

Clinical Practice Study

Outcomes of transconjunctival sutureless 27-gauge vitrectomy for stage 4 retinopathy of prematurity

Parag K Shah, Vishma Prabhu, Venkatapathy Narendran

Parag K Shah, Vishma Prabhu, Venkatapathy Narendran, Department of Pediatric Retina and Ocular Oncology, Aravind Eye Hospital, Coimbatore, Tamil Nadu 641014, India

ORCID number: Parag K Shah (0000-0002-5014-6599); Vishma Prabhu (0000-0001-6845-0209); Venkatapathy Narendran (0000-0003-1201-9801).

Author contributions: All authors equally contributed to this paper with regard to study conception and design, literature review and analysis, manuscript drafting, critical revision, and editing, and final approval of the final version.

Institutional review board statement: This study was reviewed and approved by the Ethics Committee of Aravind Eye Hospital.

Informed consent statement: The participating patients provided informed consent and gave permission for publication.

Conflict-of-interest statement: We have no financial relationships to disclose.

Open-Access: This article is an open-access article which was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

Manuscript source: Invited Manuscript

Correspondence to: Dr. Parag K Shah, DNB, Department of Pediatric Retina and Ocular Oncology, Aravind Eye Hospital, Avinashi Road and Postgraduate Institute of Ophthalmology, Coimbatore, Tamil Nadu 641014, India. parag@cbe.aravind.org
Telephone: +91-422-4360400
Fax: +91-422-2593030

Received: November 2, 2017

Peer-review started: November 2, 2017

First decision: November 30, 2017

Revised: December 3, 2017

Accepted: December 14, 2017

Article in press: December 14, 2017

Published online: February 8, 2018

Abstract**AIM**

To report our initial experience with lens-sparing vitrectomy for stage 4 retinopathy of prematurity using the 27-gauge (G) system.

METHODS

This retrospective case series involved nine eyes of five babies with active stage 4 ROP, who underwent 27-G lens-sparing vitrectomy. Surgery was done using 27-G valved cannulas and sclerotomies were made 1.5 mm from the limbus. Bilateral sequential vitrectomy was done in eight eyes.

RESULTS

At one-year follow-up, anatomical outcome was favourable in all nine (100%) eyes. High-speed cutting and smaller sclerotomies were helpful in reducing the intra and post-operative complications.

CONCLUSION

27-G vitrectomy is well suited for stage 4 ROP surgeries.

Key words: Vitrectomy; Retinopathy of prematurity; 27-gauge

© **The Author(s) 2018.** Published by Baishideng Publishing Group Inc. All rights reserved.

Core tip: This is a retrospective study of nine eyes of five children with active stage 4 retinopathy of prematurity, who underwent 27-gauge microincision vitrectomy surgeries with excellent outcomes.

Shah PK, Prabhu V, Narendran V. Outcomes of transconjunctival sutureless 27-gauge vitrectomy for stage 4 retinopathy of prematurity. *World J Clin Pediatr* 2018; 7(1): 62-66 Available from: URL: <http://www.wjgnet.com/2219-2808/full/v7/i1/62.htm> DOI: <http://dx.doi.org/10.5409/wjcp.v7.i1.62>

INTRODUCTION

Pars plana vitrectomy was first described by Machemer *et al*^[1] in 1971 when he used the 17-gauge (G) system, where the incisions were > 1 mm in size. Later, O'Malley and Heintz^[2] introduced the 20-G system where the incisions became 0.9 mm in size. Although this remained the standard of care for almost three decades, the disadvantage was that apart from relatively bulky instruments, it needed conjunctival dissection followed by applications of sutures. A big change was seen when small-gauge (23 and 25 G) instrumentation was introduced^[3,4]. With thinner instrumentation, the incision became smaller to 0.5 mm and transconjunctival entry with sutureless closure became a possibility. This is extremely beneficial in the small eyes of premature neonates^[5]. The advent of 27-G vitrectomy with only 0.4 mm incisions has led to a new beginning of transconjunctival microincision vitrectomy surgery (MIVS) in retinopathy of prematurity (ROP). Oshima *et al*^[6] in 2010 were the first to describe the safety and feasibility of the 27-gauge MIVS system with excellent visual and anatomical outcomes in adults. However, its use in ROP is still not established, although Yonekawa *et al*^[7] have reported 25 and 27-G hybrid vitrectomy in complex surgeries including ROP.

The purpose of this study was to describe our initial experience in using the 27-G trocar and cannula system in pediatric eyes with stage 4 ROP.

MATERIALS AND METHODS

This is a retrospective non-comparative case series of nine eyes of five babies. All eyes had stage 4 ROP. Four babies had bilateral disease and both eyes were operated sequentially under same general anesthesia^[8]. Care was taken to rescrub and re-prepare the surgical field of the second eye as done for any new case. The entire team rescrubbed and a new set of surgical instruments were used for each eye. Written informed consent was obtained from their parents for the procedure. Ethics committee approval was obtained for this study. The study was conducted from January 2015 to December 2015. Patient records were reviewed and the data collected were date of birth, gestational age, birth weight, postconceptional age, postnatal age, and intra and post-operative status of each eye.

All surgeries were performed under general anesthesia by a single surgeon, using the Constellation Vision System with Vitrectomy 27-G Total Plus Vitrectomy Pak system (Alcon Laboratories, Texas, United States).




Figure 1 Intraoperative picture of a left eye. The three sclerotomies 1.5 mm from the limbus are labeled as 1 (inferotemporal port), 2 (superotemporal port) and 3 (superonasal port) of the 27-gauge vitrectomy system (silicon band pieces are not shown in this representative photograph). The white arrow shows the infusion tube through which balanced salt solution flows and maintains intraocular pressure. The black arrow shows the trocar handle inserting the last superonasal port.

Valved 27-G cannulas were used. The intraocular portions of the 27-G cannula, which are designed for adult eyes were shortened for these small pediatric eyes, similar to what has been described by Babu *et al*^[9]. We took a 42 silicon band, which has a width of 4 mm and a thickness of 1.25 mm. The band was divided at 2 mm and again cut in the center to get two small pieces of 2 mm x 2 mm each. The trocar and cannula were passed through these pieces till the hub with the help of a toothed forceps. Cannulas were inserted at inferotemporal, superotemporal, and superonasal quadrants, 1.5 mm away from the limbus (Figure 1). The conjunctiva and Tenon's capsule were displaced over the sclera to avoid communication between conjunctival and scleral entry sites.

Trocar cannulas were inserted *via* a "straight in" (perpendicular to the sclera) or angled (less than 90 degrees to the sclera) approach in a 1-step procedure. First, a core vitrectomy was performed using a cutting speed of 5000 cuts per minute and a suction of 150 mmHg in core mode. Peripheral vitrectomy was done using 7500 cuts per minute with the same suction in shave mode of the machine. After the vitrectomy, partial fluid air exchange was done, the cannulas were removed, and sclerotomies were left sutureless after thorough examination for any leakage. Babies were examined on postoperative days 1, 14, and after one month.

RESULTS

Nine eyes of five babies were operated during the study period. Mean gestational age of these babies was 29.2 wk (range, 28-31 wk). Mean birth weight was 1177 g (range, 950-1850 g). Mean postconceptional age at the time of surgery was 39 wk (range, 36-43 wk), and mean postnatal age was 9.6 wk (range, 6-12 wk). Out of nine eyes, seven had stage 4a and two had 4b disease. Two eyes of case 1 had ROP in zone 1, while the rest of the

Table 1 Baseline characteristics and final outcome of the cases

Case No	GA (wk)	BW (gm)	PCA (wk)	Eye	Zone	Stage	Preoperative laser	Intraoperative anti-VEGF	Final outcome
1	30	1100	36	RE	1	4a	Y	Y	F
				LE	1	4a	Y	Y	F
2	31	950	43	RE	2	4a	Y	N	F
				LE	2	4a	Y	N	F
3	28	1000	37	RE	2	4b	Y	N	F
				LE	2	4a	Y	N	F
4	28	1850	38	RE	2	4a	Y	N	F
				LE	2	4a	Y	N	F
5	29	985	41	RE	2	4b	N	N	F

GA: Gestational age; BW: Birth weight; PCA: Postconceptional age; RE: Right eye; LE: Left eye; F: Favourable.




Figure 2 Anatomical success was achieved in all ten eyes at 4-mo follow-up. A: Preoperative picture of left eyes showing stage 4A ROP with partial retinal detachment (black arrows), and the optic disc is shown by the thin white arrow; B: Postoperative picture of the same eye showing settled retinal detachment with residual fibrous tissue (black arrows). The optic disc and fovea are shown by the white arrows, respectively.

eyes had ROP in zone 2. Eight eyes of four cases had undergone laser therapy prior to vitrectomy (Table 1). Two eyes of case 1 underwent vitrectomy along with intravitreal injection of ranibizumab as this case had excessive vascularity. At 4-mo follow-up, anatomical success was seen in all ten (100%) eyes (Figure 2). We did not encounter any postoperative hypotony in any of our cases. No other complications were noted in the follow-up period. At an average follow-up of 12 mo, all eyes showed stable regressed ROP.

DISCUSSION

The introduction of 27-G vitrectomy had major concerns like reduced endoillumination, instrument fragility, and reduced flow rate during surgery^[10]. Oshima *et al*^[6] in their study describe the various steps taken to improve the MIVS. This study done in 2010 with 31 adult eyes undergoing vitreoretinal surgeries for epiretinal membrane, macular hole, vitreous hemorrhage, focal tractional retinal detachment, and vitreomacular traction also analyzed the duty cycles of cutter, infusion and aspiration rates, and clinical outcome of 27-G MIVS. Brighter light sources such as xenon and mercury vapour bulb were used to increase the endoillumination. The smaller gauge did increase the fragility of the cutter during vitrectomy and to avoid this, the shaft of the cutter was reduced from 32 mm to 25 mm, which gives a good

rigidity similar to that of a conventional 25-G vitrectomy.

High cutting rates of 27-G vitrectomy was proved to be a major advantage as it reduced the risk of iatrogenic damage to the retinal surface and intraoperative retinal tears by preventing uncut vitreous fibers entering the cutter port. Clinically and experimentally it has been proved that high cutting rates reduce retinal traction and thereby retinal breaks^[11,12].

Postoperative complications seen with 23 and 25-gauge surgery like poor wound sealing causing leakage, hypotony, and endophthalmitis^[13-15] were serious concerns, but one-step insertion technique in 27-G vitrectomy surgeries avoided these problems. Opening and closing procedures are simplified with the one-step insertion technique, consequently shortening the total operative time. Similarly, Rizzo and colleagues^[16] in 2012 described 27-G vitrectomy in 16 patients and achieved a good clinical outcome. Additional surgical indications included were rhegmatogenous retinal detachment and tractional retinal detachment. No intraoperative and postoperative complications were encountered and no cases were converted to the 23 or 25-G system.

A retrospective case series of 95 adult eyes by Khan *et al*^[17] has been extremely valuable as it includes a large sample size of 27-G MIVS operated till date. This study evaluated the change in visual acuity, postoperative intraocular pressure (IOP), and the mean operative time by diagnosis. They reported an initial fall in IOP over

the first week, which started to increase after day 30. A recently published study of cases with 20, 23, and 25-G surgeries showed a decrease in baseline IOP over a period of 3 mo and then gradual regain^[18]. Thus, they concluded that IOP regains faster to the baseline value in MIVS.

In our study, 27-G MIVS performed in pediatric eyes is the first of its kind and has not been reported till date. The problem of bending of the shaft is minimal as the plus system has a small cuff of metal supporting the base and ROP vitrectomy just needs a good core vitrectomy with keeping the instruments more perpendicular compared to adults in whom base shaving may be required. None of our case had hypotony in the postoperative period as compared to 5% in a study by Khan *et al*^[17]. However, our sample size is much smaller compared to their study. Singh *et al*^[19] showed that the different gauge vitrectomy systems are equally effective and safe. All of our cases were sutureless and at the end of the surgery none developed any wound leak, which was seen in about 0%-7.1% of cases of 23 and 25-G vitrectomy^[20-22]. No sclerotomy-related retinal tears were noted while with 23 and 25-gauge vitrectomy 0%-3.1% have been reported^[22-25].

In one case during vitrectomy, the cannula came out along with the vitrectomy cutter, and this could have been avoided by carefully removing the cutter. This could be more common in the 27-G system compared to the 25-G system, as Alcon 27-G cannulas are valved and hence get snugly attached to the instruments. In four babies, bilateral simultaneous vitrectomy was performed as all these babies had bilateral acute disease. The anatomical outcome was 100% in our study, which is better than previously reported ROP vitrectomies with larger gauges^[8,26-28].

Limitation of our study is small sample size. Thus to summarize, the 27-G system could be favourable for pediatric cases, since this technique has a favorable wound sealing structure with fewer postoperative complications and better surgical outcomes. However, studies with a larger sample size are needed to substantiate this.

ARTICLE HIGHLIGHTS

Research background

Retinopathy of prematurity (ROP) vitrectomy is challenging due to the altered and more compact structures in a pediatric eye. Hence, there is a need to invent smaller vitrectomy instruments to make the outcome of this surgery better. 27-gauge vitrectomy being the smallest gauge available commercially could have the most benefit in the pediatric age group.

Research motivation

27-gauge instruments are smaller in size and have capability of high speed cutting, which is ideal for pediatric eyes. Hence, this study was conducted to examine the feasibility of this instrument in ROP surgery.

Research objectives

The main research objective was to assess the feasibility of 27-gauge vitrectomy for ROP and to examine if the ease of surgery and surgical

outcomes could be bettered.

Research methods

27-gauge vitrectomy has been reported mainly in adult eyes. Very few studies have explored its use in the pediatric age group and especially in ROP. This is one of the few studies which tried 27-gauge vitrectomy exclusively in stage 4 ROP.

Research results

The results of this study show that sutureless transconjunctival 27-gauge vitrectomy has good anatomical outcome in stage 4 ROP.

Research conclusions

27-gauge vitrectomy is beneficial for pediatric eyes. It is safe and effective. This study shows that smaller gauge instruments are most suitable for lens-sparing vitrectomy in ROP where the surgical space is very limited with the ever looming danger of damaging the lens anteriorly and the retinal posteriorly. Even with the limited surgical space, it become easier to maneuver with 27-gauge instruments without damaging the critical structures.

Research perspectives

In the future, 27-gauge vitrectomy has the potential to become the standard of care for all ROP-related lens-sparing vitrectomies.

REFERENCES

- 1 **Machemer R**, Buettner H, Norton EW, Parel JM. Vitrectomy: a pars plana approach. *Trans Am Acad Ophthalmol Otolaryngol* 1971; **75**: 813-820 [PMID: 5566980]
- 2 **O'Malley C**, Heintz RM Sr. Vitrectomy with an alternative instrument system. *Ann Ophthalmol* 1975; **7**: 585-588, 591-594 [PMID: 1147502]
- 3 **Eckardt C**. Transconjunctival sutureless 23-gauge vitrectomy. *Retina* 2005; **25**: 208-211 [PMID: 15689813 DOI: 10.1097/00006982-200502000-00015]
- 4 **Fujii GY**, De Juan E Jr, Humayun MS, Chang TS, Pieramici DJ, Barnes A, Kent D. Initial experience using the transconjunctival sutureless vitrectomy system for vitreoretinal surgery. *Ophthalmology* 2002; **109**: 1814-1820 [PMID: 12359600 DOI: 10.1016/S0161-6420(02)01119-3]
- 5 **Gonzales CR**, Boshra J, Schwartz SD. 25-Gauge pars plicata vitrectomy for stage 4 and 5 retinopathy of prematurity. *Retina* 2006; **26**: S42-S46 [PMID: 16946678 DOI: 10.1097/01.iae.0000244288.63757.be]
- 6 **Oshima Y**, Wakabayashi T, Sato T, Ohji M, Tano Y. A 27-gauge instrument system for transconjunctival sutureless microincision vitrectomy surgery. *Ophthalmology* 2010; **117**: 93-102 [PMID: 19880185 DOI: 10.1016/j.ophtha.2009.06.043]
- 7 **Yonekawa Y**, Thanos A, Abbey AM, Thomas BJ, Todorich B, Faia LJ, Williams GA, Capone A Jr, Wolfe JD, Hassan TS. Hybrid 25- and 27-Gauge Vitrectomy for Complex Vitreoretinal Surgery. *Ophthalmic Surg Lasers Imaging Retina* 2016; **47**: 352-355 [PMID: 27065375 DOI: 10.3928/23258160-20160324-08]
- 8 **Shah PK**, Narendran V, Kalpana N. Safety and efficacy of simultaneous bilateral 25-gauge lens-sparing vitrectomy for vascularly active stage 4 retinopathy of prematurity. *Eye (Lond)* 2015; **29**: 1046-1050 [PMID: 25998945 DOI: 10.1038/eye.2015.78]
- 9 **Babu N**, Shah PK, Narendran V, Kalpana N, Kim R. An easy method to raise the 25-gauge trocar and cannula system for retinopathy of prematurity-related lens-sparing vitrectomy. *Retina* 2014; **34**: 1014-1015 [PMID: 24695066 DOI: 10.1097/IAE.0000000000000128]
- 10 **Thompson JT**. Advantages and limitations of small gauge vitrectomy. *Surv Ophthalmol* 2011; **56**: 162-172 [PMID: 21236459 DOI: 10.1016/j.survophthal.2010.08.003]
- 11 **Rizzo S**, Genovesi-Ebert F, Belting C. Comparative study between a standard 25-gauge vitrectomy system and a new ultrahigh-speed 25-gauge system with duty cycle control in the treatment of

- various vitreoretinal diseases. *Retina* 2011; **31**: 2007-2013 [PMID: 21685823 DOI: 10.1097/IAE.0b013e318213623a]
- 12 **Teixeira A**, Chong LP, Matsuoka N, Arana L, Kerns R, Bhadri P, Humayun M. Vitreoretinal traction created by conventional cutters during vitrectomy. *Ophthalmology* 2010; **117**: 1387-1392 [PMID: 20176400 DOI: 10.1016/j.ophtha.2009.11.004]
 - 13 **Woo SJ**, Park KH, Hwang JM, Kim JH, Yu YS, Chung H. Risk factors associated with sclerotomy leakage and postoperative hypotony after 23-gauge transconjunctival sutureless vitrectomy. *Retina* 2009; **29**: 456-463 [PMID: 19174725 DOI: 10.1097/IAE.0b013e318195cb28]
 - 14 **Shimada H**, Nakashizuka H, Hattori T, Mori R, Mizutani Y, Yuzawa M. Incidence of endophthalmitis after 20- and 25-gauge vitrectomy causes and prevention. *Ophthalmology* 2008; **115**: 2215-2220 [PMID: 18930557 DOI: 10.1016/j.ophtha.2008.07.015]
 - 15 **Byeon SH**, Lew YJ, Kim M, Kwon OW. Wound leakage and hypotony after 25-gauge sutureless vitrectomy: factors affecting postoperative intraocular pressure. *Ophthalmic Surg Lasers Imaging* 2008; **39**: 94-99 [PMID: 18435331 DOI: 10.3928/15428877-20080301-04]
 - 16 **Rizzo S**, Barca F, Caporossi T, Mariotti C. Twenty-seven-gauge vitrectomy for various vitreoretinal diseases. *Retina* 2015; **35**: 1273-1278 [PMID: 25946694 DOI: 10.1097/IAE.0000000000000545]
 - 17 **Khan MA**, Shahlaee A, Toussaint B, Hsu J, Sivalingam A, Dugel PU, Lakhanpal RR, Riemann CD, Berrocal MH, Regillo CD, Ho AC. Outcomes of 27 Gauge Microincision Vitrectomy Surgery for Posterior Segment Disease. *Am J Ophthalmol* 2016; **161**: 36-43 [PMID: 26429584 DOI: 10.1016/j.ajo.2015.09.024]
 - 18 **Mi CW**, Thompson JT. Long-term follow-up of intraocular pressure after vitrectomy in eyes without preexisting glaucoma. *Retina* 2015; **35**: 2543-2551 [PMID: 26035511 DOI: 10.1097/IAE.0000000000000641]
 - 19 **Singh RP**, Bando H, Brasil OF, Williams DR, Kaiser PK. Evaluation of wound closure using different incision techniques with 23-gauge and 25-gauge microincision vitrectomy systems. *Retina* 2008; **28**: 242-248 [PMID: 18301029 DOI: 10.1097/IAE.0b013e318156dea3]
 - 20 **Lakhanpal RR**, Humayun MS, de Juan E Jr, Lim JI, Chong LP, Chang TS, Javaheri M, Fujii GY, Barnes AC, Alexandrou TJ. Outcomes of 140 consecutive cases of 25-gauge transconjunctival surgery for posterior segment disease. *Ophthalmology* 2005; **112**: 817-824 [PMID: 15878061 DOI: 10.1016/j.ophtha.2004.11.053]
 - 21 **Ibarra MS**, Hermel M, Prenner JL, Hassan TS. Longer-term outcomes of transconjunctival sutureless 25-gauge vitrectomy. *Am J Ophthalmol* 2005; **139**: 831-836 [PMID: 15860288 DOI: 10.1016/j.ajo.2004.12.002]
 - 22 **Fine HF**, Iranmanesh R, Iturralde D, Spaide RF. Outcomes of 77 consecutive cases of 23-gauge transconjunctival vitrectomy surgery for posterior segment disease. *Ophthalmology* 2007; **114**: 1197-1200 [PMID: 17544779 DOI: 10.1016/j.ophtha.2007.02.020]
 - 23 **Gupta OP**, Ho AC, Kaiser PK, Regillo CD, Chen S, Dyer DS, Dugel PU, Gupta S, Pollack JS. Short-term outcomes of 23-gauge pars plana vitrectomy. *Am J Ophthalmol* 2008; **146**: 193-197 [PMID: 18547538 DOI: 10.1016/j.ajo.2008.04.010]
 - 24 **Jalil A**, Ho WO, Charles S, Dhawahir-Scala F, Patton N. Iatrogenic retinal breaks in 20-G versus 23-G pars plana vitrectomy. *Graefes Arch Clin Exp Ophthalmol* 2013; **251**: 1463-1467 [PMID: 23504085 DOI: 10.1007/s00417-013-2299-2]
 - 25 **Scartozzi R**, Bessa AS, Gupta OP, Regillo CD. Intraoperative sclerotomy-related retinal breaks for macular surgery, 20- vs 25-gauge vitrectomy systems. *Am J Ophthalmol* 2007; **143**: 155-156 [PMID: 17188054 DOI: 10.1016/j.ajo.2006.07.038]
 - 26 **Bhende P**, Gopal L, Sharma T, Verma A, Biswas RK. Functional and anatomical outcomes after primary lens-sparing pars plana vitrectomy for Stage 4 retinopathy of prematurity. *Indian J Ophthalmol* 2009; **57**: 267-271 [PMID: 19574693 DOI: 10.4103/0301-4738.53050]
 - 27 **Kychenthal A**, Dorta P. 25-gauge lens-sparing vitrectomy for stage 4A retinopathy of prematurity. *Retina* 2008; **28**: S65-S68 [PMID: 18317348 DOI: 10.1097/IAE.0b013e318159ec49]
 - 28 **El Rayes EN**, Vinekar A, Capone A Jr. Three-year anatomic and visual outcomes after vitrectomy for stage 4B retinopathy of prematurity. *Retina* 2008; **28**: 568-572 [PMID: 18398359 DOI: 10.1097/IAE.0b013e3181610f97]

P- Reviewer: Stewart MW **S- Editor:** Cui LJ **L- Editor:** Wang TQ
E- Editor: Li RF

