

CORRECTION

Correction: Sensitivity of self-reported opioid use in case-control studies: Healthy individuals versus hospitalized patients

Hamideh Rashidian, Maryam Hadji, Maryam Marzban, Mahin Gholipour, Afarin Rahimi-Movaghar, Farin Kamangar, Reza Malekzadeh, Elisabete Weiderpass, Abbas Rezaianzadeh, Abdolvahab Moradi, Nima Babbadi-Ashar, Reza Ghiasvand, Hossein Khavari-Daneshvar, Ali Akbar Haghdoost, Kazem Zendehdel

The affiliation for the third author is incorrect. Maryam Marzban is not affiliated with #3 but with the Department of Public Health, School of Public Health, Bushehr University of Medical Science, Bushehr, Iran.

Reference

- Rashidian H, Hadji M, Marzban M, Gholipour M, Rahimi-Movaghar A, Kamangar F, et al. (2017) Sensitivity of self-reported opioid use in case-control studies: Healthy individuals versus hospitalized patients. PLoS ONE 12(8): e0183017. <https://doi.org/10.1371/journal.pone.0183017> PMID: 28854228



OPEN ACCESS

Citation: Rashidian H, Hadji M, Marzban M, Gholipour M, Rahimi-Movaghar A, Kamangar F, et al. (2018) Correction: Sensitivity of self-reported opioid use in case-control studies: Healthy individuals versus hospitalized patients. PLoS ONE 13(2): e0192814. <https://doi.org/10.1371/journal.pone.0192814>

Published: February 8, 2018

Copyright: © 2018 Rashidian et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.