



Published in final edited form as:

*J Affect Disord.* 2018 March 01; 228: 248–253. doi:10.1016/j.jad.2017.12.014.

## Incidence, recurrence and comorbidity of anxiety disorders in four major developmental stages

Cecilia A. Essau<sup>1</sup>, Peter M. Lewinsohn<sup>2</sup>, Jie Xin Lim<sup>3</sup>, Ringo Ho Moon-Ho<sup>3</sup>, and Paul Rohde<sup>2</sup>

<sup>1</sup>Department of Psychology, University of Roehampton, Whitelands College, Holybourne Avenue, London SW15 4JD, UK

<sup>2</sup>Oregon Research Institute, 1776 Millrace Drive, Eugene, OR 97403-1983, USA

<sup>3</sup>Psychology, School of Social Sciences, Nanyang Technological University, 50 Nanyang Avenue, Singapore 639798

### Abstract

**Background**—Anxiety disorders are common in childhood, adolescence, and adulthood, and frequently comorbid with other mental disorders.

**Objective**—The main aim of the present study was to examine the incidence, recurrence and comorbidity rates of anxiety disorders across four developmental periods, namely, during childhood (5 – 12.9 years), adolescence (13 – 17.9 years), emerging adulthood (18 – 23.9 years), and adulthood (24 – 30 years).

**Method**—Eight hundred and sixteen participants from a large community sample were interviewed twice during adolescence, at age 24, and at age 30. They completed self-report measures of psychosocial functioning and semi-structured diagnostic interviews during adolescence and adulthood.

**Results**—The result showed first incidence of anxiety disorders to be significantly higher in childhood and adulthood than in adolescence and emerging adulthood. Female gender was associated with first incidence, but not with recurrence. The magnitude of comorbidity was comparable across the four developmental stages, except for Substance use disorders (SUD) during adolescence. The presence of anxiety disorders during childhood and adolescence was found to significantly increase the probability of having an anxiety disorder during emerging adulthood.

**Limitations**—The participants are ethnically and geographically homogenous.

---

**Publisher's Disclaimer:** This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

#### Contributors

Cecilia A Essau wrote the manuscript. Peter M Lewinsohn and Paul Rohde designed the study and wrote the protocol. Jie Xin Lim and Ringo Ho Moon-Ho undertook the statistical analysis. All authors contributed to and have approved the final manuscript.

#### Conflict of Interest

The authors have no conflict of interest to report in relation to the research presented in this manuscript.

**Conclusion**—Incidence and recurrence rates of anxiety disorders differed across the developmental periods, but no difference was found for the comorbidity rates.

### Keywords

Developmental periods; anxiety disorders; emerging adults; adolescents

---

## 1. Introduction

There is persuasive evidence that anxiety disorders are the most frequent psychiatric disorders in childhood, adolescence, and adulthood. Up to 12% of children and up to 32% of adolescents and adults in the community are estimated to have been diagnosed with any anxiety disorders (Essau, 2005; Feehan et al., 1993; Lewinsohn et al., 1997; Merikangas et al., 2010; Wittchen et al., 1998). In addition to being highly prevalent, anxiety disorders tend to co-occur frequently both among themselves and with numerous other psychiatric disorders such as with depressive and substance use disorders (Essau et al., 2000; Feehan et al., 1993; Lewinsohn et al., 1997; Wittchen et al., 1998). Studies have also reported that more females than males meet the diagnosis of anxiety disorders, with gender differences occurring as early as childhood and reaching ratios of 2:1 to 3:1 in adolescence.

Most anxiety disorders tend to have an early onset, generally in childhood or early adolescence. If left untreated, anxiety disorders that begin early in life can become chronic (Essau et al., 2014; Feehan et al., 1993; Ferdinand and Verhulst, 1995; Keller et al., 1992; Letcher et al., 2012; Pine et al., 1998) and are associated with a high probability of recurrence (Bruce et al., 2005). The presence of anxiety disorders during adolescence also predicted a two to three-fold increased risk for anxiety disorders in adulthood (Pine et al., 1998). A recent study by Copeland et al. (2014) showed that 22.7% of their participants had met criteria for any anxiety disorders by early adulthood; this prevalence rate increased to 27.0% if overanxious disorder was included in the analyses. Their result also showed childhood anxiety disorders to be associated with adverse functioning in at least one life domain during young adulthood. Essau and colleagues (2014) recently examined the associations between anxiety disorders with an onset in two developmental periods (during childhood and adolescence) and psychosocial outcomes at age 30. In that report, adolescent-onset anxiety disorder was associated with more negative course and outcome than childhood-onset anxiety disorder. Specifically, adolescent-onset anxiety disorder predicted poorer total adjustment, poorer adjustment at work, poorer family relationships, problems with the family unit, less life satisfaction, poorer coping skills, and more chronic stress in adulthood than those with childhood anxiety. Adolescent-onset anxiety disorder also predicted substance (SUD), alcohol abuse/dependence (AUD), and anxiety disorder in adulthood.

Although large longitudinal studies have enhanced our knowledge regarding the prevalence, comorbidity, and course and outcome of anxiety disorders in children, adolescents and adults, little is known about these characteristics of anxiety in emerging adults. Emerging adulthood is a developmental stage that spans between adolescence and adulthood (Arnett, 2000), which is characterized by significant psychological changes (e.g., autonomy, identity

exploration) and distinct psychosocial events (e.g., residence [i.e., independent living], social role [e.g., getting married], and career pursuits [i.e., employment]). These changes are associated with stressors that are associated with both risk and opportunities for growth, which in turn could have a significant impact in mental health (Schulenberg et al., 2004).

On the basis of this general background, the present study reports the results of a 16-year longitudinal study and compares the incidence, recurrence and comorbidity rates of anxiety disorders in childhood, adolescence, emerging adulthood, and adulthood. The more specific aims are to address the following four questions. In each of these developmental stages: (a) what are the incidence and recurrence rates of anxiety disorders?; (b) what are the effects of gender on anxiety disorders?; (c) how frequently do anxiety disorders co-occur with depressive and substance use disorders?; and (d) do anxiety disorders that are present in a prior developmental stage predict subsequent anxiety?

The hypotheses to be tested in this study were as follows: First, the highest incidence rate is hypothesized to be found during emerging adulthood; recurrence rate is expected to be the highest during adulthood (Scholten et al., 2016; Scholten et al., 2013). Second, females compared to males are expected to have significantly higher incidence and recurrence rates of anxiety disorders. Female gender is also expected to predict the recurrence of anxiety in any developmental stages (McLean et al., 2011). Third, in all developmental stages, there will be a strong association between anxiety and major depression, and between anxiety and substance use disorders (Essau et al., 2000; Lewinsohn et al., 1997; Wittchen et al., 1998). Fourth, the presence of an anxiety disorder at one developmental disorder will increase the risk of having an anxiety disorder in the subsequent developmental period (Essau et al., 2014; Pine et al., 1998).

To our knowledge, this is the first study that has compared the incidence, recurrence and comorbidity of anxiety disorders in childhood, adolescence, emerging adulthood, and adulthood with the same sample of individuals.

## 2. Methods

### 2.1. Participants

The present study used data from the Oregon Adolescent Depression Project (OADP; Lewinsohn et al., 1993), a longitudinal study of a large cohort of high school students who were randomly selected from nine high schools in western Oregon; detailed information about the OADP had been reported elsewhere (Lewinsohn et al., 2003; Pettit et al., 2006; Rohde et al., 2007). A total of 1709 adolescents (ages 14–18; mean age 16.6, SD=1.2) completed the initial (T1) assessments who were invited to participate in the second assessment. About a year later, 1507 adolescents participated at time 2 (T2; 88%). For the third assessment (T3), all adolescents with a history of a depressive disorder by T2 (n=360) or a history of non-mood disorders (n=284), and a random sample of adolescents with no history of psychopathology by T2 (n=457) were invited to participate in a third (T3) evaluation. In addition, all non-white T2 participants were retained in the T3 sample to maximize ethnic diversity. A total of 1101 young adults participated at time 3 (T3), with a mean age of 24.2 years. At age 30, all T3 participants were asked to complete another

interview assessment (mean age=30.45, SD=.70). Of the 941 who participated in the T3 assessment, 816 (87%) completed the T4 assessment.

Of these 816 individuals, 58.8% of the sample were female, with a mean age at T4 of 30.45 years (SD=0.70). Most participants were White (85.9%), others were African American (1%), Hispanic (3%), American Indian (3%), Asian (3%), and “other” (2%). About half of the sample was married (56.2%), and 41% had a bachelor’s degree or higher by T4.

## 2.2. Measures

**2.2.1. Diagnostic measures**—Participants were interviewed with the Schedule for Affective Disorders and Schizophrenia for School-Age Children (K-SADS; Orvaschel et al., 1982) at T1 and T2. For the follow-up assessments at T2 and T3, the participants were also administered with the Longitudinal Interval Follow-Up Evaluation (LIFE; Keller et al., 1987). The K-SADS/LIFE was used to collect information regarding the onset and course of disorders since the previous interview. The T4 interview consisted of a joint administration of the LIFE and the Structured Clinical Interview for DSM-IV (SCID; First et al., 1996) to collect information for new or continuing episodes since T3. Diagnoses were based on DSM-III-R criteria for T1 and T2 and Diagnostic and Statistical Manual of Mental Disorders, 4th edition (DSM-IV; American Psychiatric Association, 1994) criteria for T3 and T4.

The anxiety disorder diagnostic category included generalized anxiety disorder, overanxious disorder, post-traumatic stress disorder, panic disorder, agoraphobia, social phobia, simple phobia, obsessive–compulsive disorder, and separation anxiety disorder. Substance use disorders (SUD) included alcohol abuse or dependence (AUD) and sedative/hypnotic/anxiolytic, cannabis, stimulant, opioid, cocaine, and hallucinogen/PCP abuse or dependence, and polydrug dependence. Disruptive disorder included attention deficit and hyperactivity disorder, oppositional defiant disorder and conduct disorder. The interrater reliability for depressive disorders (MDD or dysthymia) in the four time assessments, with Kappa ranging from 0.82 to 1.00; for anxiety disorders it ranged from 0.76 to 0.87; and for SUD from 0.81 to 0.96.

**2.2.2. Statistical procedures**—First incidence, recurrence, and total incidence were calculated using descriptive statistics and corresponding 95% confidence intervals. To investigate the extent to which gender (1=female, 0=male) predicted rates of anxiety, binary logistic regression models were used. Binary regression models were also used to examine the comorbidity between anxiety with depression and SUD. Because of the frequent comorbidity between SUD and disruptive behaviour disorders, comorbidity with SUD were both unadjusted and adjusted for the presence of disruptive behaviour disorders.

## 3. Result

### 3.1. Incidence and recurrence rates of anxiety disorders

Table 1 shows the first incidence (i.e., number of participants with first episodes of anxiety at each developmental stage), recurrent (i.e., number of participants with previous anxiety disorders in the previous developmental period and who have a new anxiety episode), and

total incidence (i.e., number of cases in a developmental stage with either a first incidence or a recurrence). Following Rohde et al. (2012), the dominator for each of the developmental stage was calculated as follows: childhood (5.0 – 12.9 years) included all the participants (N=816), during the adolescence (13.0 – 17.9 years) there were 741 participants (816 minus 75), during emerging adulthood (18.0 – 23.9 years) there were 707 participants (816 minus 109), and during adulthood (24- 30 years), the number of participants were 671 (816 minus 145).

Significant differences in the incidence of anxiety disorders emerged across the four developmental stages. First incidence rates of anxiety disorders were significantly higher in childhood and adulthood than in adolescence and emerging adulthood. This higher incidence rate was related to separation anxiety in childhood, and to panic disorder in adulthood. Recurrence of anxiety disorders were found in adolescence, emerging adulthood, and adulthood, with comparable rates across these three developmental stages. The total incidence was significantly higher in adulthood than the other three developmental stages where comparable rates were found.

### 3.2. Gender and anxiety disorders

The next step of our analysis was to examine the extent to which gender predict the rates of anxiety disorders across the four developmental periods, using univariate logistic regression. As shown on Table 2, female gender was associated with first and total incidence in childhood, emerging adulthood and adulthood, but not in adolescence.

The odds ratios for the recurrence of anxiety disorders were not significant in adolescence and emerging adulthood, suggesting that female gender did not predict recurrence during these two developmental stages.

### 3.3. Comorbidity of anxiety disorders

Table 3 shows the comorbidity between anxiety disorders with MDD and SUD. The comorbidity between anxiety disorders and SUD was examined first irrespective of the presence of disruptive behavioural disorder, and then by controlling for the presence of disruptive behavioural disorders. Significant comorbidity was found between anxiety disorders and MDD across developmental periods; the magnitude of comorbidity across periods was comparable, and indicated a four-fold increase in risk. The comorbidity between anxiety and SUD was significant across three of the four developmental periods (non-significant in the adolescent period), regardless of whether or not the presence of disruptive behavioural disorder was being controlled.

### 3.4. Impact of prior anxiety disorders

The degree to which the presence of anxiety disorders during an earlier developmental period influenced the risk of having an anxiety during a later developmental period was also examined. Following Rohde et al. (2013), we examined the extent to which the rate of anxiety disorders in adolescence was a function of MDD in childhood; emerging adulthood was a function of MDD in either childhood or adolescence; and adulthood was a function of MDD in childhood, adolescence, or emerging adulthood. During adolescence, the

probability of having an anxiety disorder given childhood anxiety was 8% (OR=2.26, 95% CI [0.91, 5.62]). During emerging adulthood, the probability of having an anxiety disorder given an anxiety during childhood or adolescence was 9% (OR=2.20, 95% CI [1.05-4.63]). During adulthood, the probability of having an anxiety disorder given an anxiety during childhood, adolescence, or emerging adulthood was 8% (OR=1.13, 95% CI [0.62-2.09]).

#### 4. Discussion

The present study makes a significant contribution to our understanding of anxiety disorders by providing information on their incidence and recurrence rates and comorbidity patterns in childhood, adolescence, emerging adulthood, and adulthood in a large non-patient sample. This was made possible by following a large cohort of adolescents into adulthood. Four major findings were obtained.

First, the incidence rates of anxiety disorders were significantly higher in childhood and adulthood than the other two developmental stages. The higher incidence rate was related to the presence of separation anxiety in childhood, and to panic disorder in adulthood. The higher incidence of separation anxiety in childhood compared to other developmental stages supports previous studies that showed this disorder to have its first onset in childhood and is the earliest onset of all anxiety disorders (Beesdo et al., 2009); it also provides support to the finding that separation anxiety disorder is the most frequent anxiety disorder among children, with prevalence rates ranging from 2.8% to 8% (Beesdo et al., 2009; Bowen et al., 1990; Pine et al., 1998). Furthermore, the higher incidence of panic disorder in adulthood was in line with previous studies that reported this disorder to have its first onset in adulthood (de Graaf et al., 2003; Kessler et al., 2005).

Our hypothesis that emerging adulthood would have the highest incidence of anxiety disorders was not supported. This was surprising given major life changes and stressors (e.g., relationship instability, lack of economic independence) that are often reported during this developmental stage which could put individuals at this stage at risk of developing an anxiety disorder (Schulenberg et al., 2004). However, it is worth noting that the concept of emerging adulthood is controversial, with some authors considering this concept to be better conceptualized in terms of economic conditions instead of a psychological developmental period (Côté and Bynner, 2008).

Second, the incidence of anxiety disorders is significantly higher in females than males during childhood, emerging adult, and adulthood. This finding replicates previous studies (Essau et al., 2000; Hale et al., 2008; Letcher et al., 2012; Merikangas et al., 2010; Pine et al., 1998; Reinherz et al., 1989) showing higher rate of anxiety disorders among females than males; previous studies have reported gender differences during childhood and then increase with age which reach ratios of 2:1 to 3:1 in adolescence (Pine et al., 1998). Interestingly, no gender differences were found for first incidence of anxiety disorders during adolescence which is inconsistent with previous studies (Pine et al., 1998). While it is beyond the scope of the present study to explore the reason for this non-significant gender difference during this specific developmental stage, it could be related to the biological changes related to puberty which have similar impact on boys and girls.

Third, in agreement with previous studies (Essau et al., 2000; Kashani and Orvaschel, 1990; Last et al., 1997; Lewinsohn et al., 1997; Rohde et al., 1991), the degree of comorbidity between anxiety disorders and MDD, as well as between anxiety and SUD were significant; to our knowledge the present study was the first to have examined the comorbidity between anxiety, MDD and SUD across the four developmental periods where comparable comorbidity rates were found, except for SUD during adolescence. Despite the high comorbidity rates between anxiety and other psychiatric disorders, the meaning of comorbidity for psychopathology issues remains unclear. One explanation could be related to our current classification and assessment of psychiatric disorders (Merikangas, 1990). Another explanation is related to the etiology of disorders. As argued by Merikangas (1990), the co-occurrence of disorders could be etiologic in that one disorder causes the second disorder, or that the two disorders are manifestations of the same underlying etiologic factors.

Fourth, the presence of anxiety disorders during childhood and adolescence was found to significantly increase the probability of having an anxiety during emerging adulthood. Indeed, several longitudinal studies have shown that young people with anxiety disorders compared with those without this disorder are at increased risk to have the same disorder (Bittner et al., 2007; Newman et al., 1996; Pine et al., 1998) at follow-up (“homotypic continuity”). Moreover, studies among adults have reported that those with anxiety disorders often had these same disorders during adolescence (Gregory et al., 2007).

## 5. Limitations

The study has some limitations which should be considered. First, the extent to which our findings are generalizable to other populations is unknown because the sample was restricted to one geographical location of the United States. Related to this limitation is the fact that the sample was predominantly of European-American descent. Second, given differences in prevalence of anxiety disorders across ethnic groups, the generalizability of the present findings may be limited (Asnaani et al., 2010). Third, the OADP is a 16-year longitudinal study, and as such changes in the diagnostic criteria and assessment approaches used are inevitable; furthermore, there are some attrition at each assessment. Fourth, the information related to the diagnoses were assessed solely via self-report. Finally, similar to other longitudinal studies (e.g., Essau et al., 2002), the drop-out rate was high which was not surprising because the participants were followed over the period of 16 years.

## 6. Conclusions

The first incidence rates of anxiety disorders differ across the four developmental stages, and the presence of anxiety disorders during an earlier developmental period tended to increase the probability of having an anxiety during a later developmental period. Studies are needed to elucidate the social, psychological, and physiological processes that give rise to the differences in the incidence rates across developmental stages. A better understanding of these processes will likely play a crucial role in designing effective prevention programs for individuals in specific developmental stages.

## Acknowledgments

The authors are grateful to all those who participated in the study.

### Role of Funding Source

This research was supported in part by National Institute of Mental Health awards MH40501 and MH50522 (Dr Lewinsohn). The NIMH had no further role in study design; in collection, analysis and interpretation of data; in the writing of the report; and in the decision to submit the paper for publication.

## References

- American Psychiatric Association. Diagnostic and Statistical Manual. fourth. American Psychiatric Association; Washington: 1994.
- Arnett JJ. Emerging adulthood: A theory of development from the late teens through the twenties. *American Psychologist*. 2000; 55:469–480. [PubMed: 10842426]
- Asnaani A, Richey JA, Dimaite R, Hinton DE, Hofmann SG. A cross-ethnic comparison of lifetime prevalence rates of anxiety disorders. *Journal of Nervous and Mental Disease*. 2010; 198:551–555. [PubMed: 20699719]
- Beesdo K, Knappe S, Pine DS. Anxiety and anxiety disorders in children and adolescents: developmental issues and implications for DSM-V. *Psychiatric Clinics of North America*. 2009; 32:483–524. [PubMed: 19716988]
- Bittner A, Egger HL, Erkanli A, Jane Costello E, Foley DL, Angold A. What do childhood anxiety disorders predict? *Journal of Child Psychology and Psychiatry*. 2007; 48:1174–1183. [PubMed: 18093022]
- Bowen RC, Offord DR, Boyle MH. The prevalence of overanxious disorder and separation anxiety disorder: Results from the Ontario Child Health Study. *Journal of the American Academy of Child & Adolescent Psychiatry*. 1990; 29:753–758. [PubMed: 2228929]
- Bruce SE, Yonkers KA, Otto MW, Eisen JL, Weisberg RB, Pagano M, Shea MT, Keller MB. Influence of psychiatric comorbidity on recovery and recurrence in generalized anxiety disorder, social phobia, and panic disorder: A 12-year prospective study. *American Journal of Psychiatry*. 2005; 162:1179–1187. [PubMed: 15930067]
- Copeland WE, Angold A, Shanahan L, Costello EJ. Longitudinal patterns of anxiety from childhood to adulthood: The Great Smoky Mountains Study. *Journal of the American Academy of Child & Adolescent Psychiatry*. 2014; 53:21–33. [PubMed: 24342383]
- Côté J, Bynner JM. Changes in the transition to adulthood in the UK and Canada: The role of structure and agency in emerging adulthood. *Journal of Youth Studies*. 2008; 11:251–268.
- de Graaf R, Bijl R, Spijker J, Beekman A, Vollebergh W. Temporal sequencing of lifetime mood disorders in relation to comorbid anxiety and substance use disorders. *Social Psychiatry and Psychiatric Epidemiology*. 2003; 38:1–11. [PubMed: 12563553]
- Essau CA. Frequency and patterns of mental health services utilization among adolescents with anxiety and depressive disorders. *Depression and Anxiety*. 2005; 22:130–137. [PubMed: 16175563]
- Essau CA, Conradt J, Petermann F. Frequency, comorbidity, and psychosocial impairment of anxiety disorders in German adolescents. *Journal of Anxiety Disorders*. 2000; 14:263–279. [PubMed: 10868984]
- Essau CA, Conradt J, Petermann F. Course and outcome of anxiety disorders in adolescents. *Journal of Anxiety Disorders*. 2002; 16:67–81. [PubMed: 12171214]
- Essau CA, Lewinsohn PM, Olaya B, Seeley JR. Anxiety disorders in adolescents and psychosocial outcomes at age 30. *Journal of Affective Disorders*. 2014; 163:125–132. [PubMed: 24456837]
- Feehan M, McGee R, Williams SM. Mental health disorders from age 15 to age 18 years. *Journal of the American Academy of Child & Adolescent Psychiatry*. 1993; 32:1118–1126. [PubMed: 8282655]
- Ferdinand RF, Verhulst FC. Psychopathology from adolescence into young adulthood: An 8-year follow-up study. *American Journal of Psychiatry*. 1995; 152:1586. [PubMed: 7485620]



- First, MB., Gibbon, M., Spitzer, RL., Williams, JBW. Structured Clinical Interview for DSM-IV Axis I Disorders: Nonpatient Edition (SCID-I, Version 2.0). Biometrics Research, New York State Psychiatric Institute; New York: 1996.
- Gregory AM, Caspi A, Moffitt TE, Koenen K, Eley TC, Poulton R. Juvenile mental health histories of adults with anxiety disorders. *American Journal of Psychiatry*. 2007; 164:301–308. [PubMed: 17267794]
- Hale WW III, Raaijmakers Q, Muris P, van Hoof A, Meeus WIM. Developmental trajectories of adolescent anxiety disorder symptoms: A 5-Year prospective community study. *Journal of the American Academy of Child & Adolescent Psychiatry*. 2008; 47:556–564. [PubMed: 18356762]
- Kashani JH, Orvaschel H. A community study of anxiety in children and adolescents. *American Journal of Psychiatry*. 1990; 147:313–318. [PubMed: 2309948]
- Keller MB, Lavori PW, Friedman B, Nielsen E, Endicott J, McDonald-Scott P, Andreasen NC. The longitudinal interval follow-up evaluation: A comprehensive method for assessing outcome in prospective longitudinal studies. *Archives of General Psychiatry*. 1987; 44:540–548. [PubMed: 3579500]
- Keller MB, Lavori PW, Wunder J, Beardslee WR, Schwartz CE, Roth J. Chronic course of anxiety disorders in children and adolescents. *Journal of the American Academy of Child & Adolescent Psychiatry*. 1992; 31:595–599. [PubMed: 1644719]
- Kessler RC, Berglund P, Demler O, Jin R, Merikangas KR, Walters EE. Lifetime prevalence and age-of-onset distributions of DSM-IV disorders in the national comorbidity survey replication. *Archives of General Psychiatry*. 2005; 62:593–602. [PubMed: 15939837]
- Last CG, Hansen C, Franco N. Anxious children in adulthood: A prospective study of adjustment. *Journal of the American Academy of Child & Adolescent Psychiatry*. 1997; 36:645–652. [PubMed: 9136499]
- Letcher P, Sanson A, Smart D, Toumbourou JW. Precursors and correlates of anxiety trajectories from late childhood to late adolescence. *Journal of Clinical Child & Adolescent Psychology*. 2012; 41:417–432. [PubMed: 22551395]
- Lewinsohn PM, Hops H, Roberts RE, Seeley JR, Andrews JA. Adolescent psychopathology: I. Prevalence and incidence of depression and other DSM-III—R disorders in high school students. *Journal of Abnormal Psychology*. 1993; 102:133–144. [PubMed: 8436689]
- Lewinsohn PM, Petit JW, Joiner TE Jr, Seeley JR. The symptomatic expression of major depressive disorder in adolescents and young adults. *Journal of Abnormal Psychology*. 2003; 112:244–252. [PubMed: 12784834]
- Lewinsohn PM, Zinbarg R, Seeley JR, Lewinsohn M, Sack WH. Lifetime comorbidity among anxiety disorders and between anxiety disorders and other mental disorders in adolescents. *Journal of Anxiety Disorders*. 1997; 11:377–394. [PubMed: 9276783]
- McLean CP, Asnaani A, Litz BT, Hofmann SG. Gender differences in anxiety disorders: Prevalence, course of illness, comorbidity and burden of illness. *Journal of psychiatric research*. 2011; 45:1027–1035. [PubMed: 21439576]
- Merikangas, KR. Comorbidity for anxiety and depression: Review of family and genetic studies. In: Maser, JD., Cloninger, CR., editors. *Comorbidity of Mood and Anxiety Disorders*. American Psychiatric Press; Washington, DC: 1990. p. 331–348.
- Merikangas KR, He JP, Burstein M, Swanson SA, Avenevoli S, Cui L, Benjet C, Georgiades K, Swendsen J. Lifetime prevalence of mental disorders in U.S. adolescents: Results from the National Comorbidity Study-Adolescent Supplement (NCS-A). *Journal of the American Academy of Child & Adolescent Psychiatry*. 2010; 49:980–989. [PubMed: 20855043]
- Newman DL, Moffitt TE, Caspi A, Magdol L, Silva PA, Stanton WR. Psychiatric disorder in a birth cohort of young adults: Prevalence, comorbidity, clinical significance, and new case incidence from ages 11 to 21. *Journal of Consulting and Clinical Psychology*. 1996; 64:552–562. [PubMed: 8698949]
- Orvaschel H, Puig-Antich J, Chambers W, Tabrizi MA, Johnson R. Retrospective assessment of prepubertal major depression with the Kiddie-SADS-E. *Journal of the American Academy of Child Psychiatry*. 1982; 21:392–397. [PubMed: 7119313]

- Pettit JW, Lewinsohn PM, Joiner TE Jr. Propagation of major depressive disorder: Relationship between first episode symptoms and recurrence. *Psychiatry Research*. 2006; 141:271–278. [PubMed: 16497387]
- Pine DS, Cohen P, Gurley D, Brook J, Ma Y. The risk for early-adulthood anxiety and depressive disorders in adolescents with anxiety and depressive disorders. *Archives of General Psychiatry*. 1998; 55:56–64. [PubMed: 9435761]
- Reinherz HZ, Stewart-Berghauer G, Pakiz B, Frost AK, Moeykens BA, Holmes WM. The relationship of early risk and current mediators to depressive symptomatology in adolescence. *Journal of the American Academy of Child & Adolescent Psychiatry*. 1989; 28:942–947. [PubMed: 2808267]
- Rohde P, Lewinsohn PM, Klein DN, Seeley JR, Gau JM. Key characteristics of major depressive disorder occurring in childhood, adolescence, emerging adulthood, adulthood. *Clinical Psychological Science*. 2013; 1:41–53.
- Rohde P, Lewinsohn PM, Seeley JR. Comorbidity of unipolar depression: II. Comorbidity with other mental disorders in adolescents and adults. *Journal of Abnormal Psychology*. 1991; 100:214–222. [PubMed: 2040773]
- Rohde P, Lewinsohn PM, Seeley JR, Klein DN, Andrews JA, Small JW. Psychosocial functioning of adults who experienced substance use disorders as adolescents. *Psychology of Addictive Behaviors*. 2007; 21:155–164. [PubMed: 17563135]
- Scholten WD, Batelaan NM, Penninx BWJH, van Balkom AJLM, Smit JH, Schoevers RA, van Oppen P. Diagnostic instability of recurrence and the impact on recurrence rates in depressive and anxiety disorders. *Journal of Affective Disorders*. 2016; 195:185–190. [PubMed: 26896812]
- Scholten WD, Batelaan NM, van Balkom AJLM, Penninx WB, Smit JH, van Oppen P. Recurrence of anxiety disorders and its predictors. *Journal of Affective Disorders*. 2013; 147:180–185. [PubMed: 23218248]
- Schulenberg JE, Bryant AL, O'Malley PM. Taking hold of some kind of life: How developmental tasks relate to trajectories of well-being during the transition to adulthood. *Development and Psychopathology*. 2004; 16:1119–1140. [PubMed: 15704830]
- Wittchen HU, Nelson CB, Lachner G. Prevalence of mental disorders and psychosocial impairments in adolescents and young adults. *Psychological Medicine*. 1998; 28:109–126. [PubMed: 9483687]

### Highlights

- A 16-year longitudinal study on the incidence, recurrence and comorbidity rates of anxiety disorders during childhood, adolescence, emerging adulthood, and adulthood.
- The incidence of anxiety disorders was significantly higher in childhood and adulthood than in adolescence and emerging adulthood.
- Female gender was associated with first incidence.
- The presence of anxiety disorders during childhood and adolescence significantly increased the probability of having an anxiety during emerging adulthood.

**Table 1**

Rates of specific types of anxiety disorders during the four developmental periods

	Child		Adolescent		Emerging adult		Adulthood	
	N (%)	[95% CI]	N (%)	[95% CI]	N (%)	[95% CI]	N (%)	[95% CI]
<b>First incidence</b>								
(total anxiety)	75 (7)	[5.2, 8.3]	34 (4)	[2.3, 4.8]	36 (4)	[2.7, 5.5]	46 (7)	[4.6, 8.4]
SepAnx	34 (0.03)	[0.02, 0.04]	2 (0.002)	[0.00, 0.004]	0 (00)	[0.0, 00]	0 (00)	[0.0, 0.0]
Phobia	23 (0.02)	[0.01, 0.03]	16 (0.02)	[0.01, 0.03]	10 (0.01)	[0.004, 0.02]	16 (0.02)	[0.01, 0.03]
Agoraphobia	1 (0.00)	[0.00, 0.00]	0 (00)	[0.0, 0.0]	4 (0.001)	[0.0, 0.01]	0 (00)	[0.0, 0.0]
Social phobia	14 (0.012)	[0.01, 0.02]	6 (0.01)	[0.001, 0.01]	5 (0.005)	[0.00, 0.01]	10 (0.01)	[0.004, 0.02]
Specific phobia	0 (00)	[0.0, 0.0]	7 (0.01)	[0.001, 0.01]	3 (0.003)	[0.0, 0.01]	11 (0.01)	[0.001, 0.02]
GAD	0 (00)	[0.0, 0.0]	0 (00)	[0.0, 0.0]	1 (0.001)	[0.00, 0.00]	17 (0.02)	[0.01, 0.03]
OCD	3 (0.003)	[0.00, 0.01]	3 (0.003)	[0.00, 0.01]	0.00 (00)	[0.00, 0.00]	5 (0.004)	[0.001, 0.01]
Panic	4 (0.35)	[0.00, 0.70]	9 (0.88)	[0.30, 1.50]	17 (1.60)	[0.80, 2.40]	27 (3.10)	[1.90, 4.30]
Panic with Agora	2 (0.002)	[0.0, 0.004]	4 (0.004)	[0.0, 0.01]	3 (0.003)	[0.0, 0.01]	10 (0.01)	[0.004, 0.02]
Panic without Agora	2 (0.002)	[0.0, 0.004]	5 (0.004)	[0.001, 0.01]	14 (0.01)	[0.01, 0.02]	17 (0.02)	[0.01, 0.03]
PTSD	18 (0.02)	[0.01, 0.03]	11 (0.01)	[0.004, 0.02]	29 (0.03)	[0.02, 0.04]	12 (0.01)	[0.01, 0.02]
Acute stress	0 (00)	[0.0, 0.0]	0 (00)	[0.0, 0.0]	1 (0.001)	[0.0, 0.003]	0 (00)	[0.0, 0.0]
<b>Recurrence</b>								
(total anxiety)	0 (00)	[0.0, 0.0]	6 (8)	[1.7, 13.8]	10 (9)	[3.5, 13.9]	12 (8)	[3.4, 11.8]
SepAnx	0 (00)	[0.0, 0.0]	1 (0.03)	[0.00, 0.09]	0 (00)	[0.0, 0.0]	0 (00)	[0.0, 0.0]
Phobia	0 (00)	[0.0, 0.0]	0 (00)	[0.0, 0.0]	0 (00)	[0.0, 0.0]	0 (00)	[0.0, 0.0]

	Child		Adolescent		Emerging adult		Adulthood	
	N (%)	[95% CI]	N (%)	[95% CI]	N (%)	[95% CI]	N (%)	[95% CI]
Agoraphobia	0 (00)	[0.0, 0.0]	0 (00)	[0.0, 0.0]	0 (00)	[0.0, 0.0]	0 (00)	[0.0, 0.0]
Social phobia	0 (00)	[0.0, 0.0]	0 (00)	[0.0, 0.0]	0 (00)	[0.0, 0.0]	0 (00)	[0.0, 0.0]
Specific phobia	0 (00)	[0.0, 0.0]	0 (00)	[0.0, 0.0]	0 (00)	[0.0, 0.0]	0 (00)	[0.0, 0.0]
GAD	0 (00)	[0.0, 0.0]	0 (00)	[0.0, 0.0]	0 (00)	[0.0, 0.0]	0 (00)	[0.0, 0.0]
OCD	0 (00)	[0.0, 0.0]	0 (00)	[0.0, 0.0]	0 (00)	[0.0, 0.0]	0 (00)	[0.0, 0.0]
Panic	0 (00)	[0.0, 0.0]	0 (00)	[0.0, 0.0]	1 (7.12)	[0.00, 21.2]	1 (3.12)	[0.00, 9.20]
Panic with Agora	0 (00)	[0.0, 0.0]	0 (00)	[0.0, 0.0]	1 (0.14)	[0.0, 0.43]	0 (00)	[0.0, 0.0]
Panic without Agora	0 (00)	[0.0, 0.0]	0 (00)	[0.0, 0.0]	0 (00)	[0.0, 0.0]	1 (0.05)	[0.00, 0.14]
PTSD	0 (00)	[0.0, 0.0]	1 (0.05)	[0.00, 0.15]	1 (0.03)	[0.00, 0.09]	1 (0.02)	[0.00, 0.05]
Acute stress	0 (00)	[0.0, 0.0]	0 (00)	[0.0, 0.0]	0 (00)	[0.0, 0.0]	0 (00)	[0.0, 0.0]
<b>Total incidence</b>								
(total anxiety)	75 (7)	[5.2, 8.3]	40 (4)	[2.6, 5.1]	46 (5)	[3.2, 6]	58 (7)	[4.9, 8.4]
SepAnx	34 (0.03)	[0.02, 0.04]	3 (0.003)	[0.00, 0.006]	0 (00)	[0.0, 0.0]	0 (00)	[0.0, 0.0]
Phobia	23 (0.02)	[0.01, 0.03]	16 (0.02)	[0.01, 0.03]	10 (0.01)	[0.00, 0.02]	16 (0.02)	[0.01, 0.03]
Agoraphobia	1 (00)	[0.0, 0.0]	0 (00)	[0.0, 0.0]	4 (00)	[0.0, 0.01]	0 (00)	[0.0, 0.0]
Social phobia	14 (0.01)	[0.01, 0.02]	6 (0.01)	[0.001, 0.01]	5 (0.01)	[0.0, 0.01]	10 (0.01)	[0.004, 0.02]
Specific phobia	0 (00)	[0.0, 0.0]	7 (0.01)	[0.001, 0.01]	3 (0.003)	[0.0, 0.01]	11 (0.01)	[0.004, 0.02]
GAD	0 (00)	[0.0, 0.0]	0 (00)	[0.0, 0.0]	1 (0.001)	[0.00, 0.00]	17 (0.02)	[0.01, 0.03]
OCD	3 (0.003)	[0.00, 0.01]	3 (0.003)	[0.00, 0.01]	0 (00)	[0.0, 0.0]	5 (0.004)	[0.001, 0.01]

	Child		Adolescent		Emerging adult		Adulthood	
	N (%)	[95% CI]	N (%)	[95% CI]	N (%)	[95% CI]	N (%)	[95% CI]
Panic	4 (0.35)	[0.00, 0.70]	9 (0.88)	[0.30, 1.50]	18 (1.67)	[0.90, 2.50]	28 (3.10)	[1.90, 4.30]
Panic with Agora	2 (0.002)	[0.0, 0.004]	4 (0.004)	[0.0, 0.01]	4 (0.00)	[0.0, 0.01]	10 (0.01)	[0.01, 0.02]
Panic without Agora	2 (0.002)	[0.0, 0.01]	5 (0.004)	[0.001, 0.01]	14 (0.01)	[0.01, 0.02]	18 (0.02)	[0.01, 0.03]
PTSD	18 (0.02)	[0.01, 0.03]	12 (0.01)	[0.01, 0.02]	30 (0.03)	[0.02, 0.04]	13 (0.01)	[0.01, 0.02]
Acute stress	0 (00)	[0.0, 0.0]	0 (00)	[0.0, 0.0]	1 (0.001)	[00, 0.003]	0 (00)	[0.0, 0.0]

Note: SepAnx = Separation Anxiety Disorder; GAD = Generalised Anxiety Disorder; OCD = Obsessive-Compulsive Disorder; Panic with Agora = Panic disorder with Agoraphobia; Panic without Agora = Panic disorder without Agoraphobia; PTSD = Post-traumatic Stress Disorder; OR = odds ratio; CI = confidence interval.

**Table 2**

Gender predicting anxiety disorders during the four developmental periods

Disorders	Child		Adolescent		Emerging adult		Adulthood	
	OR	[95% CI]	OR	[95% CI]	OR	[95% CI]	OR	[95% CI]
First incidence	2.95	[1.67, 5.10]***	2.07	[0.89, 4.85]	2.48	[1.10, 5.61]*	2.15	[1.08, 4.28]*
Recurrence	-		1.45	[0.16, 13.36]	1.32	[0.26, 6.70]	-	
Total incidence	2.95	[1.67, 5.18]***	2.12	[0.96, 4.69]	2.43	[1.17, 5.03]*	2.42	[1.26, 4.68]**

Note: OR = odds ratio; CI = confidence interval. For child, there is no recurrence, thus, the OR cannot be computed.

For adult, none of the male participants had any recurrence, thus the OR cannot be computed.

**Table 3**

Period comorbidity of anxiety disorders during the four developmental periods

Disorders	Child		Adolescent		Emerging adult		Adulthood		
	%	OR	%	OR	%	OR	%	OR	
	[95% CI]		[95% CI]		[95% CI]		[95% CI]		[95% CI]
Major depression	16 v 4	4.59	64 v 29	4.31	57 v 28	3.53	60 v 24	4.74	
	[2.29, 9.20]		[2.61, 7.4]		[1.98, 6.30]		[2.77, 8.10]		
Substance disorder <sup>a</sup>	5 v 1	4.40	28 v 22	1.35	42 v 28	1.90	34 v 18	2.31	
	[1.37, 14.15]		[0.80, 2.29]		[1.08, 3.38]		[1.33, 4.01]		
Substance disorder <sup>b</sup>	4 v 1	4.59	26 v 20	1.43	44 v 47	1.87	35 v 17	2.34	
	[1.43, 14.66]		[0.83, 2.47]		[1.04, 3.34]		[1.34, 4.10]		

Note: Childhood, 5.0–12.9 years of age; adolescence, 13.0–17.9 years of age; emerging adulthood, 18.0–23.9 years of age; adulthood, 24–30 years of age.

Percentages (%) represent anxiety disorders vs. no anxiety disorders. All tests significant at  $p < .001$ . OR = odds ratio; CI = confidence interval.

<sup>a</sup>Unadjusted associations.

<sup>b</sup>Adjusted for the lifetime presence of disruptive behavior disorders.