



## Corrigendum: A high-flow nasal cannula system set at relatively low flow effectively washes out CO<sub>2</sub> from the anatomical dead space of a respiratory-system model

Yu Onodera, Ryo Akimoto, Hiroto Suzuki, Masaki Nakane, and Kaneyuki Kawamae

*Department of Anesthesiology, Yamagata University Faculty of Medicine, Yamagata, Japan*

Korean J Anesthesiol 2017 February 70(1): 105-106

<https://doi.org/10.4097/kjae.2017.70.1.105>

The article by Onodera Y, et al. entitled, “A high-flow nasal cannula system set at relatively low flow effectively washes out CO<sub>2</sub> from the anatomical dead space of a respiratory-system model” (Korean J Anesthesiol 2017 Feb 70(1): 105-106) contained an error in author’s name.

In the original publication of this paper the name of the fourth author was incorrectly rendered as Nakane Masaki. The given name and family name were inadvertently inverted. The correct order is: Masaki Nakane.

---

Korean J Anesthesiol 2018 February 71(1): 75  
<https://doi.org/10.4097/kjae.2018.71.1.75>

© This is an open-access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>), which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Copyright © The Korean Society of Anesthesiologists, 2018

Online access in <http://ekja.org>