



SYSTEMATIC REVIEW

Postural disorders in mouth breathing children: a systematic review



Patricia Dayrell Neiva ^{a,b}, Renata Noce Kirkwood ^{c,*}, Polyana Leite Mendes ^d,
Karl Zabjek ^d, Helena Gonçalves Becker ^a, Sunita Mathur ^d

^a Universidade Federal de Minas Gerais (UFMG), Programa de Pós-graduação Ciências da Saúde: Saúde da Criança e do Adolescente, Departamento de Otorrinolaringologia, Belo Horizonte, MG, Brazil

^b Pontifícia Universidade Católica de Minas Gerais (PUC-MG), Departamento de Fisioterapia, Belo Horizonte, MG, Brazil

^c Universidade Federal de Minas Gerais (UFMG), Programa de Pós-graduação em Ciências da Reabilitação, Belo Horizonte, MG, Brazil

^d University of Toronto, Department of Toronto, Toronto, Ontario, Canada

Received 26 July 2016; received in revised form 9 September 2016; accepted 21 November 2016

Available online 5 July 2017

KEYWORDS

Mouth breathing syndrome;
Oral breathing;
Children;
Posture;
Photography;
Kinematics

Abstract

Background: Mouth breathing syndrome can cause sleep disturbances that compromise the performance of children in school. It might also cause postural abnormalities involving the head and cervical spine; however, the association between postural abnormalities and mouth breathing in children is unclear.

Objective: To assess the methodological quality of studies and determine if there is an association between mouth breathing and postural disorders in children.

Methods: Databases comprised MEDLINE, CINAHL, PEDro, LILACS, EMBASE and Cochrane Central Registrar of Controlled Trials. Searches were until March 2016 and included studies that evaluated postural disorders in children diagnosed with mouth breathing. The Downs and Black checklist was used to evaluate the quality of the evidences.

Results: Ten studies were included totaling 417 children from 5 to 14 years. Two studies used the New York State Postural Rating Scale, seven used photography and one used motion capture to measure posture. The methods used to analyze the data included the Postural Analysis Software (SAPO), Fisiometer, ALCimagem and routines in MATLAB program. Quality assessment resulted in low scores (<14) for all the studies. The main areas of weakness were a clear description of the participants, of the methods used to access posture, of the principal confounders and

* Corresponding author at: Universidade Federal de Minas Gerais – UFMG, Avenida Antônio Carlos, 6627, CEP 31270-901, Belo Horizonte, MG, Brazil.

E-mail: renata.kirkwood@gmail.com (R.N. Kirkwood).

lack of power analysis. External and internal validity were also threatened by the lack of a representative sample and blinding of the participants and assessors, respectively.

Conclusions: The review provides low evidence that mouth-breathing pattern in children between the ages 5–14 years is associated with postural deviations.

© 2017 Associação Brasileira de Pesquisa e Pós-Graduação em Fisioterapia. Published by Elsevier Editora Ltda. All rights reserved.

Introduction

The mouth-breathing syndrome (MBS) is a clinical condition among school-aged children characterized by mixed or mouth supplementary breathing that replaces an exclusively nasal breathing pattern.¹ MBS is associated with disorders of the speech organs and is generally combined with facial deformities, with the potential to progress to cardiorespiratory, endocrine diseases and sleep disturbances that can compromise the behavior and performance of the children in school.² Furthermore, this syndrome has been related to genetic factors, unhealthy oral habits and nasal obstructions of varying degrees of severity and duration and has also been associated with pediatric allergy-related otorhinolaryngology complaints.²

Studies have also correlated this condition with postural abnormalities^{3,4} and suggested that the obstruction or narrowing of the pharyngeal air space would lead to forward head projection to improve the airflow trajectory to the lower respiratory tract, causing further postural adaptations.^{5,6} With altered and sustained head positions, adaptations in the scalene, sternocleidomastoid and upper trapezius muscles might occur, which are normally associated with overuse in response to the nasal obstruction necessary to facilitate oral inspiration.⁷ Despite reports that MBS can cause postural abnormalities,^{6,8,9} the association between postural deviations and MBS is unclear. Some studies have shown that children with mouth breathing present changes in the posture of the head and cervical spine; however, other studies were unable to support such relationships.^{4,10,11}

In addition, different tools have been used to evaluate posture in mouth breathing children that vary from a simple observational chart posture, to photographic analysis or to advanced motion capture systems that used passive markers attached to the segments to determine body angles and distances.^{3,4,10,12} Different methods and marker positioning may lead to discrepancies between study findings. Moreover, forward head position has been described in most studies, however changes in postural alignment beyond the head and neck have also been considered in some studies,^{13,14} which can provide insights on the effects of MBS on the whole body postural alignment. For physical therapists, providing information on the association between mouth-breathing condition and postural changes in children is important for clinical decision-making, and this information is lacking on the literature.

Thus, we carried out a systematic review to assess the methodological quality of studies that evaluated postural disorders in children with mouth breathing and determine if there is an association between mouth breathing and postural disorders. The information from this review could help the development of rehabilitation programs focused on helping these children.

Methods

Search strategy

The following electronic databases were searched: MEDLINE, Cumulative Index to Nursing and Allied Health Literature (CINAHL), Physiotherapy Evidence Database (PEDro), LILACS database (Literatura Latino-Americana e do Caribe em Ciências da Saúde) and Excerpta Medica Database (EMBASE), including Cochrane Central Registrar of Controlled Trials. Randomized and non-randomized controlled trials were included in the review. Searches were performed by one of the authors (PN) and the strategy is described in details in [Appendix 1](#). The search results were limited to studies in children (<18 years), reported in English or Portuguese and include all published studies until March 2016.

Study selection

Two authors (PN and PM) independently reviewed the titles and abstracts retrieved from all databases and determined if the studies met the inclusion criteria. A third author (SM) decided on any disagreement. Studies were included if reported participants with mouth breathing and at least one element of postural assessment obtained with photographs, or by a software program or even a postural observational analysis that considered either the head, neck, shoulder, scapula, spine, pelvis, knee and/or foot. Inclusion criteria for mouth breathing children were based on clinical evaluation and video laryngoscopy. Signs and symptoms included the absence of active lip seal, facial features, presence of deleterious oral habits, breathing pattern during the day and overnight, tongue tonus, associated allergies and difficulty concentrating in school. Studies were excluded if reported the use of X-rays due to its invasiveness, presence of associated syndromes, or if the assessment protocol was not described. Due to the limited literature in this area all studies designs were included.

Table 1 Downs and Black¹⁶ quality assessment scores for all studies (*n* = 10).

Studies and research designs	Reporting (max 10)	External validity (max 3)	Bias (max 7)	Confounding (max 6)	Power (max 1)	Total (maximal score = 27)
Conti et al. ¹¹ Cross-sectional	6	3	1	2	0	12
Bolzan et al. ⁴ Cross-sectional	8	0	2	3	0	13
Okuro et al. ¹² Cross-sectional	6	3	2	1	0	12
Silveira et al. ³ Cross-sectional	4	1	3	0	0	8
Roggia et al. ¹⁴ Cross-sectional	4	0	2	1	0	7
Neiva et al. ¹⁰ Cross-sectional	6	1	2	1	0	10
Yi et al. ¹³ Cross-sectional	5	3	0	2	0	10
Corrêa and Berzin ⁷ Non-randomized clinical trial	6	1	2	0	0	9
Lima et al. ²⁰ Cross-sectional	2	1	2	2	0	7
Krakauer and Gulherme ²¹ Cross-sectional	3	2	1	0	0	6

Data extraction and quality assessment

The PRISMA¹⁵ checklist was used to document the studies and included information on study design, sample size, subject characteristics, etiology of oral breathing, type of postural assessment, measurement tools and outcomes measures and others. The methodological quality of the studies was conducted using the Downs and Black¹⁶ checklist. The Downs and Black¹⁶ consists of five sub-scales that examine the reporting to check if the paper has sufficient information for the reader to make and unbiased assessment of the findings, external validity of the findings, study bias that access biases in the measurements of the intervention and outcomes, confounding that access bias in the subject selection, and power analysis. Most of the scores ranged from 0 to 1, except for item five (confounders) at the reporting sub-scale that ranges from 0 to 2. The total score is 27 (Table 1). A higher score indicates better methodological quality. The following cut-off points have been suggested and adopted in this study to categorize studies by quality: excellent (26–27), good (20–25), fair (15–19) and poor (≤ 14).¹⁷ The Downs and Black checklist has been used previously in systematic reviews with studies of various designs^{17–19} and presented high internal consistency (0.89), test-retest ($r: 0.88$) and inter-rater ($r: 0.75$) reliability.¹⁶

Data extraction and study quality were reviewed independently by two investigators (PN and PM or SM), and disagreements were resolved by consensus. The kappa coefficient test used to verify the agreement between reviewers for the final classification of the studies was 0.90 (95% CI = 0.73–1.00) demonstrating an excellent level of agreement between reviewers.

Results

Study selection and characteristics

Of the 771 articles found, after deleting the duplicates, 81 remained and were retrieved electronically. Three potentially eligible studies were excluded due to the difficulty in retrieving them and failure of the authors to respond the emails. Therefore, 29 full-text articles were assessed for eligibility, and 10 met the criteria for inclusion (Fig. 1). The study designs investigated included one non-randomized clinical trial⁷ and nine cross sectional studies.^{3,4,10–14,20,21} A total of 622 mouth breathing children and 369 healthy control children were included.

The participants had a mean age ranging from 5 to 14 years, 56% of the participants were males ($n = 227$) and 44% females ($n = 190$). Three studies failed to report the gender of the participants.^{11,20,21} The clinical trial included only mouth breathing children,⁷ seven studies included mouth breathing children and a control group of nasal breather children,^{3,10–14,21} and two studies further divided their sample of mouth breathing in obstructive mouth breathing and functional mouth breathing children.^{4,20}

Methodological quality

The Downs and Black¹⁶ checklist for all selected studies ranged from six to 13, below the cut-off point of 14; therefore, the studies presented low quality (Table 1). The lower scores were a result of poor description of the participants in five studies,^{3,7,13,14,20} poor postural

measurement description in seven studies^{3,11–14,20,21}, lack of description of the confounders, such as age, sex and types of mouth breathing condition in eight studies^{3,7,10,12–14,20,21} and if the confounders were similar between the samples in five studies.^{4,7,14,20,21} Two questions from the reporting and four from the internal validity subscales were specific for randomized controlled trials, and the only clinical trial selected for this review failed to meet the criteria. In addition, sample size calculation was not reported in all studies resulting in a zero score for all studies in this subscale. Other relevant problems were absence of blinding of participants and assessors that affect the internal validity of the study, and lack of estimates in the data, such as standard errors and confidence intervals, which prevented a pooled estimate.

Postural assessment tools

Two studies used the New York State Postural Rating Scale (NYPR)^{11,12}, six used photogrammetry^{3,4,7,13,14} and one used a motion capture system (MOCAP).¹⁰ The methods used to analyze images included the Postural Analysis Software (SAPO v0.68°), Fisiometer 3.0° (Fisiometer Ltda, RJ, Brazil), ALCimagem 2-2000 and customer designed program using MATLAB (R2011a). Only one study used a Polaroid Camera²¹ with no specific method to interpret the data. None of the studies investigated the reliability and accuracy of the postural measures proposed. The psychometric properties described below are based on findings from other studies found in the literature.

The studies by Okuro et al.¹² and Conti et al.¹¹ used the New York State Postural Rating Scale (NYPR), which is an observational chart analysis associated with a plumb line to evaluate the alignment of 13 segments of the body.²² Each body segment can be scored as severe, moderate and normal giving a final score of altered or normal general body posture or neck posture. Reliability of the NYPR has been reported for intra- (0.86) and inter-rater (0.92) reliability of forward head posture, however, for other postural measurements the scale is questionable.²²

The study by Krakauer and Guilherme²¹ analyzed observationally photographs taken from a Polaroid camera, without the aid of any method. Photographic images are an important technique to assess posture, but it is known that accuracy increases even further with the use of bony landmarks.²³

The studies by Bolzan et al.⁴, Roggia et al.¹⁴ and Yi et al.¹³ used photographs associated with the Postural Software Analysis (SAPO),²⁴ which is a quantitative method that provides measurements of the alignment of body segments based on photographs of bony landmarks. The method has an excellent inter-rater agreement (>0.90) for measuring forward head posture and scapular symmetries as reported by Santos et al.²⁵ In addition, Ferreira et al.²⁴ reported inter-rater agreement below 0.75 for forward head posture and scapular symmetries with an average error analysis for all the angular and distances measurements of 0.11° and 1.8 mm, respectively. Both studies agreed that the SAPO is a reliable method to obtain postural measurements.

Similarly, Silveira et al.³ applied photography and had the images analyzed by the software Fisiometer 3.0°²⁶ and Corrêa et al.⁷ and Lima et al.²⁰ by the software

ALCimagem 2-2000.²⁷ The information available on the Fisiometer is scarce with one report on a web site stating that the method has excellent intra- and inter-rater reliability for marker positioning at the temporomandibular joint and cervical lordosis.²⁶ For the ALCimagem, Iunes et al.²⁷ reported lower reliability of the method to follow postural changes.

Only the study by Neiva et al.¹⁰ applied the Qualisys Pro-Reflex (Medical AB, Gothenburg, Sweden), which is a motion capture system (MOCAP) with high-resolution cameras and passive markers to assess posture. The data was analyzed by a costume designed program in MATLAB. The accuracy and repeatability of the MOCAP system has been reported elsewhere with excellent test-retest reliability and very high accuracy to measure distances and angles.²³ For more detailed information the reader should refer to recent published reviews.^{28,29}

Neck and head posture

Neck, head posture or forward head projection were the terms used to report head position. Using the SAPO postural program and with landmarks over the tragus of the ear, seventh cervical vertebrae (C7) and a horizontal line, Roggia et al.¹⁴ reported greater forward head position (smaller angle: 46.95°) in a group of female mouth breathing children compared to nose breathing female children (larger angle: 49.49°). Using the same instrument but with markers over the tragus, C7 and the acromion, Yi et al.¹³ also reported greater forward head posture in mouth breathing children (larger angle: 60.36°) associated with reduced cervical lordosis compared to nose breathing children (52.27°).

Assessing posture with the ALCimagem software and using the glabella, external acoustic meatus and the manubrium as the landmarks, Lima et al.²⁰ reported that head projection was significantly different between groups but did not report the values. Corrêa et al.⁷ reported increased forward head for the mouth breathing children (9.21°) at baseline compared to after the treatment (5.99°) using the same tool, but the angle was formed between a plumb line and a line passing over the ear lobe. Silveira et al.³ reported significantly higher head projection on the mouth breathing children, of 14.3 cm, compared to 11.7 cm of nose breathing children using the Fisiometer tool by measuring the distance from a vertical line to a marker placed over the temporomandibular joint. Observationally, using the NYPR scale, Okuro et al.¹² reported odds ratios of 4.27 of mouth breathing children to have forward head posture.

Shoulder and scapular position

Shoulder position was reported quantitatively in three studies using the Fisiometer,³ ALCimagem⁷ and the MOCAP,¹⁰ but none of the studies reported altered shoulder position in mouth breathing children. Based on observation of photographs, Krakauer et al.²¹ reported 60% of shoulder asymmetry in mouth breathing children and 44.4% in nose breathing children between 5 and 8 years; and in 95% and 33.3% in mouth breathing and nose breathing children, respectively, between the ages of 8.1 and 10 years.




Figure 1 PRISMA flow chart of the study.

Neiva et al.,¹⁰ using the MOCAP, investigated the position of the scapulae thoroughly with landmarks at the scapula, clavicle and spine and reported significantly right (mouth breathing $72.39 \times$ nose breathing 81.0 mm) and left (mouth breathing 75.24 mm \times nose breathing 82.64 mm) scapular elevation in mouth breathing children. Using the ALCimagem software and using the angle formed between the scapular prominence (not specified) and a vertical line, Corrêa et al.⁷ reported 9.74° of an abducted scapular angle in mouth breathing children, measured at baseline, and a significant decrease in this angle after physical therapy intervention (8.82°).

Spinal alignment

Silveira et al.³ reported significantly greater neck lordosis in mouth breathing children (7.3 cm) compared to nose breathing children (5.4 cm) assessed with the Fisiometer

by measuring the distance from a straight line to markers placed over the neck and lumbar spine. Lima et al.²⁰ using the same instrument, but with markers located at the greater cervical, thoracic and lumbar curvatures reported significantly lateral deviations of the cervical and thoracic spines in mouth breathing children but failed to report the values. Yi et al.¹³ using the SAPO postural program reported significantly greater lumbar lordosis angles in the mouth breathing group (smaller angle: 102.52°) compared to the nose breathing group (119.84°) through an angle formed between the first lumbar vertebrae, the anterior superior iliac spine and the greater trochanter.

Thoracic kyphosis was reported significantly in two studies^{10,13,20} that used the SAPO postural program. Yi et al.¹³ measured the angle formed between the acromion and first lumbar and seventh thoracic vertebrae, and reported increased thoracic kyphosis in mouth breathing children (45.89°) compared to nose breathing (41.33°).

Table 2 Summary of the studies included in the systematic review ($n=10$).

Studies	Sample and type of diagnosis	Instrument	Postural variables measured	Results	Conclusions
Okuro et al., ¹² Cross-sectional	8–11 years 45 Mouth breathing 16 F–29 M 62 Nose breathing 23 M–39 F Clinical diagnosis	New York Postural Rating (NYPR)	Neck posture: anterior head position with flexion of the lower portion and extension of the upper portion of the spine Alignment of the head: anterior tilting of the head	Mouth breathing: 36 (80%) and nose breathing: 30 (48.4%) OR = 4.27 [95% CI: 1.63–11.42], $p < 0.001$ of having altered head posture. Mouth breathing: 18 (40%) and nose breathing: 33 (53.2%) OR = 0.59 [95% CI: 0.25–1.37], $p = 0.176$ of having altered body posture	Mouth breathing children presented cervical postural changes compared to nose breathing children
Conti et al., ¹¹ Cross-sectional	5–14 years 323 Mouth breathing 124 Nose breathing Clinical diagnosis	New York Postural Rating (NYPR) assessed observationally of 13 body segments and scored as severe, moderate and normal	Full body alignment: posterior and lateral view neck, shoulders, spine, abdomen, hips, feet and arches Detailed description of the measurements were not provided	Mouth breathing: 60.74% normal posture, 29.63% moderate, 9.63% severe postural abnormalities Nose breathing: 56.20% normal posture, 42.98% moderate posture, 0.83% severe postural abnormalities Posterior view: head, shoulder, feet and arches significantly different between groups ($p \leq 0.0002$) Lateral view: thorax, shoulder, spine, trunk and abdomen significantly different between groups ($p \leq 0.0003$). Neck posture did not differ between groups in the lateral view ($p = 0.260$) Odds ratio (95% CI) of the mouth breathing group for the score of full body alignment: Severe × Normal: odds ratio of 24.99 (95% CI: 3.127–194.127) Moderate × Normal: odds ratio of 2.911 (95% CI: 1.775–4.774)	The study showed an association between mouth breathing and body posture in children and adolescents
Bolzan et al., ⁴ Cross-sectional	8–11 years 22 Obstructive mouth breathing 10 F–10 M 15 Habitual mouth breathing 9 F–13 M 15 Nose breathing 13 F–2 M Diagnosis by video nasofibroscopy	Photography Postural Software Analysis (SAPo v0.68 [®])	Lateral view: Head posture measured through the angle formed by the tragus, seventh cervical vertebra and the horizontal line	Head posture: Nose breathing: $46.15^\circ \pm 4.27^\circ$ Obstructive mouth breathing: $45.71^\circ \pm 4.34^\circ$ Habitual mouth breathing: $45.09^\circ \pm 5.42^\circ$ No difference among groups ($p = 0.45$)	The study showed that head posture seems to be not influenced by breathing pattern

Table 2 (Continued)

Studies	Sample and type of diagnosis	Instrument	Postural variables measured	Results	Conclusions
Silveira et al., ³ Cross-sectional	8–12 years 17 Mouth breathing 7 F–10 M 17 Nose breathing 9 F–11 M Clinical diagnosis	Photography Postural Software Fisiometer®	Sagittal view: Head projection: distance from a straight line drawn perpendicular to the ground and passing behind and near to the body to markers placed at temporomandibular joint Shoulder projection: distance from a straight line drawn perpendicular to the ground and passing behind and near to the body to markers placed at the acromioclavicular joint Neck lordosis: distance from a straight line drawn perpendicular to the ground and passing behind and near to the body to markers placed, not clearly specified, on the neck region. Lumbar lordosis: distance from a straight line perpendicular to the ground and passing behind and near to the body to markers placed, not clearly specified, on the lumbar region.	Head projection: Greater distance between the posterior line and the marker at the temporomandibular joint in the mouth breathing group (14.3 cm) \times nose breathing (11.7 cm) group, $p=0.005$ Neck lordosis: Greater distance between the posterior line and the neck in the mouth breathing (7.3 cm) \times nose breathing (5.4 cm) group, $p=0.016$ No differences were found for the shoulder projection distance 13.5 cm \times nose breathing 11.1 cm , $p=0.2$ and for the lumbar lordosis distance (mouth breathing 6.3 cm \times nose breathing 5.9 cm , $p=0.49$	Mouth breathing children have head projection and neck hyperlordosis
Roggia et al., ¹⁴ Cross-sectional	8–12 years 51 Mouth breathing 20 F–31 M 58 Nose breathing 34 F–24 M Clinical diagnosis	Photography Postural Software Analysis (SAPo v0.68 [®])	Full body alignment comparing the right and left sides of the head, pelvis, line of gravity, knee and ankle segments of boys and girls. Horizontal alignment of the head: angle between the tragus, C7 spinous process and the horizontal. The smaller the angle, the greater the forward head posture. Vertical alignment of the body: the angle between acromion, vertical and lateral malleolus. Positive angular measure: body leaning forward, and negative, backward. Horizontal alignment of the pelvis: the angle between the anterior superior iliac spine, posterior-superior and horizontal. Negative angular measurement: concavity close to zero: rectification, less negative closer to normal. Knee angle: angle between the greater trochanter, the knee joint line and lateral malleolus Ankle angle: angle between the knee joint line, lateral malleolus and horizontal. Measure angle $>90^\circ$: tibia tilted back, if $<90^\circ$: tibia bending forward	Full Body alignment left \times right views: knee angle statistically different between right and left side views of mouth breathing boys (R side: 3.70 ± 6.28 and L side: -0.41 ± 6.03 , $p=0.0143$); no difference found in the group of nose breathing boys and mouth breathing and nose breathing girls Comparisons between groups: Females: significantly difference in the horizontal head alignment angle, with greater forward head posture of the female mouth breathing group ($46.95 \pm 5.65^\circ$) \times Nose breathing group ($49.49 \pm 6.01^\circ$), $p=0.0486$; but no difference in the other measurements Males: significantly difference in ankle angle between males of the Mouth breathing \times nose breathing (mouth breathing: $84.74^\circ \pm 3.12^\circ$ and nose breathing: $83.1^\circ \pm 2.74^\circ$, $p=0.0034$); but no difference in the other measurements	Mouth breathing school boys presented hyperextended knee and decreased ankle angle compared to mouth breathing boys, suggesting an adaptation due to the excess of knee extension Mouth breathing school girls presented forward head posture compared to nose breathing girls

Table 2 (Continued)

Studies	Sample and type of diagnosis	Instrument	Postural variables measured	Results	Conclusions
Neiva et al., ¹⁰ Cross-sectional	8–12 years 21 Mouth breathing 21 Nose breathing Diagnosis by Video nasofibroscopy	Qualisys ProReflex system MATLAB for data analysis	Scapular elevation: obtained from vertical distance in millimeters from the marker positioned over C7 to the centroid point of the scapula. Greater scapular elevation indicated a lesser distance between markers Protrusion of the shoulders: angle obtained from the intersection of a straight line passing through the posterior angle of the acromion and C7 with a straight horizontal line intercepting the posterior acromion on the sagittal plane. An increased angle indicates that the shoulder is projected forward in relation to C7 Anterior tilt of the scapula: angle obtained from the intersection of a straight line passing through C7 and T7 markers and a straight line passing through markers on the posterior angle of the acromion and centroid of the scapula Scapular abduction: corresponds to the horizontal distance in millimeters from the centroid point of the scapula to the spinal column Forward head position: obtained from the angle formed by the intersection of a straight line passing through the marker on the tragus of the ear and C7 and a straight horizontal line intercepting C7 on the sagittal plane. This angle described the position of the head in relation to C7 Thoracic kyphosis: measured as the sum of the angles formed by the intersection of a straight line passing through T2 and the marker positioned 4.5 cm below T2 with a vertical axis and the angle formed by the intersection of a straight line passing through T12 and a marker positioned 4.5 cm above T12 with the same vertical axis Upward rotation of the scapula: angle obtained from the intersection of a straight line passing through the C7 and T7 markers and a line passing through the markers over the medial edge of the root of the scapular spine and the inferior angle of the scapula Internal rotation of the scapula: obtained from the intersection of a horizontal rod positioned in the abdominal region (frontal plane) with a straight line passing through markers located on the root of the scapula spine and the posterior angle of the acromion.	MB children presented significantly increased superior scapular position of the right (72.39 ± 10.71 mm) and left scapula (75.24 ± 10.45 mm) compared to the right (81.0 ± 9.85 mm) and left scapular position (82.64 ± 8.40 mm) of the nose breathing group ($p < 0.05$) Shoulder protrusion: no difference in the right ($141.54^\circ \pm 11.2^\circ$) and left ($138.31^\circ \pm 13.74^\circ$) of the mouth breathing group compared to the right ($139.02^\circ \pm 11.2^\circ$) and left ($132.63^\circ \pm 9.87^\circ$) of the nose breathing group ($p > 0.05$) Anterior tilt of the scapula: no difference in the right ($49.56^\circ \pm 6.42^\circ$) and left ($49.27^\circ \pm 7.34^\circ$) of the mouth breathing group compared to the right ($50.8^\circ \pm 7.39^\circ$) and left ($48.46^\circ \pm 9.95^\circ$) of the nose breathing group ($p > 0.05$) Scapular abduction: no difference in the right (95.11 ± 7.46 mm) and left (96.13 ± 6.03 mm) of the mouth breathing group compared to the right (96.3 ± 6.3 mm) and left (101.09 ± 10.96 mm) of the nose breathing group ($p > 0.05$) Forward head position: no difference in the right ($48.9^\circ \pm 4.4^\circ$) and left ($48.10^\circ \pm 6.8^\circ$) of the mouth breathing group compared to the right ($47.59^\circ \pm 4.6^\circ$) and left ($48.50^\circ \pm 6.30^\circ$) of the nose breathing group ($p > 0.05$) Thoracic kyphosis: no difference in the mouth breathing group ($31.96^\circ \pm 10.97$) compared to the nose breathing groups ($30.82^\circ \pm 16.93^\circ$) ($p > 0.05$) Upward rotation of the scapula: no difference in the right ($1.01^\circ \pm 6.84^\circ$) and left ($-3.42^\circ \pm 6.15^\circ$) of the mouth breathing group compared to the right ($2.98^\circ \pm 3.91^\circ$) and left ($-2.61^\circ \pm 5.86^\circ$) of the nose breathing group ($p > 0.05$) Internal rotation of the scapula: no difference in the right ($35.24^\circ \pm 4.2^\circ$) and left ($33.81^\circ \pm 5.69^\circ$) of the mouth breathing group compared to the right ($38.43^\circ \pm 6.01^\circ$) and left ($35.55^\circ \pm 6.72^\circ$) of the nose breathing group ($p > 0.05$)	Increased superior scapular position may be due to the forward head, clinically observed in mouth breathing children, that leads to altered position of the mandible

Table 2 (Continued)

Studies	Sample and type of diagnosis	Instrument	Postural variables measured	Results	Conclusions
Yi et al., ¹³ Cross-sectional	5–12 years 30 Mouth breathing 22 Nose breathing Diagnosis by video nasofibroscopy	Photography Postural Software Analysis (SAPO) v0.68°	Cervical lordosis: angle formed by the tragus of the ear, the acromion and C7. The larger the angle, the further forward the position of the head and the lower the degree of cervical lordosis. Thoracic kyphosis: angle formed by the acromion to L1 and from L1 to T7, where L1 was the apex of the angle. The larger the angle, the greater the degree of thoracic kyphosis. Lumbar lordosis: angle drawn between three anatomic points: L1, ASIS and the greater trochanters, where ASIS was the apex of the angle. The smaller the angle, the greater the degree of lumbar lordosis. Pelvic position: angle drawn between the ASIS, the midpoint of the knee joint on the lateral face and the greater trochanters where the midpoint of the joint line was the apex of the angle. The greater the angle, the greater the pelvic tilt.	Cervical lordosis: significantly difference between mouth breathing ($60.36^\circ \pm 9.54^\circ$) × nose breathing ($52.27^\circ \pm 8.58^\circ$), $p = 0.003$, with reduced cervical lordosis found on the mouth breathing group. Lumbar lordosis: significantly difference between mouth breathing ($102.52^\circ \pm 9.67^\circ$) × nose breathing ($119.84^\circ \pm 5.35^\circ$), $p = 0.001$, with increased lumbar lordosis found on the mouth breathing group. Thoracic kyphosis: significantly difference between mouth breathing ($45.89^\circ \pm 5.26^\circ$) × nose breathing ($41.33^\circ \pm 4.64^\circ$), $p = 0.002$, with increased thoracic kyphosis found on the mouth breathing group. Pelvic position: significantly difference between mouth breathing ($9.98^\circ \pm 1.49^\circ$) × nose breathing ($6.93^\circ \pm 1.00^\circ$), $p = 0.001$, with increased anterior tilt of the pelvis found on the mouth breathing group.	Mouth breathing children project their heads forward to facilitate airflow. In addition, the children presented decreased cervical lordosis due to the forward projection of the head, increased lumbar lordosis, thoracic kyphosis and anterior pelvic tilt
Corrêa & Bezin ⁷ Non-randomized clinical trial	9–11 years 19 Mouth breathing 8 F-11 M Diagnosed by video nasofibroscopy	Photography Postural Software ALCImagem	Lateral view: Forward head: angle formed between the plumb line and a line passing the ear lobe Forward shoulder: angle formed between a plumb line a line passing over the acromion Flexion/extension head: angle formed between the plumb line and a line passing over the mentum Scapula abduction or winged scapula: angle formed between the plumb line and a line passing over the scapular prominence Frontal view: Head Tilt right/left: angle formed between a vertical line drawn through the manubrium and a line passing through the left/right ear lobe Shoulder elevation right/left: angle formed between a vertical line drawn through the manubrium and a line passing through the left/right coracoid process Posterior view: Scapular abduction left/right: angle formed between a line passing the superior scapular angle, C7 and superior scapular angle	Baseline × Post treatment Forward head: $9.21^\circ \pm 4.51^\circ \times 5.99^\circ \pm 1.49^\circ$, $p = 0.003$ Forward shoulder: $4.18^\circ \pm 2.53^\circ \times 2.9^\circ \pm 3.13^\circ$, $p = 0.121$ Flexion/extension head: $17.68^\circ \pm 4.1^\circ \times 15.83^\circ \pm 3.04^\circ$, $p = 0.1027$ Abducted/winged scapula: $9.74^\circ \pm 2.25^\circ \times 8.82^\circ \pm 2.49^\circ$, $p = 0.0458$ Head tilt right: $30.19^\circ \pm 2.6^\circ \times 29.66^\circ \pm 3.5^\circ$, $p = 0.379$ Head tilt left: $30.96^\circ \pm 3.2^\circ \times 31.33^\circ \pm 2.6^\circ$, $p = 0.454$ Shoulder elevation right: $84.73^\circ \pm 3.22^\circ \times 86.37^\circ \pm 3.53^\circ$, $p = 0.236$ Shoulder elevation left: $85.56^\circ \pm 3.14^\circ \times 87.07^\circ \pm 4.29^\circ$, $p = 0.113$ Scapular abduction right: $47.09^\circ \pm 6.61^\circ \times 44.14^\circ \pm 5.98^\circ$, $p = 0.0158$ Scapular abduction left: $103.89^\circ \pm 12.92^\circ \times 103.42^\circ \pm 10.12^\circ$, $p = 0.873$	The exercise program was effective in reducing the forward head posture, scapular abduction, shoulder elevation and internal rotation of the scapula on the right side of mouth breathing children

Table 2 (Continued)

Studies	Sample and type of diagnosis	Instrument	Postural variables measured	Results	Conclusions
Lima et al. ²⁰ Cross-sectional	8–10 years 17 Obstructive mouth breathing 26 Functional mouth breathing 19 Nose breathing Clinical Diagnosis	Photography Postural Software ALCimagem	Lateral view: Chin retraction: angle formed between the glabella, external acoustic meatus and mentum Head forward projection: angle formed between the glabella, external acoustic meatus and manubrium sterni Lateral deviation of the cervical, thoracic and lumbar spine: markers located at greatest cervical, thoracic and lumbar curvatures Pelvis and knee alignment: markers at ASIS, gluteal line, and lateral condyles of the femur Frontal View: Horizontalization of the collarbones: coracoid process, condylar angle and a horizontal line Posterior view: Scapular position: angle between the superior and inferior angle of the scapular and T12	Chin retraction ($p=0.014$) and head forward projection ($p=0.0036$) were significantly different between the obstructive mouth breathing \times nose breathing group. The obstructive mouth breathing group showed more chin and head deviations. Cervical deviations were significantly different between obstructive mouth breathing \times nose breathing ($p=0.0004$) and obstructive mouth breathing \times functional mouth breathing ($p=0.0148$). Thoracic deviations were significantly different between obstructive mouth breathing \times nose breathing ($p=0.0009$) and obstructive mouth breathing \times functional mouth breathing ($p=0.0073$). The authors classified it at altered thoracic convexity. All the other measurements were not significantly different between the three groups. The values of the angles were not reported.	Obstructive mouth breathing and Functional mouth breathing children presented different spine postural alterations; therefore should be treated differently
Krakauer & Guilherme ²¹ Cross-sectional	5–10 years 30 Nose breathing Clinical diagnosis	Polaroid Camera Observational analysis of the photos	Frontal view: symmetry/asymmetry of the shoulder in relation to the ground. Lateral view: head position (anterior, posterior, or normal) in relation to the neck Dorsal view: symmetry/asymmetry of the scapulae in relation to a perpendicular axis passing through the body.	Frontal view: asymmetry observed in 60% mouth breathing children and 44.4% nose breathing children between 5 and 8 years; and 95% and 33.3% of mouth breathing and nose breathing children, respectively, between the ages of 8.1 and 10 years Lateral view: alteration observed in 80% mouth breathing children and 33.3% nose breathing children between 5 and 8 years; and 100% and 25% of mouth breathing and nose breathing children, respectively, between the ages of 8.1 and 10 years Dorsal view: asymmetry observed in 83.3% mouth breathing children and 44.4% nose breathing children between 5 and 8 years; and 70% and 41.7% of mouth breathing and nose breathing children, respectively, between the ages of 8.1 and 10 years	Children with nose breathing, age 8 and above present better posture than those who continue mouth breathing beyond age 8

M, male; F, female; ASIS, anterior superior iliac spine; T12, twelve thoracic vertebrae; C7, seventh cervical vertebrae.

Lima et al.²⁰ reported significant difference in thoracic convexity between the obstructive mouth breathing and nose breathing groups as well as between the functional mouth breathing and nose breathing groups, but failed to report the landmarks used and the values obtained (Table 2).

Full body, knee and pelvis alignment

Severe and moderate full body alignment was observed qualitatively using the NYPR scale by Conti et al.¹¹ in 9.63% (323/31) and 29.63% (323/96) of mouth breathing children, respectively, compared to 0.83% (124/1) and 42.98% (124/53) of severe and moderate body alignment, respectively, in nose breathing children. The authors also reported significant difference in the posterior and lateral views alignment between groups, except for the neck alignment.¹¹

Tibial and pelvis alignment were assessed with the SAPO postural software with an angle formed by a line between the knee joint line, lateral malleolus and a horizontal. Roggia et al.¹⁴ reported an increased forward bending of the tibia (of 84.74° meaning the ankle joint was more dorsiflexed) in mouth breathing compared to nose breathing children (83.1°). Yi et al.¹³ found an increased anterior tilt of the pelvis in mouth breathing children (9.98°) compared to nose breathing (6.93°). The angle was formed by a line passing over the anterior superior iliac spine, midpoint of the knee joint and greater trochanter.

Discussion

This systematic review assessed the association between mouth breathing patterns in children and postural alignment. Ten studies were found that addressed our question, of those, six studies found forward head posture in mouth breathing children, six found a higher percentage of full body misalignment and two studies reported scapular deviations. However, all the studies scored less than 14 on the Downs and Black¹⁶ checklist. Therefore, there is low quality evidence suggesting that mouth breathing pattern is associated with postural misalignment in children between the ages of 5–14 years. In addition, the review also highlights the lack of standardized terminology, bone landmarks, reference values and the reliability and validity of the methods and measurement tools used to evaluate posture in mouth breathing children.

The main areas of weakness were a clear description of the participants, of the methods used to access posture and the principal confounders. External validity was threatened by the lack of representative samples in half of the studies as well as internal validity, since none of the studies attempted to blind the participants or assessors. Another important area of concern was the adjustment of confounders for the postural analysis. For example, there are many factors that influence postural assessment such as age, gender, body mass and height, postural coordination and the presence of pain.³⁰ Furthermore, growth spurts occur around 9–12 year and may cause alterations in body shape and dimensions that could affect muscle flexibility.^{31,30} As a consequence, postural misalignments would be expected in this age group. Penha et al.³² reported a high frequency of postural alterations in healthy children from 7 to 10 years of age. In

addition, Ludwig et al.³¹ and McEvoy and Grimmer,³⁰ showed the effect of age on standing posture. Therefore, age should be considered when evaluating posture in mouth breathing children.

Measurement errors, such as marker positioning and marker movement are also another important source of errors that should be considered. So far, there is no standard recommendation to measure posture. Photogrammetry has been the choice of measurement being more precise than observational postural assessment.³³ However, it is paramount to increase measurement reliability and accuracy. Training of the evaluators for landmark palpation and marker positioning as well as knowledge of the potential device errors are essential. Standardization of the body position and the type of information provided to the participant is also helpful to improve reliability. According to Zabjek et al.³⁴ data collection duration is another important aspect of reliability. The lack of reporting the reliability of the measurements threatens the studies internal validity and credibility of the studies.

The literature supports that forward head posture is a strategy adopted by mouth breathing children to facilitate and accelerate airflow.³⁵ Forward head position is a combination of extension of the upper cervical spine, flexion of the lower cervical and thoracic spine leading to an increase in cervical lordosis.³⁶ Prolonged head forward posture causes increased strain on the extensor muscles of the head and stretches the infrahyoid muscles creating an inferior and posterior traction the hyoid bone. As a consequence, the mandible is pulled in a direction of retraction and depression.³⁶ Therefore, measurements should include bony landmarks at the lower cervical spine or upper thoracic spine and at the mandible to reflect a true forward head posture. For example, an angle formed by the ear lobe, C7 and the mandible would reflect a true head forward, since it would also include mandible retraction. Measurements with landmarks at the trunk and shoulder, as presented by Yi et al.¹³ and Lima et al.²⁰ might be influenced by the position of these structures. Shoulder elevation would place the acromion slightly higher and would affect the head position angle as proposed by Yi et al.¹³ On the other hand, trunk rotation would also affect the marker positioning over the manubrium as proposed by Lima et al.²⁰ In addition, measurements that rely on perpendicular lines^{3,7} are subject to errors due to the lack of consistent bony landmarks and body motion. Therefore, markers placed over segments that reflect the true head forward measurement might improve the reliability and facilitate comparisons among studies.

Limitations of this review included the heterogeneity of the measurements and tools used to access posture, lack of power and internal and external validity of the studies. In addition, studies in languages other than English or Portuguese were also excluded. There is no evidence that non-English written papers are substantially better than the ones published in English.^{37,38} Methodological quality can be improved in future studies by considering effect size and power analysis, controlled designs, appropriate analysis of the reliability and validity of the instruments and landmarks used to access posture, blinding of the examiners, random sampling, and eligibility criteria of the participants by controlling the types of mouth breathing patterns. Understanding the prime alterations of posture in mouth breathing

children and being able to reflect on that earlier is important to increase the quality of life of these children.

Conclusion

The result of this review provides low evidence that mouth-breathing pattern in children between the ages 5–14 years is associated with postural deviations. Future studies should consider the methodological issues discussed and attempt to develop standardized methods to assess postural measurements in mouth breathing children.

Conflicts of interest

The authors declare no conflicts of interest.

Acknowledgements

The Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES), a Brazilian funding agency, supported the primary author during her partial doctoral program in Canada.

Appendix 1.

The following combination of concepts and Boolean operators were used to retrieve the studies:

- 01 "mouth breathing"[MeSH Terms] OR mouth breathing[Text Word]
 - 02 oral breathing
 - 03 oral respiration
 - 04 breathing mouth
 - 05 1 or 2 or 3 or 4
 - 06 postural disorders
 - 07 posture
 - 08 postural
 - 09 postural abnormalities
 - 10 kinesis
 - 11 positioning
 - 12 orthostatic
 - 13 6 or 7 or 8 or 9 or 10 or 11
 - 14 05 and 13
 - 15 "child"[MeSH Terms] OR children[Text Word]
 - 16 preschool children
 - 17 adolescent
 - 18 15 or 16 or 17
 - 19 14 and 18
 - 20 research design
 - 21 comparative study
 - 22 20 or 21
 - 23 19 and 22
-
2. Felcar JM, Bueno IR, Massan AC, Torezan RP, Cardoso JR. Prevalence of mouth breathing in children from an elementary school. *Cien Saude Colet.* 2010;15:437–444.
 3. Silveira WD, Mello FC, Guimaraes FS, Menezes SL. Postural alterations and pulmonary function of mouth-breathing children. *Braz J Otorhinolaryngol.* 2010;76:683–686.
 4. Bolzan G, Souza J, Boton L, Silva A, Corrêa E. Facila type and head posture of nasal and mouth-breathing children. *J Soc Brasil Fonoaudiol.* 2011;23:315–320.
 5. Kamper SJ, Henschke N, Hestbaek L, Dunn KM, Williams CM. Musculoskeletal pain in children and adolescents. *Braz J Phys Ther.* 2016;20:275–284.
 6. Cuccia AM, Lotti M, Caradonna D. Oral breathing and head posture. *Angle Orthod.* 2008;78:77–82.
 7. Correa EC, Berzin F. Efficacy of physical therapy on cervical muscle activity and on body posture in school-age mouth breathing children. *Int J Pediatr Otorhinolaryngol.* 2007;71:1527–1535.
 8. Chaves TC, de Andrade e Silva TS, Monteiro SA, Watanabe PC, Oliveira AS, Grossi DB. Craniocervical posture and hyoid bone position in children with mild and moderate asthma and mouth breathing. *Int J Pediatr Otorhinolaryngol.* 2010;74:1021–1027.
 9. Malhotra S, Pandey RK, Nagar A, Agarwal SP, Gupta VK. The effect of mouth breathing on dentofacial morphology of growing child. *J Indian Soc Pedod Prev Dent.* 2012;30:27–31.
 10. Neiva PD, Kirkwood RN, Godinho R. Orientation and position of head posture, scapula and thoracic spine in mouth-breathing children. *Int J Pediatr Otorhinolaryngol.* 2009;73:227–236.
 11. Conti PB, Sakano E, Ribeiro MA, Schivinski CI, Ribeiro JD. Assessment of the body posture of mouth-breathing children and adolescents. *J Pediatr (Rio J).* 2011;87:357–363.
 12. Okuro RT, Morcillo AM, Sakano E, Schivinski CI, Ribeiro MA, Ribeiro JD. Exercise capacity, respiratory mechanics and posture in mouth breathers. *Braz J Otorhinolaryngol.* 2011;77:656–662.
 13. Yi LC, Jardim JR, Inoue DP, Pignatari SS. The relationship between excursion of the diaphragm and curvatures of the spinal column in mouth breathing children. *J Pediatr (Rio J).* 2008;84:171–177.
 14. Roggia B, Correa B, Pranke GI, Facco R, Rossi AG. Postural control of mouth breathing school aged children regarding gender. *Pro Fono.* 2010;22:433–438.
 15. Liberati A, Altman DG, Tetzlaff J, et al. The PRISMA statement for reporting systematic reviews and meta-analyses of studies that evaluate health care interventions: explanation and elaboration. *Ann Intern Med.* 2009;151:W65–W94.
 16. Downs SH, Black N. The feasibility of creating a checklist for the assessment of the methodological quality both of randomised and non-randomised studies of health care interventions. *J Epidemiol Community Health.* 1998;52:377–384.
 17. Silverman SR, Schertz LA, Yuen HK, Lowman JD, Bickel CS. Systematic review of the methodological quality and outcome measures utilized in exercise interventions for adults with spinal cord injury. *Spinal Cord.* 2012;50:718–727.
 18. Crow J, Pizzari T, Buttifant D. Muscle onset can be improved by therapeutic exercise: a systematic review. *Phys Ther Sport.* 2011;12:199–209.
 19. Huffer D, Hing W, Newton R, Clair M. Strength training for plantar fasciitis and the intrinsic foot musculature: a systematic review. *Phys Ther Sport.* 2016, <http://dx.doi.org/10.1016/j.ptsp.2016.08.008>, pii:S1466-853X(16)30075-X.
 20. Lima LC, Barauna MA, Sologuren MJ, Canto RS, Gastaldi AC. Postural alterations in children with mouth breathing assessed by computerized biophotogrammetry. *J Appl Oral Sci.* 2004;12:232–237.

References

1. Barros JR, Becker HM, Pinto JA. Evaluation of atopy among mouth-breathing pediatric patients referred for treatment to a tertiary care center. *J Pediatr (Rio J).* 2006;82:458–464.

21. Krakauer LH, Guilherme A. Relationship between mouth breathing and postural alterations of children: a descriptive analysis. *Int J Orofacial Myol.* 2000;26:13–23.
22. Arnold CM, Beatty B, Harrison EL, Olszynski W. The reliability of five clinical postural alignment measures for women with osteoporosis. *Physiother Canada.* 2000;52:286–294.
23. Liu H, Holt C, Evans S. Accuracy and repeatability of an optical motion analysis system for measuring small deformations of biological tissues. *J Biomech.* 2007;40:210–214.
24. Ferreira EA, Duarte M, Maldonado EP, Burke TN, Marques AP. Postural assessment software (PAS/SAPO): validation and reliability. *Clinics (Sao Paulo).* 2010;65:675–681.
25. Santos MM, Silva MPC, Sanada LS, Alves CRJ. Photogrammetric postural analysis on healthy seven to ten-year-old children: interrater reliability. *Rev Bras Fisioter.* 2009;13: 350–355.
26. Dias F, Antonio H. Relatório de confiabilidade do Fisiometer. In: *Baseado na dissertação de Waleska da Silveira Venturelli.*; 2006:1–22.
27. Iunes DH, Castro FA, Salgado HS, Moura IC, Oliveira AS, Bevilacqua-Grossi D. Confiabilidade intra e interexaminadores e repetibilidade da avaliação postural pelo fotogrametria. *Rev Bras Fisioter.* 2005;9:327–334.
28. Fortin C, Feldman DE, Cheriet F, Labelle H. Clinical methods for quantifying body segment posture: a literature review. *Disabil Rehabil.* 2011;33:367–383.
29. Dunk NM, Lalonde J, Callaghan JP. Implications for the use of postural analysis as a clinical diagnostic tool: reliability of quantifying upright standing spinal postures from photographic images. *J Manip Physiol Ther.* 2005;28: 386–392.
30. McEvoy MP, Grimmer K. Reliability of upright posture measurements in primary school children. *BMC Musculoskelet Disord.* 2005;6:35.
31. Ludwig O, Mazet C, Mazet D, Hammes A, Schmitt E. Age-dependency of posture parameters in children and adolescents. *J Phys Ther Sci.* 2016;28:1607–1610.
32. Penha PJ, Baldini M, Joao SM. Spinal postural alignment variance according to sex and age in 7- and 8-year-old children. *J Manip Physiol Ther.* 2009;32:154–159.
33. Iunes DH, Grossi DB, Oliveira AS, Castro FA, Salgado HS. Comparative analysis between visual and computerized photogrammetry postural assessment. *Rev Bras Fisioter.* 2009;13:308–315.
34. Zabjek KF, Leroux MA, Coillard C, Rivard CH, Prince F. Evaluation of segmental postural characteristics during quiet standing in control and Idiopathic Scoliosis patients. *Clin Biomech (Bristol, Avon).* 2005;20:483–490.
35. Solow B. Growth changes in head posture related to craniofacial development. *Am J Orthod Dentofac Orthop.* 1986;89:132–140.
36. Neumann DA. Kinesiology of the musculoskeletal system. *Foundations for physical rehabilitation.* 1 vol, 1st ed. St. Louis: Mosby, Inc.; 2002:1–597.
37. Mancini MC, Cardoso JR, Sampaio RF, Costa LC, Cabral CM, Costa LO. Tutorial for writing systematic reviews for the Brazilian Journal of Physical Therapy (BJPT). *Rev Bras Fisioter.* 2014;18:471–480.
38. Shiwa SR, Moseley AM, Maher CG, Pena Costa LO. Language of publication has a small influence on the quality of reports of controlled trials of physiotherapy interventions. *J Clin Epidemiol.* 2013;66:78–84.