



Published in final edited form as:

Adv Synth Catal. 2017 November 23; 359(22): 4062–4066. doi:10.1002/adsc.201701179.

Gold-catalyzed Fluorination of Alkynyl Esters and Ketones: Efficient Access to Fluorinated 1,3-Dicarbonyl Compounds

Xiaojun Zeng^a, Zhichao Lu^b, Shiwen Liu^a, Gerald B. Hammond^b, and Bo Xu^a


^aCollege of Chemistry, Chemical Engineering and Biotechnology, Donghua University, 2999 North Renmin Lu, Shanghai 201620, China

^bDepartment of Chemistry, University of Louisville, Louisville, KY 40292, USA

Abstract

We developed an efficient synthesis of 2-fluoro-1,3-dicarbonyl compounds using readily available alkynyl ketones or esters as starting material. The key step is the insertion of hydrogen fluoride (HF) to the gold carbene intermediate generated from cationic gold catalyzed addition of *N*-oxides to alkynyl ketones or esters. This method gives excellent chemical yields and regioselectivity with good functional group tolerance.

Graphical abstract



Keywords

Gold Catalysis; Fluorination; 1,3-Dicarbonyl Compounds; Alkynyl Esters and Ketones

Due to fluorine's unique properties such as small size and metabolically resistant C-F bond, the selective substitution of hydrogen by fluorine constitutes a key strategy in pharmaceuticals, agrochemicals and materials research.^[1] More specifically, fluorinated 1,3-dicarbonyl compounds are important fluorine building blocks and motifs in medicine and materials.^[2] Literature syntheses of fluorinated 1,3-dicarbonyl compounds are generally based on the electrophilic fluorination of 1,3-dicarbonyl compound precursors (Scheme 1, a–e). For examples, Shibata and coworkers reported the fluorination of β -keto-esters using methyl NFSI as fluorination agent under Lewis acid-catalysis (Scheme 1a).^[3] Many other electrophilic fluorination systems like Selectfluor in imidazolium ionic liquid (Scheme 1b),

Correspondence to: Gerald B. Hammond; Bo Xu.

Supporting information for this article is available on the WWW under <http://dx.doi.org/10.1002/adsc.201701179>.

[4] Selectfluor in continuous-flow microreactors, [5] Selectfluor/ZnCl₂ (Scheme 1c) have been developed. [6] The fluorination of 1,3-dicarbonyl compound precursors can also be achieved using Ar-I/HF-pyridine/mCPBA (Scheme 1d) [7] and HF/PhIO fluorination system (Scheme 1e). [8] In addition, the Hayes and Moody group reported a nucleophilic fluorination of α -diazo- β -ketoesters with HBF₄ as a fluorine-carrier (Scheme 1f). [9] These methods do have some drawbacks though, among them: 1) relatively limited scope; 2) non-atom-economic fluorination systems; 3) poor functional group tolerance caused by the use of strong oxidants; and 4) difficulty in controlling overreaction (i.e., difluorination). [6], [10]


Because alkynyl ketones or esters are readily available compounds, and because the alkynyl group can be considered as a masked ketone, the task of converting alkynyl ketones or esters to fluorinated 1,3-dicarbonyl compounds is a straightforward synthetic method. Hydrogen fluoride (HF) is one the most atom-economic fluorination reagent. Furthermore, HF can be regarded as compatible with cationic gold catalysts; this was demonstrated by the synthesis of fluoroalkenes through the gold catalyzed addition of HF to alkynes [11] and the synthesis of α -fluoroketone *via* HF insertion to a gold carbene intermediate. [12] These methods proved efficient when a terminal alkyne were utilized as substrate; internal alkynes showed low or no reactivity, and the regioselectivity depended on steric and electronic biases at either end of the alkyne. We proposed that the ester or ketone group could function as an activator and director in the fluorination of an alkyne. Herein, we report a widely applicable, highly efficient and divergent synthesis of 2-fluoro-1,3-dicarbonyl compounds from readily available alkynyl ketones/esters (Scheme 1g).

We carried out the fluorination of alkynyl ketone **1a** as our model reaction (Table 1). The oxidant we use was an *N*-oxide, which is mild, [13], [14] hence, we expected good functional group tolerance. We recently demonstrated that excess amounts of silver salt in gold catalysis almost always had a deleterious effect on the reactivity of gold catalysts and that a preformed gold catalyst like L-Au-NTf₂, usually exhibited the best reactivity. [15] So, initially we used the preformed gold catalyst JohnPhos-AuNTf₂. [16] We screened the effects of *N*-oxides **2** (Table 1, entries 1–4), and found that *N*-oxide **2d** gave the best result (Table 1, entry 4). The screening of various solvents indicated that PhCF₃ was the optimal solvent (Table 1, entries 4–7). We proceeded to investigate the effects of the ligands (Table 1, entries 6–10). [17] All the ligands tested gave reasonably good results, but, among them, JohnPhos was slightly better (Table 1, entries 6–10). We also confirmed that a silver salt alone was not able to catalyze this reaction (Table 1, entry 11).

With the optimized conditions in hand, we explored the substrate scope and functional group tolerance for the fluorination of alkynyl ketones (Table 2). We evaluated the effect of different [R¹, R²] combinations in the alkynyl ketones **1** (Table 2). The substitution pattern (*meta*, *para*) and electronic properties of aromatic substituents (electron deficient or rich) of R¹ played a small role; good yields were obtained regardless (Table 2, 3a–3f). Various functional groups such as nitrile (Table 2, 3b), ether (Table 2, 3c), halide (Table 2, 3d), ketone (Table 2, 3e), ester (Table 2, 3f, 3h), and sulfonate (Table 2, 3g) were well tolerated. It should be noted that when both R¹, R² were aromatics, no reaction took place (Table 2, 3i). Lack of reactivity in this case may be caused by the fact that the triple bond and two phenyl rings form a highly delocalized system which weaken the polarization of triple bond.

Our reaction conditions could also be applied to the fluorination of alkynyl esters **4**, leading to the synthesis of 2-fluoro-3-ketoester **5** (Table 3). First, we evaluated the effect of R¹ substitution of the 3-ketoester **4**: electron rich and electron deficient substituents only had a very small effect on the chemical yields (Table 3, **5a–5l**). The substitution pattern (*ortho*, *meta*, *para*) of R¹ only played a small role; good yields were obtained regardless (Table 3, **5a–5l**). And various functional groups such as nitro group (Table 2, **5c**), chloride (Table 3, **5d**, **5j**, **5k**), ester (Table 3, **5f**), sulfonamide (Table 3, **5g**), benzyl ether (Table 3, **5h**), fluoride (Table 3, **5i**) and alkene (Table 3, **5n**) were well tolerated. We also evaluated more [R¹, R²] combinations, such as [R¹ = alkyl, R² = alkyl] (Table 3, **5m–5n**), [R¹ = alkyl, R² = aryl] (Table 3, **5o**). These combinations gave good yields of product **5**. It should be noted that an activated ester (phenol ester, **5o**) survived our conditions. Activated esters like **5o** can be used for further transformations like amide formation.

The proposed mechanism is illustrated in Scheme 2. First, addition of *N*-oxides **2** to the cationic gold activated alkyne starting material gives vinyl gold intermediate **A**, which then transforms into the reactive gold carbene intermediate **B**.^[18] The hydrogen bonding interaction between HF and **A** may facilitate the formation of gold carbene **B**. Finally, insertion of HF gold carbene **B** gives products **3** or **5**.^[12]



(eq-1)

Our protocol could be easily scaled up without affecting the chemical yield (eq 1). The versatility of fluorinated 1,3-dicarbonyl compounds **3** and **5** in organic synthesis is clearly underscored in Scheme 3. For examples, reaction of **3a** with hydrazine leads to a facile synthesis of fluorinated pyrazole **6**.^[19] Iridium-catalyzed asymmetric allylic alkylation of **5a** produces the highly functionalized fluoro-building block **7**.^[20] And the guanidine-catalyzed Michael addition of **5a** gives **8** in good yield and selectivity.^[21] [3+3] Cyclization of **3a** with 1,3-bis(silyl enol ethers) leads to an efficient synthesis of fluorinated phenols **9**.^[22] Asymmetric Mannich reaction of **5a** with a bifunctional thiourea catalyst furnishes **10**.^[23]

Similarly, a highly enantioselective Michael addition of **3a** to nitrostyrene yields **11** with tertiary stereocenters.^[24] **3** can also be used in the synthesis of fluorinated pyrimidine scaffold **12**.^[25]

In summary, we have developed a widely applicable synthesis of 2-fluoro-1,3-dicarbonyl compounds starting with readily available alkynyl ketones or esters. This method gave excellent chemical yields and regioselectivity with good functional group tolerance. And the reaction can be conducted under ambient atmosphere. Other fluorination systems based on HF/gold combinations are currently being investigated in our laboratories.

Experimental Section

General procedure for fluorination of **1** or **4**

A polypropylene vial was charged with alkynone **1** or alkynoate **4** (0.2 mmol, 1 equiv), 2,6-dibromopyridine *N*-oxide **2d** (0.28 mmol, 70 mg), and HF-pyridine (HF content 70 wt/wt%, 0.8 mmol, 27 mg, 4 equiv) and PhCF₃ (0.4 mL). The reaction was commenced by the addition of JohnPhosAuNTf₂ (5 mol%, 7.75 mg) and the reaction mixture was heated to 40 °C. The progress of the reaction was monitored by TLC. The reaction typically took 4 h or overnight to complete. Upon completion, the mixture was quenched with saturated sodium bicarbonate. The mixture was extracted with DCM and washed with 1N HCl. The organic layers were collected, dried over anhydrous MgSO₄ and filtered. The solvent was removed under reduced pressure and the residue was subjected to flash column chromatography purification (eluent: hexanes/ethyl acetate) to give 2-fluoro-1,3-dicarbonyl compounds **3** or **5**.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments


We are grateful to the National Science Foundation of China for financial support (NSFC-21472018) and China Recruitment Program of Global Experts for financial support. G.B.H. is grateful to the National Institutes of Health (1R01GM121660-01). XZ is grateful to the China Scholarship Council.

References


1. a) Chambers, RD. Fluorine in organic chemistry. Blackwell Publishing Ltd./CRC Press; Boca Raton, FL: 2004. b) Hiyama, T. Organofluorine compounds, chemistry and applications. Springer-Verlag; Berlin: 2000. c) Kirsch, P. Modern fluoroorganic chemistry. Wiley-VCH; Weinheim: 2004. d) Muller K, Faeh C, Diederich F. Science. 2007; 317:1881–1886. [PubMed: 17901324] e) Schlosser M. Angew Chem Int Ed. 1998; 37:1496.f) Soloshonok, VA. Fluorine-containing synthons, ACS symposium series 911. Oxford University Press; Washington, D.C: 2005. g) Uneyama, K. Organofluorine Chemistry. Blackwell publishing; Oxford: 2006.
2. a) Guo Y, Twamley B, Shreeve J n M. Org Biomol Chem. 2009; 7:1716–1722. [PubMed: 19343262] b) Kim K-Y, Kim BC, Lee HB, Shin H. J Org Chem. 2008; 73:8106–8108. [PubMed: 18808183] c) Prakash GKS, Beier P. Angew Chem Int Ed. 2006; 45:2172–2174.d) Burger EC, Barron BR, Tunge JA. Synlett. 2006:2824–2826.e) Davis FA, Han W, Murphy CK. J Org Chem. 1995; 60:4730–4737.f) Dolbier WR Jr, Lee SK, Phanstiel OIV. Tetrahedron. 1991; 47:2065–2072.
3. Fukushi K, Suzuki S, Kamo T, Tokunaga E, Sumii Y, Kagawa T, Kawada K, Shibata N. Green Chem. 2016; 18:1864–1868.
4. Reddy AS, Laali KK. Tetrahedron Lett. 2015; 56:5495–5499.
5. Baumann M, Baxendale IR, Martin LJ, Ley SV. Tetrahedron. 2009; 65:6611–6625.
6. Jiang F, Zhao Y, Hu J. Organic Chemistry Frontiers. 2014; 1:625–629.
7. Kitamura T, Kuriki S, Morshed MH, Hori Y. Org Lett. 2011; 13:2392–2394. [PubMed: 21473585]
8. Suzuki S, Kamo T, Fukushi K, Hiramatsu T, Tokunaga E, Dohi T, Kita Y, Shibata N. Chem Sci. 2014; 5:2754–2760.
9. Pasceri R, Bartrum HE, Hayes CJ, Moody CJ. Chem Commun. 2012; 48:12077–12079.
10. Leng DJ, Black CM, Pattison G. Org Biomol Chem. 2016; 14:1531–1535. [PubMed: 26695741]
11. a) Akana JA, Bhattacharyya KX, Mueller P, Sadighi JP. J Am Chem Soc. 2007; 129:7736–7737. [PubMed: 17547409] b) Gorske BC, Mbofana CT, Miller SJ. Org Lett. 2009; 11:4318–4321.

- [PubMed: 19711904] c) He G, Qiu S, Huang H, Zhu G, Zhang D, Zhang R, Zhu H. *Org Lett.* 2016; 18:1856–1859. [PubMed: 27045165] d) Nahra F, Patrick SR, Bello D, Brill M, Obled A, Cordes DB, Slawin AMZ, O'Hagan D, Nolan SP. *ChemCatChem.* 2015; 7:240–244. [PubMed: 26236406] e) Okoromoba OE, Han J, Hammond GB, Xu B. *J Am Chem Soc.* 2014; 136:14381–14384. [PubMed: 25260170]
12. Zeng X, Liu S, Shi Z, Liu G, Xu B. *Angew Chem Int Ed.* 2016; 55:10032–10036.
13. a) Wang Y, Muratore ME, Echavarren AM. *Chem Eur J.* 2015; 21:7332–7339. [PubMed: 25786384] b) Xi Y, Su Y, Yu Z, Dong B, McClain EJ, Lan Y, Shi X. *Angew Chem Int Ed.* 2014; 53:9817–9821. c) Liu Y, Yu Z, Zhang JZ, Liu L, Xia F, Zhang J. *Chem Sci.* 2016; 7:1988–1995. d) Liu L, Zhang J. *Chem Soc Rev.* 2016; 45:506–516. [PubMed: 26658761] e) Nunes dos Santos Comprido L, Klein JEMN, Knizia G, Kästner J, Hashmi ASK. *Angew Chem Int Ed.* 2015; 54:10336–10340. f) Hashmi, AS., Toste, FD. *Modern gold catalyzed synthesis.* Wiley-VCH; Weinheim, Germany: 2012. g) Toste, FD., Michelet, V. *Gold catalysis: an homogeneous approach.* Imperial College Press; London: 2014.
14. a) Luo Y, Ji K, Li Y, Zhang L. *J Am Chem Soc.* 2012; 134:17412–17415. [PubMed: 23039251] b) He W, Li C, Zhang L. *J Am Chem Soc.* 2011; 133:8482–8485. [PubMed: 21563762] c) Ye L, He W, Zhang L. *J Am Chem Soc.* 2010; 132:8550–8551. [PubMed: 20521793] d) Ye L, Cui L, Zhang G, Zhang L. *J Am Chem Soc.* 2010; 132:3258–3259. [PubMed: 20166668] e) Lu B, Li C, Zhang L. *J Am Chem Soc.* 2010; 132:14070–14072. [PubMed: 20853846] f) He W, Xie L, Xu Y, Xiang J, Zhang L. *Org Biomol Chem.* 2012; 10:3168–3171. [PubMed: 22419033] g) Wang Y, Ji K, Lan S, Zhang L. *Angew Chem Int Ed.* 2012; 51:1915–1918. h) Chen H, Zhang L. *Angew Chem Int Ed.* 2015; 54:11775–11779. i) Wang Y, Zheng Z, Zhang L. *J Am Chem Soc.* 2015; 137:5316–5319. [PubMed: 25835372] j) Kawade RK, Liu R-S. *Org Lett.* 2013; 15:4094–4097. [PubMed: 23905777] k) Hung H-H, Liao Y-C, Liu R-S. *J Org Chem.* 2013; 78:7970–7976. [PubMed: 23875583] l) Ghorpade S, Su M-D, Liu R-S. *Angew Chem Int Ed.* 2013; 52:4229–4234. m) Talbot EPA, Richardson M, McKenna JM, Toste FD. *Adv Synth Catal.* 2014; 356:687–691. [PubMed: 24839436] n) Wang L, Xie X, Liu Y. *Angew Chem Int Ed.* 2013; 52:13302–13306. o) Qian D, Hu H, Liu F, Tang B, Ye W, Wang Y, Zhang J. *Angew Chem Int Ed.* 2014; 53:13751–13755. p) Zhao J, Liu J, Xie X, Li S, Liu Y. *Org Lett.* 2015; 17:5926–5929. [PubMed: 26596136] q) Xu M, Ren T-T, Li C-Y. *Org Lett.* 2012; 14:4902–4905. [PubMed: 22954390] r) Xie L, Liang Z, Yan D, He W, Xiang J. *Synlett.* 2013; 24:1809–1812.
15. Lu Z, Han J, Hammond GB, Xu B. *Org Lett.* 2015; 17:4534–4537. [PubMed: 26335841]
16. Mézailles N, Ricard L, Gagosz F. *Org Lett.* 2005; 7:4133–4136. [PubMed: 16146370]
17. Wang W, Hammond GB, Xu B. *J Am Chem Soc.* 2012; 134:5697–5705. [PubMed: 22376128]
18. Xu Z, Chen H, Wang Z, Ying A, Zhang L. *J Am Chem Soc.* 2016; 138:5515–5518. [PubMed: 27082456]
19. Surmont R, Verniest G, Kimpe N De. *Org Lett.* 2010; 12:4648–4651. [PubMed: 20843014]
20. Liu W-B, Reeves CM, Stoltz BM. *J Am Chem Soc.* 2013; 135:17298–17301. [PubMed: 24160327]
21. Xue H, Jiang D, Jiang H, Kee CW, Hirao H, Nishimura T, Wong MW, Tan C-H. *J Org Chem.* 2015; 80:5745–5752. [PubMed: 25974273]
22. Hussain I, Yawer MA, Lau M, Pundt T, Fischer C, Görls H, Langer P. *Eur J Org Chem.* 2008; 2008:503–518.
23. Han X, Kwiatkowski J, Xue F, Huang K-W, Lu Y. *Angew Chem Int Ed.* 2009; 48:7604–7607.
24. a) Kwiatkowski J, Lu Y. *Eur J Org Chem.* 2015; 2015:320–324. b) Han X, Luo J, Liu C, Lu Y. *Chem Commun.* 2009:2044–2046.
25. Lan Y, Chen Y, Cao X, Zhang J, Wang J, Xu X, Qiu Y, Zhang T, Liu X, Liu B-F, Zhang G. *J Med Chem.* 2014; 57:10404–10423. [PubMed: 25420090]


literature examples

[F⁺]:


- a) Me-NFSI/Ti(OⁱPr)₄ F-N(SO₂Me)₂ (Me-NFSI)
toluene, reflux, 12 h
- b) Selectfluor / ionic liquid, 80 °C
- c) Selectfluor / ZnCl₂, THF, rt
- d) Ar-I/HF-pyridine/*m*-CPBA
- e) aq HF, PhIO, DCM, 40 °C




this work



Scheme 1.
Synthesis of fluorinated 1,3-dicarbonyl compounds.




Scheme 2.
Proposed mechanism for the formation of **3** and **5**.



Scheme 3.
Divergent syntheses from **3** or **5** from literature reports.

Table 1

Optimization of fluorination of alkynyl ketone.^[a]

Entry	Catalyst	N-oxide 2	Solvent	Yield ^[b] (%)
1	JohnPhosAuNTf ₂	2a	DCM	58
2	JohnPhosAuNTf ₂	2b	DCM	73
3	JohnPhosAuNTf ₂	2c	DCM	60
4	JohnPhosAuNTf ₂	2d	DCM	80
5	JohnPhosAuNTf ₂	2d	PhF	86
6	JohnPhosAuNTf ₂	2d	PhCF ₃	93
7	XPhosAuCl+AgNTf ₂	2d	PhCF ₃	91
8	Dppf(AuCl) ₂ +AgNTf ₂	2d	PhCF ₃	89
9	IPrAuCl+AgNTf ₂	2d	PhCF ₃	84
10	PPh ₃ AuCl+AgNTf ₂	2d	PhCF ₃	90
11	AgNTf ₂	2d	PhCF ₃	0
12 ^[c]	JohnPhosAuNTf ₂	2d	PhCF ₃	80
13 ^[d]	JohnPhosAuNTf ₂	2d	PhCF ₃	85
14 ^[e]	JohnPhosAuNTf ₂	2d	PhCF ₃	89

^[a] Conditions: **1a** (0.2 mmol), N-Oxide **2** (0.28 mmol), pyridine-HF (0.8 mmol HF, 4 equiv), catalyst (5 mol%), activator (5 mol%), solvent (0.4 mL), 40 °C in a sealed polypropylene tube for 8 h.


^[b] Determined by GC-MS analysis.

^[c] Pyridine-HF (2 equiv) was used.

^[d] Pyridine-HF (2.5 equiv) was used.


^[e] pyridine-HF JohnPhosAuNTf₂ (1 mol%) was used.

Table 2

Scope for the synthesis of 2-fluoro-1,3-diketone **3**.^[a]

^[a]Reaction conditions: alkyne **1** (0.2 mmol), N-Oxide **2a** (0.28 mmol), pyridine-HF (0.8 mmol HF), JohnPhosAuNTf₂ (5 mol %), PhCF₃ (0.4 mL), 40 °C in a sealed polypropylene tube for 4 h or overnight. All yields are isolated yields; **3** exists as a mixture of ketone form and enol form.

Table 3

Scope for the synthesis of 2-fluoro-3-keto-ester 5.^[a]

^[a] Reaction conditions: alkyne 4 (0.2 mmol), N-Oxide 2a (0.28 mmol), pyridine-HF (0.8 mmol HF), JohnPhosAuNTf₂ (5 mol %), PhCF₃ (0.4 mL), 40 °C in a sealed polypropylene tube for 4 h or overnight; rt. All yields are isolated yields.