

The fauna and perspective of rodentia ectoparasites in Iran relying on their roles within public health and veterinary characteristics

Mousa khosravani¹

Received: 13 May 2017 / Accepted: 22 September 2017 / Published online: 3 October 2017
© Indian Society for Parasitology 2017

Abstract Historically, rodents-ectoparasites are responsible for many diseases in human and domestic animal communities because they are well-adapted to the environment and nutritional resources. In addition, ectoparasites can affect the reproductive rates, ecological fitness and the dispersal pattern of their hosts. Data was compiled through all publications relating to ectoparasites fauna from Iranian rodents. Host specificity, rodent-parasites interaction, geographical distribution, ecological trends as well as the medical or veterinary importance of ectoparasites are considered in this review. Also, results compared with other equal global findings. According to the obtained information, the genus *Hyalomma* considered being the major vector of Crimean-Congo hemorrhagic fever. Sistan-Balouchistan, Isfahan, Fars and Khuzestan are endemic provinces of this disease in Iran. Ixodes ticks (especially *Hyalomma* spp.) commonly existed throughout the months but had high activity in spring and late autumn seasons. Muridae as the most diversified rodent family was more reiterated exploited host by all parasites in this knowledge. *Tatera indica* confirmed as a positive host for *Francisella tularensis* by serological test but the definitive reservoir is still unclear in Iran. *Meriones persicus* as a nocturnal jird has been permanently presented in semi-desert areas except Dasht-e Kavir desert. This rodent was the frequent host for Pulicid fleas followed by *Mus musculus* and *Cricetulus migratorius* (Cricetidae). *Rattus norvegicus* was specialized host for *Polyplax spinulosa* (spiny rat louse). *Pulex*

irritans from Pulicidae was spread in particularly west of Iran which known as an endemic focus of plague. Mites were high diversity than other taxa. The domestic vertebrates (sheep, goat and cattle) were primitive hosts and rodents were discovered as auxiliary hosts for ticks in Iran apparently. The available data suggested that lice had a type of model of host specificity due to their morphological traits. Meanwhile, Alborz and Zagros mountains ranges provided the specialized habitats for animals which have been surrounded by northern and western regions. Perhaps these mountainous areas can prevent gene flow between local mammalian populations. Notably, the genealogy of parasite genome can orient toward the evolutionary process into speciation point. In general, more researches are needed to decipher the whole sights of host-parasite association and the role of ectoparasites within zoonosis diseases.

Keywords Diversity · Ectoparasites · Host · Iran · Rodents

Introduction

Rodentia species are the important pest for many agricultural products as well as oil seeds within pre and post-harvest stages. Also, they are widely distributed and adapted to different ecosystems and climates due to their reproductive abilities (Parshad 1999). Commensal rodents can damage the stored foods, digging burrow under buildings and gnawing materials because of physical and behavioral features (Brooks and Jackson 1973; Brown et al. 2008). Indeed, they are important as reservoir, carrier or incidental hosts in veterinary and medical scopes also contribute to transmission of zoonosis diseases including: Bubonic Plague, Tularemia, Bartonellosis, Leptospirosis,

✉ Mousa khosravani
mousa.khosravani2000@yahoo.com

¹ Department of Medical Entomology and Vector Control, School of Health, Shiraz University of Medical Sciences, P.O. Box 71645-111, Shiraz, Fars Province, Iran

Tick-borne relapsing fever, Murine typhus, Trichinelliasis, Toxoplasmosis, Echinostomiasis, Schistosomiasis, Leishmaniasis and Hantavirus Pulmonary Syndrome (Meerburg et al. 2009). Rodents can transmit pathogens to human directly or indirectly. For example, Omsk hemorrhagic fever virus (OHFV) transmits through direct contact of humans with infected animals or ectoparasite bite (tick) (Li et al. 2004). Numerous of the rodent-borne diseases have occurred in Southeastern Asia where rodents' population were increased in accordance with rice cultivation. (WHO 1988). In addition, gastrointestinal termatodes such as *Plagiorchis muris* and *Neodiplostomum seoulense* as endoparasites have been isolated from human and rodents communities which have a significant role in human health (Albon et al. 2002; Wells et al. 2007; Lee et al. 2013). Host specificity is one of the most important parameters which describes the ecological and evolutionary patterns of both host and parasite (Rosario Robles et al. 2016). Herein fleas (*Xenopsylla conformis*) at a given stage can affect the behavioral defence of their rodents (Hawlena et al. 2007). The genus *Ornithodoros* has been presented in the colony of Palearctic rodents (e.g. *Meriones* spp., *Microtus* spp. and *Tatera* spp.) (Naddaf et al. 2012). This tick divides the two species; *Ornithodoros tholozani* and *O. erraticus*. *Borrelia persica* (Spirochaetal agent of Tick-borne relapsing fever) is associated with *O. tholozani*. In contrast, *Borrelia microti* parasitizes *O. erraticus* in southern Persia (Aghighi et al. 2007; Naddaf et al. 2015). These soft ticks were mostly found from rodents burrows located in northwestern and central Iran specially Ardabil district (Masoumi-Asl et al. 2009). Unfortunately, the plague caused a pandemics disaster in the twentieth century and millions of people died throughout the world by this disease. The oriental rat flea (*Xenopsylla cheopis*) was identified as the main vector for bubonic plague (Adjemian et al. 2007; Gage and Kosoy 2005). Afterwards, the plague caused a horrible decline in Iranian population during the nineteenth century. More than two-thirds of residents from Kermanshah province died due to an outbreak of disease during the 1800–1906 period (Seyf 1989). Besides, *Yersinia pestis* (the microbial organism of plague) circulates between the wild mammal's population such as *Meriones* spp. rodents in Kurdistan province as the main former endemic foci of plague disease, west of Iran (Hashemi-Shahraki et al. 2016). It should be noted that Leishmaniasis is prevalent with high incidence rates in different parts of Iran (Khosravani et al. 2016). Gerbillinae Subfamily has been proven as the reservoir host for Zoonotic cutaneous leishmaniasis (ZCL) (Rassi et al. 2006). Strikingly, Dr. Etemad was the eminent Iranian scientist that published the principal's information of the Rodentia order in 1977. Sciuridae, Hystricidae, Dipodidae, Muscardinidae, Muridae and Cricetidae were discovered as the families of rodents from Iran (Sedaghat

and Salahi-Moghaddam 2010a, b). The objective of present study is a complete reappraisal of the ectoparasites associated with rodents that have been recorded from Iran. This paper is the first novel in Iran that addresses all aspects of rodents-ectoparasites based on their importance to public health and veterinary scopes. Definitive species of host, local distribution, host-parasites interaction, host specification, biodiversity, the pattern of seasonal activity and ecological trends were accommodated in this review. Also, these results compared with other global studies comprehensively.

Materials and methods

Study site

Iran (32°00'N; 53°00'E) is a country in western Asia with an area of 1,648,195 km² (80,000,000 population). Azerbaijan, Armenia, Turkmenistan, Pakistan, Afghanistan, Turkey and Iraq have a common border with Iran (a common name: Persia). Zagros and Alborz mountains ranges are high altitude areas. Southern shores of Caspian is covered by forest and trees. The Dasht-e Kavir and the Dasht-e Lut are the Iran's largest deserts also two-third of this country have arid or semiarid climates. The average annual rainfall are 246 mm approximately thereby, precipitation has a significant role in the life of plants and animals in Iran.

Research tool

This descriptive study was given on earlier investigations and references that have been conducted on different rodents and their ectoparasites fauna in Iran. Parasites classified in four groups are as follows: fleas, mites, ticks and lice. All taxonomic information about ectoparasites was ranked based on recent definitions. Sometimes the scientific classification and nomenclature of parasites have overlapping meanings but considerably related (Small 1989). In this approach, local distribution of both rodent species and their ectoparasites were tabulated. Many parameters of host-parasite re-evaluated and reviewed. In addition, some characteristics of ectoparasites compared with previous researches across rodent species as follows: geographic ranges of host-parasite "host switching in parasites, the community dynamics of parasite related to its host, ecological changes, host behaviour, parasite species richness" host defence, competition in parasite, host-parasite interaction, transmission of pathogen agents to human and the medical/veterinary importance of rodents ectoparasites. All data gathered and arranged from well-known scientific databases and search engines such as

ScienceDirect, Scopus, ProQuest, Sage, Wiley Online Library, Nature, Springer, Jstor, Bio One, PubMed, Google scholar, ResearchGate, Magiran and Sid.

Results

In this review 58 arthropod specimens within four taxa (flea, mite, tick and lice) were found on rodents species from 22 regions as follows:

Order: Siphonaptera Latreille, 1825

Pulicidae

Xenopsylla buxtoni Jordan, 1949, *Xenopsylla gerbilli* Wngner, 1903, *Xenopsylla nubica* Rothschild, 1903, *Xenopsylla astia* Rothschild, 1911, *Xenopsylla cheopis* Rothschild, 1903, *Xenopsylla conformis* Rothschild, 1904, *Xenopsylla nuttalli* Loff, 1930, *Ctenocephalides felis* Bouche, 1835, *Pulex irritans* Linn, 1758

Ceratophyllidae

Nosopsyllus fasciatus Bosc, 1800, *Nosopsyllus iranensis* Wagner et Argyropulo, 1934, *Nosopsyllus medus* Jordan, 1938

Leptopsyllidae

Leptopsylla taschenbergi taschenbergi Wagner, 1898, *Leptopsylla aethiopica* Rothschild, 1908

Hystrihopsyllidae

Stenoponia tripectinata Tiraboschi, 1902, *Ctenophthalmus rettigi* Rothschild, 1908

Subclass: Acari Leach, 1817

Order: Mesostigmata

Hirstionyssidae

Hirstionyssus Fonseca, 1948

Laelapidae

Laelaps nuttalli Hirst, 1915, Hurlbut 1949, *Laelaps acuminata*, *Laelaps algericus* Hirst, 1925, *Echinolaelaps echidninus* Berlese, 1887, Hurlbut 1949, *Androlaelaps hermaphrodita* Berlese, 1887, *Haemolaelaps glasgowi* Ewing

Dermanyssidae

Dermanyssus gallina Geer, 1778, *Dermanyssus sanguineus* Hirst, 1914, *Dermanyssus americanus* Ewing, 1922, *Liponyssoides Sanguineus* Hirst

Macronyssidae

Ornithonyssus bacoti Hirst, 1931, Pippin & Shimada, 1966, *Ornithonyssus sylviarum* Canestrini and Fanzago, 1877

Pachylaelapidae

Pachylaelaps spp. Berlese, 1888

Order: Trombidiformes

Suborder: Prostigmata

Cheyletidae Leach, 1815

Cheyletus malaccensis

Myobiidae

Myobia musculi Schrank, 1781, *Radfordia affinis* Poppe, 1896

Trombiculidae Ewing, 1929

Microtrombicula spp.

Order: Sarcoptiformes

Acaridae

Acarus siro Linnaeus, 1758, *Tyrophagus putrescentiae* Schrank 1781, *Caloglyphus* Berlese, 1924

Order: Astigmata

Myocoptidae Gunther, 1942

Myocoptes musculus Koch, 1844, *Trichoecius romboutsii* Van Eynhoven, 1946

Glycyphagidae

Lepidoglyphus destructor Schrank, 1781

Order: Opilioacarida Johnston, 1968 (= Notostigmata With, 1903–1904)

Opiliacaridae With, 1902

Order: Ixodida

Ixodidae Koch, 1844

Haemaphysalis Koch, 1844, *Rhipicephalus* sp., *Hyalomma* spp., *Boophilus* spp., *Haemaphysalis punctata* Canestrini & Fanzago, 1878, *Ixodes trianguliceps* Birula, 1895, *Ixodes ricinus* Linnaeus, 1758

Subclass: Pterygota Lang, 1888

Superorder: Exopterygota

Order: Phthiraptera Haeckel, 1896

Hoplopleuridae

Neohaematopinu spp. Mjöberg, 1910, Ferris, 1942, *Hoplopleura oenomydis* Ferris 1921, *Hoplopleura captiosa* Johnson, 1960

Polyplacidae

Polyplax spinulosa Burmeister, 1839, *Polyplax stephensi* Christophers and Newstead, 1906, *Polyplax asiatica* Ferris, 1923, *Polyplax paradoxa* Johnson, 1960, *Polyplax gerbilli* Ferris, 1923, *Polyplax serrata* Burmeister, 1839, *Eulinognathus* spp.

Discussion

Fleas

All fleas insects have consisted of 16 species belong to 4 families, Pulicidae, Ceratophyllidae, Leptopsyllidae and Hystrihopsyllidae in Iran as Pulicidae contains over 57% of all species alone (Table 1). Fleas are obligatory hematophagous (male & female) with 2550 species/subspecies from 18 families that struggle in the Holarctic, Nearctic, Palearctic, Afrotropical and Oriental regions. They have

Table 1 Ectoparasites fauna of Iranian rodents in the various districts

Ectoparasite	Host	Region	Researcher
Fleas			
<i>Xenopsylla</i> spp.	<i>Rattus norvegicus</i> , <i>Nesokia indica</i>	Ahvaz, Hovizeh, Mashhad	Rahdar et al. (2015) and Moravvej et al. (2015)
<i>Xenopsylla buxtoni</i>	<i>Meriones persicus</i> , <i>Microtus socialis</i> , <i>Calomyscus bailwardi</i> , <i>Cricetulus migratorius</i> , <i>Rattus rattus</i> , <i>R. norvegicus</i> , <i>Tatera indica</i>	Khorram abad, Meshkinshahr, Bandar Abbas	Shayan and Rafinejad (2006), Zarei et al. (2015) and Hanafi-Bojd et al. (2007)
<i>Xenopsylla gerbilli</i>	<i>M. persicus</i>	Meshkinshahr	Mohebbali et al. (1997)
<i>Xenopsylla nubica</i>	<i>M. persicus</i> , <i>Mus musculus</i> , <i>C. migratorius</i> , Muridae and Cricetidae families	Meshkinshahr, Iran Shahr, Nikshahr	Zarei et al. (2015) and Nateghpour et al. (2013)
<i>Xenopsylla astia</i>	<i>M. persicus</i> , <i>M. musculus</i> , <i>C. migratorius</i>	Meshkinshahr, Iran Shahr, Nikshahr	Zarei et al. (2015) and Nateghpour et al. (2013)
<i>Xenopsylla cheopis</i>	<i>M. persicus</i> , <i>M. musculus</i> , <i>C. migratorius</i> , <i>N. indica</i>	Meshkinshahr, Mashhad	Zarei et al. (2015) and Moravvej et al. (2015)
<i>Xenopsylla conformis</i>	Muridae and Cricetidae families	Iranshahr, Nikshahr	Nateghpour et al. (2013)
<i>Xenopsylla nuttali</i>	<i>Rhombomys opimus</i>	North of Khorasan province, Kalaleh, Damghan, Shahrood, Badrood, Habibabad, Meshkinshahr	Tajedin et al. (2009) and Mohebbali et al. (1997)
<i>Pulex irritans</i>	Muridae and Cricetidae families	Sarpole-Zahab	Telmadarraiy et al. (2007)
<i>Ctenocephalides felis</i>	<i>M. musculus</i> , <i>M. persicus</i> , <i>C. migratorius</i>	Meshkinshahr	Zarei et al. (2015)
<i>Nosopsyllus</i> spp.	<i>N. indica</i> , <i>M. persicus</i>	Mashhad	Moravvej et al. (2015)
<i>Nosopsyllus fasciatus</i>	<i>M. persicus</i> , <i>M. musculus</i> , <i>C. migratorius</i> , <i>M. socialis</i> , <i>Spermophilus fulvus</i> , <i>N. indica</i> , <i>R. norvegicus</i>	Meshkinshahr, Khorram abad, Mashhad, Tehran, Turkmen Sahra	Zarei et al. (2015), Shayan and Rafinejad (2006), Moravvej et al. (2016), Moravvej et al. (2015), Pakdad et al. (2012), Mohebbali et al. (1997) and Gholipoury et al. (2016)
<i>Nosopsyllus medus</i>	Muridae and Cricetidae families	Sarpole-Zahab	Telmadarraiy et al. (2007)
<i>Nosopsyllus iranus</i>	<i>M. musculus</i> , <i>M. persicus</i> , <i>C. migratorius</i>	Meshkinshahr	Zarei et al. (2015)
<i>Leptopsylla aethiopica</i>	<i>M. musculus</i>	Semnan	Darvishi et al. (2014)
<i>Leptopsylla taschenbergi taschenbergi</i>	<i>Apodemus sylvaticus</i>	Razan plain	Yousefi et al. (2015)
<i>Ctenophthalmus rettigi</i>	<i>M. musculus</i> , <i>R. norvegicus</i> , <i>M. persicus</i> , <i>C. migratorius</i>	Mashhad, Meshkinshahr	Moravvej et al. (2016), Moravvej et al. (2015) and Zarei et al. (2015)
<i>Stenoponia tripectinata</i>	<i>M. persicus</i>	Meshkinshahr	Mohebbali et al. (1997)
Mites			
<i>Laelaps</i> spp.	<i>M. musculus</i>	Mashhad	Moravvej et al. (2015)
<i>Laelaps nuttalli</i>	Muridae and Cricetidae families, <i>R. norvegicus</i> , <i>T. indica</i> , <i>M. musculus</i>	Sarpole-Zahab, Bandar Abbas, Turkmen Sahra	Telmadarraiy et al. (2007), Hanafi-Bojd et al. (2007) and Gholipoury et al. (2016)
<i>Laelaps ciccuminata</i>	Muridae (Gerbillinae)	Iranshahr, Nikshahr	Nateghpour et al. (2013)
<i>Laelaps algericus</i>	<i>M. musculus</i>	Mashhad	Moravvej et al. (2016)
<i>Haemolaelaps</i> spp.	<i>M. musculus</i> , <i>M. persicus</i>	Mashhad	Moravvej et al. (2016) and Moravvej et al. (2015)

Table 1 continued

Ectoparasite	Host	Region	Researcher
<i>Haemolaelaps glasgowi</i>	<i>M. persicus</i> , <i>M. musculus</i> , <i>M. socialis</i> , <i>C. migratorius</i> , <i>Ellobius fuscocapillus</i> , <i>R. rattus</i> , <i>R. norvegicus</i> , <i>T. indica</i>	Khorram abad, Bandar Abbas	Shayan and Rafinejad (2006) and Hanafi-Bojd et al. (2007)
<i>Echinolaelaps echidninus</i>	<i>R. norvegicus</i> , <i>T. indica</i> , <i>R. rattus</i>	Bandar Abbas, Sari	Hanafi-Bojd et al. (2007) and Motevalli-Haghi et al. (2000)
<i>Androlaelaps hermaphrodita</i>	Muridae (Gerbillinae)	Iranshahr, Nikshahr	Nateghpour et al. (2013)
<i>Ornithonyssus</i> spp.	<i>M. musculus</i> , <i>M. persicus</i> , <i>C. migratorius</i> , <i>T. indica</i> , <i>R. norvegicus</i>	Meshkinshahr, Ahvaz, Hovizeh	Zarei et al. (2015) and Rahdar et al. (2015)
<i>Ornithonyssus sylviarum</i>	<i>M. persicus</i> , <i>C. bailwardi</i> , <i>C. migratorius</i>	Khorram abad	Shayan and Rafinejad (2006)
<i>Ornithonyssus bacoti</i>	Muridae and Cricetidae families, <i>M. musculus</i> , <i>T. indica</i> , <i>R. opimus</i> , <i>R. norvegicus</i> , <i>M. persicus</i>	Sarpole-Zahab, Northwest of Iran, Bandar Abbas, North of Khorasan, Kalaleh, Damghan, Shahrood, Badrood, Habibabad, Meshkinshahr, Tehran	Telmadarraiy et al. (2007), Allymehri et al. (2012), Hanafi-Bojd et al. (2007), Tajedin et al. (2009), Pakdad et al. (2012) and Mohebbali et al. (1997)
<i>Dermanyssus gallinae</i>	<i>M. musculus</i>	Northwest of Iran	Allymehri et al. (2012)
<i>Dermanyssus americanus</i>	<i>R. norvegicus</i> , <i>T. indica</i>	Bandar Abbas	Hanafi-Bojd et al. (2007)
<i>Dermanyssus sanguineus</i>	Muridae and Cricetidae families, <i>R. norvegicus</i>	Sarpole-Zahab, Bandar Abbas	Telmadarraiy et al. (2007) and Hanafi-Bojd et al. (2007)
<i>Liponyssoides sanguineus</i>	<i>R. norvegicus</i>	Ahvaz, Hovizeh	Rahdar et al. (2015)
<i>Acarus siro</i>	<i>M. musculus</i> , <i>R. rattus</i> , <i>R. norvegicus</i>	Tehran	Shiravi et al. (2013)
<i>Caloglyphus</i>	<i>M. musculus</i> , <i>R. rattus</i> , <i>R. norvegicus</i>	Tehran	Shiravi et al. (2013)
<i>Tyrophagus putrescentiae</i>	<i>M. musculus</i> , <i>R. rattus</i> , <i>R. norvegicus</i>	Tehran	Shiravi et al. (2013)
<i>Hirstionyssus</i> spp.	<i>Meriones libycus</i>	Mashhad	Moravvej et al. (2015)
<i>Trichoecius romboutsii</i>	<i>M. persicus</i>	Meshkinshahr	Mohebbali et al. (1997)
<i>Myocoptes musculus</i>	<i>M. musculus</i>	Northwest of Iran	Allymehri et al. (2012)
<i>Cheyletus malaccensis</i>	<i>M. musculus</i> , <i>R. rattus</i> , <i>R. norvegicus</i>	Tehran	Shiravi et al. (2013)
<i>Myobia musculi</i>	<i>M. musculus</i> , <i>R. rattus</i> , <i>R. norvegicus</i>	Tehran	Shiravi et al. (2013)
<i>Radfordia affinis</i>	<i>M. musculus</i> , <i>R. rattus</i> , <i>R. norvegicus</i>	Tehran	Shiravi et al. (2013)
<i>Lepidoglyphus destructor</i>	<i>M. musculus</i> , <i>R. rattus</i> , <i>R. norvegicus</i>	Tehran	Shiravi et al. (2013)
<i>Pachylaelaps</i> spp.	Muridae (Gerbillinae)	Iranshahr, Nikshahr	Nateghpour et al. (2013)
<i>Microtrombicula</i> spp.	<i>M. musculus</i> ,	Mashhad	Moravvej et al. (2016)
Opiliacaridae	<i>R. rattus</i> , <i>R. norvegicus</i>	Sari	Motevalli-Haghi et al. (2000)
Ticks			
<i>Haemaphysalis</i> spp.	<i>M. persicus</i> , <i>M. musculus</i> , <i>M. socialis</i> , <i>C. bailwardi</i> , <i>R. rattus</i> , <i>T. indica</i> , <i>Apodemus witherbeyi</i>	Khorram abad, Ahvaz, Hovizeh, Meshkinshahr Mashhad	Shayan and Rafinejad (2006), Rahdar et al. (2015), Mohebbali et al. (1997), Moravvej et al. (2016) and Moravvej et al. (2015)

Table 1 continued

Ectoparasite	Host	Region	Researcher
<i>Haemaphysalis punctata</i>	<i>M. musculus</i> , <i>M. persicus</i> , <i>N. indica</i>	Mashhad	Moravvej et al. (2016) and Moravvej et al. (2015)
<i>Rhipicephalus</i> spp.	Muridae and Cricetidae families, Muridae (Gerbillinae), <i>R. rattus</i> , <i>R. norvegicus</i> , <i>T. indica</i>	Sarpole-Zahab, Iranshahr, Nikshahr, Bandar Abbas, Sari, Turkmen Sahra	Telmadarraiy et al. (2007), Nateghpour et al. (2013), Hanafi-Bojd et al. (2007), Motevalli-Haghi et al. (2000) and Gholipoury et al. (2016)
<i>Hyalomma</i> spp.	Muridae and Cricetidae families, Muridae (Gerbillinae)	Sarpole-Zahab, Iranshahr, Nikshahr	Telmadarraiy et al. (2007) and Nateghpour et al. (2013)
<i>Boophilus</i> spp.	Muridae (Gerbillinae)	Iranshahr, Nikshahr,	Nateghpour et al. (2013)
<i>Ixodes trianguliceps</i>	<i>M. persicus</i>	Mashhad	Moravvej et al. (2015)
<i>Ixodes</i> spp.	<i>A. witherbeyi</i>	Mashhad	Moravvej et al. (2015)
<i>Ixodes ricinus</i>	<i>R. norvegicus</i>	Tehran	Pakdad et al. (2012)
Lice			
<i>Polyplax</i> spp.	Persian Squirrel (<i>Scuirus anomalus</i>), <i>M. persicus</i> , <i>M. musculus</i> , <i>C. migratorius</i>	Urmia, Meshkinshahr	Shirazi et al. (2013) and Zarei et al. (2015)
<i>Polyplax spinulosa</i>	Muridae and Cricetidae families, Muridae (Gerbillinae), <i>R. norvegicus</i>	Sarpole-Zahab, Tehran, Ahvaz, Hovizeh	Telmadarraiy et al. (2007), Nateghpour et al. (2013), Pakdad et al. (2012) and Rahdar et al. (2015)
<i>Polyplax stephensi</i>	<i>T. indica</i>	Ahvaz, Hovizeh	Rahdar et al. (2015)
<i>Polyplax asiatica</i>	<i>M. persicus</i> , <i>S. fulvus</i> , <i>T. indica</i>	Mashhad	Moravvej et al. (2015)
<i>Polyplax serrata</i>	<i>M. musculus</i>	North west of Iran	Allymehr et al. (2012)
<i>Polyplax paradoxa</i>	<i>M. persicus</i>	Mashhad	Moravvej et al. (2015)
<i>Polyplax gerbilli</i>	<i>R. norvegicus</i> , <i>T. indica</i>	Bandar Abbas	Hanafi-Bojd et al. (2007)
<i>Eulinognathus</i> spp.	<i>Allactaga</i> spp.	Ahvaz, Hovizeh	Rahdar et al. (2015)
<i>Neohaematopinus</i> spp.	<i>M. persicus</i> , <i>M. socialis</i>	Khorram abad	Shayan and Rafinejad (2006)
<i>Hoplopleura</i> spp.	<i>R. norvegicus</i> , <i>R. rattus</i>	Sari, Tehran	Pakdad et al. (2012) and Motevalli-Haghi et al. (2000)
<i>Hoplopleura captiosa</i>	<i>N. indica</i> , <i>M. musculus</i> , <i>R. norvegicus</i>	Mashhad, Bandar Abbas	Moravvej et al. (2015) and Hanafi-Bojd et al. (2007)
<i>Hoplopleura oenomydis</i>	<i>R. norvegicus</i>	Tehran	Pakdad et al. (2012)

adapted to ecological features of hosts due to phylogenetic events into their shape and behaviour (Medvedev 1996; Bitam et al. 2010). Also, they implicate as vector or reservoir of Bartonella which circulates between fleas/rodents communities via horizontal and vertical routes. For instance, *Bartonella rochalimae* and *Bartonella tribocorum* DNA has been identified in *Xenopsylla cheopis* fleas (Billeter et al. 2011). As described above Pulicidae was the largest family that more comprised genus *Xenopsylla*. *Xenopsylla nubica* have widely hosts such as *Meriones persicus* (Persian jird), *Mus musculus*, *Cricetulus migratorius* and Cricetidae families (uncertain species). *Xenopsylla conformis* have occurred on Cricetidae and Muridae (undetermined species). In a previous study compared with this research *Xenopsylla conformis mycerini* and *X. nubica theodori* were removed from *Gerbillus nanus*, *Acomys*

dimidiatus, *Meriones lybicus* (the Libyan jird) and *Meriones rex* but they were more exist on *M. rex* (Harrison et al. 2015). As such *Xenopsylla cheopis* (the Oriental rat flea) was dominant on Muridae (Persian jird, *M. musculus*, *Nesokia indica*) and a species from Cricetidae (*C. migratorius*) in Iran. Also, this flea was named as a parasite for *Rattus rattus* in Sri Lanka, Asia (Sumangali et al. 2012) and the Canary Islands in Spain, Europe (Feliu et al. 2012) as well as for *Rattus flavipectus* in Yunnan Province, China (Xian-guo et al. 2000). In other words, it is a broad species of genus *Xenopsylla* throughout the world. Sometimes, even more, this parasite considered to be the important vector of plague bacteria (*Yersinia pestis*) in the zoonotic areas (Eisen et al. 2007) As regards co-infection with *Rickettsia typhi* and *Rickettsia felis* have been shown in the Oriental rat flea from *M. musculus* in previous investigation

(Eremeeva et al. 2008). *Meriones persicus*, *M. musculus* and *C. migratorius* are hosts to *Xenopsylla astia*. This specimen was a vector within sylvatic and human forms of plague in India (Renapurkar 1988) besides, *Bandicota bengalensis* was the ideal host for *X. astia* in this country (Shyamal et al. 2011). *Xenopsylla astia*, *X. buxtoni*, *X. stenoponia*, *X. conformis* and *Pulex irritans* was recorded as vectors of plague on genera *Gerbillidae* from Muridae in the endemic spots (Kurdistan, Ardebil) in Iran (Nekouie et al. 2003). In view of *Xenopsylla gerbilli*, Mohebbali and his coworkers have reported the replacement of *X. gerbilli* on *M. persicus* in Meshkinshahr, northwestern Iran (Mohebbali et al. 1997) while this flea has infested great gerbil in plague focus, Central Asia (Samia et al. 2011). Another species of Pulicidae, *P. irritans* (the human flea) resided on Muridae and Cricetidae families of rodents in the west of Iran that was once the important foci of plague. This species as domestic ectoparasite was prevailing flea in the risk districts of plague in Madagascar (Ratovonjato et al. 2014) and Tanzania, Africa (Laudisoit et al. 2007). *Ctenocephalides felis* (Pulicidae) is known as cat flea was found on three species of small mammals e.g. *M. musculus*, *M. persicus* and *C. migratorius*. It was established where CSD (cat scratch disease) cases documented increasingly (Breitschwerdt and Kordick 2000). Additionally, it was confirmed as a vector of *Rickettsia felis* (the etiologic agent of rickettsial fleaborne spotted fever) (Jiang et al. 2006), plague and murine typhus (Rust 1997). It is noteworthy that infected female flea can transmit *Rickettsia typhi* (the causative agent of Murine typhus) by transovarial transmission (Ford et al. 2004). In this research, the Ceratophyllidae family contained 3 species as *Nosopsyllus fasciatus* (the northern rat flea) was extremely common parasitic arthropod on *M. persicus*, *M. musculus*, *C. migratorius*, *Microtus socialis*, *Spermophilus fulvus* (squirrel), *N. indica* (the Indian Bandicoot) and *R. norvegicus*. In Compared with this survey it has been collected from Norway rat in rural areas from dumps during the warm months (0.24 fleas per rat) also mostly from buildings in the cold season with 1.3 (Burbutis and Hansens 1955) and 2.7 fleas per rat in U.S (Eskey 1938). Furthermore, *Nosopsyllus fasciatus* transmits *Trypanosoma lewisi* among rat populations. A filarial nematode (*Dipetalonema reconditum*) can transmit by *C. canis* and *C. felis* (Ford et al. 2004). *Ctenocephalides felis canis*, *X. cheopis* and *P. irritans* have been infected as the accidental hosts by the cysticercoids of *Hymenolepis diminuta* (the rat tapeworm), *Hymenolepis nana* and *Dipylidium caninum* (Oldham 1931; Ford et al. 2004) also transmit to susceptible hosts (especially children) upon ingestion of these intermediated hosts occasionally (Craig and Ito 2007). *Leptopsylla aethiopica* and *L. taschenbergi* from Leptopsyllidae have newly been recorded from Iran (Darvishi

et al. 2014; Yousefi et al. 2015) (Table 1). *Stenoponia tripectinata* (Hystrichopsyllidae) didn't only find from Iran. This flea has been removed from *Meriones crassus*, *Jaculus jaculus* and *Gerbillus cheesman* within two genders (female & male) in Northern Saudi Arabia (Lewis 1964) whereas it was taken on *M. persicus* predominantly in Iran. *Mus musculus*, *R. norvegicus*, *M. persicus* and *C. migratorius* were the determined hosts for *Ctenophthalmus rettigi* from this family. *Ctenophthalmus* species was also described from *Apodemus agrarius* (genus of Muridae) in the Far East (Smit 1955).

Mites

In the case of mites, 25 genus/species from 12 families are known in this study. They are high biodiversity than other groups. Laelapidae, Macronyssidae and Dermanyssidae (mesostigmatid mite) more have been recorded in Iran. According to my knowledge *Laelaps nuttalli* (domestic rat mite) has been found on many of rodents from Cricetidae (unknown species) and *T. indica* (Table 1). It was also common on *R. norvegicus* (Yang et al. 2009), *R. rattus mansorius* and *Rattus exulans* in other places (Wilson 1967). *Laelaps echidninus* has appeared on Norway/black rats in Chile, South America (Lareschi and Gonzalez-Acuna 2010) thereby, it can be concluded that *Laelaps* mites initially associate with *Rattus* spp. as Indian gerbil was secondary hosts for this group. In this family, *Haemolaelaps* spp. (adult and nymphal stages) was taken from birds, nest and rodents but more tended to rodent hosts. *Haemolaelaps glasgowi* has been demonstrated on captured hosts belong to different families (Muridae and Cricetidae) as follows: *M. persicus*, *M. musculus*, *M. socialis*, *C. migratorius*, *Ellobius fuscocapillus*, *R. rattus*, *R. norvegicus* and *T. indica*. As this ectoparasite was occurred on *Microtus arvalis* and *Citellus citellus* (ground squirrel) in Turkey (Cicek et al. 2008) as well as on the Meadow Vole in Wisconsin, north-central United State (Amin 1976). *Echinolaelaps echidninus* (gamasid mite) was a scarce species that happened on genera rat naturally in the spring season (Davis 1951) also on *T. indica* in this literature. This mite is a vector of *Hepatozoon muris*, which involves white rats (Jakeman 1961). Genus *Dermanyssus* from Dermanyssidae consists of 23 species (Roy and Chauve 2007). It was determined as a vector of viral zoonoses diseases such as Saint-Louis encephalitis virus and fowl-pox. Moreover, bacterial agents (*Salmonella gallinarum*, *Erysipelothrix rhusiopathiae*) isolated from *Dermanyssus gallinae* on Flows hosts. Western equine encephalitis virus (WEE) has also been isolated in wild birds roosts from this acarion although no transovarial transmission or viral replication has been observed. In general, the role of these mites into pathogenicity process

is a controversial issue yet (Valiente Moro et al. 2005). *Dermanyssus gallinae* (the red mite of poultry) was collected solely from *M. musculus* in my study. It would infest birds/nests (Huhta 2016; Arjomandi et al. 2013) especially hen in other regions if the condition was suitable for this mite (Rosen et al. 2002). *Dermanyssus sanguineus* can parasitize birds like other dermanyssid mites (Roy and Chauve 2010). Muridae and Cricetidae families, in particular, *R. norvegicus* have been noted as the hosts for this mite based on current data. This hematophagous mite also has been occurred on house sparrows (Brown and Wilson 1975). *Liponyssoides sanguineus* (the mouse mite) from Dermanyssidae has been assigned as the main vector of *Rickettsia akari*. This agent can transmit via bite and causes Rickettsialpox disease in human. It has been recorded from Ukraine, Korea, North Carolina, South Africa (Lahey 2003) and Netherlands (Renvoise et al. 2012). In Persia, this mite first was reported on *R. norvegicus* in Ahvaz city (Rahdar et al. 2015) although the possible cases have not been diagnosed yet. Within the Macronyssidae, *Ornithonyssus bacoti* (the tropical rat mite) was parasitic on *M. musculus*, *T. indica* (Indian gerbil), *Rhombomys opimus*, *R. norvegicus* and *M. persicus* (Muridae). This species prefers Rats to other groups as the principal hosts (Reeves et al. 2007; Bhuyan and Nath 2016). *Bartonella* spp. and *Rickettsia* spp. were detected in *O. bacoti* by molecular technic although the figure of transmission of these pathogens to human or animal is still unclear (Reeves et al. 2007). On the other hand, it infests small mammals' fur also it has medical important due to causing dermatitis in human (Beck and Folster-Holst 2009; Baumstark et al. 2007; Rahdar and Vazirianzadeh 2009). Furthermore, this mite feeds on the man when other hosts are not accessible (Fischer and Walton 2014). Considerably, *Litomosoides carinii* (the filarial nematode) of rodents can parasitize *O. bacoti* as an intermediated host (Fagir and El-Rayah 2009). *Ornithonyssus sylviarum* (The northern fowl mite) was revealed on *M. persicus*, *Calomyscus bailwardi* and *C. migratorius* in the findings. It specimen served as an avian parasite on the diversity of orders of birds (e.g. Falconiformes, Passeriformes and Piciformes) (Knee and Proctor 2007; Mullens et al. 2001; Owen et al. 2009). Interestingly, *Dermanyssus gallinae* (obligatory ectoparasite), *O. bacoti* and *O. sylviarum* had analogous relationship morphologically (Beck and Folster-Holst 2009). In this overview, *Acarus siro* (the flour mite), *Lepidoglyphus destructor*, *Cheyletus malaccensis*, *Tyrophagus putrescentiae*, *Caloglyphus berlesei*, *Myobia musculi* and *Radfordia affinis* were only common on commensal rats particularly *Mus musculus* in wheat stores. The first four species have more abundant in rodents' burrows. Many records have shown allergic reactions through contact with storage mites (the flour mite, *T.*

putrescentiae and *L. destructor*) in patients that were sensitized to *Dermatophagoides pteronyssinus* (the house-dust mite) (Green and Woolcock 1978; Llerena et al. 1991; Heide et al. 1998; Johansson et al. 2001). *Tyrophagus putrescentiae* (the mold mite) was known as the causal ectoparasite of allergic manifestations among farmers (Sanchez-Ramos and Castanera 2001). *Cheyletus malaccensis* (Cheyletidae) as known predator was used into biological control in order to reduce its prey density either *A. siro* (Solomon 1962; Pekar and Hubert 2008) or *T. putrescentiae* (the cheese mite) in gain/seeds products of plants (Kuwahara et al. 1975; Palyvos et al. 2006). House mice were favorable host for *Myobia musculi* and *Radfordia* (Myobiid mites) in South Carolina. (Reeves and Cobb 2005) As they have been typically presented on the murine rodents in Europe (Bochkov and Labrzycka 2003). Two species of Myocoptid mites have been reported from Iran. *Myocoptes musculus* and *Trichoecius romboutsii* were taken from house mouse and *M. persicus* respectively. Dr. Flynn removed these species from mice (Fain and Hyland 1970). *Microtrombicula* spp. was merely common on *M. musculus*. Another chigger mite, *Leptotrombidium scutellare* (a vector of tsutsugamushi disease) has been presented on individual rodents (Wei et al. 2010) chiefly *Eothenomys miletus* (vole) male in China (Peng et al. 2015). Additionally, these groups transmit the rickettsial agents to man only in the larval sitting (Pratt 1963). Several investigations implicate that both chiggers and Laelapinae mites can transmit Hantaan virus to rodents in laboratory and field examinations. This viral pathogen causes HFRSV (hemorrhagic fever with renal syndrome) in human through inhalation of aerosols of small mammals feces although their vector competence is still unfamiliar (Yu and Tesh 2014). Among Opilioacaridae, 3 genera have been described in the new world: *Neocaracus*, *Opilioacarus* and *Caribeacarus* (Vazquez and Klompen 2009). In conducted researches, these species were observed in the regions with warm climates (Vazquez and Klompen 2015). This family was determined on commensal rats in the present publication. *Pachylaelaps* spp. (Pachylaelapidae) was removed from Murid rodents. This genus has been collected from soil and decomposing organic matter in Iran (Ahadiyat et al. 2014). Pachylaelapid mites are free-living soil mites consist of 16 genera with 230 species in throughout the world. They were found in organic substances, soil, decaying wood and plant material as well as nests of mammals and birds (Masan and Halliday 2014; Ozbek 2015; Masan et al. 2016).

Ticks

Ticks as the competent vectors transmit the infectious disease to human, domestic and wild animals including

Rocky spotted fever, Lyme disease, Babesiosis, Theileriosis, Dermatophilosis and Issyk-Kul fever virus (Fuente et al. 2008; Lindquist et al. 2016). Ixodidae transmits rickettsiae to the next generation by the transovarial mechanism. In addition, they acquire bacteria via transstadial direction from a stage to another stage during feeding practice or localization of rickettsia in the salivary glands. This approach deals with pathogen circulation between vector and host to host (Parola et al. 2005). Hard ticks array within two lineages (the Prostriata and Metastriata) belong to Ixodidae family. The Metastriata group consists of 4 subfamilies: Amblyomminae, Haemaphysalinae, Rhipicephalinae and Hyalomminae. Similarly, Rhipicephalinae includes 9 genera e.g. *Rhipicephalus*, *Dermacentor*, *Margaropus*, and *Boophilus*. (Beati and Keirans 2001). Ticks have been assigned many studies in Iran, including the Ixodidae and Argasidae families. In the case of Ixodidae, *R. norvegicus* was a unique host for *Ixodes ricinus* in Iran so far. This species is the most widespread tick in central Europe (Sekeyova et al. 2000). Besides, it was reported to be positive for *Rickettsia raoultii*, *R. monacensis* and *R. Helvetica* by real-time PCR (Obiegala et al. 2016). In Poland, *Ixodes (I.) ricinus* was more reiterative ticks pertaining to bacterial infections like human granulocytic anaplasmosis (HGA) and Lyme borreliosis (LB) (Grzeszczuka et al. 2006). Subsequently, Granulocytic ehrlichial (GE) DAN was detected by the molecular method in both *Ixodes ricinus* and wild small mammals (*Apodemus sylvaticus*, *A. flavicollis*, *Sorex araneus*) mainly *Clethrionomys glareolus* in Switzerland (Liz et al. 2000). *Babesia microti* (the rodent parasite) and *Babesia divergens* as the causal haemoprotozoan of human babesiosis were demonstrated in *I. ricinus* (nymphal and adult stages) by PCR assay in Europe (Duh et al. 2001; Skotarczak and Cichočka 2001). *B. divergens* as a zoonotic protozoan causes bovine babesiosis in cattle (Zintl et al. 2003). Other species of Ixodidae, *I. trianguliceps* was typically associated with Persian jird in the present study. Also, this tick was a vector of human anaplasmosis. In other respects, it manages the transmission of *Anaplasma phagocytophilum* and *B. microti* to rodents communities under natural conditions (Bown et al. 2008). The genera *Boophilus* currently has been categorized within the subgenus of *Rhipicephalus* which subdivides into five species as follows: *R. (Boophilus) annulatus*, *R. (B) decoloratus*, *R. (B) geigy*, *R. (B) kohlsi* and *R. (B) microplus* (Lempereur et al. 2010). These have been proven as vectors of Babesiosis in livestock and wild animals (Oliveira-Sequeira et al. 2005). *Gerbillinae* has hosted *Boophilus* ticks in the recent data. *Rhipicephalus Sanguineus* (the brown dog tick) has been indicated to be a vector/reservoir of *Rickettsia conorii* and *Ehrlichia canis* (Dantas-Torres 2008). This specimen has been confirmed as a vector of

Theileria spp., *Babesia* spp. and *Anaplasma marginale* parasites in Iranian sheep (Ranjbar-Bahadori 2003; Khayat-Nouri and Hashemzadeh-Farhang 2011; Farzinnia et al. 2012); also it has been accepted to be a significant vector of *Ehrlichia canis* (responsible for canine monocytic ehrlichiosis) (Bremer et al. 2005) as well as spotted fever group rickettsia in Nearctic zones (Burgdorfer et al. 1975). *Haemaphysalis* (166 species) is distributed throughout the world and it acts as the vector of tick-borne diseases such as Kyasanur Forest disease, tick-borne encephalitis virus (Far Eastern subtype) and Rickettsioses (Ahanfarig et al. 2011). This genus has less propensity to bite humans (Duh et al. 2006) but it has been indicated to be the vector of *Babesia major* and *B. bigemina* (protozoan parasites) in animals (Yin et al. 1996). Indeed, it found in vast species e.g. *Haemaphysalis parva*, *H. sulcata*, *H. choldokovskyi*, *H. concinna*, *H. numidiana*, *H. inermis* as well as *H. punctate* that they have infested domestic animals (cattle, sheep, goats and camels) in mountainous regions adjacent to the Caspian Sea, Iran (Rahbari et al. 2007; Hosseini-Chegeni et al. 2014; Hosseini-Vasoukolaei et al. 2010). *Haemaphysalis* ticks collected from *M. musculus*, *M. persicus* and *N. indica*. *Haemaphysalis erinacei* was often found on the hosts as follows: hedgehog, bat, rodents (*Meriones* spp.) and fox cub (Hosseini-Chegeni et al. 2014). Compared with the recent survey, house mouse and *R. norvegicus* have hosted *H. punctate* in Ukraine (Akimov and Nebogatkin 2012). The genera *Hyalomma* has been considered to be the major vector of Crimean-Congo hemorrhagic fever (CCHF) virus and *Nairovirus* transmits through close contact with contaminated blood and tissues of vertebrates (Maltezou et al. 2010). Also this ixodid tick has been gathered from domestic ruminants within variant species including: *Hyalomma dromedarii*, *H. marginatum*, *H. anatolicum*, *H. detritum*, *H. asiaticum*, *H. rufipes* and *H. excavatum* in the whole of Iran (Yakhchali and Haji hasanzadehzarza 2004; Fakoorziba et al. 2006; Nasiri et al. 2010; Salim-abadi et al. 2010; Hosseini-Chegeni et al. 2013; Mirzaei and Khedri 2014). Strikingly, *Rhipicephalus sanguineus*, *H. marginatum*, *H. anatolicum*, *H. asiaticum* and *H. dromedarii* species have been considered as the reservoirs/vectors for CCHF virus in Iran (Telmadarraiy et al. 2015). It is noteworthy that the CCHFV genome newly was isolated from *Rhipicephalus appendiculatus* by RT-PCR in Iran but more researches are needed to discriminate vectorial capacity of this species (Fakoorziba et al. 2015). CCHF is an endemic disease whereas Sistan-Balouchistan, Isfahan, Fars and Khuzestan are the more involved focus in Iran (Alavi-Nainia et al. 2006; Chinikar et al. 2010). Notably, *Tatera indica* has shown to be positive for tularemia by serological test (Pourhossein et al. 2015) coupled with the point that definitive reservoir is not explicit in Iran (Zargar et al. 2015). Ixodid ticks have been

proposed as the vector of *Francisella (F.) tularensis* for man namely *Dermacentor (D.) andersoni*, *D. variabilis*, and *Amblyomma americanum*. *Dermacentor marginatus*, *D. reticulatus*, and *Ixodes (I.) ricinus* ticks were suspected vectors for this bacteria. (Gehringer et al. 2012). On the other hand, *Francisella tularensis* serves as the most biological agent in the weapons of mass destruction (WMD) (Sjostedt 2007). In this literature Argasid soft tick wasn't observed on rodent bodies directly but *Ornithodoros* and *Argas* genera have been collected from human place, domestic animal corrals and poultry farms (Nabian et al. 2007; Enayati et al. 2012; Mohammadi et al. 2013; Aghighi et al. 2007; Telmadarrai et al. 2004; Moradi et al. 2009; Banafshi et al. 2004). Soft ticks have been clarified as certain vectors for TBRF (Tick-borne relapsing fever) infection. This disease has been described to be an endemic disease in rural regions of many provinces such as Hamadan, Zanjan, Kurdistan and Qazvin in particular Ardabil in northwestern Iran (Masoumi-Asl et al. 2009; Moemenbellah-Fard et al. 2009; Rafinejad et al. 2012).

Lice

Lice (Phthiraptera) divide into 4986 genus/species within 4 sub-orders: Anoplura, Amblycera, Ischnocera and Rhynchophthirina. They can infest mammals, birds and many classes of animals (Kenis and Roques 2010). Trench fever, Epidemic typhus and louse-borne relapsing fever (LBRF) are associated with lice (Raoult et al. 1999). In terms of the genus *Polyplax*, it had a wide host spectrum in Iran (Table 1). For instance, *Rattus norvegicus* was specialized host for *Polyplax spinulosa* (spiny rat louse) in Iran similar to other studies (Koyee et al. 2011; Sanchez-Montes et al. 2013). This louse is cosmopolitan parasitic arthropod collected from *Ratt* spp. in all continents as follows: *R. exulans*, *R. nitidus*, *R. norvegicus*, *R. argentiventer*, *R. tanezumi* and *R. rattus* (Durden and Page 1991; Durden and Musser 1994; Durden et al. 1997). It can transmit *Haemobartonella muris* from rat to rat (Crystal 1958). *Polyplax paradoxa* occurred on *M. persicus*, *P. gerbilli* on Norway rat and *T. indica*, *P. serrata* on *M. musculus*, *P. stephensi* on *T. indica*, *P. asiatica* on Persian jird, Indian jird and *Spermophilus fulvus* as well as *Polyplax* spp. on *Sciurus anomalus*. Substantially, *Polyplax serrata* can transmit *Eperythrozoon coccoides* (EC) to mice (Reeves and Cobb 2005). Remarkably, *Polyplax paradoxa* was found on *Meriones crassus* (Durden and Musser 1994). *Gerbillus pyramidum* harbored *P. gerbilli* in Egypt (Happold 1968). *Polyplax indica* appeared on *Golunda ellioti* (the Indian bush rat) (Mishra and Kulkarni 1974). *Eulinognathus* spp. on polyplacidae is ectoparasite of the desert rodent, *Al-lactaga* spp. in Iran up to now. *Eulinognathus americanus* associated with *Ctenomys talarum* (tucó–tucó) in South

America (Martino et al. 2015). *Neohaematopinus* spp. (Hoplopleuridae) was observed on *M. persicus* and *M. socialis* in this information. At least 31 species/subspecies of this genus have been recognized in Holarctic and Asian regions (Shinozaki et al. 2004). More publications have quoted that *Neohaematopinus* species was closely associated with squirrel, *N. appressus* (flying squirrels) (Johnson 1972a), and *N. sciuri* from the grey squirrel (O'Connor et al. 2005) as well as *N. callosciuri* from Pallas squirrel (*Callosciurus erythraeus*) (Shinozaki et al. 2004). Among the *Hoplopleura* species, *H. captiosa* and *H. oenomydis* (the tropical rat louse) were presented on *N. indica*, *M. musculus*, *R. norvegicus* and *R. rattus* at this point. It is apparent that the genus *Hoplopleura* has been distributed widely in geographical territories. *Hoplopleura hirsute* was isolated from cotton rats, *Sigmodon* spp. in the U.S (Kucera et al. 2007) also another species of *Hoplopleura*, *H. acanthopus* from *lasiopodomys* (Kristofik 1999), *Microtus longicaudus* (the long-tailed vole) and *Microtus montanus* (vole) (Hansen 1964). Other records have been described within various landscapes. *Hoplopleura cornata* was found on *R. cornatus* in Australia (Kim 1972), *H. pahari* on *Mus pahari* (Johnson 1972b), *H. captiosa* on *Mus cervicolor*, Thailand (Kim 1966) *H. ramgarh* on *Mus platythrix*, India (Mishra et al. 1972) *H. oenomydis* on (black and Norway rats) (Pratt and Karp 1953) and *H. pacifica* on *R. norvegicus* lastly (Gomez 1989).

Interaction of host-parasite within diversity, density, habitat, climate and host indices

Parasites affect the reproductive, survival dynamic of hosts as well as host dispersal pattern (Heeb et al. 1999) Also, they increase the mortality rate of the juvenile host rodents (Hawlena et al. 2006). As discussed above Muridae was an important group of rodent host that has been diversely adapted to the all geographical regions of Iran. For instance, *R. norvegicus* and black rat have a large-scale distribution along with coastal provinces like Mazandaran, Hormozgan and Bushehr. They are extended to the central parts recently. *Meriones persicus* as a nocturnal jird has been permanently presented in semi-desert areas except Dasht-e Kavir desert. *Tatera indica* digs burrow contiguous with the grasslands and human dwelling that was found in Southeastern Iran. *Meriones libicus* (Libyan jird) was a preponderant host in every part of Iran particularly Yazd and Fars provinces. Until recently, there is no record of this species from the Northwestern and forest regions. *Rhombomys opimus* or great gerbil was more a sociable rodent that was occurred in sandy and clay deserts. It has been reported from Northeastern (Khorasan and Golestan zones) and central districts of Iran (Sedaghat and Salahi-

Moghaddam 2010a, b; Zarei et al. 2010, Dehghani et al. 2013). Murid mammals have been determined as the natural reservoirs of Zoonotic cutaneous leishmaniasis infection (ZCL). For example, *Rhombomys opimus* is documented as the major reservoir of ZCL in central and northeast; *M. libycus*, *T. indica* and *N. indica* in the center towards southwest Iran (Mirzaei et al. 2011; Ghaffari et al. 2013). As a consequence *Meriones persicus* is the frequent host for Pulicid fleas in Iran (Maleki-Ravasan et al. 2017) followed by *M. musculus* and *C. migratorius* (Cricetidae). *Pulex irritans* from Pulicidae is spread in particularly west of Iran that is known as an endemic foci of plague. *Xenopsylla buxtoni*, *N. fasciatus* and *Haemaphysalis* (tick) had a wide host spectrum. It appears that these are heteroxenous parasites because they can shift their mammalian hosts within the life cycle repeatedly (Whiting et al. 2008). Switching from small mammals to bird hosts was also seen in fleas pertaining to ecological characteristics but up to 94% of fleas species principally parasitize mammals and insectivores (Krasnov 2008). It is postulated that fleas are may be considered as the host-opportunistic parasites into two forms; locally or high alpha-specificity (a particular district where parasites may exploit the available of host species) (Krasnov et al. 2011). Additionally, more perspective works are necessary to decipher the mechanism of host speciation in parasites. The genealogy of parasite genome can orient toward the evolutionary process into speciation point (Nieberding and Olivieri 2006). Surveys showed that the fleas population grow according to hosts propensities under the environmental condition (Rafinejad et al. 2013; Momenbellah-Fard et al. 2014). Consequently, micro climatic factors (e.g. humidity and temperature) can affect the survival of juvenile fleas in their habitat (Krasnov et al. 2001). Fleas are strongly susceptible to lack of humidity in their sites also they can spend off-host without blood meals around 89 days (Cox et al. 1999). The Pattern of parasitism is perplexed view related to sexual dimorphism, the scheme of seasonality burdens and body mass of host species (Krasnov et al. 2005; Kiffner et al. 2013). This review was shown that Sistan-Baluchistan and Khuzestan provinces had the similar ectoparasites richness may be due to the same climatic and environmental requirements of species (Rahdar et al. 2015; Nateghpour et al. 2013; Krasnov et al. 2010). Therefore, understanding the ecological models of *Hyalomma* ticks can help us to control of CCHF disease (Ansari et al. 2014) in these provinces. Accordingly, domestic vertebrates (sheep, goat, cattle and cow) was considered as the primitive hosts also Rodentia was alternative hosts for ticks in Iran (Telmadarraiy et al. 2004; Rahbari et al. 2007; Nabian et al. 2007; Salari-Lak

et al. 2008; Mohammadi et al. 2013). There is an increasing belief that Ixodes ticks (particularly *Hyalomma* spp.) commonly existed throughout the months, but more frequent in spring and late autumn while Argasid ticks exhibited exclusively in autumn to late winter in Northwestern Iran (Rahbari 1995; Salari-Lak et al. 2008; Bahman Shabestari and karimian 2011). Other ectoparasites groups such as *Polyplax spinulosa* (sucking louse) and *L. nuttalli* (mite) infested rodents in autumn and spring respectively (Telmadarraiy et al. 2007). As such, seasonal tick activity had two major peaks, one in early summer and the other one in autumn in accordance with the environmental temperature in addition, ticks were less susceptible to relative humidity fluctuations (Hussein 1980). Each kind of seasonal alternation affects the preferred host species in ticks relating to their biome (Randolph and Rogers 2007). *Boophilus annulatus* and *I. ricinus* (on domestic ruminants) are disturbed only in the regions with the highest percentage of forest cover in Iran (Nabian et al. 2007) also many tick species such as *Ornithodoros* and *Rhipicephalus* are more prevalent in the case of numbering in mountainous suburbs compared with plateau areas (Telmadarraiy et al. 2004). Alborz mountain range has steppe climates, which lies between south of the Caspian Sea and plateau (Darvish et al. 2014). Prevalence of rodents and related ectoparasites was high in some fragments containing organic supplies but they was had less parasites diversity (Hamidi et al. 2015). Access to food resources affect the evolutionary aspects within host–parasite linkage (Tschirren et al. 2007). It is remarkable that the greater degree of homogeneity within ecological components in habitat of sociable rodents reduce macroparasites diversity (Bordes et al. 2007) and increase host density (Froeschke et al. 2013). Notably, Host tendency within biased parasitism has been assigned periodically in many studies (Gorrell and Schulte-Hostedde 2008; Morand et al. 2004; Perez-Orella and Schulte-Hostedde 2005). Immunosuppression, Social reaction and stable temperature during the reproductive time induce parasite load in female than male (Christe et al. 2007). Additionally, most female ectoparasites have active selection to host gender for feeding therefore they adapt to male host greatly (Khokhlova et al. 2011). There are well known hypotheses for interpretation of this phenomenon. First, high level of androgenic hormones (e.g. testosterone) can depress immune system in male hosts and increase the predisposition to parasitism (Mougeot et al. 2006). Male host grooms less than female during the reproductive seasons because of the suppressive impact of testosterone (Mooring and Hart 1995). This hormone lead to more mobility of host within its home range

thereby, it obtain high infestation rate of parasite (Hughes and Randolph 2001). Secondary outcome focus on the sexual size dimorphism (SSD) refers to individual males that more likely harbor greater parasites as many of mammalian males have the high body mass than females (Harrison et al. 2010). There is no simple mechanism to this affair actually. Besides, difference in ectoparasitism may result from variant model of parasite behaviour based on occupied burrows between males and females (Waterman et al. 2014). In this review two independent ectoparasites (*O. bacoti* and *H. oenomydis*) coexisted on *R. norvegicus*. Perhaps host defence has the important role in the interactions between rival species and organizes co-occurrence of parasite communities (Bush and Malenke 2008). Mites were more diverse than others may be due to their small size body compared with other taxa (Walter and Proctor 2013). This finding was obviously shown that *P. spinulosa* was strictly host-specific to Rat species. In the following, Haematophagous lice have certain morphological traits within claw structure that adapted to their host body and hair size. For this reason, they are mainly host-specific and rarely attach to each other individuals hair (Cannon 2010; Zuo et al. 2011). Unfortunately, few studies have been carried out concerning to host-ectoparasite aspects in Persia thereby, the ecological, seasonality, diversity and host speciation dimensions are still equivocal approximately. Obviously, it should be employed more genetic and phylogenetic studies about the identification of ectoparasites within subfamilies or sub-species levels to incorporate a comprehensive picture of parasite-host association. Authorities importantly scrimmage with two emerging diseases. Tick-borne relapsing fever is the more predominant disease in the highland mountainous regions and CCHF cases have been more reported from southern, central and western Iran. Although, plague cases has not been reported since long time ago but related fleas species have been occurred on captured rodents in Bijar County (the earlier district of plague) frequently. Strikingly, Alborz and Zagros mountains ranges provide the specialized habitat for animals that are enclosed by northern and western regions as well as Kavir and Dasht-e Loot deserts in central plateau toward south of Iran. Besides, the Iranian mountains can prohibit gene flow between local mammalian communities thus lead to generate the populations with different gene pool. Also this event raise the genetic variation into sympatric/allopatric speciation probably. In addition to, human interventions in nature within urbanization or agricultural activities induce many uncontrolled changes in animal life history and vector-borne diseases. For instance, these modifications affect longevity, diversity, microclimate, density and adaptive host in ectoparasites.

Also they impress the habitat, abundance, spatial distribution and supply accessibility in hosts by different methods (Friggens and Beier 2010).

Acknowledgements I would like to thank Ms. Azam Rafat-panah for help to write the draft of the manuscript.

Compliance with ethical standards

Conflict of interest This paper hasn't received financial interests by any organization, corporation or university.

References

- Adjemian JZ, Foley P, Gage KL, Foley JE (2007) Initiation and spread of traveling waves of plague, *Yersinia pestis*, in the Western United States. *Am J Trop Med Hyg* 76(2):365–375. doi: [10.4269/ajtmh.2007.76.365](https://doi.org/10.4269/ajtmh.2007.76.365)
- Aghighi Z, Assmar M, Piazak N, Javadian E et al (2007) Distribution of soft ticks and their natural infection with *Borrelia* in a focus of relapsing fever in Iran. *Iran J Arthropod-Borne Dis* 1(2):14–18
- Ahadiyat A, Masan P, Cheraghali Z, Joharchi O (2014) First report of the subgenus *Pachylaelaps* (*Longipachylaelaps*) Mašán (Mesostigmata: Eviphidoidea: Pachylaelapidae) from Iran. *Persian J Acarol* 3(1):99–102
- Ahantari A, Malaisri P, Hirunkanokpun S et al (2011) Detection of *Rickettsia* and a novel *Haemaphysalis shimoga* symbiont bacterium in ticks in Thailand. *Curr Microbiol* 62:1496–1502. doi: [10.1007/s00284-011-9887-3](https://doi.org/10.1007/s00284-011-9887-3)
- Akimov IA, Nebogatkin IV (2012) Distribution of the tick *Haemaphysalis punctata* (Acari, Ixodidae) in Ukraine. *Vestnik Zoologii* 46(4):e46–e51. doi: [10.2478/v10058-012-0030-0](https://doi.org/10.2478/v10058-012-0030-0)
- Alavi-Nainia R, Moghtaderib A, Koohpayeha HR, Sharifi-Mooda B et al (2006) Crimean-Congo hemorrhagic fever in Southeast of Iran. *J Infect* 52:378–382. doi: [10.1016/j.jinf.2005.07.015](https://doi.org/10.1016/j.jinf.2005.07.015)
- Albon SD, Stien A, Irvine RJ, Langvatn R, Ropstad E, Halvorsen O (2002) The role of parasites in the dynamics of a reindeer population. *Proc R Soc Lond B Biol Sci* 269(1500):1625–1632. doi: [10.1098/rspb.2002.2064](https://doi.org/10.1098/rspb.2002.2064)
- Allymehr M, Tavassoli M, Manoochehri MH, Ardavan D (2012) Ectoparasites and gastrointestinal helminths of house mice (*Mus musculus*) from poultry houses in Northwest Iran. *Comp Parasitol* 79(2):283–287. doi: [10.1654/4534.1](https://doi.org/10.1654/4534.1)
- Amin OM (1976) Lice, mites, and ticks of southeastern Wisconsin mammals. *Great Lakes Entomol* 9(4):195–198
- Ansari H, Shahbaz B, Izadi Sh, Zeinali M et al (2014) Crimean-Congo hemorrhagic fever and its relationship with climatic factors in southeast Iran: a 13-year experience. *J Infect Dev Ctries* 8(6):749–757. doi: [10.3855/jidc.4020](https://doi.org/10.3855/jidc.4020)
- Arjomandi E, Kazemi SH, Afshari A (2013) Fauna and diversity of the manure-inhabiting Mesostigmata (Acari) in Kerman County, South Eastern Iran. *Persian J Acarol* 2(2):253–263
- Bahman Shabestari A, Karimian A (2011) Ixodidae ticks fauna of sheep at Qazvin province. *J Vet Clin Res* 2(4):241–247 (**In Persian**)
- Banafshi O, Rafinejad J, Esmail-Nasab N (2004) A study on Argasidae distribution in internal shelters and assessment of infestation rate of *Ornithodoros tholozani* to *Borrelia persica* in Bijar county, Kordestan Province, 2003. *J Kordestan Univ Med Sci* 31:50–58 (**In Persian**)
- Baumstark J, Beck W, Hofmann H (2007) Outbreak of tropical rat mite (*Ornithonyssus bacoti*) dermatitis in a home for disabled persons. *Dermatology* 215(1):66–68. doi: [10.1159/000102037](https://doi.org/10.1159/000102037)

- Beati L, Keirans JE (2001) Analysis of the systematic relationships among ticks of the genera *Rhipicephalus* and *Boophilus* (Acari: Ixodidae) based on mitochondrial 12S ribosomal DNA gene sequences and morphological characters. *J Parasitol* 87(1):32–48
- Beck W, Folster-Holst R (2009) Tropical rat mites (*Ornithonyssus bacoti*) serious ectoparasites. *JDDG* 7:667–670. doi: [10.1111/j.1610-0387.2009.07140.x](https://doi.org/10.1111/j.1610-0387.2009.07140.x)
- Bhuyan PJ, Nath AJ (2016) Record of tropical rat mite, *Ornithonyssus bacoti* (Acari: Mesostigmata: Macronyssidae) from domestic and peridomestic rodents (*Rattus rattus*) in Nilgiris, Tamil Nadu, India. *J Arthropod-Borne Dis* 10(1):98–101
- Billetter SA, Gundi VAKB, Rood MP, Kosoy MY (2011) Molecular detection and identification of *Bartonella* species in *Xenopsylla cheopis* fleas (Siphonaptera: Pulicidae) collected from *Rattus norvegicus* Rats in Los Angeles, California. *Appl Environ Microbiol* 77(21):7850–7852. doi: [10.1128/AEM.06012-11](https://doi.org/10.1128/AEM.06012-11)
- Bitam I, Dittmar K, Parola P (2010) Fleas and flea-borne diseases. *Int J Infect Dis* 14:e667–e676. doi: [10.1016/j.ijid.2009.11.011](https://doi.org/10.1016/j.ijid.2009.11.011)
- Bochkov AV, Labrzycka A (2003) A revision of the European species of the genus *Myobia* von Heyden, 1826 (Acari, Myobiidae). *Acta Parasitol* 48(1):24–40
- Bordes F, Blumstein DT, Morand S (2007) Rodent sociality and parasite diversity. *Biol Lett* 3:692–694. doi: [10.1098/rsbl.2007.0393](https://doi.org/10.1098/rsbl.2007.0393)
- Bown KJ, Lambin X, Telford GR, Ogden NH (2008) Relative Importance of *Ixodes ricinus* and *Ixodes trianguliceps* as Vectors for *Anaplasma phagocytophilum* and *Babesia microti* in Field Vole (*Microtus agrestis*) Populations. *Appl Environ Microbiol* 74(23):7118–7125. doi: [10.1128/AEM.00625-08](https://doi.org/10.1128/AEM.00625-08)
- Breitschwerdt EB, Kordick DL (2000) bartonella infection in animals: carriership, reservoir potential, pathogenicity, and zoonotic potential for human infection. *J Clin Microbiol Rev* 13(3):428–438
- Bremer WG, Schaefer JJ, Wagner ER, Ewing SA (2005) Transstadial and intrastadial experimental transmission of *Ehrlichia canis* by male *Rhipicephalus sanguineus*. *Vet Parasitol* 131:95–105. doi: [10.1016/j.vetpar.2005.04.030](https://doi.org/10.1016/j.vetpar.2005.04.030)
- Brooks JE, Jackson WB (1973) A review of commensal rodents and their control. *C R C Crit Rev Environ Control* 3(1–4):405–453. doi: [10.1080/1064338309381607](https://doi.org/10.1080/1064338309381607)
- Brown NS, Wilson GIA (1975) Comparison of the ectoparasites of the house sparrow (*Passer domesticus*) from North America and Europe. *Am Midl Nat* 94(1):154–165
- Brown PR, Yee N, Singleton GR, Kenney AJ et al (2008) Farmers' knowledge, attitudes, and practices for rodent management in Myanmar. *Int J Pest Manag* 54(1):69–76. doi: [10.1080/09670870701474047](https://doi.org/10.1080/09670870701474047)
- Burbutis PP, Hansens EJ (1955) Fleas on rats (*Rattus Norvegicus*) in New Jersey. *J N Y Entomol Soc* 63:103–106
- Burgdorfer W, Sexton DJ, Gerloff RK, Anacker RL et al (1975) *Rhipicephalus sanguineus*: vector of a new spotted fever group Rickettsia in the United States. *Infect Immun* 12(1):205–210
- Bush SE, Malenke JR (2008) Host defence mediates interspecific competition in ectoparasites. *J Anim Ecol* 77:558–564. doi: [10.1111/j.1365-2656.2007.01353.x](https://doi.org/10.1111/j.1365-2656.2007.01353.x)
- Cannon ShM (2010) Size correlations between sucking lice and their hosts including a test of harrison's rule. *Electro Theses Dissertations* 764. <http://digitalcommons.georgiasouthern.edu/etd/764>
- Chinikar S, Ghiasi SM, Hewson R, Moradi M, Haeri A (2010) Crimean-Congo hemorrhagic fever in Iran and neighboring countries. *J Clin Virol* 47:110–114. doi: [10.1016/j.jcv.2009.10.014](https://doi.org/10.1016/j.jcv.2009.10.014)
- Christe P, Glaizot O, Evanno G, Bruyndonckx N et al (2007) Host sex and ectoparasites choice: preference for, and higher survival on female hosts. *J Anim Ecol* 76:703–710. doi: [10.1111/j.1365-2656.2007.01255.x](https://doi.org/10.1111/j.1365-2656.2007.01255.x)
- Cicek H, Stanyukovich M, Yagi S (2008) Gamasine mite (Parasitiformes: Mesostigmata) infestations of small mammals (Mammalia: Rodentia, Insectivora) in Turkey. *Turkiy Parazitoloji Dergisi* 32(1):65–70
- Cox R, Stewart PD, Macdonald DW (1999) The ectoparasites of the European Badger, *Meles meles*, and the behavior of the host-specific flea, *Paraceras melis*. *J Insect Behav* 12(2):245–265
- Craig P, Ito A (2007) Intestinal cestodes. *Curr Opin Infect Dis* 20:524–532
- Crystal MM (1958) The mechanism of transmission of haemobartonella muris (Mayer) of rats by the spined Rat Louse, *Polyplax spinulosa* (Burmeister). *J Parasitol* 44(6):603–606
- Dantas-Torres F (2008) The brown dog tick, *Rhipicephalus sanguineus* (Latreille, 1806) (Acari: Ixodidae): from taxonomy to control. *Vet Parasitol* 152:173–185. doi: [10.1016/j.vetpar.2007.12.030](https://doi.org/10.1016/j.vetpar.2007.12.030)
- Darvish J, Mohammadi Z, Mahmoudi A, Siahsharvie R (2014) Faunistic and taxonomic study of Rodents from northwestern Iran. *Iran J Anim Biosyst* 10(2):119–136
- Darvishi MM, Youssefi MR, Changizi E et al (2014) A new flea from Iran. *Asian Pac J Trop Dis* 4(2):85–87. doi: [10.1016/S2222-1808\(14\)60321-2](https://doi.org/10.1016/S2222-1808(14)60321-2)
- Davis DE (1951) Observations on rat ectoparasites and typhus fever in San Antonio, Texas. *Public Health Reports (1896–1970)* 66(52):1717–1726
- Dehghani R, Seyedi HR, Dehqan S, Sharifi H (2013) Geographical distribution of mouse and mouse-borne diseases in Iran. A review article. *Feyz* 17(2):203–219 (In Persian)
- Duh D, Petrovec M, Avsic-Zupanc T (2001) Diversity of *Babesia* infecting European sheep ticks (*Ixodes ricinus*). *J Clin Microbiol* 39(9):3395–3397
- Duh D, Punda-Polic V, Trilar T (2006) Molecular identification of *Rickettsia felis*-like Bacteria in *Haemaphysalis sulcata* ticks collected from domestic animals in Southern Croatia. *Ann NY Acad Sci* 1078:347–351. doi: [10.1196/annals.1374.068](https://doi.org/10.1196/annals.1374.068)
- Durden LA, Musser GG (1994) The mammalian hosts of the sucking lice (Anoplura) of the world: a host-parasite list. *Bull Soc Vector Ecol* 19(2):130–168
- Durden LA, Page BF (1991) Ectoparasites of commensal rodents in Sulawesi Utara, Indonesia, with notes on species of medical importance. *Med Vet Entomol* 5:1–7
- Durden LA, Kollars TM Jr, Patton SH, Gerhard RR (1997) Sucking lice (Anoplura) of mammals of Tennessee. *J Vector Ecol* 22(1):71–76
- Eisen RJ, Bearden SW, Wilder AP et al (2006) Early-phase transmission of *Yersinia pestis* by unblocked fleas as a mechanism explaining rapidly spreading plague epizootics. *PNAS* 103(42):15380–15385. doi: [10.1073/pnas.0606831103](https://doi.org/10.1073/pnas.0606831103)
- Enayati AA, Asgarian F, Amouei A, Esfandiari B et al (2012) First records of *Ornithodoros* sp. soft ticks from an endemic relapsing fever region in Northern Iran. *HealthMED* 6(6):1911–1916
- Eremeeva ME, Warashina WR, Sturgeon MM et al (2008) *Rickettsia typhi* and *R. felis* in Rat Fleas (*Xenopsylla cheopis*), Oahu, Hawaii. *Emerg Infect Dis* 14(10):1613–1615. doi: [10.3201/eid1410.080571](https://doi.org/10.3201/eid1410.080571)
- Eskey CR (1938) Flea infestation of domestic rats in San Francisco, Calif. *Public Health Reports (1896–1970)* 53(23):948–951
- Fagir DM, El-Rayah EA (2009) Parasites of the Nile rat in rural and urban regions of Sudan. *Integr Zool* 4:179–187. doi: [10.1111/j.1749-4877.2009.00148.x](https://doi.org/10.1111/j.1749-4877.2009.00148.x)
- Fain A, Hyland K (1970) Notes on the myocoptidae of North America with description of a new species on the Eastern Chipmunk, *Tamias striatus* Linnaeus. *J N Y Entomol Soc* 78(2):80–87

- Fakoorziba MR, Neghab M, Alipour H, Moemenbellah-Fard MD (2006) Tick borne crimean-congo hemorrhagic fever in Fars province, Southern Iran: epidemiologic characteristics and vector surveillance. *Pak J Biol Sci* 9(14):2681–2684
- Fakoorziba MR, Naddaf-Sani AA, Moemenbellah-Fard MD (2015) First phylogenetic analysis of a Crimean-Congo hemorrhagic fever virus genome in naturally infected *Rhipicephalus appendiculatus* ticks (Acari: Ixodidae). *Arch Virol* 160:1197–1209. doi:10.1007/s00705-015-2379-1
- Farzinnia B, Saghafipour A, Telmadarreyi Z (2012) Survey of tick species distribution in sheep and camel in Qom city, Iran, 2010–2011. *J North Khorasan Uni Med Sci* 4(3):391–398 (In Persian)
- Feliu C, Lopez M, Gomez M, Torres J et al (2012) Parasite fauna of rodents (Murinae) from El Hierro (Canary Islands, Spain): a multidisciplinary approach. *Acta Parasitol* 57(2):171–178. doi:10.2478/s11686-012-0016-7
- Fischer K, Walton S (2014) Parasitic mites of medical and veterinary importance—is there a common research agenda? *Int J Parasitol* doi:10.1016/j.ijpara.2014.08.003
- Ford PL, Fagerlund RA, Duszynski DW, Polechla PJ (2004) Fleas and lice of mammals in New Mexico. Gen. Tech. Rep. RMRS-GTR-123. Fort Collins, CO: U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station
- Friggens MM, Beier P (2010) Anthropogenic disturbance and the risk of flea-borne disease transmission. *Oecologia* 164:809–820. doi:10.1007/s00442-010-1747-5
- Froeschke G, Luther van der Mescht Lvd, McGeoch M, Matthee S (2013) Life history strategy influences parasite responses to habitat fragmentation. *Int J Parasitol* doi:10.1016/j.ijpara.2013.07.003
- Fuente JDL, Estrada-Pena A, Venzal JM et al (2008) Overview: ticks as vectors of pathogens that cause disease in humans and animals. *Front Biosci* 13:6938–6946
- Gage KL, Kosoy MY (2005) Natural history of plague: perspectives from more than a century of research. *Annu Rev Entomol* 50:505–528. doi:10.1146/annurev.ento.50.071803.130337
- Gehring H, Schacht E, Maylaender N (2012) Presence of an emerging subclone of *Francisella tularensis holarctica* in *Ixodes ricinus* ticks from south-western Germany. *Ticks Tick-borne Dis*. doi:10.1016/j.ttbdis.2012.09.001
- Ghaffari D, Hakimi Parizi M, Yaghoobi Ershadi MR, Sharifi I, Akhavan AA (2013) A survey of reservoir hosts in two foci of cutaneous leishmaniasis in Kerman province, southeast of Iran. *J Parasit Dis*. doi:10.1007/s12639-013-0275-8
- Gholipoury M, Rezaei HR, Namroodi S, Arab-Khzaeli F (2016) Zoonotic and non-zoonotic parasites of wild rodents in turk-man Sahra, Northeastern Iran. *Iran J Parasitol* 11(3):350–357
- Gomez MS (1989) The occurrence of *Hoplopleura pacifica* (Anoplura, Hoplopleuridae) on *Rattus norvegicus* of Barcelona. *Ann Parasitol Hum Comp* 64(6):516–517
- Gorrell JC, Schulte-Hostedde AI (2008) Patterns of parasitism and body size in red squirrel (*Tamiasciurus hudsonicus*). *Can J Zool* 86:99–107. doi:10.1139/Z07-123
- Green WF, Woolcock AJ (1978) *Tyrophagus putrescentiae*: an allergenically important mite. *Clin Allergy* 8:135–144
- Grzeszczuka A, Ziarkob S, Kovalchuk O, Stanczak J (2006) Etiology of tick-borne febrile illnesses in adult residents of north-eastern Poland: report from a prospective clinical study. *Int J Med Microbiol* 296(S1):242–249. doi:10.1016/j.ijmm.2006.01.007
- Hamidi K, Nourani L, Moravvej Gh (2015) The relationship of ectoparasite prevalence to the capturing season, locality and species of the murine rodent hosts in Iran. *Persian J Acarol* 4(4):409–423
- Hanafi-Bojd AA, Shahi M, Baghaii M et al (2007) A study on rodent ectoparasites in Bandar Abbas: the main economic southern seaport of Iran. *Iran J Environ Health Sci Eng* 4(3):173–176
- Hansen CG (1964) Ectoparasites of mammals from Oregon. *Great Basin Nat* 24(2):75–81
- Happold DCD (1968) Observations on *Gerbillus pyramidum* (Gerbillinae, Rodentia) at Khartoum. *Sudan Mammalia* 32(1):44–53. doi:10.1515/mamm.1968.32.1.44
- Harrison A, Scantlebury M, Montgomery WL (2010) Body mass and sex-biased parasitism in wood mice *Apodemus sylvaticus*. *Oikos* 119(7):1099–1104. doi:10.1111/j.1600-0706.2009.18072.x
- Harrison A, Robbb GN, Alagaili AN (2015) Ectoparasite fauna of rodents collected from two wildlife research centres in Saudi Arabia with discussion on the implications for disease transmission. *Acta Trop*. doi:10.1016/j.actatropica.2015.03.022
- Hashemi-Shahraki A, Carniel E, Mostafavi E (2016) Plague in Iran: its history and current status. *Epidemiol Health* 38, ID: e2016033
- Hawlena H, Abramsky Z, Krasnov BR (2006) Ectoparasites and age-dependent survival in a desert rodent. *Oecologia* 148:30–39. doi:10.1007/s00442-005-0345-4
- Hawlena H, Abramsky Z, Krasnov BR, Saltz D (2007) Host defence versus intraspecific competition in the regulation of infrapopulations of the flea *Xenopsylla conformis* on its rodent host *Meriones crassus*. *Int J Parasitol* 37:919–925. doi:10.1016/j.ijpara.2007.01.015
- Heeb Ph, Werner I, Mateman AC, Koelliker M et al (1999) Ectoparasite infestation and sex-biased local recruitment of hosts. *Nature* 400(1):63–65
- Heide SVD, Niemeijer NR, Hovenga H et al (1998) Prevalence of sensitization to the storage mites *Acarus siro*, *Tyrophagus putrescentiae*, and *Lepidoglyphus destructor* in allergic patients with different degrees of sensitization to the house-dust mite *Dermatophagoides pteronyssinus*. *Allergy* 53:426–430
- Hosseini-Chegeni A, Hosseini R, Tavakoli M, Telmadarraiy Z, Abdigoudarzi M (2013) The Iranian *Hyalomma* (Acari: Ixodidae) with a key to the identification of male species. *Persian J Acarol* 2(3):503–529
- Hosseini-Chegeni A, Telmadarraiy Z, Salimi M, Arzamani K, Banafshi O (2014) A record of *Haemaphysalis erinacei* (Acari: Ixodidae) collected from Hedgehog and an identification key for the species of *Haemaphysalis* occurring in Iran. *Persian J Acarol* 3(3):203–215
- Hosseini-Vasoukolaei N, Telmadarraiy Z, Vatandoost H et al (2010) Survey of tick species parasiting domestic ruminants in Ghaemshahr county, Mazandaran province, Iran. *Asian Pacific J Trop Med* pp 804–806
- Hughes VL, Randolph SE (2001) Testosterone increases the transmission potential of tick-borne parasites. *Parasitology* 123:365–371. doi:10.1017/S0031182001008599
- Huhta V (2016) Catalogue of the mesostigmata mites in Finland. *Memoranda Soc Fauna Flora Fennica* 92:129–148
- Hussein HS (1980) *Ixodes trianguliceps*: seasonal abundance and role in the epidemiology of *Babesia microti* infection in north-western England. *Ann Trop Med Parasitol* 74(5):531–539. doi:10.1080/00034983.1980.11687381
- Jakeman LAR (1961) The internal anatomy of the spiny rat mite, *Echinolaelaps echidninus* (Berlese). *J Parasitol* 47(2):329–349
- Jiang J, Soeatmadji DW, Henry KM et al (2006) *Rickettsia felis* in *Xenopsylla cheopis*, Java, Indonesia. *Emerg Infect Dis* 12(8):1281–1283
- Johansson E, Aponno M, Lundberg M, Hage-Hamsten MV (2001) Allergenic cross-reactivity between the nematode *Anisakis simplex* and the dust mites *Acarus siro*, *Lepidoglyphus destructor*, *Tyrophagus putrescentiae*, and *Dermatophagoides pteronyssinus*. *Allergy* 56:660–666

- Johnson PT (1972a) *Neohaematopinus appressus*, a new species of sucking louse from an Aslan tree squirrel (Anoplura). *Pacific Insects* 14(2):389–392
- Johnson PT (1972b) Two new species of *Hoplopleura* Enderlein from Laotian Murids (Anoplura). *Pacific Insects* 14(3):607–611
- Kenis M, Roques A (2010) Lice and Fleas (Phthiraptera and Siphonaptera). Chapter 13.4. In: Roques A et al (eds) *Alien terrestrial arthropods of Europe*. *BioRisk* 4(1): 833–849. doi: [10.3897/biorisk.4.65](https://doi.org/10.3897/biorisk.4.65)
- Khayat-Nouri MH, Hashemzadeh-Farhang H (2011) Survey on species diversity and season changes of Ixodidae ticks in Tabriz in 2009. *Vet. J Islamic Azad Uni Tabriz Branch* 5(3): 1273–1279 (In Persian)
- Khokhlova IS, Serobyana V, Degen AA, Krasnov BR (2011) Discrimination of host sex by a haematophagous ectoparasite. *Anim Behav* 81:275–281. doi: [10.1016/j.anbehav.2010.10.018](https://doi.org/10.1016/j.anbehav.2010.10.018)
- Khosravani M, Nasiri Z, Keshavarz D, Rafat-Panah A (2016) Epidemiological trend of cutaneous leishmaniasis in two endemic focus of disease, south of Iran. *J Parasite Dis* 40:1609–1613. doi: [10.1007/s12639-015-0740-7](https://doi.org/10.1007/s12639-015-0740-7)
- Kiffner Ch, Stanko M, Morand S, Khokhlova IS et al (2013) Sex-biased parasitism is not universal: evidence from rodent–flea associations from three biomes. *Oecologia*. doi: [10.1007/s00442-013-2664-1](https://doi.org/10.1007/s00442-013-2664-1)
- Kim KCh (1966) A new species of *Hoplopleura* from Thailand, with notes and description of nymphal stages of *Hoplopleura captiosa* Johnson (Anoplura). *Parasitology* 56:603–612
- Kim KCh (1972) A new species of *Hoplopleura* (Anoplura: Hoplopleuridae) from an Australian rat. *Pacific Insects* 14(4):675–678
- Knee W, Proctor H (2007) Host records for *Ornithonyssus sylviarum* (Mesostigmata: Macronyssidae) from Birds of North America (Canada, United States, and Mexico). *J Med Entomol* 44(4):709–713. doi: [10.1603/0022-2585\(2007\)44\[709:HRFOSM\]2.0.CO;2](https://doi.org/10.1603/0022-2585(2007)44[709:HRFOSM]2.0.CO;2)
- Koyee QMK, Ahmed RK, Aziz KK, Ahmed HSh, Abdula AH (2011) Infestation rate with polyplax spinulosa (burmeister, 1839) among certain laboratory albino rats (*Rattus norvegicus*) in relation to different washing agents in two Erbil city universities. *Zanco J Med Sci* 15(1):47–52
- Krasnov BR (2008) *Functional and evolutionary ecology of fleas: a model for ecological parasitology*. Cambridge University Press, Cambridge
- Krasnov BR, Khokhlova IS, Fielden LJ, Burdelova NV (2001) Effect of air temperature and humidity on the survival of pre-imaginal stages of two flea species (Siphonaptera: Pulicidae). *J Med Entomol* 38(5):629–637. doi: [10.1603/0022-2585-38.5.629](https://doi.org/10.1603/0022-2585-38.5.629)
- Krasnov BR, Morand S, Hawlena H, Khokhlova IS, Shenbrot GI (2005) Sex-biased parasitism, seasonality and sexual size dimorphism in desert rodents. *Oecologia* 146:209–217. doi: [10.1007/s00442-005-0189-y](https://doi.org/10.1007/s00442-005-0189-y)
- Krasnov BR, Mouillot D, Shenbrot GI, Khokhlova IS et al (2010) Similarity in ectoparasite faunas of Palaearctic rodents as a function of host phylogenetic, geographic or environmental distances: which matters the most? *Int J Parasitol* 40:807–817. doi: [10.1016/j.ijpara.2009.12.002](https://doi.org/10.1016/j.ijpara.2009.12.002)
- Krasnov BR, Mouillot D, Shenbrot GI, Khokhlova IS, Poulin R (2011) Beta-specificity: the turnover of host species in space and another way to measure host specificity. *Int J Parasitol* 41:33–41. doi: [10.1016/j.ijpara.2010.06.001](https://doi.org/10.1016/j.ijpara.2010.06.001)
- Kristofik J (1999) Sucking lice (Phthiraptera) from Mongolian mammals. *Biol Bratislava* 54(2):143–149
- Kucera JR, Durden LA, Kim KC (2007) New records of sucking lice (Phthiraptera: Anoplura) from the western United States. *J Vector Ecol* 32(2):366–370. doi: [10.3376/1081-1710\(2007\)32\[366:NROSLP\]2.0.CO;2](https://doi.org/10.3376/1081-1710(2007)32[366:NROSLP]2.0.CO;2)
- Kuwahara Y, Ishii S, Fukami H (1975) Neryl formate: alarm pheromone of the cheese mite, *Tyrophagus putrescentiae* (Schrank) (Acarina, Acaridae). *Experientia* 31(10):1115–1116
- Lahey T (2003) Rickettsialpox in Turkey. *Emerg Infect Dis* 9(11):1496–1498
- Lareschi M, Gonzalez-Acuna D (2010) Acari, Laelapidae (ectoparasitic mites), central and southern Chile. *Check List* 6(4):546–548
- Laudisoit A, Leirs H, Makundi RH et al (2007) Plague and the Human Flea, Tanzania. *Emerg Infect Dis* 13(5):687–693
- Lee Y-I, Pyeon H-J, Seo M (2013) Intestinal parasites among wild rodents in Northern Gangwon-do, Korea. *Korean J Parasitol* 51(5):603–606. doi: [10.3347/kjp.2013.51.5.603](https://doi.org/10.3347/kjp.2013.51.5.603)
- Lempereur L, Geysen D, Madder M (2010) Development and validation of a PCR–RFLP test to identify African Rhipicephalus (Boophilus) ticks. *Acta Trop* 114:55–58. doi: [10.1016/j.actatropica.2010.01.004](https://doi.org/10.1016/j.actatropica.2010.01.004)
- Lewis RE (1964) A collection of fleas (Siphonaptera) from Northern Saudi Arabia. *J Parasitol* 50(2):313–318
- Li L, Rollin PE, Nichol ST, Shope RE, Barrett ADT, Holbrook MR (2004) Molecular determinants of antigenicity of two subtypes of the tick-borne flavivirus Omsk haemorrhagic fever virus. *J Gen Virol* 85(6):1619–1624
- Lindquist EE, Galloway TD, Artsob H (2016) *A handbook to the ticks of Canada (Ixodida: Ixodidae, Argasidae)*. Biological Survey of Canada Monograph Series No. 7. doi: [10.3752/9780968932186](https://doi.org/10.3752/9780968932186)
- Liz JS, Anderes L, Sumner JW, Massung RF et al (2000) PCR detection of granulocytic ehrlichiae in *Ixodes ricinus* ticks and wild small mammals in western Switzerland. *J Clin Microbiol* 38(3):1002–1007
- Llerena LP, Ferruindez-Caldas E, Gracia LRC, Lackey RF (1991) Sensitization to *Bbmia tropicalis* and *Lepidoglyphus destructor* in *Dermatopiragrides* spp-attergic individuals. *J Allergy Clin Immunol* 88(6):943–950
- Maleki-Ravasan N, Solhjoui-Fard S, Beaucournu J-C, Laudisoit A, Mostafavi E (2017) The Fleas (Siphonaptera) in Iran: diversity, host range, and medical importance. *PLoS Negl Trop Dis* 11(1):e0005260. doi: [10.1371/journal.pntd.0005260](https://doi.org/10.1371/journal.pntd.0005260)
- Maltezou HC, Andonova L, Andraghetti R, Bouloy M et al (2010) Crimean-Congo hemorrhagic fever in Europe: current situation calls for preparedness. *Euro Surveill*. 15(10):19504. <http://www.eurosurveillance.org/ViewArticle.aspx?ArticleId=19504>
- Martino NS, Romero MD, Malizia AI (2015) Parasitism underground: lice (Insecta: Phthiraptera) from *Ctenomys talarum* (Rodentia: Ctenomyidae) along its coastal distribution in Argentina. *Acta Parasitol* 60(1):154–157. doi: [10.1515/ap-2015-0021](https://doi.org/10.1515/ap-2015-0021)
- Masan P, Halliday B (2014) A new genus and species of soil mite (Acari: Pachylaelapidae: Pachylaelapinae) from the Canary Islands. *J Natl Hist* 48(7–8):403–412. doi: [10.1080/00222933.2013.791950](https://doi.org/10.1080/00222933.2013.791950)
- Masan P, Ozbek HH, Fend'a P (2016) Two new species of *Pachylaelaps* Berlese, 1888 from the Iberian Peninsula, with a key to European species (Acari, Gamasida, Pachylaelapidae). *ZooKeys* 603:71–95. doi: [10.3897/zookeys.603.9038](https://doi.org/10.3897/zookeys.603.9038)
- Masoumi-Asl H, Goya MM, Vatandoost H, Zahraei SM et al (2009) The epidemiology of tick-borne relapsing fever in Iran during 1997–2006. *Travel Med Infect Dis* 7:160–164. doi: [10.1016/j.tmaid.2009.01.0](https://doi.org/10.1016/j.tmaid.2009.01.0)
- Medvedev SG (1996) Geographical distribution of families of Fleas (Siphonaptera). *Entomol Rev* 76(8):978–992
- Meerburg BG, Singleton GR, Kijlstra A (2009) Rodent-borne diseases and their risks for public health. *Crit Rev Microbiol* 35(3):221–270. doi: [10.1080/10408410902989837](https://doi.org/10.1080/10408410902989837)
- Mirzaei M, Khedri J (2014) Ixodidae ticks in cattle and sheep in Sistan and Baluchestan Province (Iran). *Vet Ital* 50(1):65–68. doi: [10.12834/VetIt.1018.00](https://doi.org/10.12834/VetIt.1018.00)

- Mirzaei A, Rouhani S, Taherkhani H, Farahmand M et al (2011) Isolation and detection of *Leishmania* species among naturally infected *Rhombomys opimus*, a reservoir host of zoonotic cutaneous leishmaniasis in Turkemen Sahara, North East of Iran. *Exp Parasitol* 129:375–380. doi:[10.1016/j.exppara.2011.08.020](https://doi.org/10.1016/j.exppara.2011.08.020)
- Mishra AC, Kulkarni SM (1974) *Polyplax Indica*, sp. nov. (Anoplura: Hoplopleuridae), parasitizing *Golunda ellioti* in India. *Orient Insects* 8(1):89–94. doi:[10.1080/00305316.1974.10434447](https://doi.org/10.1080/00305316.1974.10434447)
- Mishra AC, Bhat HR, Kulkarni SM (1972) *Hoplopleura ramgarh* sp.nov. and *Hoplopleura sinharh* sp.nov. (Anoplura: Hoplopleuridae), parasitizing *Mus* spp. (Rodentia: Muridae) in India. *Parasitology* 65:11–21
- Moemenbellah-Fard MD, Banafshi O, Rafinejad J, Ashraf H (2009) Tick-borne relapsing fever in a new highland endemic focus of western Iran. *Ann Trop Med Parasitol* 103(6):529–537. doi:[10.1179/136485909X451852](https://doi.org/10.1179/136485909X451852)
- Moemenbellah-Fard MD, Shahriari B, Azizi K, Fakoorziba MR et al (2014) Faunal distribution of fleas and their blood-feeding preferences using enzyme-linked immunosorbent assays from farm animals and human shelters in a new rural region of southern Iran. *J Parasit Dis*. doi:[10.1007/s12639-014-0471-1](https://doi.org/10.1007/s12639-014-0471-1)
- Mohammadi M, Ghavami MB, Mohammadi J (2013) Study of phenotypic variability and borrelia infectivity in different populations of *Ornithodoros Tholozani* in Zanjan Province. *J Zanjan Uni Med Sci* 21(87):83–93 (In Persian)
- Mohebbali M, Rezai H, Farahnak A, Kanani Nootash A (1997) A survey on parasitic fauna (Helminths and Ectoparasites) of the rodents in Meshkinshahr district North West Iran. *J Fac Vet Med Univ Tehran* 25(3):23–25 (In Persian)
- Mooring MS, Hart BL (1995) Differential grooming rate and tick load of territorial male and female impala, *Aepyceros melampus*. *Behav Ecol* 6(1):94–101
- Moradi A, Telmadarraiy Z, Moradi A (2009) The survey on infestation rate of sheep to ticks and their distribution in Bahar county. *J Vet* 83:26–28 (In Persian)
- Morand S, Bellocq JGD, Stanko M, Miklisova D (2004) Is sex-biased ectoparasitism related to sexual size dimorphism in small mammals of Central Europe? *Parasitology* 129:505–510. doi:[10.1017/S0031182004005840](https://doi.org/10.1017/S0031182004005840)
- Moravvej G, Hamidi K, Nourani L, Bannazade H (2015) Occurrence of ectoparasitic arthropods (Siphonaptera, Acarina, and Anoplura) on rodents of Khorasan Razavi Province, northeast of Iran. *Asian Pac J Trop Dis* 5(9):716–720. doi:[10.1016/S2222-1808\(15\)60919-7](https://doi.org/10.1016/S2222-1808(15)60919-7)
- Moravvej G, Hamidi K, Nourani L (2016) Relationship between the sex and age of *Mus musculus* (Rodentia: Muridae) with ectoparasites prevalence in northeast of Iran. *Persian J Acarol* 5(1):51–62
- Motevalli-Haghi F, Shirzad G, Mahdi S, Mohammad-Mahdi S, Behzad P (2000) The survey on rodental ectoparasites in Sari and its central areas (1997–1998). *J Mazandaran Uni Med Sci* 10(27):1–7 (In Persian)
- Mougeot F, Redpath SM, Piertyny SB (2006) Elevated spring testosterone increase parasite intensity in male red grouse. *Behav Ecol* 17(1):117–125. doi:[10.1093/beheco/arj005](https://doi.org/10.1093/beheco/arj005)
- Mullens BA, Hinkle NC, Robinson LJ, Szijj CE (2001) Dispersal of northern fowl mites, *Ornithonyssus sylviarum* among hens in experimental poultry house. *J Appl Poultry Res* 10:60–64
- Nabian S, Rahbari S, Shayan P, Haddadzadeh HR (2007) Current status of tick fauna in North of Iran. *Iran J Parasitol* 2(1):12–17
- Naddaf SR, Ghazinezhad B, Bahramali G, Cutler SJ (2012) Phylogenetic analysis of the spirochete *Borrelia microti*, a potential agent of relapsing fever in Iran. *J Clin Microbiol* 50(9):2873–2876
- Naddaf SR, Ghazinezhad B, Sedaghat MM, Masoumi-Asl H, Cutler SJ (2015) Tickborne Relapsing Fever in Southern Iran, 2011–2013. *Emerg Infect Dis*. doi:[10.3201/eid2106.141715](https://doi.org/10.3201/eid2106.141715)
- Nasiri A, Telmadarraiy Z, Vatandoost H et al (2010) Tick infestation rate of sheep and their distribution in Abdanan County, Ilam Province, Iran, 2007–2008. *Iran J Arthropod-Borne Dis* 4(2):56–60
- Nateghpour M, Akhavan AA, Hanafi-Bojd AA, Telmadarraiy Z et al (2013) Wild rodents and their ectoparasites in Baluchistan area, southeast of Iran. *Trop Biomed* 30(1):72–77
- Nekouie H, Razavi MR, Seyedipour GH (2003) Investigation of *Yersinia Pestis* in *Xenopsylla astia*. *Southeast Asian J Trop Med Public Health* 34(Suppl 2):158–161
- Nieberding CM, Olivieri I (2006) Parasites: proxies for host genealogy and ecology? *Trends Ecol Evol* 22(3):156–165. doi:[10.1016/j.tree.2006.11.012](https://doi.org/10.1016/j.tree.2006.11.012)
- Obiegala A, Oltersdorf C, Silaghi C et al (2016) Rickettsia spp. in small mammals and their parasitizing ectoparasites from Saxony, Germany. *Vet Parasitol* 5:19–24. doi:[10.1016/j.vprsr.2016.08.008](https://doi.org/10.1016/j.vprsr.2016.08.008)
- O'Connor JP, Sleeman DP, Butler FT (2005) A review of the Irish Anoplura (Insecta: Phthiraptera). *Ir Nat J* 28(2):62–67
- Oldham JN (1931) On the Arthropod Intermediate Hosts of *Hymenolepis diminuta* (Rudolphi 1819). *J Helminthol* IX 1:21–28
- Oliveira-Sequeira TCG, Oliveira MCS, Araujo JP Jr, Amarante AFT (2005) PCR-based detection of *Babesia bovis* and *Babesia bigemina* in their natural host *Boophilus microplus* and cattle. *Int J Parasitol* 35:105–111. doi:[10.1016/j.ijpara.2004.09.002](https://doi.org/10.1016/j.ijpara.2004.09.002)
- Owen JP, Delany ME, Cardona CJ et al (2009) Host inflammatory response governs fitness in an avian ectoparasite, the northern fowl mite (*Ornithonyssus sylviarum*). *Int J Parasitol* 39:789–799. doi:[10.1016/j.ijpara.2008.12.008](https://doi.org/10.1016/j.ijpara.2008.12.008)
- Ozbeka HH (2015) An unusual new species of Pachylaelaps (Acari: Pachylaelapidae) from Turkey. *Int J Acarol* 41(1):67–70. doi:[10.1080/01647954.2014.988645](https://doi.org/10.1080/01647954.2014.988645)
- Pakdad K, Ahmadi NA, Aminalroaya R et al (2012) A study on rodent ectoparasites in the North district of Tehran, Iran during 2007–2009. *J Paramed Sci (JPS)* 3(1):42–46
- Palyvos NE, Athanassiou CG, Kavallieratos NG (2006) Acaricidal effect of a diatomaceous earth formulation against *Tyrophagus putrescentiae* (Astigmata: Acaridae) and its predator *Cheyletus malaccensis* (Prostigmata: Cheyletidae) in four grain commodities. *J Econ Entomol* 99(1):229–236. doi:[10.1603/0022-0493\(2006\)099\[0229:AEOADE\]2.0.CO;2](https://doi.org/10.1603/0022-0493(2006)099[0229:AEOADE]2.0.CO;2)
- Parola P, Paddock CD, Raoult D (2005) Tick-borne rickettsioses around the world: emerging diseases challenging old concepts. *Clin Microbiol Rev* 18(4):719–756. doi:[10.1128/CMR](https://doi.org/10.1128/CMR)
- Parshad VR (1999) Rodent control in India. *Integr Pest Manag Rev* 4(2):97–126. doi:[10.1023/A:1009622109901](https://doi.org/10.1023/A:1009622109901)
- Pekar S, Hubert J (2008) Assessing biological control of *Acarus siro* by *Cheyletus malaccensis* under laboratory conditions: effect of temperatures and prey density. *J Stored Prod Res* 44:335–340. doi:[10.1016/j.jspr.2008.02.011](https://doi.org/10.1016/j.jspr.2008.02.011)
- Peng P-Y, Guo X-G, Song W-Y (2015) Analysis of ectoparasites (chigger mites, gamasid mites, fleas and sucking lice) of the Yunnan red-backed vole (*Eothenomys miletus*) sampled throughout its range in southwest China. *Med Vet Entomol* 29:403–415. doi:[10.1111/mve.12134](https://doi.org/10.1111/mve.12134)
- Perez-Orella C, Schulte-Hostedde AI (2005) Effects of sex and body size on ectoparasite loads in the northern flying squirrel (*Glaucomys sabrinus*). *Can J Zool* 83:1381–1385. doi:[10.1139/Z05-126](https://doi.org/10.1139/Z05-126)
- Pourhossein B, Esmaeili S, Gyuranecz M, Mostafavi E (2015) Tularemia and plague survey in rodents in an earthquake zone in southeastern Iran. *Epidemiol Health* 37:e2015050
- Pratt HD (1963) Mites of public health importance and their control, training guide-insect control series. U.S. Department of health, Education, and Welfare Public Health Service, Communicable Disease Center, Atlanta, Georgia

- Pratt HD, Karp H (1953) Notes on the Rat Lice *Polyplax spinulosa* (Burmeister) and *Hoplopleura oenomydis* Ferris. *J Parasitol* 39(5):495–504
- Rafinejad J, Shemshad K, Banafshi O (2012) Epidemiological study on tick-borne (Acari: Argasidae) relapsing fever in Kurdistan Province, Iran, 2000–2004. *Florida Entomol* 95(3):758–763. doi: [10.1653/024.095.0327](https://doi.org/10.1653/024.095.0327)
- Rafinejad J, Piazak N, Dehghan A, Shemshad Kh, Basseri HR (2013) Effect of some environmental parameters on fleas density in human and animal shelters. *Am J Res Commun*. www.usa-journals.com, ISSN: 2325-4076
- Rahbari S (1995) Studies on some ecological aspects of Tick West Azarbidjan, Iran. *J Appl Anim Res* 7:189–194
- Rahbari S, Nabian S, Shayan P, Haddadzadeh HR (2007) Status of *Haemaphysalis* tick infestation in domestic ruminants in Iran. *Korean J Parasitol* 45(2):129–132
- Rahdar M, Vazirianzadeh B (2009) A case report of tropical rat mite infestation *Ornithonyssus bacoti* (Dermanyssidae: Acarina) in Ahvaz, SW, Iran. *Jundishapur J Microbiol* 2(2):78–80
- Rahdar M, Vazirianzadeh B, Roiantan ES, Amraei K (2015) Identification of collected ectoparasites of rodents in the west of Khuzestan Province (Ahvaz and Hovizeh), southwest of Iran. *Asian Pac J Trop Dis* 5(8):627–631. doi: [10.1016/S2222-1808\(15\)60902-1](https://doi.org/10.1016/S2222-1808(15)60902-1)
- Randolph SE, Rogers DJ (2007) Ecology of tick-borne disease and the role of climate. In: Ergonul O, Whitehouse CA (eds) *Crimean-congo hemorrhagic fever*. Springer, pp 167–186
- Ranjbar-Bahadori Sh (2003) Study of species diversity of animal ticks in Garmsar. *Vet Med Univ Tehran* 58(1):11–14 (**In Persian**)
- Raoult D, Birtles RJ, Montoya M et al (1999) Survey of three bacterial louse-associated diseases among rural Andean communities in Peru: prevalence of epidemic typhus, trench fever, and relapsing fever. *Clin Infect Dis* 29:434–436
- Rassi Y, Javadian E, Amin M et al (2006) *Meriones libycus* is the main reservoir zoonotic cutaneous leishmaniasis in south Islamic Republic of Iran. *East Mediterr Health J* 12:474–477
- Ratovonjato J, Rajerison M, Rahelinirina S, Boyer S (2014) *Yersinia pestis* in pulex irritans fleas during plague outbreak, madagascar. *Emerg Infect Dis* 20(8):1414–1415. doi: [10.3201/eid2008.130629](https://doi.org/10.3201/eid2008.130629)
- Reeves WK, Cobb KD (2005) Ectoparasites of House Mice (*Mus musculus*) from Pet Stores in South Carolina, USA. *Comp Parasitol* 72(2):193–195. doi: [10.1654/4178](https://doi.org/10.1654/4178)
- Reeves WK, Loftis AD, Szumlas DE et al (2007) Rickettsial pathogens in the tropical rat mite *Ornithonyssus bacoti* (Acari: Macronyssidae) from Egyptian rats (*Rattus* spp.). *Exp Appl Acarol* 41:101–107. doi: [10.1007/s10493-006-9040-3](https://doi.org/10.1007/s10493-006-9040-3)
- Renapurkar DM (1988) Distribution and susceptibility of *Xenopsylla astia* to DDT in Maharashtra State, India. *Int J Trop Insect Sci* 9(3):377–380
- Renvoise A, Wount JW, Schroeff JGVD et al (2012) A case of Rickettsialpox in Northern Europe. *Int J Infect Dis* 16:e221–e222. doi: [10.1016/j.ijid.2011.11.009](https://doi.org/10.1016/j.ijid.2011.11.009)
- Rosario Robles MD, Kinsella JM, Galliari C, Navone GT (2016) New host, geographic records, and histopathologic studies of *Angiostrongylus* spp (Nematoda: Angiostrongylidae) in rodents from Argentina with updated summary of records from rodent hosts and host specificity assessment. *Mem Inst Oswaldo Cruz Rio de Janeiro* 111(3):181–191. doi: [10.1590/0074-02760150371](https://doi.org/10.1590/0074-02760150371)
- Rosen S, Yeruham Y, Braverman Y (2002) Dermatitis in humans associated with the mite *Pyemotes tritici*, *Dermanyssus gallinae*, *Ornithonyssus bacoti* and *Androlaelaps casalis* in Israel. *Med Vet Entomol* 16:442–444
- Roy L, Chauve CM (2007) Historical review of the genus *Dermanyssus Duges*, 1834 (Acari: Mesostigmata: Dermanyssidae). *Parasite* 14:87–100. doi: [10.1051/paraite/2007142087](https://doi.org/10.1051/paraite/2007142087)
- Roy L, Chauve C (2010) The genus *Dermanyssus* (Mesostigmata: Dermanyssidae): history and species characterization. In: Sabelis MW, Bruin J (eds) *Trends in acarology: proceedings of the 12th international congress*, doi: [10.1007/978-90-481-9837-5_8](https://doi.org/10.1007/978-90-481-9837-5_8)
- Rust MK (1997) The biology, ecology, and management of the Cat flea. *Annu Rev Entomol* 42:451–473
- Salari-Lak Sh, Vatandoost H, Telmadarraiy Z, Mahdi RE, Kia EB (2008) Seasonal activity of ticks and their importance in tick-borne infectious diseases in West Azerbaijan, Iran. *Iran J Arthropod-Borne Dis* 2(2):28–34
- Salim-abadi Y, Telmadarraiy Z, Vatandoost H (2010) Hard ticks on domestic ruminants and their seasonal population dynamics in Yazd Province, Iran. *Iran J Arthropod-Borne Dis* 4(1):66–71
- Samia NI, Kausrud KL, Heesterbeek H et al (2011) Dynamics of the plague-wildlife-human system in Central Asia are controlled by two epidemiological thresholds. *PNAS* 108(35):14527–14532
- Sanchez-Montes S, Guzman-Cornejo C, Leon-Paniagua L, Rivas G (2013) A checklist of sucking lice (Insecta: Phthiraptera: Anoplura) associated with Mexican wild mammals, including geographical records and a host-parasite list. *Zootaxa* 3722(2):183–203. doi: [10.11646/zootaxa.3722.2.4](https://doi.org/10.11646/zootaxa.3722.2.4)
- Sanchez-Ramos I, Castanera P (2001) Acaricidal activity of natural monoterpenes on *Tyrophagus putrescentiae* (Schrank), a mite of stored food. *J Stored Prod Res* 37:93–101
- Sedaghat MM, Salahi-Moghaddam A (2010) Mapping the distribution of the important rodents reservoir in Iran. *J Aja Univ Med Sci (JAUMS)* 8(3) (**In Persian**)
- Sedaghat MM, Salahi-Moghaddam A (2010b) Mapping the distribution of the important rodents reservoir in Iran. *JAUMS* 8(3):210–223 (**In Persian**)
- Sekeyova Z, Fournier PE, Rehacek J, Raoult D (2000) Characterization of a new spotted fever group rickettsia detected in *Ixodes ricinus* (Acari: Ixodidae) Collected in Slovakia. *J Med Entomol* 37(5):707–713. doi: [10.1603/0022-2585-37.5.707](https://doi.org/10.1603/0022-2585-37.5.707)
- Seyf A (1989) Iran and the Great Plague, 1830–1831. *Maisonneuve and Larose. Studia Islamica* 69:151–165. <http://www.jstor.org/stable/1596071>
- Shayan A, Rafinejad J (2006) Arthropod parasites of Rodents in Khorram Abbad district, Lorestan Province of Iran. *Iran J Publ Health* 35(3):70–76
- Shinozaki Y, Yoshizawa K, Murata K, Shiibashi T (2004) The first record of sucking louse, *Neohaematopinus collosciuri*, infesting Pallas squirrels in Japan. *J Vet Med Sci* 66(3):333–335
- Shiravi AH, Ardeshir F, Jabbarpour S, Hojati V (2013) The study of mites fauna in the nests and bodies of the rodents in wheat stores in Tehran Province, Iran. *Iran J Biol* 26(3):314–325 (**In Persian**)
- Shirazi Sh, Bahadori F, Mostafaei T-S, Ronaghi H (2013) First report of *Polyplax* sp. in a Persian Squirrel (*Sciurus anomalus*) in Tabriz, Northwest of Iran. *Turkiye Parazitoloj Derg* 37:299–301. doi: [10.5152/tpd.2013.3085](https://doi.org/10.5152/tpd.2013.3085)
- Shyamal B, Sohan L, Veena M et al (2011) Detection of enzootic Plague foci in peninsular India. *J Commun Dis* 43(3):169–176
- Sjostedt A (2007) Tularemia: history, epidemiology, pathogen physiology, and clinical manifestations. *Ann NY Acad Sci* 1105:1–29. doi: [10.1196/annals.1409.009](https://doi.org/10.1196/annals.1409.009)
- Skotarczak B, Cichocka A (2001) Isolation and amplification by polymerase chain reaction DNA of *Babesia microti* and *Babesia divergens* in ticks in Poland. *Ann Agric Environ Med* 8:187–189
- Small E (1989) Systematics of biological systematics (or, Taxonomy of Taxonomy). *JSTOR Taxon* 38(3):335–356. doi: [10.2307/1222265](https://doi.org/10.2307/1222265)
- Smit FGAM (1955) Three subspecies of *Ctenophthalmus congener* Rothschild from the Far east (Siphonaptera: Hystrichopsyllidae). *Entomol Soc Lond B Taxon* 24(3–4):69–75. doi: [10.1111/j.1365-3113.1955.tb01478.x](https://doi.org/10.1111/j.1365-3113.1955.tb01478.x)
- Solomon ME (1962) Ecology of the flour mite, *Acarus siro* L. (= *Tyroglyphus farfnue* DeG.) *Ann. appl. BWL* 50:178–184
- Sumangali K, Rajapakse RPVJ, Rajakaruna RS (2012) Urban rodents as potential reservoirs of zoonoses: a parasitic survey in two

- selected areas in Kandy district. *Ceylon J Sci* 41(1):71–77. doi: [10.4038/cjsbs.v41i1.4539](https://doi.org/10.4038/cjsbs.v41i1.4539)
- Tajedin L, Rassi Y, Oshaghi MA, Telmadarraiy Z et al (2009) Study on ectoparasites of *Rhombomys opimus*, the main reservoir of zoonotic cutaneous leishmaniasis in endemic foci in Iran. *Iran J Arthropod-Borne Dis* 3(1):41–45
- Telmadarraiy Z, Bahrami A, Vatandoost H (2004) A survey on fauna of ticks in West Azerbaijan Province, Iran. *Iran J Publ Health* 33(4):65–69
- Telmadarraiy Z, Vatandoost H, Mohammadi S et al (2007) Determination of rodent ectoparasite fauna in Sarpole-Zahab District, Kermanshah Province, Iran, 2004–2005. *Iran J Arthropod-Borne Dis* 1(1):58–62
- Telmadarraiy Z, Chinikar S, Vatandoost H et al (2015) Vectors of Crimean Congo hemorrhagic fever virus in Iran. *J Arthropod-Borne Dis* 9(2):137–147
- Tschirren B, Bischoff LL, Saladin V, Richner H et al (2007) Host condition and host immunity affect parasite fitness in a bird-ectoparasite system. *Funct Ecol* 21:372–378. doi: [10.1111/j.1365-2435.2007.01235.x](https://doi.org/10.1111/j.1365-2435.2007.01235.x)
- Valiente Moro C, Chauve C, Zenner L (2005) Vectorial role of some Dermansoid mites (Acari, Mesostigmata, Dermansoidea). *Parasite* 12(2):99–109. doi: [10.1051/parasite/200512209](https://doi.org/10.1051/parasite/200512209)
- Vazquez MAM, Klompen H (2009) New species of new world opilioacaridae (Acari: Parasitiformes) with the description of a new genus from the Caribbean region. *Zootaxa* 2061:23–44
- Vazquez MAM, Klompen H (2015) The family Opilioacaridae (Parasitiformes: Opilioacarida) in Mexico, description of two new species and notes on biology and geographical distribution. *Zootaxa* 3957(5):535–552
- Walter DE, Proctor HC (2013) *Mites: Ecology, Evolution & Behaviour: Life at a Microscale*. Springer Dordrecht Heidelberg New York London. doi: [10.1007/978-94-007-7164-2](https://doi.org/10.1007/978-94-007-7164-2)
- Waterman JM, Macklin GF, Enright C (2014) Sex-biased parasitism in Richardson's ground squirrels (*Urocitellus richardsonii*) depends on the parasite examined. *Can J Zool* 92:73–79. doi: [10.1139/cjz-2013-0151](https://doi.org/10.1139/cjz-2013-0151)
- Wei L, Wang X, Wang CH, He H (2010) A survey of ectoparasites from wild rodents and *Anourosorex squamipes* in Sichuan Province, Southwest China. *J Ecol Nat Environ* 2(8):160–166
- Wells K, Smales LR, Kalko EKV, Pfeiffer M (2007) Impact of rain-forest logging on helminth assemblages in small mammals (Muridae, Tupaiidae) from Borneo. *J Trop Ecol* 23(1):35–43. doi: [10.1017/S0266467406003804](https://doi.org/10.1017/S0266467406003804)
- Whiting MF, Whiting AS, Hastriter MW, Dittmar K (2008) A molecular phylogeny of fleas (Insecta: Siphonaptera): origins. *Cladistics* 24:677–707. doi: [10.1111/j.1096-0031.2008.00211.x](https://doi.org/10.1111/j.1096-0031.2008.00211.x)
- WHO (1988) *Vector-borne Disease Control in Humans through Rice Agroecosystem Management*. International Rice Research Institute in collaboration with The WHO/FAO/UNEP Panel of Experts on Environmental Management for Vector Control
- Wilson N (1967) Acarina: mesostigmata dermanyssoidea, laelapidae, spinturnicidae parasitic on vertebrates. *Insects of Micronesia* 3(5):133–148
- Xian-guo G, Zheng-da G, Ti-jun Q et al (2000) Spatial pattern analysis of *Xenopsylla cheopis* (Siphonaptera: Pulicidae) on its dominant rat host, *Rattus flavipectus* in the foci of human plague in Yunnan, China. *Entomol Sin* 7(1):47–52
- Yakhchali M, Haji Hasanzadehzarza Sh (2004) Study on some ecological aspects and prevalence of different species of hard ticks (Acarina: Ixodidae) on cattle, buffalo, and sheep in Oshnavieh suburb. *Pajouhesh Sazandegi* 63:30–35 (**In Persian**)
- Yang P, Oshiro S, Warashina W (2009) Ectoparasitic arthropods occurring on *Rattus norvegicus* and *Rattus rattus* collected from two properties on the Island of Oahu, Hawaii (Acarina, Siphonaptera, and Anoplura). *Hawaiian Entomol Soc* 41:53–56
- Yin H, Lu W, Luo J et al (1996) Experiments on the transmission of *Babesia major* and *Babesia bigemina* by *Haemaphysalis punctata*. *Vet Parasitol* 67:89–98
- Yousefi A, Chaechi Nosrati MR, Karimi A, Naisi S (2015) *Leptopsylla taschenbergi taschenbergi* (Siphonaptera: Leptopsyllidae), new flea from Iran. *Asian Pac J Trop Dis* 5(8):606–607. doi: [10.1016/S2222-1808\(15\)60897-0](https://doi.org/10.1016/S2222-1808(15)60897-0)
- Yu X, Tesh RB (2014) The role of mites in the transmission and maintenance of hantaan virus (Hantavirus: Bunyaviridae). *J Infect Dis* 210:1693–1699. doi: [10.1093/infdis/jiu336](https://doi.org/10.1093/infdis/jiu336)
- Zarei R, Darvish J, Esmaeili HR, Tarahomi M (2010) A biosystematic survey of Shiraz (Central Part) rodents. *Iran J Biol* 23(4):573–583 (**In Persian**)
- Zarei Z, Mohebbali M, Heidari Z et al (2017) Wild rodent ectoparasites collected from Northwestern Iran. *J Arthropod-Borne Dis* 11(1):36–41
- Zargar A, Maurin M, Mostafavi E (2015) Tularemia, a re-emerging infectious disease in Iran and neighboring countries. *Epidemiol Health* 37:1–6. doi: [10.4178/epih/e2015011](https://doi.org/10.4178/epih/e2015011)
- Zintl A, Mulcahy G, Skerrett HE, Taylor SM, Gray JS (2003) *Babesia divergens*, a bovine blood parasite of veterinary and zoonotic importance. *Clin Microbiol Rev* 16(4):622–636. doi: [10.1128/CMR.16.4.622-636.2003](https://doi.org/10.1128/CMR.16.4.622-636.2003)
- Zuo X-H, Guo X-G, Zhan Y-Z, Wu D et al (2011) Host selection and niche differentiation in sucking lice (Insecta: Anoplura) among small mammals in southwestern China. *Parasitol Res* 108:1243–1251. doi: [10.1007/s00436-010-2173-7](https://doi.org/10.1007/s00436-010-2173-7)