



Published in final edited form as:

J Cell Biochem. 2018 April ; 119(4): 3586–3597. doi:10.1002/jcb.26552.

Mechanoactivation of the angiotensin II type 1 receptor induces β -arrestin-biased signaling through $G\alpha_i$ coupling

Jialu Wang^{1,2}, Kenji Hanada², Clarice Gareri², and Howard A. Rockman^{1,2,3}

¹Department of Cell Biology, Duke University Medical Center, Durham, North Carolina

²Department of Medicine, Duke University Medical Center, Durham, North Carolina

³Department of Molecular Genetics and Microbiology, Duke University Medical Center, Durham, North Carolina

Abstract

Ligand activation of the angiotensin II type 1 receptor (AT1R), a member of the G protein-coupled receptor (GPCR) family, stimulates intracellular signaling to mediate a variety of physiological responses. The AT1R is also known to be a mechanical sensor. When activated by mechanical stretch, the AT1R can signal via the multifunctional adaptor protein β -arrestin, rather than through classical heterotrimeric G protein pathways. To date, the AT1R conformation induced by membrane stretch in the absence of ligand was thought to be the same as that induced by β -arrestin-biased agonists, which selectively engage β -arrestin thereby preventing G protein coupling. Here, we show that in contrast to the β -arrestin-biased agonists TRV120023 and TRV120026, membrane stretch uniquely promotes the coupling of the inhibitory G protein ($G\alpha_i$) to the AT1R to transduce signaling. Stretch-triggered AT1R- $G\alpha_i$ coupling is required for the recruitment of β -arrestin2 and activation of downstream signaling pathways, such as EGFR transactivation and ERK phosphorylation. Our findings demonstrate additional complexity in the mechanism of receptor bias in which the recruitment of $G\alpha_i$ is required for allosteric mechanoactivation of the AT1R-induced β -arrestin-biased signaling.

Keywords

β -arrestin; angiotensin II; G protein; mechanotransduction

Correspondence: Dr. Howard A. Rockman, Department of Medicine, Duke University Medical Center, DUMC 3104, 226 CARL Building, Durham, NC 27710. h.rockman@duke.edu.

Current address for Kenji Hanada is Department of Cardiology, Hirosaki University Graduate School of Medicine, Hirosaki, Aomori, Japan.

ORCID

Howard A. Rockman <http://orcid.org/0000-0003-2921-1584>

Jialu Wang and Kenji Hanada authors contributed equally.

SUPPORTING INFORMATION

Additional Supporting Information may be found online in the supporting information tab for this article.

CONFLICT OF INTEREST

The authors declare that they have no conflicts of interest with the contents of this article.

1 | INTRODUCTION

The angiotensin II type 1 receptor (AT1R), a member of the G protein-coupled receptor (GPCR) superfamily, is a well-known regulator of blood pressure and cardiac function.¹ In response to the balanced agonist angiotensin II (AngII), AT1Rs couple to the heterotrimeric G-protein triggering the dissociation of $G\alpha_q$ from $G\beta$, which subsequently activates the signaling effector phospholipase C (PLC) to promote generation of the second messengers inositol-1,4,5-trisphosphate (IP3) and diacylglycerol (DAG).¹ This G protein-mediated signaling is rapidly turned off by the receptor desensitization, a process that involves phosphorylation of the C-terminal tail of the AT1R mediated by G protein-coupled receptor kinases (GRKs) and recruitment of the multifunctional scaffold protein β -arrestin that impairs further G protein coupling.²⁻⁴ It is now known that β -arrestins also function as signal transducers in their own right, regulating a large network of cellular responses.⁵ Several AT1R ligands, such as [Sar1, Ile4, Ile8]-AngII (SII) and TRV120023, have been shown to selectively activate β -arrestin-mediated pathways and therefore termed as β -arrestin-biased agonists.^{6,7}

Our current understanding for ligand stimulated AT1R signaling is based on a mechanism whereby ligand binding stabilizes distinct receptor conformations allowing them to engage signal transducers such as G proteins or β -arrestin to affect activation of an array of signaling pathways, a concept known as functional selectivity.^{1,6,8-12} Interestingly, activation of AT1Rs can be mechanosensitive, whereby membrane stretch allosterically stabilizes a distinct β -arrestin-biased AT1R conformation triggering ligand-independent β -arrestin-mediated signaling.^{13,14} The precise molecular mechanism by which mechanical stress activates AT1R signaling is not clearly understood.

It is now well appreciated that AT1Rs may couple to multiple signaling transducers with different efficacy and kinetics depending on cellular systems and stimuli.^{15,16} For example, in proximal tubule cells, the AngII-activated AT1R recruits $G\alpha_i$ leading to the inhibition of adenylate cyclase activity.¹⁷ In cardiac fibroblasts, AngII activates the AT1R-mediated ERK signaling through the $G\beta\gamma$ subunits of $G\alpha_i$ ¹⁸ and the weak β -arrestin-biased agonist [Sar¹, Ile⁴, Ile⁸]-angiotensin II (SII) can promote AT1R coupling to both $G\alpha_q$ and $G\alpha_i$.¹⁵

We, therefore, set out to determine whether $G\alpha_i$ plays a role as a signal transducer for the AT1R under conditions of mechanoactivation. Indeed, we show that membrane stretch promotes the recruitment of $G\alpha_i$ to the AT1R, which is required for the activation of β -arrestin-biased signaling as measured by epidermal growth factor receptor (EGFR) internalization and extracellular signal-regulated kinases (ERK) phosphorylation. In contrast, $G\alpha_i$ is not required for signaling induced by the balanced agonist AngII or the β -arrestin-biased agonist TRV120023 and TRV120026. Notably, our findings demonstrate mechanistic heterogeneity for AT1R-mediated β -arrestin-biased signaling induced by mechanoactivation versus biased ligands and support a concept that membrane stretch may stabilize a biased AT1R conformation that is distinct from that stabilized by a β -arrestin-biased ligand.

2 | MATERIALS AND METHODS

2.1 | Cell culture

HEK 293 cells stably expressing hemagglutinin (HA)-tagged AT1Rs were cultured in minimum essential medium (MEM) supplemented with 10% FBS, 100 U/mL penicillin and 100 µg/mL streptomycin (ThermoFisher Scientific, Waltham, MA), and 100 µg/mL of zeocin (Invitrogen, Carlsbad, CA) at 37°C in a humidified environment with 5% CO₂ as previously described.¹³ HEK 293 cells stably expressing cyan fluorescent protein (CFP)-tagged AT1R and yellow fluorescent protein (YFP)-tagged β-arrestin2 were cultured in MEM supplemented with 10% FBS, 100 U/mL penicillin and 100 µg/mL streptomycin, 300 µg/mL of zeocin, and 150 µg/mL of Geneticin (ThermoFisher Scientific).

2.2 | Generation of Gα_i knockout cell line with CRISPR-Cas9 gene editing

Plasmids carrying *S. pyogenes* Cas9 (SpCas9) next to a cloning site for guide RNA (gRNA) with EGFP (pSpCas9 (BB)-2A-GFP, Addgene 48138) were obtained from Addgene (deposited by the laboratory of Dr. F. Zhang¹⁹). Designing of the guide RNAs for Gα_i and cloning the guide RNAs into the Cas9 plasmids were performed as previously described.¹⁹ Gα_{i1} was targeted using guide sequence oligos (top: CACCGCGCCGTCCTCACGGA GGTG; bottom: AAACCAACCTCCGTGAGGACGGCGC), Gα_{i2} was targeted using guide sequence oligos (top: CACCGAG ACAACCGCCCGGTACTGC, bottom: AAACGCAGTACCG GGCGGTTGTCTC), and Gα_{i3} was targeted using guide sequence oligos (top: CACGGGACGGCTAAAGATTGA CTT; bottom: AAACAAGTCAATCTTTAGCCGTCCC). The guide sequence oligos were cloned into pSpCas9 (BB)-2A-GFP. Plasmids targeting the three Gα_i subtypes were co-transfected into HEK293 cells. GFP positive cells were selected by fluorescence-activated cell sorting, diluted for growth and single cell colonies were obtained. The Gα_i knockout were confirmed by Western blot.

2.3 | Proximity ligation assay

Proximity ligation assay (PLA) was performed following manufacture's protocol (Sigma-Aldrich, St. Louis, MO) and previously published paper.²⁰ Briefly, HEK293 cells stably expressing HA-tagged AT1R were plated on collagen-coated glass bottom cell culture dishes. After overnight incubation in serum-free medium with vehicle or 200 ng/mL pertussis toxin (List Biological Labs, Campbell, CA), cells were stimulated with 1 µM AngII (Sigma-Aldrich), 1 µM TRV120023 (synthesized by GenScript, Inc., Piscataway, NJ), or osmotic stretch (by adding same volume of double-distilled H₂O to the medium as previously described¹⁴) for 10 min at 37°C in a humidified environment with 5% CO₂, and fixed with 4% PFA. Fixed cells were permeabilized with 0.2% Triton-X100 for 10 min, blocked with manufacture's blocking solution for 30 min, and incubated overnight with anti-Gα_i mouse monoclonal antibody (NewEast Biosciences, King of Prussia, PA) and anti-HA rabbit antibody (Sigma-Aldrich) at 4°C. The next day, cells were subsequently incubated with Duolink In Situ PLA Probe Anti-Mouse MINUS and Anti-Rabbit PLUS for 1 h, ligation buffer for 30 min, amplification buffer (Duolink In Situ Detection Reagents Green) for 100 min, and mounted with Duolink In Situ Mounting Medium with DAPI. Images were recorded using the Zeiss Axio Observer Z1 confocal microscope. Quantification of PLA

signal representing the AT1R-G α_i interaction was performed by averaging the intensity of the green fluorescence per cell and subtracting the non-specific background. Background signal was determined as green fluorescence intensity per cell in wildtype HEK293 cells without HA-tagged AT1R overexpression.

2.4 | Co-immunoprecipitation

HEK293 cells stably expressing HA-tagged AT1Rs were transfected with G α_{i2} for the detection of AT1R-G α_i co-IP, and not transfected for the AT1R-G α_q co-IP experiments. Cells were incubated in serum-free medium with vehicle or 200 ng/mL PTX for overnight, and then stimulated with 1 μ M AngII, 10 μ M TRV120023 or osmotic stretch for 10 min. After stimulation, cells were lysed in 0.5% MNG lysis buffer (20 mM HEPES, 150 mM NaCl, 0.5% Lauryl Maltose Neopentyl Glycol, 2 mM sodium orthovanadate, 1 mM PMSF, 10 mM sodium fluoride, 10 μ g/ml aprotinin, 5 μ g/mL leupeptin and phosphatase inhibitors) with gentle agitation at 4°C for 1 h, and then centrifuged for 15 min at 13 200 rpm. 0.5–1 mg of protein was incubated with 30 μ L of anti-HA magnetic beads (Pierce, Waltham, MA) for overnight. Immunoprecipitates were then separated by SDS-PAGE, transferred to PVDF membrane (Bio-Rad, Hercules, CA) and subjected to immunoblotting with various primary antibodies. Immunoblots were detected using enhanced chemiluminescence (Thermo Fisher Scientific) and analyzed with ImageJ software.

2.5 | Confocal microscopy

In the β -arrestin translocation assay, HEK 293 cells stably expressing CFP-tagged AT1R and YFP-tagged β -arrestin2 were transiently transfected with GRK5 using Fugene 6 (Roche, Basel, Switzerland) and then plated in collagen-coated glass bottom cell culture dishes. After overnight serum-starvation with vehicle or 200 ng/mL PTX, cells were stimulated with 1 μ M AngII, 1 μ M or 10 μ M TRV120023, 1 μ M or 10 μ M TRV120026, or osmotic stretch for 10 min, and then fixed with 4% PFA. In the EGFR internalization assay, HEK293 cells stably expressing HA-tagged AT1R were transiently transfected with green fluorescent protein (GFP)-tagged EGFR and GRK5 plasmids, and stimulated and fixed as described above. Cells were visualized using a Zeiss Axio Observer Z1 confocal microscope. Quantification of β -arrestin translocation was performed by counting the percentage of the cells showing intracellular aggregates of β -arrestin. Similarly, quantification of EGFR internalization was performed by counting the percentage of cells showing EGFR internalization.

2.6 | ERK phosphorylation assay

We performed immunoblotting as previously described.¹³ In brief, HEK293 cells stably expressing HA-tagged AT1Rs were plated on collagen-coated six-well dishes at a confluence of 60–70%. Cells were serum-starved for overnight and pretreated with vehicle or inhibitors (PTX: 200 ng/mL for overnight; Ro318425 (Sigma-Aldrich): 1 μ M for 30 min; U73122 (Sigma-Aldrich): 10 μ M for 30 min; AG1478 (Sigma-Aldrich): 1 μ M for 30 min), and then stimulated with 1 μ M AngII, 1 μ M or 10 μ M TRV120023, 1 μ M or 10 μ M TRV120026, or osmotic stretch for 10 min. After stimulation, cells were lysed in 1% NP-40 lysis buffer (20 mM Tris, pH7.4, 137 mM NaCl, 20% glycerol, 1% Nonidet P-40, 2 mM sodium orthovanadate, 1 mM PMSF, 10 mM sodium fluoride, 10 μ g/mL aprotinin, 5 μ g/mL

leupeptin and phosphatase inhibitors) with gentle agitation for 1 h at 4°C and centrifuged for 15 min at 13 200 rpm. Protein concentrations were measured with Bio-Rad protein assay reagent. Proteins were separated by SDS-PAGE, and transferred to PVDF membrane and blocked with 5% skim-milk. Membranes were incubated with phospho ERK1/2 (1:1000 Cell Signaling, Danvers, MA) or total ERK1/2 (1:2000, EMD Millipore, Billerica, MA) antibody overnight with gentle agitation at 4°C, and then Horseradish peroxidase-conjugated secondary antibody for 1 h at room temperature. Protein bands were visualized by ECL with the Syngene Imager and quantified by densitometry using GeneSnap (Syngene, Cambridge, UK) or ImageJ software. In left ventricle (LV) of the mice, tissues were homogenized in NP-40 lysis buffer. Immunoblotting was performed using the protocol described above.

2.7 | Ex-vivo LV balloon stretch

Animal studies were carried out according to approved protocols and animal welfare regulations of Duke University Medical Center's Institutional Review Boards. We performed ex-vivo LV balloon stretch experiment as previously described.¹³ Briefly, 8–12 weeks old, gender matched wild type mice (purchased from Jackson laboratory, Bar Harbor, ME) were pretreated with 25 µg/kg of PTX or vehicle through intraperitoneal (IP) injection at 48 h before experiment. LV balloon was made by stretching a polypropylene membrane into the shape of the LV cavity. Mice were anesthetized with a mixture of ketamine (100 mg/kg) and xylazine (2.5 mg/kg), and heparinized by IP injection. After 10 min, chest was opened and heart was excised. Aorta was cannulated to the needle and balloon was inserted to LV through mitral valve. Heart was retrograde perfused with buffer containing 118 mM NaCl, 4.7 mM KCl, 1.2 mM MgSO₄, 1.2 mM KH₂PO₄, 2.5 mM CaCl₂, 25 mM NaHCO₃, 0.5 mM Na-EDTA, and 5.5 mM glucose with O₂ bubbling through Langendorff apparatus (Hugo Sachs Harvard Apparatus, Holliston, MA) set at 37°C. LV balloon was connected to a Statham P23 Db pressure transducer (Gould Statham Instruments, Westminister, CA) through Polyethylene-50 (PE-50) for pressure monitoring. Intraballoon pressure was recorded continuously with a pressure-recording system (MacLab, Millar Instruments, Houston, TX). Balloon inflation was started after 10 min perfusion, and diastolic intraballoon pressure was kept within 30–50 mmHg with minor adjustment. Control was defined as perfusion without balloon inflation for identical periods of time. After 10 min balloon inflation, heart was removed from the system, and snap frozen in liquid nitrogen after removing atrium and right ventricle. If intraballoon pulse pressure was less than 20 mmHg or heart rate was less than 100 bpm continuously, the heart was excluded from this study.

2.8 | Statistics

Data are expressed as mean ± SEM. Statistical significance was determined with a one- or two-way analysis of variance (ANOVA) with Bonferroni correction for multiple comparisons using GraphPad Prism software. A *P*-value of <0.05 was considered significant.

3 | RESULTS

3.1 | Membrane stretch promotes the recruitment of G α_i to AT1Rs

Previous work has shown that ligand stimulated AT1Rs may transduce signaling through multiple G proteins such as G α_q and G α_i , as well as the multifunctional adaptor protein β -arrestin.¹⁵ It is now appreciated that AT1Rs, in response to biomechanical membrane stretch, also activate signaling but in a G α_q -independent β -arrestin-dependent manner.^{13,14} To determine whether G α_i is involved in this stretch-induced AT1R signaling, we investigated the effect of membrane stretch on AT1R-G α_i coupling. HEK293 cells stably expressing HA-tagged AT1R were stimulated with the balanced agonist AngII, the β -arrestin-biased agonist TRV120023, or hypotonicity-induced membrane stretch, respectively. The recruitment of G α_i to AT1R was then measured with an in situ proximity ligation assay (PLA), in which protein-protein interactions can be directly visualized and quantified by the fluorescence signal. We found that osmotic stretch significantly increased the amount of G α_i recruited to AT1R, shown as the green PLA signal (Figure 1). Pretreatment with pertussis toxin (PTX), a G α_i inhibitor that catalyzes the ADP-ribosylation of G α_i and prevents its coupling to receptors, blocked the stretch-induced G α_i recruitment to AT1Rs. Notably, the balanced agonist AngII and the β -arrestin biased agonist TRV120023 did not alter the PLA signal.

To further demonstrate recruitment of G α_i to the stretch-activated AT1R, we performed co-immunoprecipitation experiments. Consistent with the PLA results, the level of G α_i bound to AT1R was increased by osmotic stretch, but not affected by AngII and TRV120023 (Figures 2A and 2B). In contrast, AngII promoted the recruitment of G α_q to AT1R, while neither TRV120023 nor stretch had any effect (Figures 2C and 2D), which is consistent with previous studies suggesting that TRV120023- and stretch-induced signaling is independent of G α_q . Thus, in contrast to the balanced agonist AngII and the β -arrestin-biased agonist TRV120023, G α_i is only recruited to the mechano activated AT1R.

3.2 | G α_i is required for membrane stretch-induced β -arrestin2 recruitment to AT1Rs

To dissect the role of G α_i in stretch-induced AT1R signaling, we tested the effect of the G α_i inhibitor PTX on β -arrestin2 recruitment using HEK293 cells stably expressing both CFP-tagged AT1R and YFP-tagged β -arrestin2. Due to low abundance in HEK293 cells of G protein-coupled receptor kinase 5 (GRK5), a kinase that mediates AT1R phosphorylation required for osmotic stretch-induced β -arrestin recruitment,¹³ cells were transfected with GRK5. In the absence of stimulation, AT1R localizes on the plasma membrane and β -arrestin2 is evenly distributed in cytoplasm (Figure 3A). When cells were stimulated with AngII, TRV120023, or osmotic stretch, β -arrestin2 translocated from cytoplasm to endocytic vesicles where it co-localized with the internalized AT1R (AngII: $87.0 \pm 1.7\%$, 1 μ M TRV120023: $73.8 \pm 2.1\%$, 10 μ M TRV120023: $73.3 \pm 6.3\%$, osmotic stretch: $36.9 \pm 2.1\%$ of the cells showed β -arrestin2 translocation, respectively, all $P < 0.0001$ compared with unstimulated cells), indicating the recruitment of β -arrestin2 to AT1R. Pretreatment with PTX significantly blocked the osmotic stretch-induced β -arrestin2 recruitment ($8.3 \pm 1.4\%$, $P < 0.0001$ compared with vehicle-pretreated cells); while without any effect on the AngII- and TRV120023-induced response (with PTX, AngII: $84.5 \pm 2.0\%$, 1 μ M TRV120023: $75.8 \pm 2.0\%$, 10 μ M TRV120023: $74.7 \pm 6.8\%$, $P =$ n.s. compared with vehicle) (Figure 3B).

We previously showed that the mechanoactivated AT1R has specificity for β -arrestin2.¹³ To test whether PTX would influence β -arrestin1 translocation to the AT1R, we stably expressed both CFP-tagged AT1Rs and YFP-tagged β -arrestin1 in HEK293 cells and performed recruitment assays. While the recruitment of β -arrestin1 to AT1R was robustly induced by AngII and TRV120023, and was not affected by PTX pretreatment, β -arrestin1 translocation was not observed in cells treated with hypotonic stretch (Supplementary Figure S1).

Taken together, these results indicate a requirement of $G\alpha_i$ for osmotic stretch-induced β -arrestin2 recruitment to the AT1R, and indicates important heterogeneity in AT1R signaling induced by a β -arrestin-biased agonist compared to that by membrane stretch.

3.3 | Mechanoactivated-AT1R induces EGFR transactivation and ERK1/2 phosphorylation in a $G\alpha_i$ -dependent manner

Both ligand stimulation and mechanoactivation of AT1R promotes the shedding of membrane-bound EGF and subsequent transactivation of EGFR in a β -arrestin-dependent manner.^{13,21} To dissect the mechanism of stretch-induced $G\alpha_i$ -dependent AT1R signaling, we tested the effect of PTX on AT1R-mediated EGFR transactivation. We transfected HEK293 cells stably expressing AT1Rs with GFP-tagged EGFRs and monitored EGFR activation by GFP internalization with confocal microscopy. In the absence of stimulation, EGFRs were located on the plasma membrane. Stimulation of AT1R induced EGFR internalization as shown by redistribution into cellular aggregates (AngII: $44.4 \pm 1.3\%$, TRV120023: $12.0 \pm 1.7\%$, or osmotic stretch: $25.6 \pm 2.6\%$ of the cells showed EGFR internalization, AngII: $P < 0.0001$, TRV120023: $P < 0.01$, osmotic stretch: $P < 0.0001$ compared with unstimulated cells) (Figures 4A and 4B). Pretreatment with PTX significantly inhibited EGFR internalization induced by osmotic stretch (with PTX, $9.6 \pm 1.6\%$, $P < 0.0001$ compared with vehicle-pretreated cells), whereas have no effect on AngII or TRV120023 (with PTX, AngII: $42.8 \pm 1.9\%$, TRV120023: $11.2 \pm 1.4\%$, $P = \text{n.s.}$ compared with vehicle) (Figures 4A and 4B). These data indicate that $G\alpha_i$ is required for AT1R-mediated EGFR transactivation promoted by osmotic stretch, but not for the response induced by AngII or the β -arrestin-biased ligand TRV120023.

We next determined the effect of PTX on AT1R-mediated ERK1/2 phosphorylation. In HEK293 cells stably expressing AT1Rs, both AngII and osmotic stretch significantly increased ERK1/2 phosphorylation, whereas TRV120023 showed a mild increase (Figures 4C and 4D). Similar to that observed for EGFR transactivation, PTX blocked the osmotic stretch-induced ERK1/2 signaling ($P < 0.001$ compared to vehicle), but did not alter the AngII response (Figures 4C and 4D).

To determine whether a similar signaling mechanism is involved in vivo, we next tested whether $G\alpha_i$ is necessary to activate mechanical stretch-induced AT1R signaling in mouse heart tissue. A polypropylene membrane balloon was placed in the left ventricle (LV) of a Langendorff-perfused mouse heart and then inflated to apply a mechanical stretch on the heart. We have previously shown that stretch of the heart induces ERK1/2 phosphorylation in a AT1R- and β -arrestin2-dependent manner.¹³ While LV balloon stretch of hearts significantly increased the phosphorylation of ERK1/2 ($P < 0.001$ compared with control),

pretreatment with PTX abolished this response ($P < 0.01$ compared with the vehicle-pretreated hearts), indicating a requirement of $G\alpha_i$ for stretch-induced ERK1/2 signaling in vivo (Figures 4E and 4F).

To determine the relative contribution of $G\alpha_q$ and EGFR transactivation on AT1R signaling, we tested the effect of the protein kinase c (PKC) inhibitor Ro318425, phospholipase C inhibitor U73122 and the EGFR inhibitor AG1478 on ERK1/2 phosphorylation. Both Ro318425 and U73122 blocked AngII-induced ERK1/2 signaling (for both inhibitors, $P < 0.0001$ compared with vehicle), while neither had effect on the osmotic stretch response (Figure 5A–D). In contrast, AG1478 blocked ERK1/2 phosphorylation induced by either AngII ($P < 0.0001$ compared with vehicle) or osmotic stretch ($P < 0.01$ compared with vehicle) (Figures 5E and 5F).

Collectively, these data support a concept that the balanced ligand AngII stimulates AT1R ERK1/2 signaling through both the $G\alpha_q$ and β -arrestin transducers, while mechanoactivated AT1R signaling is mediated through a $G\alpha_i$ -dependent, β -arrestin-dependent mechanism.

3.4 | TRV120026, another β -arrestin-biased agonist, induces β -arrestin-biased AT1R signaling independent of $G\alpha_i$

To determine whether the $G\alpha_i$ dependency for the β -arrestin-biased AT1R signaling is specifically induced by mechanical stress, we tested the effect of the $G\alpha_i$ inhibitor PTX on the AT1R signaling induced by TRV120026, another β -arrestin-biased agonist with very similar potency of β -arrestin recruitment to AT1R.^{7,22} In HEK293 cells stably expressing the CFP-tagged AT1R and YFP-tagged β -arrestin2, TRV120026 significantly induced β -arrestin2 translocation, to the same extent of TRV120023 response (1 μ M TRV120026: $62.0 \pm 2.0\%$, 10 μ M TRV120026: $72.7 \pm 2.4\%$, 10 μ M TRV120023: $72.0 \pm 2.3\%$ of cells showed β -arrestin2 translocation, all $P < 0.0001$ compared with unstimulated cells) (Figures 6A and 6B). Pretreatment of PTX have no effect on TRV120026-induced response (Figures 6A and 6B), suggesting that $G\alpha_i$ is not required for TRV120026-induced β -arrestin2 translocation. Similar to TRV120023, TRV120026 induced mild increase of ERK1/2 phosphorylation, which was not affected by PTX pretreatment (Figures 6C and 6D). Taken together, these data suggest that the β -arrestin-biased agonists TRV120023 and TRV120026 both induce AT1R signaling in a $G\alpha_i$ -independent manner, which is distinct from that induced by osmotic stretch.

3.5 | Depletion of $G\alpha_i$ by CRISPR-Cas9 gene editing blocks osmotic stretch-induced β -arrestin2 recruitment to AT1Rs

To more robustly determine the role of $G\alpha_i$ in osmotic stretch-induced AT1R signaling, we measured β -arrestin2 translocation in a $G\alpha_i$ knockout cell line²³ that is completely deficient of all three subtypes of $G\alpha_i$ generated by CRISPR/Cas9 gene editing (Figure 7A). In wild type HEK293 cells transfected with CFP-tagged AT1R and YFP-tagged β -arrestin2 (together with GRK5), stimulation with AngII, TRV120023, TRV120026, and osmotic stretch induced the translocation of β -arrestin2. In the $G\alpha_i$ knockout cells, the osmotic stretch-induced β -arrestin2 translocation was blocked, whereas the responses induced by other stimuli were not affected (Figures 7B and 7C). These data are consistent with our PTX

experiments and robustly demonstrate that $G\alpha_i$ is required for stretch-induced β -arrestin2 recruitment to the AT1R.

4 | DISCUSSION

In this study, we identify a new molecular mechanism for β -arrestin-biased AT1R signaling. Membrane stretch, which has been shown to activate β -arrestin-biased AT1R signaling without activating $G\alpha_q$, specifically promotes the recruitment of another subtype of G proteins, $G\alpha_i$, to the AT1R (Figure 8). AT1R- $G\alpha_i$ coupling is required for the recruitment of β -arrestin to the receptor and the induction of subsequent β -arrestin-dependent signaling as measured by EGFR internalization and ERK1/2 phosphorylation. In contrast, the $G\alpha_i$ recruitment and the dependence of $G\alpha_i$ for β -arrestin-mediated signaling was not observed for the balanced agonist AngII, or, remarkably, the β -arrestin-biased agonists TRV120023 and TRV120026 (Figure 8). These results indicate mechanistic divergence of AT1R-mediated β -arrestin-biased signaling triggered by the different modes of receptor activation, membrane stretch and β -arrestin-biased ligands in this case.

It is well established that GPCR can engage to multiple G protein subtypes, and the efficacy and kinetics of distinct G protein coupling are regulated by the specificity of ligands or allosteric modulators.¹⁶ Although in our study we did not observe $G\alpha_i$ recruitment to AngII-stimulated AT1Rs, previous studies have shown that AngII may promote AT1R- $G\alpha_i$ coupling to inhibit adenylyl cyclase and to regulate Ca^{2+} channels in certain tissue or cell types.²⁴⁻²⁶ In addition, a recent study reported that SII, an AT1R ligand previously described as a β -arrestin-biased agonist, can also weakly activates $G\alpha_q$ and $G\alpha_i$.¹⁵ Despite the variation of the effect of AngII on $G\alpha_i$ activation, which might due to the difference in experimental systems or assay sensitivity, it is clear that in response to different stimuli AT1Rs can engage distinct transducers to induce cellular signaling.

A full understanding of the diversity of GPCR signaling through distinct transducers activated by different stimuli remains to be elucidated. Besides their role in receptor internalization and desensitization, β -arrestins are also critical transducers for GPCR signaling that involved in a wide diversity of physiological and pathological processes. Interestingly, the G protein signaling and β -arrestin signaling are not mutually exclusive. The signaling mediated by many $G\alpha_i$ -coupled receptors requires both $G\alpha_i$ and β -arrestin.²⁷ Upon the previous understanding, the β -arrestin-mediated signaling of most $G\alpha_s$ or $G\alpha_q$ receptors appears to be independent of G proteins.²⁷ However, our study suggests that certain modes of stimulation, mechanoactivation in this case, may switch the primarily $G\alpha_q$ -coupled receptors to coupling to $G\alpha_i$. The now $G\alpha_i$ -coupled receptors then activate downstream β -arrestin signaling.

Considerable structural information is now available for GPCRs in complex with ligands or signal transducers and support the concept that receptors adopt distinct conformations to selectively regulate different arrays of downstream signaling.^{28,29} Biophysical analysis of mechanical stretch-activated AT1R suggest that the AT1R undergoes anticlockwise rotation and a shift of transmembrane domain 7 into the ligand-binding pocket, whereas AT1R blocker candesartan stabilized the receptor in the inactive conformation and prevented

stretch induced conformational change.³⁰ Our previous work suggested that membrane stretch induces a β -arrestin-biased conformational state of AT1Rs and an active β -arrestin conformation indistinguishable from that induced by the biased agonist TRV120023.^{13,14} However, we now show that membrane stretch induces a distinct AT1R- β -arrestin signaling complex through its recruitment of $G\alpha_i$, suggesting the existence of heterogeneity in biased-receptor conformations.

Given the importance of GRK-mediated receptor phosphorylation on the recruitment of β -arrestins to the receptor, the concept of a bar code of receptor phosphorylation has emerged as a potential mechanism for how ligand-specific receptor conformations selectively recruit specific transducers.^{12,31,32} According to this hypothesis, the receptor adopts distinct conformational states when bound to specific ligands, and each then recruits unique subsets of GRKs favoring distinct regions of the receptor, leading to phosphorylation of distinct amino acid residues on the c-terminal tail of the receptor, generating a phosphorylation “bar code.” These distinct GRK-mediated phosphorylation patterns direct the recruitment of signal transducers, such as β -arrestins, stabilizing them into unique active conformations and activating diversity in signaling profiles. This hypothesis is compatible with previous findings for the AT1R signaling whereby GRK5 and GRK6 are required for β -arrestin-mediated MAPK activation,¹³ whereas GRK2 and GRK3 mediate receptor internalization.³ Therefore, in our conceptual model, we speculate that membrane stretch induces a unique AT1R conformation that exposes an allosteric binding sites for $G\alpha_i$, and the bound $G\alpha_i$ in turn stabilizes a unique receptor conformation allowing receptor phosphorylation at specific residues on the c-terminal tail of the receptor by a unique subset of GRKs. The stretch-induced GRK-mediated distinctive phosphorylation pattern of the AT1R then induces a unique β -arrestin-biased receptor conformational state, leading to the activation of subsequent biased signaling. Future phosphoproteomic studies will be needed to determine the specific AT1R c-terminus phosphorylation pattern induced by balanced agonist, β -arrestin-biased agonist or membrane stretch.

AT1R plays vital roles in the regulation of cardiovascular function, and is one of the major targets for the therapeutic treatment of cardiovascular diseases such as congestive heart failure and hypertension.^{1,9,33} As $G\alpha_q$ -mediated AT1R signaling has been shown to mediate cardiomyocyte hypertrophy whereas β -arrestin-mediated signaling is thought to promote cardioprotection,^{34–36} the β -arrestin-biased agonist may provide novel clinical implications on promoting cardiac contractility and simultaneously blocking cardiac dysfunction.^{22,37–39} With our data now showing that two different β -arrestin-biased stimuli, membrane stretch and the biased ligands likely engage distinct AT1R molecular states to activate β -arrestin signaling, the potential to screen for unique biased ligands that can distinguish these different pathways may be possible.

In conclusion, we show that membrane stretch specifically promotes AT1R coupling to $G\alpha_i$, triggering activation of β -arrestin-biased signaling. These new insights for mechanoactivation of the AT1R reveal additional regulatory complexity for biased agonism.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

Funding information: National Institutes of Health, Grant numbers: HL56687, HL75443; American Heart Association, Grant number: 14PRE20480352



This work was supported by National Institutes of Health grants HL56687 and P01 HL75443 to HAR and a AHA Predoctoral Fellowship 14PRE20480352 to JW.

References

1. Rockman HA, Koch WJ, Lefkowitz RJ. Seven-transmembrane-spanning receptors and heart function. *Nature*. 2002; 415:206–212. [PubMed: 11805844]
2. Kohout TA, Lefkowitz RJ. Regulation of G protein-coupled receptor kinases and arrestins during receptor desensitization. *Mol Pharmacol*. 2003; 63:9–18. [PubMed: 12488531]
3. Kim J, Ahn S, Ren XR, et al. Functional antagonism of different G protein-coupled receptor kinases for beta-arrestin-mediated angiotensin II receptor signaling. *Proc Natl Acad Sci U S A*. 2005; 102:1442–1447. [PubMed: 15671181]
4. Moore CA, Milano SK, Benovic JL. Regulation of receptor trafficking by GRKs and arrestins. *Annu Rev Physiol*. 2007; 69:451–482. [PubMed: 17037978]
5. Shukla AK, Xiao K, Lefkowitz RJ. Emerging paradigms of beta-arrestin-dependent seven transmembrane receptor signaling. *Trends Biochem Sci*. 2011; 36:457–469. [PubMed: 21764321]
6. Rajagopal S, Rajagopal K, Lefkowitz RJ. Teaching old receptors new tricks: biasing seven-transmembrane receptors. *Nat Rev Drug Discov*. 2010; 9:373–386. [PubMed: 20431569]
7. Strachan RT, Sun JP, Rominger DH, et al. Divergent transducer-specific molecular efficacies generate biased agonism at a G protein-coupled receptor (GPCR). *J Biol Chem*. 2014; 289:14211–14224. [PubMed: 24668815]
8. Lefkowitz RJ, Shenoy SK. Transduction of receptor signals by beta-arrestins. *Science*. 2005; 308:512–517. [PubMed: 15845844]
9. Violin JD, Lefkowitz RJ. Beta-arrestin-biased ligands at seven-transmembrane receptors. *Trends Pharmacol Sci*. 2007; 28:416–422. [PubMed: 17644195]
10. Reiter E, Ahn S, Shukla AK, Lefkowitz RJ. Molecular mechanism of beta-arrestin-biased agonism at seven-transmembrane receptors. *Annu Rev Pharmacol Toxicol*. 2012; 52:179–197. [PubMed: 21942629]
11. Violin JD, Crombie AL, Soergel DG, Lark MW. Biased ligands at G-protein-coupled receptors: promise and progress. *Trends Pharmacol Sci*. 2014; 35:308–316. [PubMed: 24878326]
12. Wisler JW, Xiao K, Thomsen AR, Lefkowitz RJ. Recent developments in biased agonism. *Curr Opin Cell Biol*. 2014; 27:18–24. [PubMed: 24680426]
13. Rakesh K, Yoo B, Kim IM, Salazar N, Kim KS, Rockman HA. Beta-Arrestin-biased agonism of the angiotensin receptor induced by mechanical stress. *Sci Signal*. 2010; 3:ra46. [PubMed: 20530803]
14. Tang W, Strachan RT, Lefkowitz RJ, Rockman HA. Allosteric modulation of beta-arrestin-biased angiotensin II type 1 receptor signaling by membrane stretch. *J Biol Chem*. 2014; 289:28271–28283. [PubMed: 25170081]
15. Sauliere A, Bellot M, Paris H, et al. Deciphering biased-agonism complexity reveals a new active AT1 receptor entity. *Nat Chem Biol*. 2012; 8:622–630. [PubMed: 22634635]
16. Masuho I, Ostrovskaya O, Kramer GM, Jones CD, Xie K, Martemyanov KA. Distinct profiles of functional discrimination among G proteins determine the actions of G protein-coupled receptors. *Sci Signal*. 2015; 8:ra123. [PubMed: 26628681]

17. Crajoinas RO, Polidoro JZ, Carneiro de Morais CP, Castelo-Branco RC, Girardi AC. Angiotensin II counteracts the effects of cAMP/PKA on NHE3 activity and phosphorylation in proximal tubule cells. *Am J Physiol Cell Physiol*. 2016; 311:C768–C776. [PubMed: 27510906]
18. Zou Y, Komuro I, Yamazaki T, et al. Cell type-specific angiotensin II-evoked signal transduction pathways: critical roles of Gbetagamma subunit, Src family, and Ras in cardiac fibroblasts. *Circ Res*. 1998; 82:337–345. [PubMed: 9486662]
19. Ran FA, Hsu PD, Wright J, Agarwala V, Scott DA, Zhang F. Genome engineering using the CRISPR-Cas9 system. *Nat Protoc*. 2013; 8:2281–2308. [PubMed: 24157548]
20. Choo YY, Hagen T. Mechanism of cullin3 E3 ubiquitin ligase dimerization. *PLoS ONE*. 2012; 7:e41350. [PubMed: 22911784]
21. Kim J, Ahn S, Rajagopal K, Lefkowitz RJ. Independent beta-arrestin2 and Gq/protein kinase Czeta pathways for ERK stimulated by angiotensin type 1A receptors in vascular smooth muscle cells converge on transactivation of the epidermal growth factor receptor. *J Biol Chem*. 2009; 284:11953–11962. [PubMed: 19254952]
22. Violin JD, DeWire SM, Yamashita D, et al. Selectively engaging beta-arrestins at the angiotensin II type 1 receptor reduces blood pressure and increases cardiac performance. *J Pharmacol Exp Ther*. 2010; 335:572–579. [PubMed: 20801892]
23. Wang J, Hanada K, Staus DP, et al. Galphai is required for carvedilol-induced beta1 adrenergic receptor beta-arrestin biased signaling. *Nat Commun*. 2017; 8:1706. [PubMed: 29167435]
24. Hescheler J, Rosenthal W, Hinsch KD, Wulfern M, Trautwein W, Schultz G. Angiotensin II-induced stimulation of voltage-dependent Ca²⁺ currents in an adrenal cortical cell line. *EMBO J*. 1988; 7:619–624. [PubMed: 2456209]
25. Maturana AD, Casal AJ, Demaurex N, Vallotton MB, Capponi AM, Rossier MF. Angiotensin II negatively modulates L-type calcium channels through a pertussis toxin-sensitive G protein in adrenal glomerulosa cells. *J Biol Chem*. 1999; 274:19943–19948. [PubMed: 10391942]
26. de Gasparo M, Catt KJ, Inagami T, Wright JW, Unger T. International union of pharmacology. XXIII. The angiotensin II receptors. *Pharmacol Rev*. 2000; 52:415–472. [PubMed: 10977869]
27. Lefkowitz RJ. Arrestins come of age: a personal historical perspective. *Prog Mol Biol Transl Sci*. 2013; 118:3–18. [PubMed: 23764048]
28. Rosenbaum DM, Rasmussen SG, Kobilka BK. The structure and function of G-protein-coupled receptors. *Nature*. 2009; 459:356–363. [PubMed: 19458711]
29. Vaidehi N, Kenakin T. The role of conformational ensembles of seven transmembrane receptors in functional selectivity. *Curr Opin Pharmacol*. 2010; 10:775–781. [PubMed: 20933468]
30. Yasuda N, Miura S, Akazawa H, et al. Conformational switch of angiotensin II type 1 receptor underlying mechanical stress-induced activation. *EMBO Rep*. 2008; 9:179–186. [PubMed: 18202720]
31. Nobles KN, Xiao K, Ahn S, et al. Distinct phosphorylation sites on the beta(2)-adrenergic receptor establish a barcode that encodes differential functions of beta-arrestin. *Sci Signal*. 2011; 4:ra51. [PubMed: 21868357]
32. Zhou XE, He Y, de Waal PW, et al. Identification of phosphorylation codes for arrestin recruitment by G protein-coupled receptors. *Cell*. 2017; 170:457–469. e413. [PubMed: 28753425]
33. Pfeffer MA, Swedberg K, Granger CB, et al. Effects of candesartan on mortality and morbidity in patients with chronic heart failure: the CHARM-Overall programme. *Lancet*. 2003; 362:759–766. [PubMed: 13678868]
34. Fan G, Jiang YP, Lu Z, et al. A transgenic mouse model of heart failure using inducible Galpha q. *J Biol Chem*. 2005; 280:40337–40346. [PubMed: 16210321]
35. Benigni A, Corna D, Zoja C, et al. Disruption of the Ang II type 1 receptor promotes longevity in mice. *J Clin Invest*. 2009; 119:524–530. [PubMed: 19197138]
36. Whalen EJ, Rajagopal S, Lefkowitz RJ. Therapeutic potential of beta-arrestin- and G protein-biased agonists. *Trends Mol Med*. 2011; 17:126–139. [PubMed: 21183406]
37. Rajagopal K, Whalen EJ, Violin JD, et al. Beta-arrestin2-mediated inotropic effects of the angiotensin II type 1A receptor in isolated cardiac myocytes. *Proc Natl Acad Sci U S A*. 2006; 103:16284–16289. [PubMed: 17060617]

38. Ahn S, Kim J, Hara MR, Ren XR, Lefkowitz RJ. β -Arrestin-2 Mediates Anti-apoptotic Signaling through Regulation of BAD Phosphorylation. *J Biol Chem.* 2009; 284:8855–8865. [PubMed: 19171933]
39. Boerrigter G, Lark MW, Whalen EJ, Soergel DG, Violin JD, Burnett JC Jr. Cardiorenal actions of TRV120027, a novel ss-arrestin-biased ligand at the angiotensin II type I receptor, in healthy and heart failure canines: a novel therapeutic strategy for acute heart failure. *Circ Heart Fail.* 2011; 4:770–778. [PubMed: 21835984]

(A) Proximity Ligation Assay (PLA): HA-AT1R x G α_i 

NS: No Stimulation; TRV023: TRV120023; OSM: Osmotic Stretch

FIGURE 1.

Osmotic stretch promoted G α_i recruitment to AT1R, while the balanced agonist AngII and the β -arrestin-biased agonist TRV120023 did not have significant effect. HEK293 cells stably expressing HA-tagged AT1Rs were pretreated with vehicle or 200 ng/mL PTX for overnight, then stimulated with 1 μ M AngII, 1 μ M TRV120023 or osmotic stretch for 10 min. In proximity ligation assay (PLA), cells were immuno-stained with G α_i antibody raised in mouse and HA antibody raised in rabbit. The recruitment of G α_i to AT1R is indicated by the green fluorescence signal. A: Representative image showings osmotic stretch-induced G α_i recruitment to AT1R. Scale bar = 10 μ M. B: The PLA signal was quantified as the


average intensity of the green fluorescence per cell, and normalized by subtracting the non-specific background determined as the green fluorescence intensity per cell in wildtype HEK293 cells without HA-AT1R overexpression. Data represent the mean \pm SEM for four independent experiments. Statistical significance versus unstimulated cells was assessed using one-way ANOVA with Bonferroni correction

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript



NS: No Stimulation; TRV023: TRV120023; OSM: Osmotic Stretch

FIGURE 2.

Osmotic stretch specifically induced AT1R coupling to $G\alpha_i$, but not $G\alpha_q$. In contrast, AngII promoted AT1R- $G\alpha_q$ coupling, while TRV120023 and osmotic stretch did not have significant effect. A and B: Osmotic stretch increases the amount of $G\alpha_i$ bound to AT1R, which was blocked by the PTX pretreatment. HEK293 cells stably expressing HA-tagged AT1Rs were transfected with $G\alpha_{i2}$ 48 h before experiment, and pretreated with vehicle or 200 ng/mL PTX in serum-free medium for overnight. Cell were stimulated with 1 μ M AngII, 10 μ M TRV120023, or osmotic stretch for 10 min. $G\alpha_i$ recruitment was determined with co-immunoprecipitation assay. HA-AT1Rs were immunoprecipitated with anti-HA magnetic beads, and bound $G\alpha_{i2}$ was detected with western blot. C and D: AngII specifically induced $G\alpha_q$ recruitment to AT1R. AT1R stable cells were serum-starved for overnight, then stimulated with 1 μ M AngII, 10 μ M TRV120023 or osmotic stretch for 10 min. The AT1R- $G\alpha_q$ coupling was determined with co-immunoprecipitation. Signals were quantified by densitometry, normalized to the immunoprecipitated AT1R level, and expressed as fold change over the unstimulated cells. Data represent the mean \pm SEM for n


independent experiments as marked on the figure. Statistical significance versus unstimulated cells was assessed using one-way ANOVA with Bonferroni correction

Author Manuscript


Author Manuscript

Author Manuscript

Author Manuscript

**FIGURE 3.**

$G\alpha_i$ is required for the osmotic stretch-induced β -arrestin2 recruitment to the AT1R. HEK 293 cells stably expressing CFP-tagged AT1R and YFP-tagged β -arrestin2 were transiently transfected with GRK5. After pretreatment with vehicle or 200 ng/mL PTX for overnight, cells were stimulated with 1 μ M AngII, 1 μ M, or 10 μ M TRV120023, or osmotic stretch for 10 min. A: In response to AngII, TRV120023, or osmotic stretch, AT1R redistributed from the plasma membrane to intracellular puncta, and the cytosolic evenly distributed β -arrestin2 translocated and co-localized with the internalized AT1Rs, indicating the β -arrestin2 recruitment to AT1Rs. Pretreatment with PTX significantly inhibited osmotic stretch-induced β -arrestin2 recruitment, whereas has no effect to AngII- or TRV120023-induced response. Scale bar = 10 μ M. B: Quantification of β -arrestin2 translocation was expressed as the percentage of cells showing intracellular aggregates of β -arrestin. Data represent the mean \pm SEM for n independent experiments as marked on the figure. Statistical significance versus unstimulated cells, or between the vehicle and PTX pretreated group, was assessed using two-way ANOVA with Bonferroni correction

**FIGURE 4.**

Mechanically activated AT1R signaling is dependent on $G\alpha_i$. A and B: PTX selectively blocked AT1R-mediated EGFR internalization induced by osmotic stretch. HEK 293 cells stably expressing AT1R were transiently transfected with GFP-tagged EGFR and GRK5. Cells were pretreated with vehicle or 200 ng/mL PTX for overnight, and then stimulated with 1 μ M AngII, 1 μ M TRV120023, or osmotic stretch for 10 min. Quantification of EGFR internalization was performed by counting the percentage of the cells showing intracellular aggregates of GFP-EGFR. All three stimulations promoted EGFR internalization, whereas only the osmotic stretch-induced response is PTX-sensitive. Scale bar = 10 μ M. C and D: $G\alpha_i$ is required for osmotic stretch-induced ERK1/2 phosphorylation in HEK293 cells. AT1R stable cells pretreated with vehicle or PTX were stimulated with 1 μ M AngII, 1 μ M TRV120023 or osmotic stretch for 10 min. Cell lysates were analyzed for phosphorylated ERK1/2 (pERK1/2) and total ERK1/2 (tERK1/2) with Western blot. The pERK1/2 signals were quantified by densitometry, normalized to tERK1/2, and expressed as fold change over unstimulated cells. ERK1/2 phosphorylation was significantly increased in response to AngII or osmotic stretch. PTX pretreatment diminished osmotic stretch-induced ERK1/2 phosphorylation, whereas has no effect on AngII-induced response. E and F: PTX blocked mechanical stretch-induced ERK1/2 phosphorylation in mice heart. In ex vivo perfused wild type heart, LV balloon stretch significantly increased ERK1/2 phosphorylation. Pretreatment with PTX (25 μ g/kg) significantly inhibited balloon stretch-induced ERK1/2 phosphorylation. Quantification of pERK1/2 signal was performed as mentioned above. Data represent the mean \pm SEM for *n* independent experiments or animals as marked on the


figure. Statistical significance was assessed using two-way ANOVA with Bonferroni correction

Author Manuscript


Author Manuscript

Author Manuscript


Author Manuscript

**FIGURE 5.**


Osmotic stretch-induced ERK1/2 phosphorylation is independent of $G\alpha_q$, whereas dependent on EGFR transactivation. A and B: Osmotic stretch-induced ERK1/2 phosphorylation was not blocked by Ro318425, suggesting $G\alpha_q$ is not required for this response. AT1R stable cells were pretreated with 1 μ M PKC inhibitor Ro318425 for 30 min, then stimulated with 1 μ M AngII, 1 μ M TRV120023, or osmotic stretch for 10 min. C and D: The PLC inhibitor U73122 blocked the AngII-induced ERK1/2 phosphorylation, whereas had no effect on the osmotic stretch response. Cells were pretreated with 10 μ M U73122. E and F: AG1478 blocked ERK1/2 phosphorylation induced by AngII or osmotic stretch, suggesting the ERK1/2 activation is mediated by the transactivated EGFRs. AT1R stable cells were pretreated with 1 μ M EGFR inhibitor AG1478 for 30 min before stimulation. The pERK1/2 signals were quantified by densitometry, normalized to tERK1/2, and expressed as fold change over unstimulated cells. Data represent the mean \pm SEM for n independent experiments or animals as marked on the figure. Statistical significance was assessed using two-way ANOVA with Bonferroni correction

**FIGURE 6.**

TRV120026 induces the β -arrestin-biased AT1R signaling in a $G\alpha_i$ -independent manner, consistent with the TRV120023 response. A and B: PTX had no effect on the TRV120026-induced β -arrestin2 recruitment to AT1R. HEK 293 cells stably expressing CFP-tagged AT1R and YFP-tagged β -arrestin2 were transiently transfected with GRK5. After pretreatment with vehicle or 200 ng/mL PTX for overnight, cells were stimulated with 1 μ M AngII, 10 μ M TRV120023, 1 μ M or 10 μ M TRV120026, or osmotic stretch for 10 min. Scale bar = 10 μ M. Quantification of β -arrestin2 translocation was expressed as the percentage of cells showing intracellular aggregates of β -arrestin. C and D: similar to the effect of TRV120023, TRV120026 induced a mild increase of ERK1/2 phosphorylation, which was not PTX-sensitive. HEK293 cells stably expressing AT1R were pretreated and stimulated as mentioned above. The pERK signals were quantified by densitometry, normalized to tERK, and expressed as fold change over unstimulated cells. Data represent the mean \pm SEM for n independent experiments as marked on the figure. Statistical significance was assessed using two-way ANOVA with Bonferroni correction

**FIGURE 7.**

Osmotic stretch-induced β -arrestin2 recruitment to AT1R was blocked in $G\alpha_i$ knockout cells. A: Western blot showing the knockout of $G\alpha_i$ in HEK293 cells by CRISPR-Cas9 gene editing. B: AngII, TRV120023, and TRV120026 induced β -arrestin2 translocation in $G\alpha_i$ knockout cells to the same extent as that in wild type cells. In contrast, the osmotic stretch-induced response was blocked in $G\alpha_i$ knockout cells. $G\alpha_i$ knockout cells were transfected with CFP-tagged AT1R, YFP-tagged β -arrestin2, and GRK5. Cells were stimulated with 1 μ M AngII, 10 μ M TRV120023, 10 μ M TRV120026, or osmotic stretch for 10 min. B: Quantification of β -arrestin2 translocation was expressed as the percentage of cells showing intracellular aggregates of β -arrestin. Data represent the mean \pm SEM for n independent experiments as marked on the figure. Statistical significance was assessed using two-way ANOVA with Bonferroni correction

**FIGURE 8.**

Schematic model of AT1R signaling induced by different modes of receptor activation. Different stimulations stabilize the receptors into distinct active conformational states, allowing the recruitment of unique subsets of transducers that subsequently activates divergent arrays of downstream signaling. When induced by the balanced agonist AngII, the AT1Rs transduce signaling through both G_{α_q} and β -arrestins, whereas the biased agonists TRV120023 and TRV120026 preferentially induce β -arrestin signaling without activating G proteins. In contrast, another β -arrestin-biased receptor stimulation, mechanical stress, specifically promotes the recruitment of G_{α_i} to the receptors and activates β -arrestin signaling through a mechanism distinct from that induced by the biased ligands