



Published in final edited form as:

*Am J Prev Med.* 2017 September ; 53(3 Suppl 1): S78–S85. doi:10.1016/j.amepre.2017.03.026.

## Bars, Nightclubs, and Cancer Prevention: New Approaches to Reduce Young Adult Cigarette Smoking

Pamela M. Ling, MD, MPH<sup>1</sup>, Louisa M. Holmes, PhD<sup>1</sup>, Jeffrey W. Jordan, MA<sup>2</sup>, Nadra E. Lisha, PhD<sup>1</sup>, and Kirsten Bibbins-Domingo, MD, PhD, MAS<sup>3</sup>

<sup>1</sup>Center for Tobacco Control Research and Education, and Division of General Internal Medicine, Department of Medicine, University of California, San Francisco, San Francisco, California

<sup>2</sup>Rescue Agency, San Diego, California

<sup>3</sup>Center for Vulnerable Populations, Department of Medicine, and Department of Epidemiology and Biostatistics, University of California, San Francisco, San Francisco, California

### Abstract

**Introduction**—Tobacco contributes to multiple cancers, and it is largely preventable. As overall smoking prevalence in California declines, smoking has become concentrated among high-risk groups. Targeting social/cultural groups (i.e., “peer crowds”) that share common values, aspirations, and activities in social venues like bars and nightclubs may reach high-risk young adult smokers. Lack of population data on young adult peer crowds limits the ability to assess the potential reach of such interventions.

**Methods**—This multimodal population-based household survey included young adults residing in San Francisco and Alameda counties. Data were collected in 2014 and analyzed in 2016. Multivariable logistic regressions assessed smoking by sociodemographic factors, attitudes, self-rated health, peer crowd affiliation, and bar/nightclub attendance.

**Results**—Smoking prevalence was 15.1% overall; 35.3% of respondents sometimes or frequently attended bars. In controlled analyses, bar attendance (AOR=2.13, 95% CI=1.00, 4.53) and binge drinking (AOR=3.17, 95% CI=1.59, 6.32) were associated with greater odds of smoking, as was affiliation with “Hip Hop” (AOR=4.32, 95% CI=1.48, 12.67) and “Country” (AOR=3.13, 95% CI=1.21, 8.09) peer crowds. Multivariable models controlling for demographics estimated a high probability of smoking among bar patrons affiliating with Hip Hop (47%) and Country (52%) peer crowds.

**Conclusions**—Bar attendance and affiliation with certain peer crowds confers significantly higher smoking risk. Interventions targeting Hip Hop and Country peer crowds could efficiently reach smokers, and peer crowd–tailored interventions have been associated with decreased

---

This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Address correspondence to: Pamela M. Ling, MD, MPH, Center for Tobacco Control Research and Education, Division of General Internal Medicine, Department of Medicine, University of California, San Francisco, 530 Parnassus Ave., Suite 366, San Francisco CA 94143. [pamela.ling@ucsf.edu](mailto:pamela.ling@ucsf.edu).

This article is part of a supplement issue titled Opportunities for Cancer Prevention During Early Adulthood.

#### SUPPLEMENTAL MATERIAL

Supplemental materials associated with this article can be found in the online version at <https://doi.org/10.1016/j.amepre.2017.03.026>.

smoking and binge drinking. Targeted interventions in bars and nightclubs may be an efficient way to address these cancer risks.

---

## INTRODUCTION

Decreasing tobacco use is important for cancer prevention.<sup>1</sup> In California, smoking declined from 23.7% to 11.6% between 1988 and 2014, and lung cancer rates have fallen twice as fast as the rest of the U.S.<sup>2</sup> As smoking declines overall, it is increasingly concentrated in high-risk subgroups, including those with low SES and racial/ethnic or sexual/gender minority groups.<sup>2</sup>

Young adults are important targets for cancer prevention. During young adulthood, most tobacco experimenters either quit or progress to regular smoking and addiction.<sup>3</sup> Tobacco marketing strategies entrench assumptions about tobacco use that consumers perceive to be “natural,” such as links between drinking and smoking.<sup>4–10</sup> Tobacco marketing campaigns were designed to reinforce alcohol and tobacco co-use.<sup>11,12</sup> The tobacco industry also developed sophisticated strategies targeting young adults<sup>13</sup> with campaigns targeting their values, activities, aspirations, peer groups, and social environments.<sup>14,15</sup> Tobacco companies developed special marketing campaigns focused on young adult bar patrons because in addition to having higher smoking rates, they were regarded as key “trendsetters” influencing their peers.<sup>16,17</sup>

Tobacco marketing campaigns also increase disparities by encouraging use among vulnerable groups such as poor women<sup>18</sup>; rural men<sup>19–22</sup>; African Americans<sup>23–26</sup>; Hispanics<sup>27</sup>; Asians<sup>28</sup>; the working poor<sup>29</sup>; the military<sup>30–32</sup>; the lesbian, gay, bisexual, and transgender (LGBT) community<sup>33–35</sup>; or those with mental illness.<sup>36</sup> Additionally, tobacco companies target young people.<sup>37–42</sup> Thus, there is a critical need to counteract predatory tobacco marketing, particularly within high-risk subgroups.

One innovative tobacco prevention approach leverages targeted marketing strategies to develop targeted branded anti-tobacco interventions utilizing psychographic factors such as social affiliations, activities, and values.<sup>13</sup> Peer crowd affiliation is one way to approximate psychographic segmentation to identify high-risk subgroups of young adults and adolescents.<sup>43,44</sup> Distinct from one’s personal networks, “peer crowds” reflect common values, activities, fashion, and influencers that link different social networks. Peer crowds, described in detail elsewhere,<sup>45</sup> include “Hipsters” (who value creativity and individuality, prefer indie rock music, and set trends); “Hip Hop” (who identify with having overcome disproportionate life challenges to succeed, express strength and success, prefer hip hop music, and value strength, honor, and respect); or “Young Professionals” (who value their careers, prioritize networking).

Peer crowd affiliation is associated with tobacco use, independent of demographics.<sup>45</sup> Peer crowds have not yet been studied in population-based samples of young adults. This population-based study in San Francisco and Alameda counties in 2014 describes the size of young adult peer crowds and the relationships among peer crowd affiliation, demographic characteristics, and smoking. The main hypothesis was that there would be significant

differences in smoking behavior based on peer crowd identification, independent of sociodemographic factors. Exploratory analyses also addressed if the combination of bar attendance and peer crowd affiliation could increase the efficiency of targeted interventions.

## METHODS

### Data Sample

Young adults are difficult to reach as they frequently move households, experience changing life circumstances (e.g., going to college, taking new jobs), and are less likely to have telephone landlines. Surveys using mobile phone numbers may not reflect respondents' actual location. Most population-based health surveys including young adults are national samples lacking sufficient numbers of young adults in a particular region. The 2014 San Francisco Bay Area Young Adult Health Survey, a unique multimode probabilistic household survey stratified by race/ethnicity of young adults in San Francisco and Alameda Counties, was conducted to facilitate local planning for tobacco control programs (analysis done in 2016).

Potential respondent households were identified from marketing address lists and Census data, identifying households with approximately 40% chance of a young adult occupant (N=15,000 addresses) and stratified into eight samples accounting for Asian and Hispanic ethnic surnames, racial/ethnic composition, and one random sample from the remaining population of young adults. Next, the 2009–2013 American Community Survey and 2010 decennial Census data were used in a multistage sampling design to identify Census blocks in which 15% of residents were either Hispanic, black, or non-Hispanic white and in the eligible age range to oversample black and Hispanic young adults. The survey utilized mail/web, telephone, and face-to-face interviews. All households were first contacted by letter; followed by postcard and letter reminders (three mailings); followed by phone calls to nonresponding households; followed by face-to-face visits. Most surveys were completed on paper or online (five respondents completed by phone, skipping the peer crowd measure). Phone calls primarily determined if there were no eligible young adults in the household. A random selection of addresses ( $n=1,240$ ) from the households lacking either a mail or phone response were visited, as well as randomly selected housing units from the block sample, to attempt to complete surveys ( $n=1,636$  housing units, a total of 2,876 households), resulting in 1,363 complete surveys (30% response rate). All survey protocols were approved by the IRB at the University of California, San Francisco.

### Measures

Respondents reporting smoking cigarettes on 1–30 of the last 30 days were coded as current smokers. Age was measured continuously, subtracting respondent date of birth from date of survey administration. Sex was measured as male (1) or female (0). Race/ethnicity was measured with two items (*Are you of Hispanic, Latino, or Spanish origin?* and *What is your race?*), with 14 categories corresponding to American Community Survey.<sup>46</sup> Responses were combined to generate categories: Latino, non-Hispanic white, non-Hispanic black, non-Hispanic Asian/Pacific Islander, and non-Hispanic other race. Sexual orientation was measured with the question: *Do you think of yourself as... heterosexual or straight;*

*homosexual, gay, lesbian, or queer; bisexual; transgender, or other*, coded as LGBT (1) or straight (0). Education was measured with 16 categories ranging from *no school* to *doctorate degree*, and dichotomized as college educated (current enrollment in college or bachelor's degree or higher; 1) with others as (0).

The I-Base Survey™ was used to determine peer crowd affiliation. The measure includes a grid of images representing different peer crowds developed using extensive formative qualitative research described previously<sup>43</sup> where young people group images and describe the characteristics of different friend groups. The 72 images that were selected most consistently for each peer crowd were included on the grid. Survey respondents chose three male and three female photos that *best fit into your main group of friends*. Scoring is described in more detail elsewhere<sup>45</sup>; selected pictures scored 3, 2, or 1 points for their corresponding peer crowd based on rank. Participants also selected those who *least fit into your main group of friends*, scoring -1, -2, and -3, respectively. The total score for each peer crowd ranged from -12 to 12; respondents were categorized based on the crowd in which they scored highest. Seventy-eight (7%) respondents had tied high scores and were randomly assigned to one of the two tied peer crowds. The six peer crowds were named “Hipster,” “Country,” “Hip Hop,” “Partier,” “Homebody,” and “Young Professional”; names are used for reporting here, but not in campaigns.

Perceived smoking prevalence was measured, asking respondents to estimate the percentage of people their age they perceived to smoke tobacco. Perception that smoking reduces stress was measured using agreement with *smoking cigarettes helps to reduce stress* on a Likert scale from *not at all* (1) to *extremely* (7). As in prior studies,<sup>47</sup> anti-tobacco attitudes were measured using agreement with three items: *I want to be involved in efforts to get rid of cigarette smoking; I would like to see the cigarette companies go out of business; and Taking a stand against smoking is important to me* on a 7-point Likert scale, calculating mean response across the three items.

Bar attendance was measured with *How often do you go to bars or clubs? (never, rarely, sometimes, or frequently)*. Authors coded *sometimes* or *frequently* as bar patrons (1), and others not (0). Binge drinking was measured asking number of days in the past 30 when respondents had *five or more alcoholic drinks within a few hours*. Those drinking on at least 1 of the past 30 days, and consuming at least five drinks as specified were coded as binge drinkers (1).

Self-rated health was measured with *Would you say your health in general is... excellent; very good; good; fair; poor?* Respondents answering *excellent* or *very good* were coded dichotomously as high self-rated health (1).

## Statistical Analysis

Of 1,363 total respondents, 1,151 (84%) provided information for peer crowd scores, and 1,344 provided sufficient information for imputation (Table 1). Multiple imputation in Stata, version 14, was used to estimate the missing values, and final models with and without the imputed values were checked for bias with imputation. Bivariate relationships between peer crowd affiliation and smoking were analyzed for the full sample and among bar patrons

(Table 2). Hierarchical logistic regressions on smoking first controlled for peer crowd and sociodemographic characteristics, then added self-rated health and tobacco attitudes (Table 3). The “margins” command in Stata measured the discrete probabilities associated with being a smoker and peer crowd affiliation, holding age, sex, race/ethnicity, LGBT identification, and education at their means. All results were weighted using person-level weights constructed following data collection. All analyses used the “svyset” command to adjust for the complex sampling design and clustering.

## RESULTS

Sample characteristics are presented in Table 1. The overall smoking prevalence in the sample was 15.1%, higher than the 10.1% adult prevalence in San Francisco County.<sup>2</sup> The weighted sample had a mean age of 22.7 years, was 50% male, and racially and ethnically diverse, with approximately 32% non-Hispanic white, 28% non-Hispanic Asian, 24% Latino, and 10% non-Hispanic black individuals. Approximately 50% of respondents reported current enrollment in college, and 34% reported having a bachelor’s degree or more. The peer crowds with low smoking rates made up a large proportion of the population: 36.5% and 35.4% reported strongest affiliation with Young Professional and Homebody peer crowds.

Though 35.3% of all respondents attended bars sometimes or frequently, 57% of smokers attended bars sometimes or frequently (Table 2). Smoking was higher among bar patrons (24%) versus 10% among those who rarely/never attended bars. Differences in smoking rates were even greater taking peer crowds into account: young adult bar patrons who affiliated with the Partier, Hip Hop, and Country peer crowds reported smoking rates of between 40% and 57%, whereas <10% of Young Professionals smoked (Figure 1). A large proportion of the smokers within the Country (72%); Partier (71%); and Hip Hop (66%) peer crowds attended bars.

In multivariable logistic regressions, peer crowd affiliation was independently associated with smoking, across both the first iteration of the model, controlling for sociodemographic factors and the second iteration of the model, which added attitudinal and health-related behaviors (Table 3). Hip Hop (AOR=5.58, 95% CI=2.24, 13.88) and Country (AOR=3.55, 95% CI=1.37, 9.19) peer crowd affiliation was associated with significantly greater odds of smoking, controlling for age, gender, race/ethnicity, sexual orientation, and education. This association remained significant in Model 2 for Hip Hop (AOR=4.42, 95% CI=1.51, 12.98) and Country (AOR=3.06, 95% CI=1.17, 7.99) affiliation, controlling for tobacco-related attitudes and risk behaviors. Binge drinking was also associated with smoking. Factors associated with lower odds of smoking included non-Hispanic Asian/Pacific Islander race, more education, strong anti-tobacco attitudes, and very good/excellent health (Table 3).

After specifying the multivariable models, the discrete probabilities of smoking given affiliation with a certain peer crowd and propensity to frequent bars were estimated, holding demographic factors (age, sex, race, sexual orientation, and education) at their means. The probability of reaching smokers through bars differed considerably by peer crowd. For example, Young Professionals (37.1% of the sample) have a low (8.2%) smoking rate, but

represent a large number of smokers ( $n \approx 8,000$ ). Bars are not an efficient way to reach Young Professionals, as the probability of a Young Professional bar patron being a smoker is only 10% ( $\approx 4,200$  smokers). By contrast, the Hip Hop peer crowd (8.4% of the sample) has a much higher smoking rate (39.2%), representing approximately 8,500 smokers. This model estimated the probability of smoking among Hip Hop bar patrons at 47% ( $\approx 5,600$  smokers), suggesting that bars are an efficient way to reach Hip Hop smokers. Similarly, the probability of smoking among Country bar patrons was 52% ( $\approx 2,900$  smokers).

## DISCUSSION

This study is the first to estimate the potential reach of bar and nightclub interventions to different peer crowds within the young adult population in the San Francisco Bay Area. The results suggest that bars and nightclubs efficiently reach smokers, particularly those who identify with the Hip Hop and Country peer crowds. Additionally, a large number of young adult smokers do not frequent bars, and these reflect larger-size peer crowds with lower risk (e.g., Young Professionals). Campaigns such as the recent “Finish IT” campaign sponsored by the Truth Initiative® emphasizing low smoking prevalence and mobilizing nonsmokers to end the tobacco epidemic may appeal particularly to these lower-risk groups. However, members of peer crowds with substantially more smoking (e.g., Partier, Hip Hop, and Country) may not relate to messages highlighting low overall smoking rates. This study suggests that bars are an efficient way to reach these high-risk peer crowds, as 60%–70% of the smokers in these peer crowds attended bars and clubs. Furthermore, as bars cater to certain types of patrons that frequently align with the different peer crowds, bar-based interventions can be easily tailored to peer crowd.

Bars and nightclubs are a nexus for risk behavior among young adults. This may be due to the aggressive marketing of both tobacco and alcohol in bars,<sup>11,48,49</sup> reinforcing effects of tobacco and alcohol, which increases the risk of addiction,<sup>50,51</sup> or features of the physical and social environment that normalize risky behaviors (including other drug use, sexual risk taking).<sup>52–56</sup> Conversely, bars and clubs pose opportunities for cancer prevention, addressing the cancer risk afforded by both tobacco and alcohol use. Tobacco and alcohol co-use increases the risk of cancers of the mouth, throat, esophagus, and upper aerodigestive tract.<sup>57–59</sup> Public health interventions focused on one behavior may affect other risks: Anti-smoking interventions for young adult bar patrons have been associated with decreases in binge drinking.<sup>60–62</sup> This approach warrants further study.

More important, because peer crowds are connected to young adult social identities, lifestyles, and values, messages targeted to peer crowds may be more relevant. Adolescents who identified with the peer crowd in tailored advertisements reported stronger anti-smoking attitudes and lower levels of smoking susceptibility.<sup>63</sup> Young adult bar and nightclub interventions targeted to Hipsters,<sup>62</sup> Partiers,<sup>61</sup> and LGBT audiences<sup>64</sup> have shown significant decreases in smoking. These tailored messages differ in content and style depending on the peer crowd (Appendix Figure 1, available online). For example, messages appealing to the Hipster crowd frequently connected the issue of tobacco use, marketing, or production to social justice issues, such as the environment, tobacco production and world hunger, or animal testing. By contrast, messages for the Hip Hop peer crowd might address

how the tobacco industry has targeted communities of color with menthol or flavored little cigars/cigarillos.<sup>23,39,65</sup>

Peer crowds are also racially and ethnically diverse. In the San Francisco Bay Area Young Adult Health Survey, the Hip Hop peer crowd was 31.7% non-Hispanic black, 26.6% Hispanic, 15.7% non-Hispanic white, 15.6% non-Hispanic Asian/Pacific Islander, and 10.5% non-Hispanic other race/ethnicity. Although it may be difficult to reach Asian smokers by targeting the general population (Asians made up 27.7% of the sample, with a smoking prevalence of 8.6%, representing 7,338 smokers), by targeting the Hip Hop peer crowd one could reach approximately 1,600 (22%) Asian smokers.

Public health programs frequently need to balance the objective to reach a large part of the population with the efficiency of spending limited resources. For example, assume a hypothetical situation where the San Francisco health department has a budget of \$100,000 for a local media campaign, and for simplicity, it costs \$1 to reach a young adult one time. It would cost approximately \$96,000 to reach the approximately 8,000 Young Professional smokers. To reach almost the same number of 8,500 Hip Hop smokers, it would cost approximately \$22,000 (because of the high prevalence of smoking in this peer group). For a budget of \$100,000, one could choose to reach Young Professional smokers once, or Hip Hop smokers four to five times. By concentrating on Hip Hop bar patrons, one could spend approximately \$11,000 to reach approximately two thirds of the Hip Hop smokers, and that a budget of \$100,000 would reach this high-risk group almost ten times. In this sample, approximately 70% of the population identified with Young Professional or Homebody peer crowds, including about half of the smokers, whereas the other half of smokers were in the 30% of the population identifying with high-risk peer crowds (Partier, Hipster, Hip Hop, and Country). Addressing smoking among high-risk peer crowds reaches smokers more efficiently, as smoking rates were 20%–40% in the high-risk peer crowds versus 8%–14% in the low-risk peer crowds.

### Limitations

Though the study sampling and weighting enabled population estimates, the overall response rate for the survey was relatively low (30%) and so may include nonresponse bias. The study sample may not generalize beyond the two studied counties. The sample was not large enough to estimate both smoking and race/ethnicity within peer crowds for all groups. Pictorial peer crowd measures used in this study were not compared to prior measures, which utilize lists of crowd names.<sup>66–68</sup> More research validating peer crowd measures is needed,<sup>69</sup> but peer crowd affiliation has been recognized as a powerful strategy for tailored interventions.<sup>70</sup> All measures were self-reported without biochemical validation of smoking status.

### CONCLUSIONS

This is the first study of peer crowds in a population-based sample of young adults. Bar attendance and affiliation with high-risk peer crowds can reach smokers more efficiently, increasing the probability of reaching a smoker from 15% in the general population to 40%–50% by targeting Hip Hop, Partier, and Country bar patrons. Targeting high-risk peer crowds

can reach half of the young adult smokers in San Francisco with greater efficiency. Targeting high-risk peer crowds also preferentially reaches disproportionately affected groups of young adult smokers and may be a valuable tool to address disparities in tobacco-related cancer risk and morbidity.

## Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

## Acknowledgments

Publication of this article was supported by the U.S. Centers for Disease Control and Prevention (CDC), an Agency of the U.S. Department of Health and Human Services, under contract number: 200–2017-M-94637.

The authors wish to thank Drs. Enrico Marcelli and Anthony Roman for assistance with complex sampling, study design, and data weighting.

This study was supported by the NIH/National Institute of Minority Health and Health Disparities (5P60MD006902) and the National Cancer Institute (U01-CA154240). The content is solely the responsibility of the authors and does not necessarily represent the official views of NIH. The study sponsors had no role in study design; collection, analysis, and interpretation of data; writing the report; or the decision to submit the report for publication.

Mr. Jordan is an employee of Rescue Agency, a company that implements bar and nightclub interventions discussed in this study. No other financial disclosures were reported by the authors of this paper.

## References

1. U.S. DHHS. The Health Consequences of Smoking: 50 Years of Progress. A Report of the Surgeon General. Atlanta, GA: U.S. DHHS, CDC, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health; 2014.
2. California Department of Public Health. California Tobacco Control Program. California Tobacco Facts and Figures 2016. Sacramento, CA: California Department of Public Health; 2016.
3. Ling PM, Glantz SA. Why and how the tobacco industry sells cigarettes to young adults: evidence from industry documents. *Am J Public Health*. 2002; 92(6):908–916. <https://doi.org/10.2105/AJPH.92.6.908>. [PubMed: 12036776]
4. Nichter, M. *Lighting Up: The Rise of Social Smoking on College Campuses*. New York: New York University Press; 2015.
5. Dawson DA. Drinking as a risk factor for sustained smoking. *Drug Alcohol Depend*. 2000; 59(3): 235–249. [https://doi.org/10.1016/S0376-8716\(99\)00130-1](https://doi.org/10.1016/S0376-8716(99)00130-1). [PubMed: 10812284]
6. Little HJ. Behavioral mechanisms underlying the link between smoking and drinking. *Alcohol Res Health*. 2000; 24(4):215–224. [PubMed: 15986716]
7. John U, Meyer C, Rumpf H-J, Hapke U. Probabilities of alcohol high-risk drinking, abuse or dependence estimated on grounds of tobacco smoking and nicotine dependence. *Addiction*. 2003; 98(6):805–814. <https://doi.org/10.1046/j.1360-0443.2003.00381.x>. [PubMed: 12780369]
8. Weitzman ER, Chen Y-Y. The co-occurrence of smoking and drinking among young adults in college: national survey results from the United States. *Drug Alcohol Depend*. 2005; 80(3):377–386. <https://doi.org/10.1016/j.drugalcdep.2005.05.008>. [PubMed: 16009507]
9. Harrison ELR, Desai RA, McKee SA. Nondaily smoking and alcohol use, hazardous drinking, and alcohol diagnoses among young adults: findings from the NESARC. *Alcohol Clin Exp Res*. 2008; 32(12):2081–2087. <https://doi.org/10.1111/j.1530-0277.2008.00796.x>. [PubMed: 18828805]
10. Nichter M, Nichter M, Carkoglu A, Lloyd-Richardson E. Tobacco Etiology Research Network (TERN). Smoking and drinking among college students: “it’s a package deal”. *Drug Alcohol Depend*. 2010; 106(1):16–20. <https://doi.org/10.1016/j.drugalcdep.2009.07.025>. [PubMed: 19758771]




11. Jiang N, Ling PM. Reinforcement of smoking and drinking: tobacco marketing strategies linked with alcohol in the United States. *Am J Public Health*. 2011; 101(10):1942–1954. <https://doi.org/10.2011.300157>. [PubMed: 21852637]
12. Jiang N, Ling P. Vested interests in addiction research and policy. Alliance between tobacco and alcohol industries to shape public policy. *Addiction*. 2013; 108(5):852–864. <https://doi.org/10.1111/add.12134>. [PubMed: 23587076]
13. Ling PM, Glantz SA. Using tobacco-industry marketing research to design more effective tobacco-control campaigns. *JAMA*. 2002; 287(22):2983–2989. <https://doi.org/10.1001/jama.287.22.2983>. [PubMed: 12052128]
14. Pierce JP, Messer K, James LE, et al. Camel No. 9 cigarette-marketing campaign targeted young teenage girls. *Pediatrics*. 2010; 125(4):619–626. <https://doi.org/10.1542/peds.2009-0607>. [PubMed: 20231181]
15. R.J. Reynolds Tobacco Company. [Accessed August 19, 2008] Camel smokes. [www.camelsmokes.com](http://www.camelsmokes.com). Published 2008
16. Sepe E, Ling PM, Glantz SA. Smooth moves: bar and nightclub tobacco promotions that target young adults. *Am J Public Health*. 2002; 92(3):414–419. <https://doi.org/10.2105/AJPH.92.3.414>. [PubMed: 11867322]
17. Hendlin Y, Anderson SJ, Glantz SA. “Acceptable rebellion”: marketing hipster aesthetics to sell Camel cigarettes in the U.S. *Tob Control*. 2010; 19(3):213–222. <https://doi.org/10.1136/tc.2009.032599>. [PubMed: 20501494]
18. Brown-Johnson CG, England LJ, Glantz SA, Ling PM. Tobacco industry marketing to low socioeconomic status women in the U.S.A. *Tob Control*. 2014; 23(e2):e139–e146. <https://doi.org/10.1136/tobaccocontrol-2013-051224>. [PubMed: 24449249]
19. Ling PM, Haber LA, Wedl S. Branding the rodeo: a case study of tobacco sports sponsorship. *Am J Public Health*. 2010; 100(1):32–41. <https://doi.org/10.2105/AJPH.2008.144097>. [PubMed: 19910357]
20. Mejia AB, Ling PM. Tobacco industry consumer research on smokeless tobacco users and product development. *Am J Public Health*. 2010; 100(1):78–87. <https://doi.org/10.2105/AJPH.2008.152603>. [PubMed: 19910355]
21. Kostygina G, Ling PM. Tobacco industry use of flavourings to promote smokeless tobacco products. *Tob Control*. 2016; 25(suppl 2):ii40–ii49. <https://doi.org/10.1136/tobaccocontrol-2016-053212>. [PubMed: 27856998]
22. Cortese DK, Ling PM. Enticing the new lad: masculinity as a product of consumption in tobacco industry-developed lifestyle magazines. *Men Masc*. 2011; 14(1):4–30. <https://doi.org/10.1177/1097184X09352177>. [PubMed: 21687813]
23. Balbach ED, Gasior RJ, Barbeau EM. R.J. Reynolds’ targeting of African Americans: 1988–2000. *Am J Public Health*. 2003; 93(5):822–827. <https://doi.org/10.2105/AJPH.93.5.822>. [PubMed: 12721151]
24. Yerger VB, Malone RE. African American leadership groups: smoking with the enemy. *Tob Control*. 2002; 11(4):336–345. <https://doi.org/10.1136/tc.11.4.336>. [PubMed: 12432159]
25. Yerger VB, Przewoznik J, Malone RE. Racialized geography, corporate activity, and health disparities: tobacco industry targeting of inner cities. *J Health Care Poor Underserved*. 2007; 18(4 suppl):10–38. <https://doi.org/10.1353/hpu.2007.0120>. [PubMed: 18065850]
26. Yerger VB, McCandless PM. Menthol sensory qualities and smoking topography: a review of tobacco industry documents. *Tob Control*. 2011; 20(suppl 2):ii37–ii43. <https://doi.org/10.1136/tc.2010.041988>. [PubMed: 21504930]
27. Iglesias-Rios L, Parascandola M. A historical review of R.J. Reynolds’ strategies for marketing tobacco to Hispanics in the United States. *Am J Public Health*. 2013; 103(5):e15–e27. <https://doi.org/10.2105/AJPH.2013.301256>.
28. Muggli ME, Pollay RW, Lew R, Joseph AM. Targeting of Asian Americans and Pacific Islanders by the tobacco industry: results from the Minnesota Tobacco Document Depository. *Tob Control*. 2002; 11(3):201–209. <https://doi.org/10.1136/tc.11.3.201>. [PubMed: 12198269]

29. Barbeau EM, Leavy-Sperounis A, Balbach ED. Smoking, social class, and gender: what can public health learn from the tobacco industry about disparities in smoking? *Tob Control*. 2004; 13(2): 115–120. <https://doi.org/10.1136/tc.2003.006098>. [PubMed: 15175523]
30. Smith EA, Malone RE. Tobacco promotion to military personnel: “the plums are here to be plucked”. *Mil Med*. 2009; 174(8):797–806. <https://doi.org/10.7205/MILMED-D-04-4108>. [PubMed: 19743733]
31. Smith EA, Malone RE. “Everywhere the soldier will be”: wartime tobacco promotion in the U.S. military. *Am J Public Health*. 2009; 99(9):1595–1602. <https://doi.org/10.2105/AJPH.2008.152983>. [PubMed: 19608945]
32. Joseph AM, Muggli M, Pearson KC, Lando H. The cigarette manufacturers’ efforts to promote tobacco to the U.S. military. *Mil Med*. 2005; 170(10):874–880. <https://doi.org/10.7205/MILMED.170.10.874>. [PubMed: 16435763]
33. Stevens P, Carlson LM, Hinman JM. An analysis of tobacco industry marketing to lesbian, gay, bisexual, and transgender (LGBT) populations: strategies for mainstream tobacco control and prevention. *Health Promot Pract*. 2004; 5(3 suppl):129s–134s. <https://doi.org/10.1177/1524839904264617>. [PubMed: 15231106]
34. Smith EA, Thomson K, Offen N, Malone RE. “If you know you exist, it’s just marketing poison”: meanings of tobacco industry targeting in the lesbian, gay, bisexual, and transgender community. *Am J Public Health*. 2008; 98(6):996–1003. <https://doi.org/10.2105/AJPH.2007.118174>. [PubMed: 18445800]
35. Smith EA, Malone RE. The outing of Philip Morris: advertising tobacco to gay men. *Am J Public Health*. 2003; 93(6):988–993. <https://doi.org/10.2105/AJPH.93.6.988>. [PubMed: 12773366]
36. Apollonio DE, Malone RE. Marketing to the marginalised: tobacco industry targeting of the homeless and mentally ill. *Tob Control*. 2005; 14(6):409–415. <https://doi.org/10.1136/tc.2005.011890>. [PubMed: 16319365]
37. Lambert A, Sargent JD, Glantz SA, Ling PM. How Philip Morris unlocked the Japanese cigarette market: lessons for global tobacco control. *Tob Control*. 2004; 13(4):379–387. <https://doi.org/10.1136/tc.2004.008441>. [PubMed: 15564622]
38. Bansal R, John S, Ling PM. Cigarette advertising in Mumbai, India: targeting different socioeconomic groups, women, and youth. *Tob Control*. 2005; 14(3):201–206. <https://doi.org/10.1136/tc.2004.010173>. [PubMed: 15923471]
39. Hafez N, Ling PM. Finding the Kool Mixx: how Brown & Williamson used music marketing to sell cigarettes. *Tob Control*. 2006; 15(5):359–366. <https://doi.org/10.1136/tc.2005.014258>. [PubMed: 16998169]
40. Braun S, Mejia R, Ling PM, Perez-Stable EJ. Tobacco industry targeting youth in Argentina. *Tob Control*. 2008; 17(2):111–117. <https://doi.org/10.1136/tc.2006.018481>. [PubMed: 18299308]
41. Hafez N, Ling PM. How Philip Morris built Marlboro into a global brand for young adults: implications for international tobacco control. *Tob Control*. 2005; 14(4):262–271. <https://doi.org/10.1136/tc.2005.011189>. [PubMed: 16046690]
42. Lee S, Ling PM, Glantz SA. The vector of the tobacco epidemic: tobacco industry practices in low and middle-income countries. *Cancer Causes Control*. 2012; 23(suppl 1):117–129. <https://doi.org/10.1007/s10552-012-9914-0>. [PubMed: 22370696]
43. Lee YO, Jordan JW, Djakaria M, Ling PM. Using peer crowds to segment Black youth for smoking intervention. *Health Promot Pract*. 2014; 15(4):530–537. <https://doi.org/10.1177/1524839913484470>. [PubMed: 23628591]
44. Sussman S, Pokhrel P, Ashmore RD, Brown BB. Adolescent peer group identification and characteristics: a review of the literature. *Addict Behav*. 2007; 32(8):1602–1627. <https://doi.org/10.1016/j.addbeh.2006.11.018>. [PubMed: 17188815]
45. Lisha NE, Jordan JW, Ling PM. Peer crowd affiliation as a segmentation tool for young adult tobacco use. *Tob Control*. 2016; 25(suppl 1):i83–i89. <https://doi.org/10.1136/tobaccocontrol-2016-053086>. [PubMed: 27697952]
46. U.S. Department of Commerce, Economics and Statistics Administration, U.S. Census Bureau. [Accessed October 31, 2016] The American Community Survey. [www2.census.gov/programs-surveys/acs/methodology/questionnaires/2014/quest14.pdf](http://www2.census.gov/programs-surveys/acs/methodology/questionnaires/2014/quest14.pdf) Published 2014

47. Ling PM, Neilands TB, Glantz SA. Young adult smoking behavior: a national survey. *Am J Prev Med.* 2009; 36(5):389–394. <https://doi.org/10.1016/j.amepre.2009.01.028>. [PubMed: 19269128]
48. Belstock SA, Connolly GN, Carpenter CM, Tucker L. Using alcohol to sell cigarettes to young adults: a content analysis of cigarette advertisements. *J Am Coll Health.* 2008; 56(4):383–389. <https://doi.org/10.3200/JACH.56.44.383-390>. [PubMed: 18316281]
49. Jiang N, Ling PM. Impact of alcohol use and bar attendance on smoking and quit attempts among young adult bar patrons. *Am J Public Health.* 2013; 103(5):e53–e61. <https://doi.org/10.2105/AJPH.2012.301014>. [PubMed: 23488485]
50. Lopez-Quintero C, Perez de los Cobos J, Hasin DS, et al. Probability and predictors of transition from first use to dependence on nicotine, alcohol, cannabis, and cocaine: results of the National Epidemiologic Survey on Alcohol and Related Conditions (NESARC). *Drug Alcohol Depend.* 2011; 115(1–2):120–130. <https://doi.org/10.1016/j.drugalcdep.2010.11.004>. [PubMed: 21145178]
51. Hymowitz N, Cummings KM, Hyland A, Lynn WR, Pechacek TF, Hartwell TD. Predictors of smoking cessation in a cohort of adult smokers followed for five years. *Tob Control.* 1997; 6(suppl 2):S57–S62. [https://doi.org/10.1136/tc.6.suppl\\_2.S57](https://doi.org/10.1136/tc.6.suppl_2.S57).
52. Lindsay J. Young Australians and the staging of intoxication and self-control. *J Youth Stud.* 2009; 12(4):371–384. <https://doi.org/10.1080/13676260902866520>.
53. Palamar JJ, Acosta P, Sherman S, Ompad DC, Cleland CM. Self-reported use of novel psychoactive substances among attendees of electronic dance music venues. *Am J Drug Alcohol Abuse.* 2016; 42(6):624–632. <https://doi.org/10.1080/00952990.2016.1181179>. [PubMed: 27315522]
54. Droste N, Miller P, Pennay A, Zinkiewicz L, Lubman DI. Environmental contexts of combined alcohol and energy drink use: associations with intoxication in licensed venues. *Int J Drug Policy.* 2016; 36:58–66. <https://doi.org/10.1016/j.drugpo.2016.06.016>. [PubMed: 27497238]
55. Perry MJ, Solomon LJ, Winett RA, et al. High risk sexual behavior and alcohol consumption among bar-going gay men. *AIDS.* 1994; 8(9):1321–1324. <https://doi.org/10.1097/00002030-199409000-00016>. [PubMed: 7802987]
56. Casswell S, Pledger M, Pratap S. Trajectories of drinking from 18 to 26 years: identification and prediction. *Addiction.* 2002; 97(11):1427–1437. <https://doi.org/10.1046/j.1360-0443.2002.00220.x>. [PubMed: 12410783]
57. Zeka A, Gore R, Kriebel D. Effects of alcohol and tobacco on aerodigestive cancer risks: a meta-regression analysis. *Cancer Causes Control.* 2003; 14(9):897–906. <https://doi.org/10.1023/B:CACO.0000003854.34221.a8>. [PubMed: 14682447]
58. Prabhu A, Obi KO, Rubenstein JH. The synergistic effects of alcohol and tobacco consumption on the risk of esophageal squamous cell carcinoma: a meta-analysis. *Am J Gastroenterol.* 2014; 109(6):822–827. <https://doi.org/10.1038/ajg.2014.71>. [PubMed: 24751582]
59. Pelucchi C, Gallus S, Garavello W, Bosetti C, La Vecchia C. Alcohol and tobacco use, and cancer risk for upper aerodigestive tract and liver. *Eur J Cancer Prev.* 2008; 17(4):340–344. <https://doi.org/10.1097/CEJ.0b013e3282f75e91>. [PubMed: 18562959]
60. Kalkhoran S, Lisha NE, Neilands TB, Jordan JW, Ling PM. Evaluation of bar and nightclub intervention to decrease young adult smoking in New Mexico. *J Adolesc Health.* 2016; 59(2):222–229. <https://doi.org/10.1016/j.jadohealth.2016.04.003>. [PubMed: 27265423]
61. Fallin A, Neilands TB, Jordan JW, Hong JS, Ling PM. Wreaking “havoc” on smoking: social branding to reach young adult “partiers” in Oklahoma. *Am J Prev Med.* 2015; 48(1 suppl 1):S78–S85. <https://doi.org/10.1016/j.amepre.2014.09.008>. [PubMed: 25528713]
62. Ling PM, Lee YO, Hong J, Neilands TB, Jordan JW, Glantz SA. Social branding to decrease smoking among young adults in bars. *Am J Public Health.* 2014; 104(4):751–760. <https://doi.org/10.2105/AJPH.2013.301666>. [PubMed: 24524502]
63. Moran MB, Sussman S. Changing attitudes toward smoking and smoking susceptibility through peer crowd targeting: more evidence from a controlled study. *Health Commun.* 2015; 30(5):521–524. <https://doi.org/10.1080/10410236.2014.902008>. [PubMed: 25204200]
64. Fallin A, Neilands TB, Jordan JW, Ling PM. Social branding to decrease lesbian, gay, bisexual, and transgender young adult smoking. *Nicotine Tob Res.* 2015; 17(8):983–989. <https://doi.org/10.1093/ntr/ntu265>. [PubMed: 26180223]

65. Gardiner PS. The African Americanization of menthol cigarette use in the United States. *Nicotine Tob Res.* 2004; 6(suppl 1):S55–S65. <https://doi.org/10.1080/14622200310001649478>. [PubMed: 14982709]
66. Sessa FM. Peer crowds in a commuter college sample: the relation between self-reported alcohol use and perceived peer crowd norms. *J Psychol.* 2007; 141(3):293–305. <https://doi.org/10.3200/JRLP.141.3.293-306>. [PubMed: 17564259]
67. Daddis C. Adolescent peer crowds and patterns of belief in the boundaries of personal authority. *J Adolesc.* 2010; 33(5):699–708. <https://doi.org/10.1016/j.adolescence.2009.11.001>. [PubMed: 19939443]
68. Van Zalk N, Van Zalk MH, Kerr M. Socialization of social anxiety in adolescent crowds. *J Abnorm Child Psychol.* 2011; 39(8):1239–1249. <https://doi.org/10.1007/s10802-011-9533-3>. [PubMed: 21695445]
69. Cross JR, Fletcher KL. The challenge of adolescent crowd research: defining the crowd. *J Youth Adolesc.* 2009; 38(6):747–764. <https://doi.org/10.1007/s10964-008-9307-6>. [PubMed: 19636778]
70. Pokhrel P, Brown BB, Moran MB, Sussman S. Comments on adolescent peer crowd affiliation: a response to Cross and Fletcher (2009). *J Youth Adolesc.* 2010; 39(2):213–216. <https://doi.org/10.1007/s10964-009-9454-4>. [PubMed: 19774454]



**Figure 1.** Smoking prevalence by peer crowd in total sample and among bar patrons.

**Table 1**

Sample Characteristics, 2014 Bay Area Young Adult Health Survey

Variable	% (SD) or M
Unweighted population, n	1,344
Weighted population, n	306,625
Weighted population with peer crowd scores, n	259,407
Bar-going population	107,274
Current smoker	15.2 (0.36)
Peer crowds	
Young Professional (ref)	36.5 (0.48)
Hipster	8.0 (0.27)
Partier	8.9 (0.28)
Hip Hop	8.2 (0.27)
Homebody	35.4 (0.48)
Country	5.8 (0.23)
Age	22.7 (2.47)
Male	50.0 (0.50)
Race/ethnicity	
NH white (ref)	31.9 (0.47)
Hispanic/Latino	23.8 (0.43)
NH black	10.1 (0.30)
NH API	27.7 (0.45)
NH other	6.5 (0.25)
LGBT	10.3 (0.30)
College (currently enrolled or graduate)	83.2 (0.37)
Perceived prevalence of smoking	34.1 (0.21)
Agrees that smoking reduces stress	2.4 (1.72)
Anti-industry attitudes	4.1 (1.86)
Attends bars sometimes/frequently	35.3 (0.48)
Binge drinking in past month	30.0 (0.46)
Very good/excellent self-rated health	59.4 (0.49)

API, Asian/Pacific Islander; LGBT, lesbian, gay, bisexual, transgender; NH, non-Hispanic.

Population Represented by Each Peer Crowd in the Total Sample and Among Bar Patrons

Table 2

Peer crowd	Estimated proportion of population represented by each peer crowd among total population				Estimated proportion of population by peer crowd among bar patrons				
	%	Population representation	Smoking rate, %	Smokers	%	Population representation	Smoking rate, %	Smokers	Smokers attending bars, %
Young Professional	37.1	96,181	8.2	7,839	43.0	40,219	10.3	4,155	53.0
Homebody	34.2	88,608	14.2	12,565	21.4	20,038	23.4	4,697	37.4
Partier	8.9	23,022	22.6	5,196	9.8	9,135	40.1	3,667	70.6
Hip Hop	8.4	21,779	39.2	8,546	12.0	11,201	50.3	5,632	65.9
Hipster	7.1	18,407	22.8	4,200	8.4	7,799	24.7	1,924	45.8
Country	4.4	11,410	35.0	3,996	5.4	5,058	56.8	2,874	71.9
Total	100.0	259,407		42,342	100.0	93,450		22,948	54.2

**Table 3**

Peer Crowd Affiliation and Current Smoking Among Bay Area Young Adults, 2014

Variable	Logistic regression results, current smoker (imputed sample, n=1,344)			
	Model 1		Model 2	
	AOR (95% CI)	SE	AOR (95% CI)	SE
Peer crowd affiliation				
Young professional (ref)			—	—
Hipster	2.56 (0.98, 6.72)	1.26	2.12 (0.75, 6.01)	1.13
Partier	2.34 (0.90, 6.06)	1.13	1.04 (0.40, 2.73)	0.51
Hip Hop	<b>5.58 (2.24, 13.88***)</b>	<b>2.59</b>	<b>4.42 (1.51, 12.98**)</b>	<b>2.43</b>
Homebody	1.57 (0.80, 3.10)	0.54	1.71 (0.82, 3.55)	0.64
Country	<b>3.55 (1.37, 9.19**)</b>	<b>1.72</b>	<b>3.06 (1.17, 7.99*)</b>	<b>1.50</b>
Sociodemographic characteristics				
Age	0.99 (0.89, 1.12)	0.06	0.89 (0.78, 1.02)	0.06
Male	<b>1.67 (1.02, 2.73*)</b>	<b>0.42</b>	1.10 (0.58, 2.12)	0.37
Race/ethnicity				
NH white (ref)			—	—
Hispanic/Latino	0.74 (0.39, 1.39)	0.24	0.55 (0.27, 1.09)	0.19
NH black	0.75 (0.30, 1.87)	0.35	0.63 (0.19, 2.08)	0.38
NH API	<b>0.41 (0.21, 0.80**)</b>	<b>0.14</b>	<b>0.41 (0.19, 0.89*)</b>	<b>0.16</b>
NH other	0.59 (0.13, 2.64)	0.45	0.26 (0.03, 1.97)	0.27
LGBT	1.07 (0.50, 2.30)	0.42	0.60 (0.21, 1.69)	0.32
College	<b>0.47 (0.27, 0.84*)</b>	<b>0.14</b>	<b>0.45 (0.24, 0.85*)</b>	<b>0.15</b>
Tobacco attitudes and health status				
Perceived prevalence of smoking			1.77 (0.54, 5.77)	1.07
Agrees that smoking reduces stress			<b>1.59 (1.36, 1.86***)</b>	<b>0.13</b>
Anti-industry attitudes			<b>0.76 (0.63, 0.91**)</b>	<b>0.07</b>
Attends bars sometimes/frequently			2.08 (0.98, 4.43)	0.80
Binge drinking in past month			<b>3.17 (1.59, 6.32***)</b>	<b>1.12</b>



Logistic regression results, current smoker (imputed sample, $n=1,344$ )				
Variable	Model 1		Model 2	
	AOR (95% CI)	SE	AOR (95% CI)	SE
Very good/excellent self-rated health			<b>0.33 (0.18, 0.58***)</b>	<b>0.10</b>

Note: Boldface indicates statistical significance (\* $p<0.05$ ; \*\* $p<0.01$ ; \*\*\* $p<0.001$ ).

API, Asian/Pacific Islander; LGBT, lesbian, gay, bisexual, transgender; NH, non-Hispanic.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript