



## RESEARCH PAPER

# Competition between cyanobacteria and green algae at low versus elevated CO<sub>2</sub>: who will win, and why?

Xing Ji, Jolanda M.H. Verspagen, Maayke Stomp and Jef Huisman\*

Department of Aquatic Microbiology, Institute for Biodiversity and Ecosystem Dynamics, University of Amsterdam, PO Box 94248, 1090 GE Amsterdam, The Netherlands

\* Correspondence: [j.huisman@uva.nl](mailto:j.huisman@uva.nl)

Received 7 November 2016; Editorial decision 11 January 2017; Accepted 12 January 2017

Editor: Howard Griffiths, University of Cambridge

## Abstract

Traditionally, it has often been hypothesized that cyanobacteria are superior competitors at low CO<sub>2</sub> and high pH in comparison with eukaryotic algae, owing to their effective CO<sub>2</sub>-concentrating mechanism (CCM). However, recent work indicates that green algae can also have a sophisticated CCM tuned to low CO<sub>2</sub> levels. Conversely, cyanobacteria with the high-flux bicarbonate uptake system BicA appear well adapted to high inorganic carbon concentrations. To investigate these ideas we studied competition between three species of green algae and a *bicA* strain of the harmful cyanobacterium *Microcystis aeruginosa* at low (100 ppm) and high (2000 ppm) CO<sub>2</sub>. Two of the green algae were competitively superior to the cyanobacterium at low CO<sub>2</sub>, whereas the cyanobacterium increased its competitive ability with respect to the green algae at high CO<sub>2</sub>. The experiments were supported by a resource competition model linking the population dynamics of the phytoplankton species with dynamic changes in carbon speciation, pH and light. Our results show (i) that competition between phytoplankton species at different CO<sub>2</sub> levels can be predicted from species traits in monoculture, (ii) that green algae can be strong competitors under CO<sub>2</sub>-depleted conditions, and (iii) that bloom-forming cyanobacteria with high-flux bicarbonate uptake systems will benefit from elevated CO<sub>2</sub> concentrations.

**Key words:** Carbon dioxide, climate change, competition model, CO<sub>2</sub>-concentrating mechanism, cyanobacteria, green algae, harmful algal blooms, lakes, *Microcystis aeruginosa*.

## Introduction

Cyanobacterial blooms have become a major water quality problem in many eutrophic lakes worldwide (Chorus and Bartram, 1999; Verspagen *et al.*, 2006; Guo, 2007; Michalak *et al.*, 2013). They produce taste and odor compounds that may interfere with the recreational function of lakes and the preparation of drinking water (Chorus and Bartram, 1999; Watson *et al.*, 2008). Moreover, cyanobacteria can produce a variety of toxins, causing liver, digestive and neurological diseases when ingested by waterfowl, pets, cattle, and humans

(Carmichael, 2001; Codd *et al.*, 2005; Huisman *et al.*, 2005; Merel *et al.*, 2013). Hence, an improved understanding of the environmental conditions that favor the dominance of cyanobacteria over eukaryotic phytoplankton species is desirable.

Dense cyanobacterial blooms often deplete the dissolved CO<sub>2</sub> concentration in surface waters, sometimes down to less than 0.1 μmol l<sup>-1</sup> corresponding to pCO<sub>2</sub> less than 3 parts per million (ppm) (Lazzarino *et al.*, 2009; Balmer and Downing, 2011). CO<sub>2</sub> depletion by dense blooms induces

Abbreviations: CCM, CO<sub>2</sub>-concentrating mechanism; C<sub>i</sub>, inorganic carbon; CA, carbonic anhydrase; DIC, dissolved inorganic carbon; RuBisCO, ribulose-1,5-bisphosphate carboxylase/oxygenase; ppm, parts per million.

© The Author 2017. Published by Oxford University Press on behalf of the Society for Experimental Biology.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted reuse, distribution, and reproduction in any medium, provided the original work is properly cited.

high pH values, above 9 or even 10 (Talling, 1976; Ibelings and Maberly, 1998; Verspagen *et al.*, 2014b). At these pH values, most dissolved inorganic carbon (DIC) is in the form of bicarbonate, and with increasing pH an increasing fraction of DIC is converted to carbonate. Under these conditions, CO<sub>2</sub> availability can become an important limiting factor for photosynthesis. Cyanobacteria have developed a highly efficient CO<sub>2</sub>-concentrating mechanism (CCM) to take up CO<sub>2</sub> and bicarbonate as inorganic carbon (C<sub>i</sub>) source, and to augment the intracellular CO<sub>2</sub> level around the ribulose-1,5-bisphosphate carboxylase/oxygenase (RuBisCO) enzyme responsible for carbon fixation (Price *et al.*, 2008; Raven *et al.*, 2012; Burnap *et al.*, 2015). It has therefore been hypothesized that cyanobacteria are superior competitors at low CO<sub>2</sub> levels, and will dominate waters in which the dissolved CO<sub>2</sub> concentration has been depleted (Shapiro, 1990, 1997). Conversely, eukaryotic phytoplankton might be better competitors at high CO<sub>2</sub> levels. This classic paradigm has received support from several competition experiments between cyanobacteria and green algae (Caraco and Miller, 1998; Low-Décarie *et al.*, 2011, 2015; but see Verschoor *et al.*, 2013). If this paradigm is true, the logical corollary is that rising CO<sub>2</sub> levels will particularly benefit eukaryotic phytoplankton species at the expense of cyanobacteria.

Yet, recent insights indicate that this classic paradigm might be too simple. Although the details of the eukaryotic CCM are not yet fully understood, green algae can also deploy a sophisticated CCM well adapted to low CO<sub>2</sub> levels (Moroney and Ynalvez, 2007; Wang *et al.*, 2011; Meyer and Griffiths, 2013). Furthermore, recent studies have revealed a striking genetic and phenotypic diversity in the CCM of harmful cyanobacteria (Sandrini *et al.*, 2014, 2015b; Visser *et al.*, 2016). All harmful freshwater cyanobacteria investigated so far contain two CO<sub>2</sub> uptake systems and the ATP-dependent bicarbonate uptake system BCT1. In addition, however, some cyanobacteria deploy the high-affinity but low-flux bicarbonate uptake system SbtA, whereas other cyanobacteria deploy the low-affinity but high-flux bicarbonate uptake system BicA (Sandrini *et al.*, 2014). Therefore, three C<sub>i</sub> uptake genotypes can be distinguished: (i) *sbtA* strains (high-affinity specialists), (ii) *bicA* strains (high-flux specialists), and (iii) *bicA+sbtA* strains (CCM generalists). These three genotypes were first described for the genus *Microcystis* (Sandrini *et al.*, 2014, 2016), but similar genetic diversity also exists within other harmful cyanobacterial genera such as *Dolichospermum* (formerly known as *Anabaena*) and *Planktothrix* (Visser *et al.*, 2016).

Laboratory selection experiments with mixtures of several *Microcystis* strains found that the strain composition shifted from *bicA+sbtA* strains at low pCO<sub>2</sub> to *bicA* strains at high pCO<sub>2</sub> (Sandrini *et al.*, 2016). Similarly, in a eutrophic lake, *bicA+sbtA* strains were dominant when C<sub>i</sub> concentrations in the lake were depleted during a dense cyanobacterial bloom, but were replaced by *bicA* strains when C<sub>i</sub> concentrations increased later in the season (Sandrini *et al.*, 2016). These results show that the genetic composition of cyanobacterial communities adapts to changes in C<sub>i</sub> availability by means of natural selection, favoring CCM generalists at low CO<sub>2</sub> levels

while favoring high-flux specialists at high CO<sub>2</sub>. In natural waters, however, cyanobacteria compete not only amongst each other, but also against eukaryotic phytoplankton. How the competitive abilities of different cyanobacterial C<sub>i</sub> uptake genotypes perform against eukaryotic species such as green algae has not yet been investigated.

Resource competition theory provides a theoretical framework to understand and predict how changes in resource availability may affect the species composition (Tilman, 1982; Huisman and Weissing, 1994; Grover, 1997). This body of theory uses the kinetic traits of species measured in monoculture to predict the dynamics and outcome of competition for limiting resources in species mixtures. Resource competition models have been successfully applied to predict competition for nutrients and light, both in qualitative and quantitative terms (Tilman, 1977; Sommer, 1985; Huisman *et al.*, 1999; Litchman *et al.*, 2004; Stomp *et al.*, 2004; Passarge *et al.*, 2006). Competition for inorganic carbon is conceptually more complicated, however, because the species compete for two resources (CO<sub>2</sub> and bicarbonate) whose concentrations depend not only on resource uptake but also on pH and alkalinity. Moreover, pH and alkalinity depend in turn on a variety of biogeochemical processes and also change in response to the carbon and nutrient uptake activity by the phytoplankton community (Talling, 1976; Wolf-Gladrow *et al.*, 2007; Verspagen *et al.*, 2014b).

This study investigated competition between a harmful cyanobacterium (*Microcystis* PCC 7806) and three species of green algae (*Monoraphidium griffithii*, *Scenedesmus obliquus*, and *Chlorella vulgaris*) at low and at high CO<sub>2</sub> concentrations. *Microcystis* PCC 7806 is a *bicA* strain (Sandrini *et al.*, 2014), and is therefore expected to be a relatively weak competitor at low CO<sub>2</sub> levels but a stronger competitor at high CO<sub>2</sub>. To investigate this hypothesis, we ran monoculture experiments of each species at both low and high CO<sub>2</sub> levels (100 and 2000 ppm). The results of these monoculture experiments were used to parameterize a resource competition model, which predicted the competitive interactions between the species based on dynamic changes in inorganic carbon chemistry, alkalinity and pH induced by the growing phytoplankton populations. Next, competition experiments were carried out to test the model predictions. Together, the theory and experiments may help in understanding shifts in phytoplankton community composition in response to rising CO<sub>2</sub> levels.

## Competition model: theory and development

We first developed a model to investigate competition for inorganic carbon and light among phytoplankton species. The model combined previous theoretical work on growth and competition under light-limited (Huisman and Weissing, 1994; Huisman *et al.*, 1999; Passarge *et al.*, 2006) and carbon-limited conditions (Van de Waal *et al.*, 2011; Verspagen *et al.*, 2014a, b).

The model considers a well-mixed vertical water column, where depth  $z$  runs from 0 at the water surface to  $z_{\max}$  at the

bottom of the water column. The population dynamics of the phytoplankton species are governed by their light-dependent assimilation of carbon dioxide and bicarbonate. The model assumes eutrophic conditions, in which all nutrients are in ample supply and hence do not limit phytoplankton growth. Uptake of inorganic carbon and nutrients induces dynamic changes in pH and alkalinity. These changes in pH and alkalinity affect the availability of the different inorganic carbon species, which feeds back on phytoplankton growth. The growing phytoplankton populations also increase the turbidity of the water column, thereby diminishing the light available for further photosynthesis and growth.

### Species dynamics

The model assumes that the specific growth rates of the species depend on their carbon assimilation. Let  $X_i$  denote the population density of phytoplankton species  $i$ , and let  $Q_i$  denote its cellular carbon content (also known as carbon quota; sensu [Droop, 1973](#)). The population dynamics of a number of  $n$  competing species can then be written as:

$$\frac{dX_i}{dt} = \mu_i(Q_i)X_i - m_iX_i \quad i = 1, \dots, n \quad (1)$$

where  $\mu_i(Q_i)$  is the specific growth rate of species  $i$  as an increasing function of its carbon content, and  $m_i$  is its loss rate. We assume that each species requires a minimum cellular carbon content in order to function, and reaches its maximum specific growth rate when cells are satiated with carbon (see Supplementary Model S1 at *JXB* online). In our application, the loss rates of the species will be governed by the dilution rate of the chemostat (i.e.  $m_i = D$ ).

The carbon contents of the species increase through uptake of carbon dioxide ( $u_{\text{CO}_2,i}$ ) and bicarbonate ( $u_{\text{HCO}_3,i}$ ), and decrease through respiration ( $r_i$ ) and dilution of the carbon content by growth:

$$\frac{dQ_i}{dt} = u_{\text{CO}_2,i} + u_{\text{HCO}_3,i} - r_i - \mu_i(Q_i)Q_i \quad i = 1, \dots, n \quad (2)$$

We assume that uptake rates of  $\text{CO}_2$  ( $u_{\text{CO}_2,i}$ ) and bicarbonate ( $u_{\text{HCO}_3,i}$ ) are increasing functions of the ambient  $\text{CO}_2$  and bicarbonate concentration according to Michaelis–Menten kinetics. Carbon uptake and assimilation require energy from the light reactions of photosynthesis, and therefore the carbon uptake rates also depend on the photosynthetic activity of the cells and hence on light availability. Furthermore, we incorporated a simple negative feedback loop in which carbon uptake rates decrease with increasing cellular carbon content, such that carbon uptake systems are active under carbon-limiting conditions ([Eisenhut et al., 2007](#); [Burnap et al., 2015](#); [Wang et al., 2015](#)) and down-regulated when cells become satiated with carbon ([Beardall and Giordano, 2002](#); [Sandrini et al., 2015a](#)). Respiration rates ( $r_i$ ) of the species increase with their cellular carbon content, approaching maximum values when cells become satiated with carbon. The mathematical equations describing these relationships are presented in Supplementary Model S1.

### Light conditions

The underwater light gradient is described by the Lambert–Beer law ([Huisman et al., 1999](#)):

$$I(z) = I_{\text{in}} \exp(-K_{\text{bg}}z - \sum_{i=1}^n k_i X_i z) \quad (3)$$

where  $I(z)$  is the light intensity at depth  $z$ ,  $I_{\text{in}}$  is the incident light intensity,  $K_{\text{bg}}$  is the background turbidity of the water itself, and  $k_i$  is the specific light attenuation coefficient of phytoplankton species  $i$ . We note that the light gradient changes dynamically, because the light intensity at a given depth decreases with increasing phytoplankton densities. We define  $I_{\text{out}}$  as the light intensity reaching the bottom of the water column (i.e.  $I_{\text{out}} = I(z_{\text{max}})$ ).

### Dissolved inorganic carbon

Changes in the concentration of total dissolved inorganic carbon, [DIC], are described by ([Verspagen et al., 2014b](#)):

$$\frac{d[\text{DIC}]}{dt} = D([\text{DIC}]_{\text{in}} - [\text{DIC}]) + \frac{g_{\text{CO}_2}}{z_{\text{max}}} - \sum_{i=1}^n (u_{\text{CO}_2,i} + u_{\text{HCO}_3,i})X_i + \sum_{i=1}^n r_i X_i \quad (4)$$

The first term on the right-hand side of this equation describes changes through the influx ( $[\text{DIC}]_{\text{in}}$ ) and efflux of water containing DIC. The second term describes  $\text{CO}_2$  exchange with the atmosphere, where  $g_{\text{CO}_2}$  is the  $\text{CO}_2$  flux across the air–water interface and division by  $z_{\text{max}}$  converts the flux per unit surface area into the corresponding change in DIC concentration. The third term describes uptake of dissolved  $\text{CO}_2$  and bicarbonate by the photosynthetic activity of the phytoplankton community. Finally, the fourth term describes  $\text{CO}_2$  release by respiration of the phytoplankton species.

The  $\text{CO}_2$  flux across the air–water interface,  $g_{\text{CO}_2}$ , depends on the difference in partial pressure. More specifically,  $g_{\text{CO}_2}$  depends on the difference between the expected concentration of dissolved  $\text{CO}_2$  in water if in equilibrium with the partial pressure in the atmosphere and the actual dissolved  $\text{CO}_2$  concentration ([Siegenthaler and Sarmiento, 1993](#); [Cole et al., 2010](#)):

$$g_{\text{CO}_2} = v(K_0 \text{pCO}_2 - [\text{CO}_2]) \quad (5)$$

where  $v$  is the gas transfer velocity (also known as piston velocity),  $K_0$  is the solubility of  $\text{CO}_2$  gas in water (also known as Henry’s constant),  $\text{pCO}_2$  is the partial pressure of  $\text{CO}_2$  in the atmosphere, and  $[\text{CO}_2]$  is the dissolved  $\text{CO}_2$  concentration. In chemostats, gas transfer will depend on the gas flow rate ( $a$ ). We therefore assume that  $v = ba$ , where  $b$  is a constant of proportionality.

Dissolved  $\text{CO}_2$ , bicarbonate and carbonate concentrations, and pH were calculated from [DIC] and alkalinity ([Stumm and Morgan, 1996](#)). Assimilation of nitrate, phosphate and sulfate by phytoplankton involves proton consumption, thus increasing alkalinity ([Wolf-Gladrow et al., 2007](#); [Verspagen et al., 2014b](#)). Therefore, the model treats alkalinity as a dynamic variable (see Supplementary Model S1 for details).

## Materials and methods

### Species

We performed monoculture and competition experiments with four freshwater phytoplankton species: the green algae *Monoraphidium griffithii* (Berk.) Kom.-Legn. (strain CCAP 202/15A), *Scenedesmus obliquus* (strain CCAP 276/3A) and *Chlorella vulgaris* Beyerinck (strain UTEX 259), and the cyanobacterium *Microcystis aeruginosa* (strain PCC 7806). *Microcystis* PCC 7806 produces the hepatotoxins microcystin-LR and [Asp<sup>3</sup>]-microcystin-LR and the potential neurotoxins cyanopeptolin A, C and 970 (Tonk et al., 2009; Faltermann et al., 2014). All four species were unialgal but not axenic. Regular microscopic inspection confirmed that concentrations of heterotrophic bacteria remained low throughout the experiments (<1% of the total biovolume measured with a CASY TTC cell counter; OLS OMNI Life Science, Bremen, Germany).

### Chemostat experiments

All experiments were conducted in laboratory-built chemostats, specifically designed to study the population dynamics of phytoplankton species (Huisman et al., 2002; Passarge et al., 2006; Verspagen et al., 2014b). The chemostats allowed full control of light conditions, temperature, pCO<sub>2</sub> in the gas flow, and nutrient concentrations in the mineral medium. Each chemostat consisted of a flat culture vessel with an optimal path length ('mixing depth') of  $z_{\max}=5$  cm and a working volume of ~1.7 l. The vessel was illuminated from one side to create a unidirectional light gradient, using a constant incident light intensity ( $I_{\text{in}}$ ) of  $50 \pm 1$   $\mu\text{mol photons m}^{-2} \text{s}^{-1}$  provided by white fluorescent tubes (Philips PL-L 24W/840/4P, Philips Lighting, Eindhoven, The Netherlands). The temperature was maintained at  $20 \pm 1$  °C with a stainless steel cooling finger inside each chemostat and connected to a Colora thermocryostat. To avoid nutrient limitation, the chemostats were provided with a very nutrient-rich mineral medium (Van de Waal et al., 2009; Verspagen et al., 2014a). The dilution rate was maintained at  $D=0.125 \text{ d}^{-1}$ .

### CO<sub>2</sub> supply

We applied two CO<sub>2</sub> treatments. The chemostats were bubbled with gas containing a CO<sub>2</sub> concentration of either 100 ppm ('low pCO<sub>2</sub>') or 2000 ppm ('high pCO<sub>2</sub>') (Table 1). The gas was prepared as a mixture of pressurized air from which the variable CO<sub>2</sub> concentration was completely removed using a CO<sub>2</sub> scrubber (Ecodry K-MT6; Parker Zander, Lancaster, NY, USA) and subsequently a

defined amount of pure CO<sub>2</sub> gas was added to obtain the desired concentration using mass flow controllers (GT 1355R-2-15-A316 SS and 5850S, Brooks Instrument, Hatfield, PA, USA). Before entering the chemostats, the mixed gas was filter sterilized (0.2  $\mu\text{m}$  Midisart 2000 Filter, Sartorius Stedim Biotech GmbH, Göttingen, Germany) and moisturized with Milli-Q water to suppress evaporation from the chemostats. The gas was dispersed as fine bubbles supplied from the bottom of the chemostat vessels at a constant flow rate of  $a=25 \text{ l h}^{-1}$ , which also ensured homogeneous mixing of the phytoplankton populations. We checked the CO<sub>2</sub> concentration in the gas flow regularly using an environmental gas monitor (EGM-4; PP Systems, Amesbury, MA, USA).

### Experimental measurements

The experiments were sampled at least every other day. The incident light intensity ( $I_{\text{in}}$ ) was measured with a LI-COR LI-250 quantum photometer (LI-COR Biosciences, Lincoln, NE, USA) at ten randomly chosen positions at the front surface of the chemostat vessel. Likewise, the light intensity transmitted through the chemostat ( $I_{\text{out}}$ ) was measured at the back surface of the chemostat vessel (Huisman et al., 2002).

Population densities and biovolumes in samples of the monoculture experiments were measured in triplicate with a CASY TTC automated cell counter (OLS OMNI Life Science, Bremen, Germany) using a 60  $\mu\text{m}$  capillary. The cell counter could not distinguish between the different species. Therefore, population densities in the competition experiments were counted on an Accuri C6 flow cytometer (Accuri Cytometers Inc., Ann Arbor, MI, USA), which distinguished the different species in these experiments on the basis of differences in pigmentation and cell size (side scatter). We did not perform competition experiments between *Scenedesmus* and *Chlorella*, because the flow cytometer could not adequately distinguish between these two chlorophytes.

Temperature and pH were measured immediately after sampling, using a SCHOTT pH meter (SCHOTT AG, Mainz, Germany). For DIC analysis, 35 ml samples were transferred to 50 ml falcon tubes, centrifuged for 15 min at 600 g, immediately filtered over 0.45  $\mu\text{m}$  polyethersulfone membrane filters (Sartorius Stedim Biotech GmbH, Göttingen, Germany), transferred to gas-tight urine tubes (Terumo Europe NV, Leuven, Belgium) and stored at 4 °C until analysis. DIC was analysed by phosphoric acid addition using a Model 700 TOC Analyzer (OI Corp., College Station, TX, USA). Concentrations of CO<sub>2</sub>(aq), bicarbonate and carbonate were calculated from DIC and pH, based on the dissociation constants of inorganic carbon corrected for temperature and salinity (Stumm and Morgan, 1996; Verspagen et al., 2014b).

To determine the cellular carbon, nitrogen and sulfur content, pellets from the 50 ml falcon tube were transferred into 2 ml Eppendorf tubes, washed three times with a nutrient-free NaCl solution with a salinity equal to our medium, and stored at -20 °C. Subsequently, the pellets were freeze-dried and weighted, and the carbon, nitrogen and sulfur content of homogenized freeze-dried cell powder were analysed with a Vario EL Elemental Analyzer (Elementar Analysensysteme GmbH, Hanau, Germany).

### Model parameterization

The model parameters consisted of system parameters and species parameters. System parameters are under experimental control and were regularly measured during the experiments. We had already specified several of the system parameters, such as the incident light intensity ( $I_{\text{in}}$ ), pCO<sub>2</sub> level in the gas flow, and dilution rate ( $D$ ) of the chemostats. A complete list of all system parameters is provided in Table 1.

Species parameters describe the traits of the species. Some species parameters were measured experimentally. The maximum specific growth rate ( $\mu_{\max,i}$ ) and minimum cellular carbon content ( $Q_{\min,i}$ ) were determined in batch cultures at an  $I_{\text{in}}$  of  $50 \pm 1$   $\mu\text{mol photons}$

**Table 1.** System parameters applied in the experiments

Parameter	Description	Value	Units
$D$	Dilution rate	0.125	$\text{d}^{-1}$
$z_{\max}$	Mixing depth	0.05	m
$T$	Temperature	20	°C
$I_{\text{in}}$	Incident light intensity	50	$\mu\text{mol photons m}^{-2} \text{s}^{-1}$
$K_{\text{bg}}$	Background turbidity <sup>a</sup>	7–11	$\text{m}^{-1}$
$\text{DIC}_{\text{in}}$	DIC concentration at influx <sup>b</sup>	0.5 and 2.0	$\text{mmol l}^{-1}$
$\text{ALK}_{\text{in}}$	Alkalinity at influx <sup>b</sup>	0.79 and 2.29	$\text{mEq l}^{-1}$
sal	Salinity <sup>b</sup>	1.23 and 1.36	$\text{g l}^{-1}$
pCO <sub>2</sub>	CO <sub>2</sub> concentration in gas flow <sup>b</sup>	100 and 2000	ppm
$a$	Gas flow rate	25	$\text{l h}^{-1}$
$v$	Gas transfer velocity <sup>b</sup>	0.24 and 0.68	$\text{m h}^{-1}$

<sup>a</sup> Background turbidity varied among the chemostat vessels.

<sup>b</sup> The first value refers to the low pCO<sub>2</sub> experiments and the second value to the high pCO<sub>2</sub> experiments.

$\text{m}^{-2} \text{s}^{-1}$  and a temperature of  $20 \pm 1^\circ \text{C}$ . Maximum specific growth rate was measured in batch cultures aerated with gas containing a saturating  $\text{CO}_2$  concentration of 10 000 ppm. The minimum cellular carbon content was measured as the cellular carbon content in unaerated dense batch cultures that were first grown for a day in medium to which no DIC was added, and were subsequently incubated overnight in the dark. The cellular N:C and S:C ratios ( $c_{\text{N},i}$  and  $c_{\text{S},i}$ ) were calculated from the cellular carbon, nitrogen and sulfur contents measured in the steady-state monocultures of the species. The specific light attenuation coefficients ( $k_i$ ) of the species were estimated from the monoculture experiments using the Lambert–Beer law. For monocultures, Eqn (3) can be written as  $\ln(I_{\text{in}}/I_{\text{out}})/z_{\text{max}} = K_{\text{bg}} + k_i X_i$ . Hence, the specific light attenuation coefficient ( $k_i$ ) was estimated as the slope of a linear regression of  $\ln(I_{\text{in}}/I_{\text{out}})/z_{\text{max}}$  versus the population density  $X_i$ , and the background turbidity ( $K_{\text{bg}}$ ) was estimated as the intercept.

All other species parameters were estimated by fitting the model predictions to the observed dynamics in the monoculture experiments. More specifically, we fitted the time courses of population density, light transmission ( $I_{\text{out}}$ ), pH and inorganic carbon concentrations predicted by the model to the time courses measured in the monoculture experiments, following the same methodology as in earlier studies (Huisman *et al.*, 1999; Passarge *et al.*, 2006; Verspagen *et al.*, 2014b). To avoid overfitting, the model was fitted simultaneously to both the low  $\text{pCO}_2$  and high  $\text{pCO}_2$  monocultures, resulting in eight parameter estimates per species.


The species parameters obtained from the monoculture experiments were combined with the system parameters to predict the population dynamics and inorganic carbon chemistry in the competition experiments.

In the competition experiments, we calculated the rate of competitive displacement (RCD) from the slope of the linear regression of  $\ln(X_1/X_2)$  versus time, where  $X_1$  and  $X_2$  are the population densities of the two competing species (Grover, 1991; Passarge *et al.*, 2006).

## Results

### Monoculture experiments at low $\text{pCO}_2$

At low  $\text{pCO}_2$ , the phytoplankton species increased until a steady state was reached with population densities (expressed as biovolumes) ranging from  $370 \text{ mm}^3 \text{ l}^{-1}$  for *Chlorella* to  $720 \text{ mm}^3 \text{ l}^{-1}$  for *Scenedesmus* (Fig. 1A–D; Supplementary Table S1). The growing phytoplankton populations reduced the light intensity penetrating through the chemostats ( $I_{\text{out}}$ ) to  $6.2\text{--}9.3 \mu\text{mol photons m}^{-2} \text{ s}^{-1}$  and depleted the dissolved  $\text{CO}_2$  concentration by several orders of magnitude to  $<0.01 \mu\text{mol l}^{-1}$  (Fig. 1E–H). Bicarbonate concentrations decreased about one order of magnitude, and were offset by a similar increase of the carbonate concentrations, such that the total DIC concentration in the chemostats remained more or less constant. Alkalinity increased to  $1.3\text{--}2.0 \text{ mEq l}^{-1}$  depending on the species (see Supplementary Table S1 at JXB online). The pH increased by more than two units, from initial values of  $\sim 8$  to steady state values of  $\sim 10.7$  (Fig. 1I–L). The relatively high values of  $I_{\text{out}}$  in combination with severe  $\text{CO}_2$  depletion and a



**Fig. 1.** Monoculture experiments at low  $\text{pCO}_2$  (100 ppm). (A–D) Population density (expressed as biovolume) and light intensity  $I_{\text{out}}$  penetrating through the chemostat. (E–H)  $\text{CO}_2(\text{aq})$ , bicarbonate and carbonate concentrations. (I–L) Dissolved inorganic carbon (DIC) and pH. Different panels represent different species: (A, E, I) *Monoraphidium*; (B, F, J) *Microcystis*; (C, G, K) *Scenedesmus*; (D, H, L) *Chlorella*. Symbols indicate experimental data, and lines indicate model fits. Parameter values of the model are provided in Table 1 and Table 2.

**Table 2.** Species parameters estimated from the monoculture experiments

Parameter	Description	<i>Monoraphidium</i>	<i>Microcystis</i>	<i>Scenedesmus</i>	<i>Chlorella</i>	Units
$\mu_{\max}$	Maximum specific growth rate <sup>a</sup>	1.27 ± 0.05	1.04 ± 0.02	1.39 ± 0.05	1.28 ± 0.03	d <sup>-1</sup>
$k$	Specific light attenuation coefficient <sup>a</sup>	$6 \times 10^{-5} \pm 0.7 \times 10^{-6}$	$8 \times 10^{-5} \pm 1.0 \times 10^{-6}$	$6 \times 10^{-5} \pm 1.0 \times 10^{-6}$	$6 \times 10^{-5} \pm 1.0 \times 10^{-6}$	m <sup>2</sup> mm <sup>-3</sup>
$H$	Half-saturation constant for light <sup>b</sup>	50	11	30	48	μmol photons m <sup>-2</sup> s <sup>-1</sup>
$H_{\text{CO}_2}$	Half-saturation constant for CO <sub>2</sub> <sup>b</sup>	12.5	1.0	2.0	2.5	μmol l <sup>-1</sup>
$H_{\text{HCO}_3}$	Half-saturation constant for HCO <sub>3</sub> <sup>-b</sup>	80	20	30	40	μmol l <sup>-1</sup>
$r_{\max}$	Maximum respiration rate <sup>b</sup>	2.2	0.9	2.2	2.2	μmol mm <sup>-3</sup> d <sup>-1</sup>
$u_{\max, \text{CO}_2}$	Maximum uptake rate of CO <sub>2</sub> <sup>b</sup>	17.1	20.1	14.7	22.5	μmol mm <sup>-3</sup> d <sup>-1</sup>
$u_{\max, \text{HCO}_3}$	Maximum uptake rate of HCO <sub>3</sub> <sup>-b</sup>	14.0	10.8	14.7	15.7	μmol mm <sup>-3</sup> d <sup>-1</sup>
$Q_{\min}$	Minimum cellular carbon content <sup>a</sup>	14.0 ± 1.1	9.0 ± 2.7	12.5 ± 1.0	11.5 ± 0.8	μmol mm <sup>-3</sup>
$Q_{\max}$	Maximum cellular carbon content <sup>b</sup>	24.5	29.9	21.1	29.9	μmol mm <sup>-3</sup>
$C_N$	Cellular N:C ratio <sup>a</sup>	0.114 ± 0.002	0.130 ± 0.003	0.125 ± 0.003	0.100 ± 0.002	Molar ratio
$C_P$	Cellular P:C ratio <sup>b</sup>	$7.1 \times 10^{-3}$	$8.1 \times 10^{-3}$	$7.8 \times 10^{-3}$	$6.3 \times 10^{-3}$	Molar ratio
$C_S$	Cellular S:C ratio <sup>a</sup>	$3.8 \times 10^{-3} \pm 0.2 \times 10^{-3}$	$7.5 \times 10^{-3} \pm 0.3 \times 10^{-3}$	$6.0 \times 10^{-3} \pm 0.3 \times 10^{-3}$	$8.2 \times 10^{-3} \pm 0.3 \times 10^{-3}$	Molar ratio

<sup>a</sup> Parameter values measured experimentally, given as mean ± standard error.

<sup>b</sup> Parameter values estimated by fitting the model predictions to time courses of the experiments.

high pH indicate that the phytoplankton growth rates in these experiments were carbon limited.

#### Monoculture experiments at high pCO<sub>2</sub>


At high pCO<sub>2</sub>, the phytoplankton species reached population densities ranging from ~2000 mm<sup>3</sup> l<sup>-1</sup> for *Monoraphidium* to ~2500 mm<sup>3</sup> l<sup>-1</sup> for *Scenedesmus* (Fig. 2A–D; Supplementary Table S1). These values are 3.5–6 times higher than in the low pCO<sub>2</sub> experiments (Fig. 1A–D), thus confirming that phytoplankton growth in the low pCO<sub>2</sub> experiments was indeed carbon limited. The dense phytoplankton populations absorbed almost all incident light, reducing the light intensity penetrating through the chemostats ( $I_{\text{out}}$ ) to <0.2 μmol photons m<sup>-2</sup> s<sup>-1</sup> (Fig. 2A–D). The dissolved CO<sub>2</sub> concentration was only slightly reduced to ~30 μmol l<sup>-1</sup> (Fig. 2E–H). By contrast, the bicarbonate and carbonate concentration and hence also the total DIC concentration increased during the experiments (Fig. 2I–L). The DIC increase was enabled by a rise in alkalinity, from 2.3 mEq l<sup>-1</sup> at the start of the experiments to 6.2–8.4 mEq l<sup>-1</sup> at steady state (see Supplementary Table S1). Alkalinity increased during the experiments, because the high uptake rates of nitrate, phosphate, and sulfate by the growing phytoplankton populations are accompanied by proton consumption to maintain charge balance (Goldman and Brewer, 1980; Wolf-Gladrow et al., 2007). A similar increase in DIC and alkalinity induced by dense phytoplankton populations was also observed in earlier chemostat studies (Verspagen et al., 2014b). The pH increased only slightly from ~8 to ~8.6

(Fig. 2I–L). The very low  $I_{\text{out}}$  values in combination with high dissolved CO<sub>2</sub> and bicarbonate concentrations indicate that the phytoplankton growth rates in these experiments were light limited.

#### Predictions derived from the monoculture experiments

The monoculture experiments showed dynamic changes in population abundances, light conditions, carbon speciation, alkalinity, and pH, which caused concomitant changes in the growth rates of the species. We tried to capture these dynamics by the development of a mathematical model. The results show that the model generally fitted well to the monoculture data, in the experiments at both low pCO<sub>2</sub> (Fig. 1) and high pCO<sub>2</sub> (Fig. 2). The species parameters estimated from the monoculture experiments are summarized in Table 2, and will be used to predict the dynamics of the competition experiments.

Resource competition theory can provide some further insights. Consider several species competing for a single limiting resource. Each species has its own critical  $R^*$ , defined as the resource availability at which the specific growth rate of a species equals its loss rate. During competition, resource availability diminishes as the resource is consumed by the species. One by one, species start to decline when resource availability is depleted below their  $R^*$  values. This process continues, until eventually the species with lowest  $R^*$  has competitively displaced all other species. Hence, resource competition theory predicts that the species with lowest  $R^*$  will be the superior competitor (Tilman, 1982; Grover, 1997).



**Fig. 2.** Monoculture experiments at high  $p\text{CO}_2$  (2000 ppm). (A–D) Population density (expressed as biovolume) and light intensity  $I_{\text{out}}$  penetrating through the chemostat. (E–H)  $\text{CO}_2(\text{aq})$ , bicarbonate and carbonate concentrations. (I–L) Dissolved inorganic carbon (DIC) and pH. Different panels represent different species: (A, E, I) *Monoraphidium*; (B, F, J) *Microcystis*; (C, G, K) *Scenedesmus*; (D, H, L) *Chlorella*. Symbols indicate experimental data, and lines indicate model fits. Parameter values of the model are provided in Table 1 and Table 2.

In many applications, the  $R^*$  value of each species is measured as the steady-state concentration of the limiting resource in monoculture (e.g. Tilman, 1981; Passarge *et al.*, 2006; Wilson *et al.*, 2007). In our application, however,  $\text{CO}_2$  and bicarbonate provide two alternative inorganic carbon sources that are rapidly interconverted by the chemical reaction of  $\text{CO}_2$  with water, which makes it difficult to measure the  $R^*$  for  $\text{CO}_2$  and  $R^*$  for bicarbonate independently. Nevertheless, the steady-state concentrations of  $\text{CO}_2$  and bicarbonate in carbon-limited monoculture may provide useful information on the competitive abilities for inorganic carbon of the species. That is, if we assume that a low steady-state  $\text{CO}_2$  concentration in monoculture implies a high competitive ability for  $\text{CO}_2$ , then the species can be ranked according to their competitive ability for  $\text{CO}_2$  as (Fig. 3A): *Scenedesmus* > *Chlorella* > *Microcystis* > *Monoraphidium*. Similarly, based on the steady-state bicarbonate concentrations, the species can be ranked according to their competitive ability for bicarbonate as (Fig. 3B): *Scenedesmus* > *Chlorella* > *Microcystis* > *Monoraphidium*. Accordingly, the ranking of the species is the same for both  $\text{CO}_2$  and bicarbonate, indicating that *Scenedesmus* will be the best competitor for inorganic carbon, followed by *Chlorella* and then *Microcystis*, while *Monoraphidium* is the worst competitor for inorganic carbon.


At high  $\text{CO}_2$  levels,  $\text{CO}_2$  and bicarbonate were in ample supply, but light becomes a limiting resource (Fig. 2A–D).

Analogous to  $R^*$ , competition theory predicts that the species with lowest critical light intensity ( $I_{\text{out}}^*$ ) is the superior competitor for light (Huisman and Weissing, 1994; Huisman *et al.*, 1999). The critical light intensities of the species were measured as the steady-state values of  $I_{\text{out}}$  in the monoculture experiments at high  $p\text{CO}_2$ . Based on their critical light intensities, the species can be ranked according to their competitive ability for light as (Fig. 3C): *Microcystis*  $\approx$  *Scenedesmus* > *Chlorella* > *Monoraphidium*. Hence, from the monoculture data, *Microcystis* and *Scenedesmus* are predicted to be the best competitors for light with an approximately similar competitive ability, followed by *Chlorella*, while *Monoraphidium* is the worst competitor for light.

Comparison of the above species rankings indicates that *Microcystis* will become a stronger competitor when the species interactions shift from competition for inorganic carbon at low  $p\text{CO}_2$  to competition for light at high  $p\text{CO}_2$ . The relative ranking among the three green algae remains unaltered with rising  $p\text{CO}_2$ .

#### Competition experiments

The competition experiments largely confirmed the predictions derived from the monoculture experiments. For instance, in the competition experiment at low  $p\text{CO}_2$  between *Monoraphidium* and *Scenedesmus*, both species increased during the first 6 d



**Fig. 3.**  $R^*$  values of the species. (A, B)  $R^*$  values for  $\text{CO}_2(\text{aq})$  (A) and bicarbonate (B) of the species, indicative of their competitive abilities for inorganic carbon.  $R^*$  values were estimated from the  $\text{CO}_2$  and bicarbonate concentrations measured in the steady-state monocultures at low  $\text{pCO}_2$ . (C) Critical light intensities ( $I_{\text{out}}^*$ ) of the species, indicative of their competitive abilities for light. Critical light intensities were estimated from the light intensities penetrating through the steady-state monocultures at high  $\text{pCO}_2$ . All estimates are based on the mean  $\pm$  SD of the last five data points of each monoculture experiment. Sc., *Scenedesmus*; Chl, *Chlorella*; Mic, *Microcystis*; Mon, *Monoraphidium*.

(Fig. 4A). Meanwhile, pH increased to 10.8 (see Supplementary Fig. S1A), the dissolved  $\text{CO}_2$  concentration was depleted to  $<0.004 \mu\text{mol l}^{-1}$  and the bicarbonate concentration decreased to  $\sim 100 \mu\text{mol l}^{-1}$  (Supplementary Fig. S2A). These concentrations are below the  $R^*$  values for  $\text{CO}_2$  and bicarbonate of *Monoraphidium* (Fig. 3A, B). Hence, as predicted, *Monoraphidium* started to decline after the first week, and was competitively displaced by *Scenedesmus* (Fig. 4A).

At low  $\text{pCO}_2$ , *Monoraphidium* was also outcompeted by *Chlorella* (Fig. 4C) and *Microcystis* (Fig. 4E). Hence, *Monoraphidium* was the weakest competitor. Furthermore, in line with expectation, *Scenedesmus* competitively displaced *Microcystis* (Fig. 4G). Contrary to expectation, *Chlorella* and *Microcystis* appeared to coexist at low  $\text{pCO}_2$  (Fig. 4I). During the last 20 d of this experiment, however, *Chlorella* tended to increase slowly at the cost of *Microcystis*, although the

experiment didn't last long enough to witness the final outcome. The latter result is in agreement with the rate of competitive displacement (Table 3), which indicated that *Chlorella* was indeed slowly displacing *Microcystis*. According to the competition experiments, the competitive ranking of the species at low  $\text{pCO}_2$  can thus be summarized as follows: *Scenedesmus*  $>$  *Chlorella*  $\geq$  *Microcystis*  $>$  *Monoraphidium*. This is in good agreement with the rankings based on the  $R^*$  values for  $\text{CO}_2$  and bicarbonate estimated in the monoculture experiments.

In their competition experiment at high  $\text{pCO}_2$ , both *Monoraphidium* and *Scenedesmus* increased during the first 20 d (Fig. 4B). Meanwhile, pH stabilized at  $\sim 8.5$  (see Supplementary Fig. S1B at JXB online), the dissolved  $\text{CO}_2$  concentration slightly decreased to  $\sim 20 \mu\text{mol l}^{-1}$  and the bicarbonate concentration increased to  $>3,000 \mu\text{mol l}^{-1}$  (Supplementary Fig. S2B). However, the incident light was almost completely absorbed by the dense species mixture, with  $<0.1 \mu\text{mol photons m}^{-2} \text{s}^{-1}$  penetrating through the chemostat (Supplementary Fig. S1B). This value is below the critical light intensity of *Monoraphidium* (Fig. 3C). Hence, after the first 20 d, *Monoraphidium* declined, and was gradually displaced by *Scenedesmus* (Fig. 4B).

At high  $\text{pCO}_2$ , *Monoraphidium* was also outcompeted by *Chlorella* (Fig. 4D) and *Microcystis* (Fig. 4F), and hence it was again the weakest competitor. Interestingly, *Scenedesmus* and *Microcystis* appeared to coexist (Fig. 4H), in agreement with the similar critical light intensities of these two species (Fig. 3C). Their coexistence at high  $\text{pCO}_2$  was confirmed by the rate of competitive displacement, which did not differ significantly from zero for this species pair (Table 3). In line with expectation, *Microcystis* competitively displaced *Chlorella* (Fig. 4J). Hence, according to the competition experiments, the competitive abilities of the species at high  $\text{pCO}_2$  can be ranked as follows: *Microcystis*  $\approx$  *Scenedesmus*  $>$  *Chlorella*  $>$  *Monoraphidium*. This matches the species ranking based on their critical light intensities in monoculture.


Also quantitatively, the population dynamics predicted by the competition model agreed well with the results of the competition experiments (compare lines versus symbols in Fig. 4, and Supplementary Figs S1 and S2), both at low  $\text{pCO}_2$  and at high  $\text{pCO}_2$ .

## Discussion

### *Cyanobacteria versus green algae*

Both our model predictions and our experimental results contradict the classic view (Shapiro, 1990; Caraco and Miller, 1998; Low-Décarie et al., 2011, 2015) that cyanobacteria are strong competitors at low  $\text{CO}_2$  levels, whereas eukaryotic phytoplankton such as green algae are better competitors at elevated  $\text{CO}_2$ . We found the opposite. At low  $\text{CO}_2$  levels, the cyanobacterium *Microcystis* was a relatively poor competitor. It lost in competition with *Scenedesmus*, was slowly replaced by *Chlorella*, and won only against *Monoraphidium*. At high  $\text{CO}_2$  levels, *Microcystis* was a stronger competitor. It won in competition with both *Monoraphidium* and *Chlorella*, and





**Fig. 4.** Competition experiments. The competition experiments were performed at low  $p\text{CO}_2$  (100 ppm; left panels) and high  $p\text{CO}_2$  (2000 ppm; right panels). (A, B) Competition between *Monoraphidium* and *Scenedesmus*. (C, D) Competition between *Monoraphidium* and *Chlorella*. (E, F) Competition between *Monoraphidium* and *Microcystis*. (G, H) Competition between *Microcystis* and *Scenedesmus*. (I, J) Competition between *Microcystis* and *Chlorella*. Dynamic changes in light conditions, carbon speciation and pH during the competition experiments are presented in Supplementary Figs S1 and S2. Symbols indicate experimental data, and lines indicate model predictions. Parameter values of the model are provided in Tables 1 and 2.

coexisted with *Scenedesmus*. *Microcystis* was the only species that increased its competitive ranking at elevated  $\text{CO}_2$ ; the relative ranking among the three species of green algae did not change.

#### Competition at low $\text{CO}_2$

In the competition experiments at low  $p\text{CO}_2$ , the growing phytoplankton populations depleted the dissolved  $\text{CO}_2$  concentration within the first 1–3 weeks of the experiments and also

the bicarbonate concentration declined. As a consequence, the growth rates of the phytoplankton populations slowed down, and one of the species in the experiments started to displace the other. Interestingly, both the ranking of the  $R^*$  values estimated from the monocultures and the competitive replacements observed in the competition experiments show that the green algae *Scenedesmus* and *Chlorella* were stronger competitors for inorganic carbon than the cyanobacterium *Microcystis*.

These results can to a large extent be explained by the CCMs of the species. In particular, the cyanobacterium

**Table 3.** Rates of competitive displacement (RCD±standard error) in the competition experiments

RCD was calculated as the slope of a linear regression of  $\ln(X_1/X_2)$  versus time, where  $X_1$  and  $X_2$  are the population densities of species 1 and species 2. The coefficient of determination ( $R^2$ ), number of data points ( $n$ ) and significance ( $P$ ) of the linear regression are indicated.

Competition experiment	RCD (d <sup>-1</sup> )	R <sup>2</sup>	n	P
<i>Monoraphidium</i> vs <i>Microcystis</i>				
Low pCO <sub>2</sub>	-0.250 ± 0.008	0.99	14	<0.001
High pCO <sub>2</sub>	-0.043 ± 0.005	0.88	11	<0.001
<i>Monoraphidium</i> vs <i>Scenedesmus</i>				
Low pCO <sub>2</sub>	-0.292 ± 0.021	0.97	8	<0.001
High pCO <sub>2</sub>	-0.021 ± 0.002	0.84	20	<0.001
<i>Monoraphidium</i> vs <i>Chlorella</i>				
Low pCO <sub>2</sub>	-0.100 ± 0.008	0.93	16	<0.001
High pCO <sub>2</sub>	-0.062 ± 0.008	0.82	15	<0.001
<i>Scenedesmus</i> vs <i>Microcystis</i>				
Low pCO <sub>2</sub>	+0.654 ± 0.071	0.95	6	<0.001
High pCO <sub>2</sub>	0.000 ± 0.003	0.00	11	n.s.
<i>Microcystis</i> vs <i>Chlorella</i>				
Low pCO <sub>2</sub>	-0.039 ± 0.071	0.96	11	<0.001
High pCO <sub>2</sub>	+0.197 ± 0.025	0.92	11	<0.001

*Microcystis* PCC 7806 used in this study is a *bicA* strain (*sensu* Sandrini et al., 2014). It contains the bicarbonate uptake systems BicA and BCT1, but lacks the high-affinity bicarbonate uptake system SbtA. Batch experiments have shown that *Microcystis* PCC 7806 has a lower growth rate at low C<sub>i</sub> levels than strains that do have SbtA (Sandrini et al., 2014). BCT1 is induced when *Microcystis* PCC 7806 grows at low CO<sub>2</sub> levels (Sandrini et al., 2015a, b), but this uptake system appears to have a slightly lower affinity for bicarbonate than SbtA, at least in the cyanobacteria *Synechocystis* PCC 6803 and *Synechococcus* PCC 7002 (Price et al., 2004, 2008). Moreover, bicarbonate uptake by BCT1 is ATP dependent and therefore energetically quite expensive. The Na<sup>+</sup>-dependent bicarbonate uptake system BicA has a high flux rate but low affinity for bicarbonate (Price et al., 2004). Hence, the lack of SbtA offers a plausible explanation of why *Microcystis* PCC 7806 had a selective disadvantage under C<sub>i</sub>-limited conditions. Previous experiments have indeed shown that such *bicA* strains were selectively displaced by *bicA+sbtA* strains under C<sub>i</sub>-limited conditions (Sandrini et al., 2016). Our results show that cyanobacteria that lack the high-affinity bicarbonate uptake system SbtA are relatively poor competitors under C<sub>i</sub>-limited conditions, not only in comparison with other cyanobacteria but also in comparison with green algae such as *Scenedesmus* and *Chlorella*.

Our results do not of course imply that cyanobacteria are generally poor competitors under C<sub>i</sub>-limited conditions. *Microcystis* strains containing the high-affinity bicarbonate uptake system SbtA sustain higher growth rates at low C<sub>i</sub> concentrations than *bicA* strains such as *Microcystis* PCC 7806 (Sandrini et al., 2014). Moreover, recent selection experiments and lake data show that *bicA+sbtA* strains have a competitive advantage over *bicA* strains at low CO<sub>2</sub> levels

(Sandrini et al., 2016). Whether SbtA-containing cyanobacteria or green algae such as *Scenedesmus* and *Chlorella* are better competitors at low CO<sub>2</sub> levels thus remains an interesting open question.

Compared with cyanobacteria, less is known about the functioning of the CCMs in green algae. Our current understanding of eukaryotic CCMs comes largely from studies with the model organism *Chlamydomonas reinhardtii* (Moroney and Ynalvez 2007; Spalding, 2008; Wang et al., 2011; Meyer and Griffiths, 2013). Briefly, the key components of the eukaryotic CCM are quite similar to the cyanobacterial CCM and involve active CO<sub>2</sub> and bicarbonate uptake, interconversion between these two C<sub>i</sub> species by carbonic anhydrases (CAs), and a microcompartment that contains RuBisCO (Wang et al., 2015). In *Chlamydomonas*, five C<sub>i</sub> transporters have been localized, including two confirmed bicarbonate transporters (HLA3 on the plasma membrane and LCIA on the chloroplast envelope) and three undefined C<sub>i</sub> transporters (LCII on the plasma membrane and CCAPI/2 on the chloroplast envelope). Inside the chloroplast, RuBisCO is densely packed in a microcompartment named the pyrenoid (Kuchitsu et al., 1988; Engel et al., 2015). Within the pyrenoid, bicarbonate is dehydrated to CO<sub>2</sub> by a CA and released to RuBisCO (Moroney and Ynalvez, 2007).

So far, there are no studies on the CCM genes of the green algal strains we used in this study, but there is some recent work on the CCM genes of a different species of *Chlorella*, *C. pyrenoidosa*, indicating that this species has a CCM similar to that of *Chlamydomonas* (Fan et al., 2016). For instance, when shifting *C. pyrenoidosa* from high CO<sub>2</sub> to low CO<sub>2</sub> conditions, CCM-related genes such as *LCIA*, *LCIB*, and *HLA3* showed increased expression, similar to the response of *C. reinhardtii* (Fan et al., 2015). Interestingly, earlier work indicates that there is substantial variation in CCM activity within the *Chlorella* genus and even between different strains of the same species of *Chlorella*. For example, *C. vulgaris* strain 11 h and strain UTEX 263 seem to utilize only CO<sub>2</sub> as a carbon source (Miyachi et al., 1983; Tu et al., 1986), whereas *C. vulgaris* strain C-3, strain UTEX 259 and *C. pyrenoidosa* can use both CO<sub>2</sub> and bicarbonate (Miyachi et al., 1983). Our results indicate that the *C. vulgaris* strain we used can utilize bicarbonate: if this green alga could only use CO<sub>2</sub>, it would not be able to grow in monoculture in our chemostats at a dilution rate of 0.125 d<sup>-1</sup> and dissolved CO<sub>2</sub> concentrations lower than 0.01 μmol l<sup>-1</sup> (Fig. 1).

Similarly, the fact that *S. obliquus* and *M. griffithii* were also able to grow at CO<sub>2</sub> concentrations below 0.01 μmol l<sup>-1</sup> in our chemostat experiments indicates that these green algae can also utilize bicarbonate as carbon source. Earlier studies confirm the presence of a functional CCM in a different strain of *S. obliquus*. WT strain D3 has increased intracellular and extracellular CA activity, as well as a higher affinity for the uptake of CO<sub>2</sub> and bicarbonate at low compared with high CO<sub>2</sub> concentrations (Palmqvist et al., 1994). Furthermore, in our experiments *S. obliquus* could deplete dissolved CO<sub>2</sub> and bicarbonate concentrations to even lower levels than *Microcystis* PCC 7806 (Fig. 3), and displace *Microcystis* in the competition experiments at low CO<sub>2</sub> (Fig. 4).

Similar results were obtained in CO<sub>2</sub>-limited mesocosm experiments by Verschoor *et al.* (2013), where the same strain of *S. obliquus* displaced *Synechocystis* PCC 6803, a cyanobacterium that contains all five C<sub>1</sub> uptake systems (Badger *et al.*, 2006). So far, very little is known about the CCM of *Monoraphidium griffithii*, but a related species, *M. braunii*, can photoactivate a blue-light-dependent bicarbonate transport system under CO<sub>2</sub>-limiting conditions (Mora *et al.* 2002).

In total, these physiological studies indicate that many green algae are well adapted to cope with C<sub>1</sub> limitation. Similar to cyanobacteria, many species of green algae are able to induce a CCM under CO<sub>2</sub>-limiting conditions (Meyer and Griffiths, 2013). Furthermore, in eutrophic lakes, not only cyanobacteria but also green algae have been observed to develop dense blooms at low CO<sub>2</sub> concentrations and high pH (Jeppesen *et al.*, 1990; Jensen *et al.*, 1994; Beklioglu and Moss, 1995). The results of our competition experiments are in agreement with these physiological studies and field observations, and demonstrate that green algae can indeed be very effective competitors at low CO<sub>2</sub> levels.

### Competition at elevated CO<sub>2</sub>

The notion that *Microcystis* PCC 7806 lacks the high-affinity uptake system SbtA explains not only why it was a poor competitor at low CO<sub>2</sub> levels, but may also help to understand why it was more successful at high CO<sub>2</sub> levels. In *bicA+sbtA* strains of *Microcystis*, the bicarbonate uptake genes *bicA* and *sbtA* are located on the same operon and are co-transcribed (Sandrini *et al.*, 2014). Transcription of *sbtA* will be inefficient and costly, however, when inorganic concentrations are high and hence the high-affinity but low-flux system SbtA is no longer needed. This reasoning is supported by recent selection experiments, which have shown that *bicA* strains have a competitive advantage over *bicA+sbtA* strains at elevated CO<sub>2</sub> levels (Sandrini *et al.*, 2016).

Yet, the presence of the high-flux bicarbonate uptake system BicA is probably not sufficient to explain its competitive success at elevated CO<sub>2</sub> levels. Elevated CO<sub>2</sub> increased the dissolved C<sub>1</sub> concentrations, but also yielded 3- to 8-fold higher population densities, which in turn generated very low light availabilities (Figs 1 and 2, and Supplementary Table S1). At these high CO<sub>2</sub> but low light levels, carbon availability is no longer a major limiting factor and bicarbonate uptake systems tend to be down-regulated (Beardall, 1991; Sandrini *et al.*, 2015a). Instead, the intense shading induced by the dense phytoplankton populations will favor species adapted to low light environments (Huisman *et al.*, 1999). A range of studies have suggested that cyanobacteria are superior competitors for light (Mur *et al.*, 1977; Reynolds *et al.*, 1987; Huisman *et al.*, 2004; Yang and Jin, 2008; Schwaderer *et al.*, 2011). Comparison of the traits of the species investigated in this study shows that *Microcystis* PCC 7806 had the lowest half-saturation constant for light-limited growth (Table 2), indicating that it can sustain a relatively high growth rate at low light levels in comparison with the other species. *Scenedesmus* also had a relatively low half-saturation constant for light-limited growth, albeit higher than *Microcystis*, and had the highest

maximum growth rate of all four species (Table 2). Indeed, the competitive ability of the cyanobacterium *Microcystis* increased at elevated CO<sub>2</sub>, and together with *Scenedesmus* it became the best competitor for light (Fig. 3).

### Model predictions

Comparison of the model predictions and experimental results shows that the impact of elevated CO<sub>2</sub> on phytoplankton competition can be quite accurately predicted under controlled laboratory conditions. First, the system parameters and several species parameters were measured in monoculture experiments, while the remaining species parameters were estimated from least-squares fits of the model predictions to the monoculture dynamics (Figs 1 and 2). Hence, the model is calibrated with the monoculture data. Subsequently, the parameter estimates from the monocultures were used to predict the time course and outcome of competition in the species mixtures. Accordingly, the model predictions are validated with the competition experiments (Fig. 4, and Supplementary Figs S1 and S2).

Our model and experiments are of course still a major simplification in comparison with the complexities of the real CCMs of cyanobacteria and green algae competing in natural waters. For instance, the model brushes over many of the physiological details involved in the regulation of CCMs in response to changes in carbon and light availability (e.g. Beardall and Giordano, 2002; Burnap *et al.*, 2015). Furthermore, the experiments were limited to only a small number of phytoplankton species under controlled laboratory conditions, in isolation from a multitude of other hydrological, biogeochemical and ecological processes that are important in lakes. Species might be superior competitors for inorganic carbon or light, but if they are also preferentially grazed by zooplankton they are still unlikely to gain dominance in natural waters. Hence, further improvement of the model predictions might be obtained by further refinement of the underlying physiological and ecological processes.

Nevertheless, the results are promising. Our model and experiments seemed to capture the basic ingredients required to predict phytoplankton competition at different CO<sub>2</sub> levels. Similar to several previous studies (e.g. Caraco and Miller, 1998; Low-Décarie *et al.*, 2011; Trimborn *et al.*, 2013), we have shown that species traits measured in monoculture can be used to predict changes in phytoplankton species composition at elevated CO<sub>2</sub>. To our knowledge, our study is the first experimental demonstration that a mathematical model can quantitatively predict dynamic changes in phytoplankton species composition, carbon speciation, pH, alkalinity and light during the competition process.

### Conclusions

Our experimental results call for a revision of the classic paradigm that cyanobacteria are superior competitors at low CO<sub>2</sub> levels and high pH, whereas eukaryotic phytoplankton such as green algae are superior competitors at elevated CO<sub>2</sub>. Such simple dichotomies do not capture the diversity of CCMs that

have been found among and within different phytoplankton taxa. First, our results demonstrate that green algae can also be very effective competitors at low CO<sub>2</sub> levels. Second, our results show that some cyanobacterial strains are relatively poor competitors when CO<sub>2</sub> is limiting but become stronger competitors at elevated CO<sub>2</sub>. We therefore urge the CCM scientific community to further elucidate and compare the performance of different CCMs, and their selective advantages and disadvantages across a wide range of different species at the molecular, physiological and ecological level. Such a comparative approach will be essential if we are to understand and predict how the species composition of natural phytoplankton communities will respond to the anticipated rise in CO<sub>2</sub> levels.

## Supplementary data

Supplementary data are available at *JXB* online.

Model S1. Detailed description of the model.

Fig. S1. Dynamic changes in light, DIC and pH during the competition experiments.

Fig. S2. Dynamic changes in carbon speciation during the competition experiments.

Table S1. Steady-state characteristics in the monoculture experiments.

## Acknowledgements

We thank Hanneke van Bree, Sander de Groot, Duke van Velden, Maureen de Wit and Josan Yauw for their help in running the chemostats, Leo Hoitinga for assistance with the TOC-V<sub>CPH</sub> Analyzer and Elemental Analyzer, Suzanne Naus-Wiezer of the NIOO-KNAW for assistance with flow cytometry, and the reviewers for their helpful comments. This research is supported by the Division of Earth and Life Sciences (ALW) of the Netherlands Organization for Scientific Research (NWO).

## References

- Badger MR, Price GD, Long BM, Woodger FJ.** 2006. The environmental plasticity and ecological genomics of the cyanobacterial CO<sub>2</sub> concentrating mechanism. *Journal of Experimental Botany* **57**, 249–265.
- Balmer MB, Downing JA.** 2011. Carbon dioxide concentrations in eutrophic lakes: undersaturation implies atmospheric uptake. *Inland Waters* **1**, 125–132.
- Beardall J.** 1991. Effects of photon flux density on the 'CO<sub>2</sub>-concentrating mechanism' of the cyanobacterium *Anabaena variabilis*. *Journal of Plankton Research* **13**, 133–141.
- Beardall J, Giordano M.** 2002. Ecological implications of microalgal and cyanobacterial CO<sub>2</sub> concentrating mechanisms, and their regulation. *Functional Plant Biology* **29**, 335–347.
- Beklioglu M, Moss B.** 1995. The impact of pH on interactions among phytoplankton algae, zooplankton and perch (*Perca fluviatilis*) in a shallow, fertile lake. *Freshwater Biology* **33**, 497–509.
- Burnap RL, Hagemann M, Kaplan A.** 2015. Regulation of CO<sub>2</sub> concentrating mechanism in cyanobacteria. *Life* **5**, 348–371.
- Caraco NF, Miller R.** 1998. Effects of CO<sub>2</sub> on competition between a cyanobacterium and eukaryotic phytoplankton. *Canadian Journal of Fisheries and Aquatic Sciences* **55**, 54–62.
- Carmichael WW.** 2001. Health effects of toxin-producing cyanobacteria: "The CyanoHABs". *Human and Ecological Risk Assessment* **7**, 1393–1407.
- Chorus I, Bartram J.** 1999. Toxic cyanobacteria in water: a guide to their public health consequences, monitoring, and management. London: E & FN Spon.
- Codd GA, Morrison LF, Metcalf JS.** 2005. Cyanobacterial toxins: risk management for health protection. *Toxicology and Applied Pharmacology* **203**, 264–272.
- Cole JJ, Bade DL, Bastviken D, Pace ML, Van de Bogert M.** 2010. Multiple approaches to estimating air-water gas exchange in small lakes. *Limnology and Oceanography Methods* **8**, 285–293.
- Droop MR.** 1973. Some thoughts on nutrient limitation in algae. *Journal of Phycology* **9**, 264–272.
- Eisenhut M, Aguirre von Wobeser EA, Jonas L, Schubert H, Ibelings BW, Bauwe H, Matthijs HCP, Hagemann M.** 2007. Long-term response toward inorganic carbon limitation in wild type and glycolate turnover mutants of the cyanobacterium *Synechocystis* sp. strain PCC 6803. *Plant Physiology* **144**, 1946–1959.
- Engel BD, Schaffer M, Cuellar LK, Villa E, Plietzko JM, Baumeister W.** 2015. Native architecture of the *Chlamydomonas* chloroplast revealed by in situ cryo-electron tomography. *eLife* **4**, e04889.
- Faltermann S, Zucchi S, Kohler E, Blom JF, Perntaler J, Fent K.** 2014. Molecular effects of the cyanobacterial toxin cyanopeptolin (CP1020) occurring in algal blooms: global transcriptome analysis in zebrafish embryos. *Aquatic Toxicology* **149**, 33–39.
- Fan J, Xu H, Li Y.** 2016. Transcriptome-based global analysis of gene expression in response to carbon dioxide deprivation in the green algae *Chlorella pyrenoidosa*. *Algal Research* **16**, 12–19.
- Fan J, Xu H, Luo Y, Wan M, Huang J, Wang W, Li Y.** 2015. Impacts of CO<sub>2</sub> concentration on growth, lipid accumulation, and carbon-concentrating-mechanism-related gene expression in oleaginous *Chlorella*. *Applied Microbiology and Biotechnology* **99**, 2451–2462.
- Goldman JC, Brewer PG.** 1980. Effect of nitrogen source and growth rate on phytoplankton-mediated changes in alkalinity. *Limnology and Oceanography* **25**, 352–357.
- Grover JP.** 1991. Dynamics of competition among microalgae in variable environments: experimental tests of alternative models. *Oikos* **62**, 231–243.
- Grover JP.** 1997. Resource competition. London: Chapman & Hall.
- Guo L.** 2007. Doing battle with the green monster of Taihu Lake. *Science* **317**, 1166.
- Huisman J, Jonker RR, Zonneveld C, Weissing FJ.** 1999. Competition for light between phytoplankton species: experimental tests of mechanistic theory. *Ecology* **80**, 211–222.
- Huisman J, Matthijs HCP, Visser PM, eds.** 2005. Harmful cyanobacteria. Dordrecht: Springer Netherlands.
- Huisman J, Matthijs HCP, Visser PM, Balke H, Sigon CAM, Passarge J, Weissing FJ, Mur LR.** 2002. Principles of the light-limited chemostat: theory and ecological applications. *Antonie van Leeuwenhoek* **81**, 117–133.
- Huisman J, Sharples J, Stroom JM, Visser PM, Kardinaal WEA, Verspagen JMH, Sommeijer B.** 2004. Changes in turbulent mixing shift competition for light between phytoplankton species. *Ecology* **85**, 2960–2970.
- Huisman J, Weissing FJ.** 1994. Light-limited growth and competition for light in well-mixed aquatic environments: an elementary model. *Ecology* **75**, 507–520.
- Ibelings BW, Maberly SC.** 1998. Photoinhibition and the availability of inorganic carbon restrict photosynthesis by surface blooms of cyanobacteria. *Limnology and Oceanography* **43**, 408–419.
- Jensen JP, Jeppesen E, Olrik K, Kristensen P.** 1994. Impact of nutrients and physical factors on the shift from cyanobacterial to chlorophyte dominance in shallow Danish lakes. *Canadian Journal of Fisheries and Aquatic Sciences* **51**, 1692–1699.
- Jeppesen E, Sondergaard M, Sortkjaer O, Mortensen E, Kristensen P.** 1990. Interactions between phytoplankton, zooplankton and fish in a shallow, hypertrophic lake – a study of phytoplankton collapses in Lake Søbygård, Denmark. *Hydrobiologia* **191**, 149–164.
- Kuchitsu K, Tsuzuki M, Miyachi S.** 1988. Characterization of the pyrenoid isolated from unicellular green-alga *Chlamydomonas reinhardtii*: particulate form of RuBisCo protein. *Protoplasma* **144**, 17–24.

- Lazzarino JK, Bachmann RW, Hoyer MV, Canfield DE Jr.** 2009. Carbon dioxide supersaturation in Florida lakes. *Hydrobiologia* **627**, 169–180.
- Litchman E, Klausmeier CA, Bossard P.** 2004. Phytoplankton nutrient competition under dynamic light regimes. *Limnology and Oceanography* **49**, 1457–1462.
- Low-Décarie E, Bell G, Fussmann GF.** 2015. CO<sub>2</sub> alters community composition and response to nutrient enrichment of freshwater phytoplankton. *Oecologia* **177**, 875–883.
- Low-Décarie E, Fussmann GF, Bell G.** 2011. The effect of elevated CO<sub>2</sub> on growth and competition in experimental phytoplankton communities. *Global Change Biology* **17**, 2525–2535.
- Merel S, Walker D, Chicana R, Snyder S, Baurès E, Thomas O.** 2013. State of knowledge and concerns on cyanobacterial blooms and cyanotoxins. *Environment International* **59**, 303–327.
- Meyer M, Griffiths H.** 2013. Origins and diversity of eukaryotic CO<sub>2</sub>-concentrating mechanisms: lessons for the future. *Journal of Experimental Botany* **64**, 769–786.
- Michalak AM, Anderson EJ, Beletsky D, et al.** 2013. Record-setting algal bloom in Lake Erie caused by agricultural and meteorological trends consistent with expected future conditions. *Proceedings of the National Academy of Sciences of the United States of America* **110**, 6448–6452.
- Miyachi S, Tsuzuki M, Avramova ST.** 1983. Utilization modes of inorganic carbon for photosynthesis in various species of *Chlorella*. *Plant and Cell Physiology* **24**, 441–451.
- Mora C, Witt FG, Aparicio PJ, Quiñones MA.** 2002. Independent induction of two blue light-dependent monovalent anion transport systems in the plasma membrane of *Monoraphidium braunii*. *Journal of Experimental Botany* **53**, 1909–1918.
- Moroney JV, Ynalvez RA.** 2007. Proposed carbon dioxide concentrating mechanism in *Chlamydomonas reinhardtii*. *Eukaryotic Cell* **6**, 1251–1259.
- Mur LR, Gons HJ, Van Liere L.** 1977. Some experiments on the competition between green algae and blue-green bacteria in light-limited environments. *FEMS Microbiology Letters* **1**, 335–338.
- Palmqvist K, Yu JW, Badger MR.** 1994. Carbonic anhydrase activity and inorganic carbon fluxes in low- and high-C<sub>i</sub> cells of *Chlamydomonas reinhardtii* and *Scenedesmus obliquus*. *Physiologia Plantarum* **90**, 537–547.
- Passarge J, Hol S, Escher M, Huisman J.** 2006. Competition for nutrients and light: stable coexistence, alternative stable states, or competitive exclusion? *Ecological Monographs* **76**, 57–72.
- Price GD, Badger MR, Woodger FJ, Long BM.** 2008. Advances in understanding the cyanobacterial CO<sub>2</sub>-concentrating-mechanism (CCM): functional components, C<sub>i</sub> transporters, diversity, genetic regulation and prospects for engineering into plants. *Journal of Experimental Botany* **59**, 1441–1461.
- Price GD, Woodger FJ, Badger MR, Howitt SM, Tucker L.** 2004. Identification of a SulP-type bicarbonate transporter in marine cyanobacteria. *Proceedings of the National Academy of Sciences of the United States of America* **101**, 18228–18233.
- Raven JA, Giordano M, Beardall J, Maberly SC.** 2012. Algal evolution in relation to atmospheric CO<sub>2</sub>: carboxylases, carbon-concentrating mechanisms and carbon oxidation cycles. *Philosophical Transactions of the Royal Society of London. Series B, Biological Sciences* **367**, 493–507.
- Reynolds CS, Oliver RL, Walsby AE.** 1987. Cyanobacterial dominance: the role of buoyancy regulation in dynamic lake environments. *New Zealand Journal of Marine and Freshwater Research* **21**, 379–390.
- Sandrini G, Cunsolo S, Schuurmans JM, Matthijs HCP, Huisman J.** 2015a. Changes in gene expression, cell physiology and toxicity of the harmful cyanobacterium *Microcystis aeruginosa* at elevated CO<sub>2</sub>. *Frontiers in Microbiology* **6**, 401.
- Sandrini G, Jakupovic D, Matthijs HCP, Huisman J.** 2015b. Strains of the harmful cyanobacterium *Microcystis aeruginosa* differ in gene expression and activity of inorganic carbon uptake systems at elevated CO<sub>2</sub> levels. *Applied and Environmental Microbiology* **81**, 7730–7739.
- Sandrini G, Matthijs HCP, Verspagen JMH, Muyzer G, Huisman J.** 2014. Genetic diversity of inorganic carbon uptake systems causes variation in CO<sub>2</sub> response of the cyanobacterium *Microcystis*. *The ISME Journal* **8**, 589–600.
- Sandrini G, Ji X, Verspagen JMH, Tann RP, Slot PC, Luimstra VL, Schuurmans JM, Matthijs HCP, Huisman J.** 2016. Rapid adaptation of harmful cyanobacteria to rising CO<sub>2</sub>. *Proceedings of the National Academy of Sciences of the United States of America* **113**, 9315–9320.
- Schwaderer AS, Yoshiyama K, De Tezanos Pinto P, Swenson NG, Klausmeier CA, Litchman E.** 2011. Eco-evolutionary differences in light utilization traits and distributions of freshwater phytoplankton. *Limnology and Oceanography* **56**, 589–598.
- Shapiro J.** 1990. Current beliefs regarding dominance of bluegreens: the case for the importance of CO<sub>2</sub> and pH. *Verhandlungen Internationale Vereinigung für Theoretische und Angewandte Limnologie* **24**, 38–54.
- Shapiro J.** 1997. The role of carbon dioxide in the initiation and maintenance of blue-green dominance in lakes. *Freshwater Biology* **37**, 307–323.
- Siegenthaler U, Sarmiento JL.** 1993. Atmospheric carbon dioxide and the ocean. *Nature* **365**, 119–125.
- Sommer U.** 1985. Comparison between steady state and non-steady state competition: experiments with natural phytoplankton. *Limnology and Oceanography* **30**, 335–346.
- Spalding MH.** 2008. Microalgal carbon-dioxide-concentrating mechanisms: *Chlamydomonas* inorganic carbon transporters. *Journal of Experimental Botany* **59**, 1463–1473.
- Stomp M, Huisman J, De Jongh F, Veraart AJ, Gerla D, Rijkeboer M, Ibelings BW, Wollenzien UIA, Stal LJ.** 2004. Adaptive divergence in pigment composition promotes phytoplankton biodiversity. *Nature* **432**, 104–107.
- Stumm W, Morgan JJ.** 1996. *Aquatic chemistry: chemical equilibria and rates in natural waters*, 3rd edn. New York: John Wiley & Sons.
- Talling JF.** 1976. The depletion of carbon dioxide from lake water by phytoplankton. *Journal of Ecology* **64**, 79–121.
- Tilman D.** 1977. Resource competition between plankton algae: an experimental and theoretical approach. *Ecology* **58**, 338–348.
- Tilman D.** 1981. Tests of resource competition theory using four species of Lake Michigan algae. *Ecology* **62**, 802–815.
- Tilman D.** 1982. *Resource competition and community structure*. Princeton: Princeton University Press.
- Tonk L, Welker M, Huisman J, Visser PM.** 2009. Production of cyanopeptolins, anabaenopeptins, and microcystins by the harmful cyanobacteria *Anabaena* 90 and *Microcystis* PCC 7806. *Harmful Algae* **8**, 219–224.
- Trimborn S, Brenneis T, Sweet E, Rost B.** 2013. Sensitivity of Antarctic phytoplankton species to ocean acidification: growth, carbon acquisition, and species interaction. *Limnology and Oceanography* **58**, 997–1007.
- Tu CK, Acevedo-Duncan M, Wynns GC, Silverman DN.** 1986. Oxygen-18 exchange as a measure of accessibility of CO<sub>2</sub> and HCO<sub>3</sub><sup>-</sup> to carbonic anhydrase in *Chlorella vulgaris* (UTEX 263). *Plant Physiology* **80**, 997–1001.
- Van de Waal DB, Verspagen JMH, Finke JF, et al.** 2011. Reversal in competitive dominance of a toxic versus non-toxic cyanobacterium in response to rising CO<sub>2</sub>. *The ISME Journal* **5**, 1438–1450.
- Van de Waal DB, Verspagen JMH, Lüring M, Van Donk E, Visser PM, Huisman J.** 2009. The ecological stoichiometry of toxins produced by harmful cyanobacteria: an experimental test of the carbon-nutrient balance hypothesis. *Ecology Letters* **12**, 1326–1335.
- Verschoor AM, Van Dijk MA, Huisman J, Van Donk E.** 2013. Elevated CO<sub>2</sub> concentrations affect the elemental stoichiometry and species composition of an experimental phytoplankton community. *Freshwater Biology* **58**, 597–611.
- Verspagen JMH, Passarge J, Jöhnk KD, Visser PM, Peperzak L, Boers P, Laanbroek HJ, Huisman J.** 2006. Water management strategies against toxic *Microcystis* blooms in the Dutch delta. *Ecological Applications* **16**, 313–327.
- Verspagen JMH, Van de Waal DB, Finke JF, Visser PM, Huisman J.** 2014a. Contrasting effects of rising CO<sub>2</sub> on primary production and ecological stoichiometry at different nutrient levels. *Ecology Letters* **17**, 951–960.
- Verspagen JMH, Van de Waal DB, Finke JF, Visser PM, Van Donk E, Huisman J.** 2014b. Rising CO<sub>2</sub> levels will intensify phytoplankton blooms in eutrophic and hypertrophic lakes. *PLoS ONE* **9**, e104325.

- Visser PM, Verspagen JMH, Sandrini G, Stal LJ, Matthijs HCP, Davis TW, Paerl HW, Huisman J.** 2016. How rising CO<sub>2</sub> and global warming may stimulate harmful cyanobacterial blooms. *Harmful Algae* **54**, 145–159.
- Wang Y, Duanmu D, Spalding MH.** 2011. Carbon dioxide concentrating mechanism in *Chlamydomonas reinhardtii*: inorganic carbon transport and CO<sub>2</sub> recapture. *Photosynthesis Research* **109**, 115–122.
- Wang Y, Stessman DJ, Spalding MH.** 2015. The CO<sub>2</sub> concentrating mechanism and photosynthetic carbon assimilation in limiting CO<sub>2</sub>: how *Chlamydomonas* works against the gradient. *The Plant Journal* **82**, 429–448.
- Watson SB, Ridal J, Boyer GL.** 2008. Taste and odour and cyanobacterial toxins: impairment, prediction, and management in the Great Lakes. *Canadian Journal of Fisheries and Aquatic Sciences* **65**, 1779–1796.
- Wilson JB, Spijkerman E, Huisman J.** 2007. Is there really insufficient support for Tilman's R\* concept? A comment on Miller *et al.* *The American Naturalist* **169**, 700–706.
- Wolf-Gladrow DA, Zeebe RE, Klaas C, Körtzinger A, Dickson AG.** 2007. Total alkalinity: the explicit conservative expression and its application to biogeochemical processes. *Marine Chemistry* **106**, 287–300.
- Yang S, Jin X.** 2008. Critical light intensities for *Microcystis aeruginosa*, *Scenedesmus quadricauda* and *Cyclotella* sp. and competitive growth patterns under different light:N:P ratios. *Journal of Freshwater Ecology* **23**, 387–396.