

ARTICLE

Open Access

Developing gene-tagged molecular markers for evaluation of genetic association of apple *SWEET* genes with fruit sugar accumulation

Qiaoling Zhen^{1,2}, Ting Fang^{1,2}, Qian Peng^{1,2}, Liao Liao¹, Li Zhao¹, Albert Owiti^{1,2} and Yuepeng Han^{1,3,4}

Abstract

Sugar content is an important component of fruit quality. Although sugar transporters are known to be crucial for sugar accumulation, the role of genes encoding *SWEET* sugar transporters in fruit sugar accumulation remains elusive. Here we report the effect of the *SWEET* genes on fruit sugar accumulation in apple. A total of 25 *MdSWEET* genes were identified in the apple genome, and 9 were highly expressed throughout fruit development. Molecular markers of these 9 *MdSWEET* genes were developed and used for genotyping of 188 apple cultivars. The association of polymorphic *MdSWEET* genes with soluble sugar content in mature fruit was analyzed. Three genes, *MdSWEET2e*, *MdSWEET9b*, and *MdSWEET15a*, were significantly associated with fruit sugar content, with *MdSWEET15a* and *MdSWEET9b* accounting for a relatively large proportion of phenotypic variation in sugar content. Moreover, both *MdSWEET9b* and *MdSWEET15a* are located on chromosomal regions harboring QTLs for sugar content. Hence, *MdSWEET9b* and *MdSWEET15a* are likely candidates regulating fruit sugar accumulation in apple. Our study not only presents an efficient way of implementing gene functional study but also provides molecular tools for genetic improvement of fruit quality in apple-breeding programs.

INTRODUCTION

Sugar is the main carbon source and energy-supplying substance in organisms, and it plays an important role in plant growth and development. Sugar is the main product of photosynthesis and carbon dioxide assimilation occurs mainly in stromal cells of chloroplast. The main assimilation product synthesized in chloroplasts is triose phosphate, most of which will be converted to either sucrose in the cytosol or starch in the chloroplast^{1,2}. Sucrose is commonly translocated to other carbon-demanding organs through the long-distance transport

occurring in phloem. Hence, sugar transportation is critical for maintaining the source-sink balance³.

Over the past two decades, various sugar transporters have been identified in living organisms, including plants, animals, humans, and fungi^{4–6}. These transporters can be categorized into four families: sodium solute symporter transporters; major facilitator superfamily transporters; phosphotransferase system transporters; and sugar will eventually be exported transporters (*SWEET*s). Among these transporters, *SWEET*s emerge as a unique and novel class of sugar transporters. *SWEET*s are conserved evolutionarily and exist widely in eukaryotes, and prokaryotes such as archaea and eubacteria. The first member of the *SWEET* gene family, *MtN3*, was identified in *Medicago truncatula*, and its expression was induced during root nodule development⁷. Later, a homolog of the *MtN3* gene, designated *Saliva*, was identified in *Drosophila* and

Correspondence: Yuepeng Han (yphan@wbjcas.cn)

¹Key Laboratory of Plant Germplasm Enhancement and Specialty Agriculture, Wuhan Botanical Garden of the Chinese Academy of Sciences, Wuhan 430074, China

²Graduate University of Chinese Academy of Sciences, 19A Yuquanlu, Beijing 100049, China

Full list of author information is available at the end of the article

© The Author(s) 2018



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

displayed salivary gland-specific expression during embryonic development⁸. Recently, MtN3/Saliva-type genes were functionally characterized as sugar transporters in both animals and plants, and thus gave the name “*SWEET*”⁹. *SWEET* proteins are characterized by the MtN3/Saliva motif (also known as the PQ-loop repeat), which comprises three alpha-helical transmembrane domains (3-TMs). Eukaryotic *SWEET*s consists of a tandem repeat of the basic 3-TM unit separated by a single transmembrane domain, which constitutes a 3-1-3 TM structure. In contrast, prokaryotic *SWEET*s, also called *SemiSWEET*s, contain only a single 3-TM unit, suggesting that eukaryotic *SWEET*s evolved through a duplication of the 3-TM unit¹⁰.

In plants, *SWEET*s function as bidirectional uniporters that mediate influx and efflux of sugars across cell membranes. *SWEET*s can be divided into four clades¹¹. Clades I, II, and IV *SWEET*s transport predominantly hexoses, whilst clade III *SWEET*s appear to be sucrose transporters (SUTs)^{12–15}. For example, the clade I *SWEET* *AtSWEET1* and the clade II *SWEET* *OsSWEET5* mediate the uptake and efflux of glucose or galactose, respectively, across the cell membrane^{9,16}. Two clade III *SWEET*s, *AtSWEET11* and *AtSWEET12*, are involved in efflux of photosynthesized sucrose from phloem parenchyma cells into intercellular space for phloem loading and long-distance translocation of sucrose¹⁷. A newly identified clade III *SWEET*, *AtSWEET9*, transports sucrose to the apoplast and plays an essential role in nectar secretion¹⁸. The clade IV *SWEET*, *AtSWEET17*, functions as a fructose-specific uniporter, with a key role in facilitating bidirectional transport of fructose across the tonoplast in leaf and root^{12,13}. In addition, *SWEET*s have been shown to affect various physiological processes, such as pollen development^{19,20}, seed filling^{21,22}, stress and senescence^{14,16,23,24}, modulating gibberellins response²⁵, and host–pathogen interaction²⁶. Usually, abiotic stress such as cold, high salinity, and drought, and biotic stress caused by fungi or bacteria result in an induction of specific *SWEET* genes^{27,28}.

The accumulation of carbohydrates in storage organs such as seeds and fruits mainly depends on the supply of photoassimilates from photosynthetic tissues, especially source leaves. In most plants, sucrose is the major carbohydrate transported over a long distance in the veins to support the growth and development of storage organs⁶. Sugar transporter genes exhibit divergent evolutionary patterns and play important roles in sugar accumulation in plants²⁹. For example, genes encoding SUT proteins are involved in sugar translocation toward storage organs^{30–32}. Recently, the *SWEET4* gene is also found to play an important role in seed filling²². In maize and rice, *SWEET4* shows high expression during seed development and contributes to seed filling by enhancing the

importation of hexoses into the endosperm. This study sheds light on our understanding on the mechanism by which sucrose is released from maternal tissues such as seed coat to support filial tissues such as embryonic tissue. Since sugar content is an important component of fruit quality, increasing attention has also been paid to investigate the *SWEET* gene family in fruit crops such as apple³³, grapevine³⁴, and banana³⁵. However, the effect of *SWEET*s on fruit sugar accumulation remains elusive.

The domesticated apple, *Malus x domestica* Borkh., is an economically important fruit crop worldwide. Apple belongs to the family Rosaceae, and the cultivated apple is a diploidized autopolyploid species with a basic chromosome number of $x = 17$. The draft genome sequence of the domesticated apple has been released, and accounts for an approximately 750 Mb per haploid³⁶. In this study, we report the identification of the *MdSWEET* genes with high expression during apple fruit development. DNA markers for these *MdSWEET*s were developed to investigate their association with fruit sugar content. Our study not only aids our better understanding of the effect of *SWEET*s on fruit sugar accumulation but it will also be helpful for genetic improvement of fruit sugar accumulation in apple-breeding programs.

MATERIALS AND METHODS

Plant material

All 188 apple cultivars (Table S1) used in this study that show a great variation in fruit sweetness are maintained at Xingcheng Institute of Pomology of the Chinese Academy of Agricultural Sciences, Xingcheng, Liaoning, China. Young leaves used for genomic DNA extraction were collected in the spring of 2015. Leaf samples were immediately frozen in liquid nitrogen, and then stored at -80°C until use. Fruits at mature stage were randomly collected in 2015 and fruit maturity was comprehensively estimated based on skin background color and blush development, seed color turning into brown, and the previous records of maturity date. Each cultivar had three replicates, consisting of nine fruits. Fruit samples were cut into small pieces, immediately frozen in liquid nitrogen, and then stored either at -40°C for sugar measurement or at -80°C for real-time PCR (RT-PCR) analysis.

Measurement of soluble sugar content

The content of soluble sugar components was measured using high-performance liquid chromatography (HPLC) according to our previous report³⁷. Briefly, fruit samples were ground into fine powder in liquid nitrogen using an A11 basic Analytical mill (IKA, Germany). One gram of powder was dissolved in 6 ml sterilized deionized water, the mixture was extracted in an ultrasonic bath for 30 min, and then centrifuged at 6000 r/s for 15 min. The supernatant was collected, purified using a

SEPC18 syringe (Supelclean ENVI C18 SPE), and subsequently filtered through a 0.22 μm Sep-Pak filter. The filtered supernatant was used for sugar content measurement using a Dionex P680 HPLC system (Dionex Corporation, CA, USA) equipped with a refractive index detector (Shodex RI-101; Shodex Munich, Germany). The separation was performed on a Transgenomic COREGET-87C column (7.8 mm \times 300 mm, 10 μm) together with a guard column Transgenomic CARB Sep Coregel 87C. The column temperature was maintained at 85 $^{\circ}\text{C}$ by using a Dionex TCC-100 thermostated column compartment. The mobile phase was set at a flow rate of 0.6 ml/min with degassed, distilled, deionized water. Peak areas were integrated with the Chromeleon chromatography data system according to external standard solution calibrations (reagents from Sigma Chemical Co., Castle Hill, NSW, Australia). Sugar concentrations were expressed on a fresh weight (FW) basis, and total sugar content was indicated by the amount of four sugars found in apple fruit, i.e., glucose, sucrose, fructose, and sorbitol. In addition, the measurement of soluble solid content (SSC) was conducted using a pocket refractometer (Atago, Tokyo, Japan).

Identification of the *SWEET* genes in apple and their phylogenetic analysis

Coding DNA sequences of the *SWEET* gene family in *Arabidopsis thaliana* were retrieved from the Arabidopsis Information Resource (TAIR, <http://www.arabidopsis.org/>). These coding sequences were used as query sequences to compare against the apple genome sequence database (GDDH13 V1.1, <https://www.rosaceae.org/blast/>) by BlastX with a cutoff *E*-value of 1.00E-10. The homologs of *Arabidopsis SWEET* genes in the apple genome were named according to their phylogenetic relationships to the founding members of the family in *Arabidopsis*⁹. Chromosome lengths and gene locations were presented according to the draft genome sequence of the doubled haploid (GDDH13 V1.1)³⁸.

Multiple alignment of amino-acid sequences of *SWEET* genes in *Arabidopsis* and apple was conducted using the integrated MUSCLE alignment program in MEGA5 (Molecular Evolutionary GenetiMd Analysis) with default parameters³⁹. The resulting data matrix was analyzed using the Neighbor-Joining method. The bootstrap consensus tree was inferred from 1000 replicates and the bootstrap values <50% were collapsed.

RNA isolation and quantitative RT-PCR

Two apple cultivars, K9 and Shizishan 2, were randomly selected for quantitative RT-PCR (qRT-PCR) analysis. Fruit samples were collected at 30, 60, and 90 days after full bloom. Each cultivar had three biological replicates, containing of nine fruits. Fruits of each replicate were cut

into small pieces, mixed, and used for total RNA extraction. RNA extraction was conducted using RNA prep Pure Plant Kit (TianGen, Beijing, China) according to the manufacturer's instructions. RNA concentration and quantity were detected and assessed with NanoDrop2000 (Thermo Scientific).

Approximately 1 μg of total RNA was used to synthesize the first strand of cDNA using TransScript One-Step gDNA Removal and cDNA Synthesis SuperMix (TRANS, Beijing, China) following the manufacturer's instructions. qRT-PCR was performed in 20 μL reaction containing 1 \times SYBR Green II Master Mix (Takara, Dalian, China), 0.2 μM of each primer, and 0.5 μL of template cDNA. The qRT-PCR amplifications were performed using the Applied Biosystems 7500 Real-Time PCR System (Applied Biosystems, USA), and the reaction program was set as follows: 95 $^{\circ}\text{C}$ for 1 min, one cycle; and 95 $^{\circ}\text{C}$ for 5 s, 60 $^{\circ}\text{C}$ for 34 s, 40 cycles. Melting curve analysis was performed at the end of 40 cycles by heating from 55 to 95 $^{\circ}\text{C}$ at a rate of 0.5 $^{\circ}\text{C}/\text{s}$. An apple polymer ubiquitin enzyme gene (*UBQ*) was selected as a constitute control⁴⁰. The relative expression level of all detected genes was calculated according to the cycle threshold $2^{-\Delta\Delta\text{CT}}$ method. All analyses were performed in triplicates. The primer sequences are listed in Table S2.

Development of gene-tagged simple sequence repeat and cleaved amplified polymorphism sequence markers

Simple sequence repeat (SSR) or cleaved amplified polymorphism sequence (CAPS) markers were development for *SWEET*s that showed high expression in fruit of apple. SSR Hunter software and dCAPS Finder 2.0 (<http://helix.wustl.edu/dcaps/dcaps.html>) were used to screen SSRs with ≥ 7 repeats or CAPS markers, respectively, in genomic DNA sequences of each *SWEET* gene, including 2 kb upstream of the start codon, the entire coding region, and 2 kb downstream of the translation stop codon. Primers flanking the SSR and CAPS loci were designed using the Primer 5 program, and polymorphism of the SSR and CAPS markers was evaluated using four apple cultivars with high sugar content and four cultivars with low sugar content.

PCR amplification was performed using the GeneAmp PCR System 9700 (ABI, USA) with the following condition: 3 min at 94 $^{\circ}\text{C}$, followed by 35 cycles consisting of 94 $^{\circ}\text{C}$ for 30 s, 60 $^{\circ}\text{C}$ for 30 s, 72 $^{\circ}\text{C}$ for 30 s, and with a final extension of 72 $^{\circ}\text{C}$ for 7 min. For the SSR markers, 3 μL of amplification products was mixed with an equal volume of formamide loading buffer (98 % formamide, 10 mM EDTA, pH 8.0, 0.025% bromophenol blue, and xylene cyanol). The mixture was denatured at 95 $^{\circ}\text{C}$ for 5 min, and then immediately put on ice for 5 min. An aliquot of 2 μL mixture was on an 8% polyacrylamide gel and electrophoresed for 1–1.5 h at 1000 V. DNA bands were

visualized after silver staining, and their sizes were estimated on the basis of a standard 25 bp DNA ladder. For the CAPS markers, amplification products were digested with restriction enzymes and then separated on 2% agarose gel. DNA bands in agarose gel were visualized under ultraviolet light after staining with ethidium bromide.

Statistical analysis

A total of 188 apple cultivars were subjected to statistical analysis. Each cultivar was genotyped for the *SWEET* loci following analysis of molecular marker profiles. The detection of association between molecular markers and fruit sugar accumulation was performed with the software package TASSEL version 3.0 according to our previous report⁴¹. The criterion for marker–trait association was set at $P \leq 0.01$. Fisher's least significant difference at $P < 0.01$ was used to compare mean soluble sugar contents between cultivars.

RESULTS

The *SWEETs* gene family in the apple genome

A total of 25 *MdSWEET* genes were identified in the apple genome. Of the 25 *MdSWEET* genes, 24 were located on five homologous pairs of chromosomes (3–11, 5–10, 4–12, 6–14, 13–16), and 1 on chromosome 17 (Fig. 1a). Genomic structural analysis showed that the majority of *MdSWEET* genes consisted of six exons, while 4 *MdSWEET* genes, *MdSWEET5b*, *MdSWEET7a*, *MdSWEET7b*, and *MdSWEET11*, contained five exons (Fig. 1b and Table S3). The open reading frames of the *MdSWEET* genes ranged from 645 to 1020 bp in length and their deduced proteins ranging from 215 to 340 amino acids in length (Table S3). The conserved domain prediction indicated that 21 *MdSWEET* genes had seven alpha-helical TMs. By contrast, 3 *MdSWEET* genes, *MdSWEET5a*, *MdSWEET9a*, and *MdSWEET5b*, had six TMs, with absence of the TM7 domain (Table S3). Interestingly, the remaining *MdSWEET11* gene had eight TMs. In addition, it is worth noting that an additional *SWEET* gene (GDR accession no. MD01G1215700) was also found in the apple genome. This *SWEET* gene was located on chromosome 1, with four exons (Fig. 1b), contained only one Mtn3/Saliva motif, and showed extremely low expression throughout fruit development. Hence, this *SWEET* gene was deemed a pseudogene and was not included in the later analysis.

Phylogenetic analysis revealed that all the *MdSWEET* genes were divided into four clades as previously defined by Chen et al.⁹, and each *MdSWEET* gene was named following its homologous genes in *Arabidopsis* (Fig. 2). Clade I, II, III, and IV contained 10, 5, 9, and 1 *MdSWEET* genes, respectively. The *MdSWEET2* genes had nine homologs, *MdSWEET2a*, *MdSWEET2b*, *MdSWEET2c*,

MdSWEET2d, *MdSWEET2e*, *MdSWEET2f*, *MdSWEET2g*, *MdSWEET2h*, and *MdSWEET2i*, while only one copy was observed for *MdSWEET1*, *MdSWEET8*, *MdSWEET11*, and *MdSWEET17*. The remaining *MdSWEET* genes, *MdSWEET2*, *MdSWEET5*, *MdSWEET7*, *MdSWEET9*, *MdSWEET10*, *MdSWEET12*, and *MdSWEET15* contained two homologs.

Expression profiling of *MdSWEET* genes in fruit at different developmental stages

To identify *SWEET* genes that are potentially involved in fruit sugar accumulation, we investigated the expression profiling of the *MdSWEET* genes in fruits of two cultivars, K9 and Shishan 2, at different developmental stages, including juvenile, expanding, and mature stages (Fig. 3). Of the 25 *MdSWEET* genes, 16 showed extremely low expression throughout fruit development, with relative expression levels ranging from 0 to 0.1. In contrast, 9 *MdSWEET* genes, *MdSWEET2a*, *MdSWEET2b*, *MdSWEET2d*, *MdSWEET2e*, *MdSWEET7a*, *MdSWEET7b*, *MdSWEET9b*, *MdSWEET12a*, and *MdSWEET15a*, were highly expressed in fruits at all the three stages tested. Hence, these 9 *MdSWEET* genes were assumed to be potential candidates related to fruit sugar accumulation, and were further subjected to develop gene-tagged markers for evaluation of their genetic association with fruit sugar content.

Development of gene-tagged markers for *SWEETs* with high expression in fruit and their polymorphisms in a collection of apple cultivars

Two types of molecular markers, SSR and CAPS, were developed for the nine *MdSWEET* genes mentioned above (Table 1). The SSR markers for two *SWEET* genes, *MdSWEET7a* and *MdSWEET9b*, were developed based on a (CT)_n microsatellite located in the second intron, whereas dinucleotide microsatellites, including (AG)_n, (GA)_n, (AT)_n, (CT)_n, and (TA)_n, located upstream of the start codon were used to develop SSR markers for six *SWEET* genes, *MdSWEET2a*, *MdSWEET2b*, *MdSWEET2d*, *MdSWEET2e*, *MdSWEET7b*, and *MdSWEET12a* (Table 1). A T/C single-nucleotide polymorphism (SNP) in the first intron of *MdSWEET15a* was successfully used to develop a CAPS marker. PCR products harboring a “C” nucleotide at the polymorphic site could be digested with the *NdeI* enzyme, producing two fragments of 438 and 203 bp in size, while no digestion for PCR products harboring a “T” nucleotide.

These nine gene-tagged markers were subsequently used to screen a collection of 188 cultivars (Fig. 4). As a result, two alleles at each polymorphic locus were detected for three *MdSWEET* genes, *MdSWEET2a*, *MdSWEET12a*, and *MdSWEET7a*, while three alleles at each polymorphic locus were observed for five *MdSWEET*




Fig. 1 Chromosomal location of the *SWEET* gene family in apple (a) and their genomic structure (b)

genes, *MdSWEET2b*, *MdSWEET2d*, *MdSWEET2e*, *MdSWEET7b*, and *MdSWEET9b*. Six genotypes derived from the random combination of three alleles were identified at each of the four gene loci, including *MdSWEET2b*, *MdSWEET2d*, *MdSWEET2e*, and

MdSWEET7b, while only four genotypes were detected at the *MdSWEET9b* locus (Table S4). Three genotypes at each polymorphic locus were identified for three *MdSWEET* genes, *MdSWEET2a*, *MdSWEET7a*, and *MdSWEET12a*. By contrast, the CAPS marker revealed




Fig. 2 Phylogenetic tree derived from the amino-acid sequences of 25 *MdSWEET* genes in apple and 17 *AtSWEET* genes in *Arabidopsis*. All the *SWEET* genes are divided into four clades (I–IV), and numbers near branches represent bootstrap values. The circles and triangles indicate *SWEET*s from *Arabidopsis* and apple, respectively





Fig. 3 Expression profiles of *MdSWEET*s in fruits of two cultivars K9 and Shishan 2 at different developmental stages. The heat map was constructed based on relative expression levels of the *MdSWEET*s. S1, S2, and S3 represent young fruitlets, expanding fruits, and mature fruits, which were collected 30, 60, and 90 days after full bloom, respectively. The *MdSWEET* genes showing high expression throughout fruit development are highlighted in red color. The different colors represent different levels of gene expression, with the red color representing the highest values of gene expression, followed by yellow, gray, light green, and dark green

Table 1 Primer sequences of molecular markers developed for nine *SWEET* genes highly expressed in fruit of apple

Gene	ID	LG	Gene tag	Type	Motif	Location	Primer sequence (5'–3')		Expected size (bp)
							Forward	Reverse	
<i>MdSWEET2b</i>	MD11G1270800	11	SSR	(AG) ₇	93 bp of USC		TGAGGCAGAAAACAATCATAAGGGTC	GAGCACGGAAITTTGAAGCTGTAAAA	257
<i>MdSWEET2a</i>	MD03G1250600	3	SSR	(GA) ₁₁	102 bp of USC		ATACCGAGGAACCTGTAGGACCAAGC	CTCCACACTAACCAACCAGAAAAGCA	335
<i>MdSWEET2d</i>	MD05G1293100	5	SSR	(AT) ₈	664 bp of USC		CATTCAATTTTATTCGACCCGGAGAC	TGGGTTTCATCCCTCACITTCACCTCA	270
<i>MdSWEET2e</i>	MD10G1269300	10	SSR	(AT) ₇	4203 bp of USC		GTGAGCCCAACTAATCCCAT	CTTGTGGGTAGGAATCCCGATA	219
<i>MdSWEET7b</i>	MD06G112000	6	SSR	(CT) ₁₇	192 bp of USC		GGGTTTTGAGAATCTTGAGGGTAGG	TTTGATGGGTTGGACTGTAACCTGC	251
<i>MdSWEET9b</i>	MD04G1236000	4	SSR	(CT) ₁₉	Second intron		GGCCCAATGTAAGACCCCTTACTTT	CTGACCTTGTCTTCTTGGATGCGTA	348
<i>MdSWEET7a</i>	MD11G1299200	11	SSR	(CT) ₁₄	Second intron		TTCTATCTCCCTTCCAAACTTCC	GCTAAACAGTGCCACTGCATAAGGT	340
<i>MdSWEET12a</i>	MD14G1151300	14	SSR	(TA) ₁₀	1902 bp of USC		ATGACAGGGCAACTTCAGGGT	CGTAATAGTCTTTGCCCTCC	253
<i>MdSWEET15a</i>	MD16G1125300	16	CAPS	T/C	First intron		ACCTACAATCTCCATCTGTC	CCATACAATAACTCACCTGCTTC	641

USC upstream of start codon



only two genotypes, T/T and T/C, and the homologous genotype C/C was not identified in all the cultivars tested.

Association between *MdsWEET* genes and fruit sugar accumulation in apple

SSC and soluble sugar content in mature fruit for all the tested cultivars are listed in Table S1 and their distributions are shown in Fig. 5. SSC and the contents of sucrose, fructose, glucose, and total sugar components showed a normal distribution, whereas the distribution of sorbitol content was skewed toward low sorbitol contents. A wide range was observed for the concentration of various sugar components, including sucrose (3.8–41.75 mg/g FW), glucose (3.86–31.80 mg/g FW), fructose (30.48–83.82 mg/g FW), sorbitol (0.14–22.62 mg/g FW), and total sugar components (47.93–142.40 mg/g FW), with an average of 21.65, 14.98, 49.07, 3.90, and 89.60 mg/g FW, respectively. SSC ranged from 7.96 to 20.02, with an average of 12.76. Overall, the apple cultivar collection has a great variation in fruit sugar content, which suggests that they are suitable for investigating genetic association of *MdsWEET* genes with fruit sugar accumulation.

Candidate gene-based association mapping was further performed to investigate association between the polymorphic loci of *MdsWEET* genes and fruit sugar

accumulation in apple. As a result, three genes, *MdsWEET2e*, *MdsWEET9b*, and *MdsWEET15a*, showed a significant association with fruit sugar accumulation, whereas no significant association was observed for the remaining six genes, *MdsWEET2a*, *MdsWEET2b*, *MdsWEET2d*, *MdsWEET7a*, *MdsWEET7b*, and *MdsWEET12a* (Table 2). Based on the presence or absence of the (AT)₁₃ allele on the *MdsWEET2e* locus, all the tested cultivars were grouped into three genotypes, (AT)_{13/13}, (AT)_{13/7} or ₁₇, and (AT)₇ or _{17/7} or ₁₇. The (AT)_{13/13} genotype had significantly higher than both (AT)_{13/7} or ₁₇ and (AT)₇ or _{17/7} or ₁₇ genotypes for the sucrose, fructose, or total sugar content (Fig. 6). Similarly, all the cultivars were divided into three genotypes, (CT)_{19/19}, (CT)_{19/23}, and (CT)_{23/23} or ₂₆, based on the presence or absence of the (CT)₁₉ allele on the *MdsWEET9b* locus. Cultivars with two (CT)₁₉ alleles had significantly higher than cultivars with one or no (CT)₁₉ allele for the fructose or total sugar content. Based on the polymorphic locus of *MdsWEET15a*, all the cultivars were assigned to two genotypes, T/T and T/C. The T/T genotype had significantly higher than the T/C genotype for SSC and the glucose, sorbitol, or total sugar content. However, the (AT)₁₃ allele of *MdsWEET2e* and the (CT)₁₉ allele of *MdsWEET9b* had no effect on the soluble solids, glucose, or sorbitol content and the soluble solids, sucrose, glucose, or sorbitol content, respectively (data not shown). Similarly, the polymorphic locus of *MdsWEET15a* had no effect on accumulation of sucrose and fructose.

Of the above three genes associated with fruit sugar accumulation, *MdsWEET15a* had relatively higher contributions to the observed phenotypic variation, and accounted for 6.4%, 6.8%, 5.7%, and 8.4% of the observed phenotypic variation in the soluble solids, glucose, sorbitol and total sugar content, respectively. The *MdsWEET9b* gene accounted for 6.6% and 2.5% of the observed phenotypic variation in the fructose and total sugar content, respectively. By contrast, *MdsWEET2e* had lower contribution, accounting for 0.7%, 2.7%, and 3.6% of the observed phenotypic variation in the sucrose, fructose, and total sugar content, respectively.

Taken together, all the above results suggest that one clade II *SWEET* gene, *MdsWEET2e*, and two clade III *SWEET* genes, *MdsWEET9b* and *MdsWEET15a*, are genetically associated with sugar content in mature apple fruit, with *MdsWEET15a* and *MdsWEET9b* showing relatively higher contribution.

DISCUSSION

Sugar transporters play a crucial role in plant growth and development as they mediate sugar uptake or release from cells or subcellular compartments²¹. *SWEET*s are newly identified sugar transporters and they are critical for sugar transportation and sink tissue





Table 2 The probability value (*P*-value) of association between polymorphic loci of *MdSWEET* genes and soluble sugar content in mature fruits of 188 apple cultivars

Gene	Average SSC ^a	Average content of soluble sugar components (mg/g FW) ^a				
		Sucrose	Glucose	Fructose	Sorbitol	Total
<i>MdSWEET7b</i>	0.3461	0.4660	0.5707	0.9866	0.4530	0.9098
<i>MdSWEET2d</i>	0.0596	0.1501	0.7462	0.4083	0.8425	0.2916
<i>MdSWEET2b</i>	0.2998	0.3007	0.0256	0.8808	0.6281	0.9957
<i>MdSWEET2e</i>	0.0177	0.0001	0.0642	0.0083	0.0111	0.0001
<i>MdSWEET12a</i>	0.8595	0.8146	0.9991	0.4798	0.1514	0.8063
<i>MdSWEET9b</i>	0.0198	0.8811	0.1550	0.0001	0.4027	0.0085
<i>MdSWEET2a</i>	0.6065	0.9280	0.8084	0.0365	0.9427	0.6298
<i>MdSWEET7a</i>	0.0838	0.0966	0.9905	0.3783	0.0110	0.1418
<i>MdSWEET15a</i>	0.0005	0.0660	0.0005	0.0109	0.0019	0.0001

^aThe *P*-values < 0.01 are highlighted in bold

development^{17,22,42}. Although preliminary analyses of the *SWEET* gene family have been reported in several fruit crops^{33–35}, little is known about the role of *SWEET*s in fruit sugar accumulation. In this study, we report for the first time genetic association of the *SWEET* genes with

fruit sugar content in apple. Our study indicates that developing gene-tagged markers is an efficient way to investigate gene functionality, and the gene-tagged markers can be directly used in breeding programs if they are associated with horticultural traits of interest.



Duplication of *SWEETs* in the apple genome

Gene duplication is a major driving force for recruitment of genes in plants. Apple is diploid, with an autopolyploidization origin^{36,43}. Here our study reveals 25 *SWEETs* in the apple genome, which is inconsistent with a previous report that identifies a total of 29 apple *SWEETs*³³. This inconsistency is likely due to the fact that our analysis is based on of the draft genome sequence of “Golden delicious” doubled-haploid tree (GDDH13 V1.1)³⁸, while the genome sequence of apple cv. Golden Delicious³⁶ was used in the study reported by Wei et al³³. All the *MdsWEET* genes except *MdsWEET8* are located on homologous pairs of chromosomes. For example, chromosomes 6 and 14 are homologous pairs and both contain one copy of each of the four *SWEET* genes, *MdsWEET7*, *MdsWEET12*, *MdsWEET10*, and *MdsWEET5*. The *MdsWEET9* gene has two homologs, which are separately located on the bottom of two homologously paired chromosomes 4 and 12. Chromosomes 5 and 10 are homologous pairs and both contain four *MdsWEET* genes. These results unambiguously

demonstrate that duplication of *SWEET* genes is related to whole-genome duplication (WGD).

Chromosome 3 is homologous to chromosome 11. Chromosome 3 contains a single *SWEET* gene *MdsWEET2a* on the bottom chromosome, whilst a cluster of two *SWEET2* genes, *MdsWEET2b* and *MdsWEET11*, are found on the bottom of chromosome 11. Similarly, chromosomes 13 and 16 are homologous pairs. A cluster of two *SWEET2* genes, *MdsWEET15b* and *MdsWEET17*, are found on the top of chromosome 13, whilst only single *SWEET* gene *MdsWEET15a* located on the homologous region of chromosome 16. These results also suggest that tandem duplication of *SWEET* genes has probably occurred on chromosomes 11 and 13. In addition, it is worth noting that one *SWEET* gene, *MdsWEET8*, is located on the bottom of chromosome 17, but no *SWEET* gene was found on chromosome 9 that is homologous to chromosome 17. Since duplicated gene copies following WGD are prone to be rapidly lost⁴⁴, it is reasonable to speculate that a *SWEET* gene on chromosome 9 may have been lost in the ancestor of apple.

Most WGD-derived duplicated genes are prone to diverge in expression⁴⁵. This case is also detected for the apple *SWEET* genes. For example, two *SWEET* homologs, *MdSWEET15a* and *MdSWEET15b*, which are located on homologous pair of chromosomes 16 and 13, respectively, have undergone divergence in expression. *MdSWEET15a* is highly expressed throughout fruit development, whilst *MdSWEET15b* shows no expression in fruit. Similarly, *MdSWEET9a* and *MdSWEET9b* are located on homologous pair of chromosomes 12 and 4, respectively. *MdSWEET9b* shows high expression throughout fruit development, but *MdSWEET9a* with no expression in fruit. In addition, expression divergence was also observed for tandem duplicated *SWEET* genes. For example, *MdSWEET2b* and *MdSWEET11* are clustered on the bottom of chromosome 11. *MdSWEET2b* is highly expressed throughout fruit development, but *MdSWEET11* shows no expression in fruit. However, it is unclear whether or not the remaining 16 *MdSWEET* genes with extremely low expression in fruit have also diverged in expression.

Taken together, all the results above suggest that the *SWEET* genes in apple have undergone polyploidization and/or segmental duplication during the process of speciation, and some duplicated *SWEET* genes have diverged in expression.

Candidate *MdSWEET*s involved in the regulation of fruit sugar accumulation in apple

Measurement of soluble sugar content in mature fruits of 188 apple cultivars reveals that the average concentration of fructose is over twofold higher than those of sucrose, glucose, and sorbitol, which confirms our previous report of fructose being the major sugar component in mature apple fruit³⁷. Genotyping of the apple cultivar collection using nine gene-tagged molecular markers further indicates that three *MdSWEET* genes, *MdSWEET2e*, *MdSWEET15a*, and *MdSWEET9b*, are associated with fruit sugar accumulation. Six genotypes produced by three alleles at single polymorphic locus were detected for four *SWEET* genes, *MdSWEET7b*, *MdSWEET2d*, *MdSWEET2b*, and *MdSWEET2e*, and three genotypes arisen from two alleles at single polymorphic locus were observed for three *SWEET* genes, *MdSWEET12a*, *MdSWEET2a*, and *MdSWEET7a*. In contrast, the (CT)_{26/26} and (CT)_{19/26} genotypes and the C/C genotype were not found at the *MdSWEET9b* or *MdSWEET15a* loci, respectively, in the apple cultivar collection. Since selection is well-known to be a directional process that results in changes in the frequency of various genotypes in the population, the *MdSWEET15a* and *MdSWEET9b* loci have probably undergone selection during the process of apple domestication.

Linkage mapping of quantitative trait loci (QTLs) for sugar content have revealed many QTLs with minor effects on all linkage groups (LGs) except LG7 and LG17^{46–49}. *MdSWEET15a* is located on the region of LG16, which contains several QTLs for Brix and the amount of sorbitol and fructose^{48,50}. Similarly, *MdSWEET9b* is located on the region of LG4 harboring a QTL for individual sugar content⁴⁹. By contrast, *MdSWEET2e* is located far away (approximately 11.2 Mb) from the region of LG10 harboring a QTL for Brix and sucrose content. In addition, *MdSWEET2e* has a relative smaller contribution to phenotypic variation in sugar content compared with both *MdSWEET15a* and *MdSWEET9b*.

Taken together, all these results above suggest that *MdSWEET15a* and *MdSWEET9b* are likely candidates involved in sugar accumulation in apple fruit. In peach, two major QTLs for fruit sugar content have been reported on the top of LG4 and the bottom of LG5, respectively⁵¹. Interestingly, these two QTL regions both contain *SWEET* genes, based on the draft genome sequence of peach cv. Lovell⁵⁰. The two QTL regions on peach LG4 and LG5 correspond to syntenic blocks on apple LG3 and LG6, respectively, which harbor QTLs for sugar content⁵². Thus, it seems that *SWEET* genes may also function as candidates for sugar accumulation in peach and other Rosaceae fruit crops.

Both *MdSWEET9b* and *MdSWEET15a* belong to the clade III *SWEET*s that are proved to be efficient SUTs¹¹. In this study, the cultivars with the T/C genotype at the *MdSWEET15a* locus have higher average sucrose content in mature fruit compared with those with the T/T genotype. Similarly, the cultivars with two (CT)₁₉ alleles have higher average sucrose content in mature fruit than those with one or no (CT)₁₉ allele. Thus, the clade III *SWEET* genes in apple, similar to clade III *SWEET*s in *Arabidopsis*⁹, may be involved in sucrose transportation. However, the difference in average sucrose content between various genotypes in the *MdSWEET9b* or *MdSWEET15a* loci does not reach a significance level ($P < 0.01$). This might be partially due to the reason that sucrose in the vacuolar has a trend to conversion into hexoses, resulting in fructose being the major sugar component in mature apple fruit³⁷.

Besides *MdSWEET15a* and *MdSWEET9b*, two additional *SWEET* genes, *MdSWEET2a* and *MdSWEET2b*, are also located on the regions of LGs 3 and 11, respectively, which contain QTLs for fruit sorbitol content in apple⁵³. However, these two *MdSWEET* genes both have no significant association with sorbitol content. This could be attributed to certain process of sugar metabolism in fruit, where sorbitol unloaded from leaves into the fruit is converted to fructose or glucose⁵⁴. It is worth noting that

gene-tagged markers developed in this study may not correspond to functional variants that account for associations with sugar content. Thus, further studies are still needed to ascertain whether *MdSWEET2a* and *MdSWEET2b* have an influence on fruit sugar accumulation in apple.

Marker-assisted selection (MAS) is a valuable tool in breeding programs of plants, particularly fruit crops. In this study, a T/C SNP in the first intron of *MdSWEET15a* accounts for 6–8% of phenotypic variation for SSC (6.4%) and total sugar content (8.4%) among apple germplasm, while a (CT)_n SSR locus in the second intron of *MdSWEET9b* explains approximately 7% of phenotypic variation for the concentration of fructose, the major sugar component in apple fruit. Since these two gene-tagged makers account for considerable phenotypic variation, they can serve as efficient tools for genetic improvement of fruit sweetness in apple-breeding programs with MAS.

In summary, our study suggests that both *MdSWEET9b* and *MdSWEET15a* are likely candidates regulating fruit sugar accumulation in apple. It is worthy of study in the future to clarify the functions of *MdSWEET9b* and *MdSWEET15a*.

Acknowledgements

This work was supported by grants from the National Natural Science Foundation of China (Grant Nos. 31420103914) and the Overseas Construction Plan for Science and Education Base, China-Africa Center for Research and Education, Chinese Academy of Sciences (Grant No. SAJC201327).

Author details

¹Key Laboratory of Plant Germplasm Enhancement and Specialty Agriculture, Wuhan Botanical Garden of the Chinese Academy of Sciences, Wuhan 430074, China. ²Graduate University of Chinese Academy of Sciences, 19A Yuquanlu, Beijing 100049, China. ³Sino-African Joint Research Center, Chinese Academy of Sciences, Wuhan 430074, China. ⁴College of Horticulture Science and Engineering, Shandong Agricultural University, Tai-An, Shandong 271018, China

Competing interests

The authors declare that they have no competing interests.

Supplementary Information accompanies this paper at <https://doi.org/10.1038/s41438-018-0024-3>.

Received: 5 October 2017 Revised: 23 January 2018 Accepted: 25 January 2018

Published online: 20 March 2018

References

- Kuhn, C., Franceschi, V. R., Schulz, A., Lemoine, R. & Frommer, W. B. Macromolecular trafficking indicated by localization and turnover of sucrose transporters in enucleate sieve elements. *Science* **275**, 1298–1300 (1997).
- Doerflinger, F. C., Miller, W. B., Nock, J. F. & Watkins, C. B. Variations in zonal fruit starch concentrations of apples - a developmental phenomenon or an indication of ripening? *Hortic. Res-Engl.* **2**, 47 (2015).
- Gottwald, J. R., Kysan, P. J., Young, J. C., Evert, R. F. & Sussman, M. R. Genetic evidence for the in planta role of phloem-specific plasma membrane sucrose transporters. *Proc. Natl Acad. Sci. USA* **97**, 13979–13984 (2000).
- Walgreen, R. A., Lin, J. T., Kinne, R. K. H. & Walle, T. Cellular uptake of dietary flavonoid quercetin 4'-beta-glucoside by sodium-dependent glucose transporter SGLT1. *J. Pharmacol. Exp. Ther.* **294**, 837–843 (2000).
- Severi, E., Hosie, A. H. F., Hawkhead, J. A. & Thomas, G. H. Characterization of a novel sialic acid transporter of the sodium solute symporter (SSS) family and in vivo comparison with known bacterial sialic acid transporters. *FEMS Microbiol. Lett.* **304**, 47–54 (2010).
- Baker, R. F., Leach, K. A. & Braun, D. M. SWEET as sugar: new sucrose effluxers in plants. *Mol. Plant* **5**, 766–768 (2012).
- Gamas, P., de Carvalho-Niebel, F., Lescure, N. & Cullimore, J. V. Use of a subtractive hybridization approach to identify new *Medicago truncatula* genes induced during root nodule development. *Mol. Plant Microbe Interact.* **9**, 233–242 (1996).
- Artero, R. D. et al. *Saliva*, a new drosophila gene expressed in the embryonic salivary glands with homologues in plants and vertebrates. *Mech. Dev.* **75**, 159–162 (1998).
- Chen, L. Q. et al. Sugar transporters for intercellular exchange and nutrition of pathogens. *Nature* **468**, 527–U199 (2010).
- Xuan, Y. H. et al. Functional role of oligomerization for bacterial and plant SWEET sugar transporter family. *Proc. Natl Acad. Sci. USA* **110**, E3685–E3694 (2013).
- Eom, J. S. et al. SWEETs, transporters for intracellular and intercellular sugar translocation. *Curr. Opin. Plant Biol.* **25**, 53–62 (2015).
- Chardon, F. et al. Leaf fructose content is controlled by the vacuolar transporter SWEET17 in *Arabidopsis*. *Curr. Biol.* **23**, 697–702 (2013).
- Guo, W. J. et al. SWEET17, a facilitative transporter, mediates fructose transport across the tonoplast of *Arabidopsis* Roots and Leaves. *Plant Physiol.* **164**, 777–789 (2014).
- Klemens, P. A. W. et al. Overexpression of the vacuolar sugar carrier *AtSWEET16* modifies germination, growth, and stress tolerance in *Arabidopsis*. *Plant Physiol.* **163**, 1338–1352 (2013).
- Chen, L. Q., Cheung, L. S., Feng, L., Tanner, W. & Frommer, W. B. Transport of sugars. *Annu. Rev. Biochem.* **84**, 865–894 (2015).
- Zhou, Y. et al. Overexpression of *OsSWEET5* in rice causes growth retardation and precocious senescence. *PLoS ONE* **9**, e94210 (2014).
- Chen, L. Q. et al. Sucrose efflux mediated by SWEET proteins as a key step for phloem transport. *Science* **335**, 207–211 (2012).
- Lin, I. W. et al. Nectar secretion requires sucrose phosphate synthases and the sugar transporter SWEET9. *Nature* **508**, 546–549 (2014).
- Guan, Y. F. et al. RUPTURED POLLEN GRAIN1, a member of the *MtN3/saliva* gene family, is crucial for exine pattern formation and cell integrity of microspores in *Arabidopsis*. *Plant Physiol.* **147**, 852–863 (2008).
- Sun, M. X., Huang, X. Y., Yang, J., Guan, Y. F. & Yang, Z. N. *Arabidopsis* RPG1 is important for primexine deposition and functions redundantly with RPG2 for plant fertility at the late reproductive stage. *Plant Reprod.* **26**, 83–91 (2013).
- Chen, L. Q. et al. A cascade of sequentially expressed sucrose transporters in the seed coat and endosperm provides nutrition for the *Arabidopsis* embryo. *Plant Cell* **27**, 607–619 (2015).
- Sosso, D. et al. Seed filling in domesticated maize and rice depends on SWEET-mediated hexose transport. *Nat. Genet.* **47**, 1489–1493 (2015).
- Seo, P. J., Park, J. M., Kang, S. K., Kim, S. G. & Park, C. M. An *Arabidopsis* senescence-associated protein SAG29 regulates cell viability under high salinity. *Planta* **233**, 189–200 (2011).
- Le Hir, R. et al. Disruption of the sugar transporters *AtSWEET11* and *AtSWEET12* affects vascular development and freezing tolerance in *Arabidopsis*. *Mol. Plant* **8**, 1687–1690 (2015).
- Kanno, Y. et al. *AtSWEET13* and *AtSWEET14* regulate gibberellin-mediated physiological processes. *Nat. Commun.* **7**, 13245 (2016).
- Chen, L. Q. SWEET sugar transporters for phloem transport and pathogen nutrition. *New Phytol.* **201**, 1150–1155 (2014).
- Verdier, V. et al. Transcription activator-like (TAL) effectors targeting *OsSWEET* genes enhance virulence on diverse rice (*Oryza sativa*) varieties when expressed individually in a TAL effector-deficient strain of *Xanthomonas oryzae*. *New Phytol.* **196**, 1197–1207 (2012).
- Chong, J. et al. The SWEET family of sugar transporters in grapevine: *VvSWEET4* is involved in the interaction with *Botrytis cinerea*. *J. Exp. Bot.* **65**, 6589–6601 (2014).
- Wang, W. et al. Divergent evolutionary pattern of sugar transporter genes is associated with the difference in sugar accumulation between grasses and eudicots. *Sci. Rep.* **6**, 29153 (2016).

30. Baud, S. et al. The AtSUC5 sucrose transporter specifically expressed in the endosperm is involved in early seed development in *Arabidopsis*. *Plant J.* **43**, 824–836 (2005).
31. Hackel, A. et al. Sucrose transporter LeSUT1 and LeSUT2 inhibition affects tomato fruit development in different ways. *Plant J.* **45**, 180–192 (2006).
32. Zhang, W. H. et al. Nutrient loading of developing seeds. *Funct. Plant Biol.* **34**, 314–331 (2007).
33. Wei, X. Y., Liu, F. L., Chen, C., Ma, F. W. & Li, M. J. The *Malus domestica* sugar transporter gene family: identifications based on genome and expression profiling related to the accumulation of fruit sugars. *Front. Plant Sci.* **5**, 569 (2014).
34. Lecourieux, F. et al. An update on sugar transport and signalling in grapevine. *J. Exp. Bot.* **65**, 821–832 (2014).
35. Miao, H. X. et al. Genome-wide analyses of SWEET family proteins reveal involvement in fruit development and abiotic/biotic stress responses in banana. *Sci. Rep.* **7**, 3536 (2017).
36. Velasco, R. et al. The genome of the domesticated apple (*Malus x domestica* Borkh.). *Nat. Genet.* **42**, 833–839 (2010).
37. Ma, B. Q. et al. Comparative assessment of sugar and malic acid composition in cultivated and wild apples. *Food Chem.* **172**, 86–91 (2015).
38. Daccord, N. et al. High-quality de novo assembly of the apple genome and methylome dynamics of early fruit development. *Nat. Genet.* **49**, 3886 (2017).
39. Tamura, K. et al. MEGA5: molecular evolutionary genetics analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods. *Mol. Biol. Evol.* **28**, 2731–2739 (2011).
40. Fang, T. et al. Variation of ascorbic acid concentration in fruits of cultivated and wild apples. *Food Chem.* **225**, 132–137 (2017).
41. Ma, B. Q. et al. Genes encoding aluminum-activated malate transporter II and their association with fruit acidity in apple. *Plant Genome* **8**, 3 (2015).
42. Patil, G. et al. Soybean (*Glycine max*) SWEET gene family: insights through comparative genomics, transcriptome profiling and whole genome re-sequencing analysis. *BMC Genomics* **16**, 520 (2015).
43. Han, Y. P. et al. Integration of physical and genetic maps in apple confirms whole-genome and segmental duplications in the apple genome. *J. Exp. Bot.* **62**, 5117–5130 (2011).
44. Maere, S. et al. Modeling gene and genome duplications in eukaryotes. *Proc Natl Acad. Sci. USA* **102**, 5454–5459 (2005).
45. Blanc, G. & Wolfe, K. H. Functional divergence of duplicated genes formed by polyploidy during *Arabidopsis* evolution. *Plant Cell* **16**, 1679–1691 (2004).
46. Liebhard, R., Kellerhals, M., Pfammatter, W., Jertmini, M. & Gessler, C. Mapping quantitative physiological traits in apple (*Malus x domestica* Borkh.). *Plant Mol. Biol.* **52**, 511–526 (2003).
47. Kenis, K., Keulemans, J. & Davey, M. W. Identification and stability of QTLs for fruit quality traits in apple. *Tree Genet. Genomes* **4**, 647–661 (2008).
48. Kuniyama, M. et al. Identification of QTLs for fruit quality traits in Japanese apples: QTLs for early ripening are tightly related to preharvest fruit drop. *Breed. Sci.* **64**, 240–251 (2014).
49. Ma, B. Q. et al. Construction of a high density linkage map and its application in the identification of QTLs for soluble sugar and organic acid components in apple. *Tree Genet. Genomes* **12**, 1 (2016).
50. Verde, I. et al. The high-quality draft genome of peach (*Prunus persica*) identifies unique patterns of genetic diversity, domestication and genome evolution. *Nat. Genet.* **45**, 487–U447 (2013).
51. Zeballos, J. L. et al. Mapping QTLs associated with fruit quality traits in peach [*Prunus persica* (L.) Batsch] using SNP maps. *Tree Genet. Genomes* **12**, 37 (2016).
52. Illa, E. et al. Comparative analysis of rosaceous genomes and the reconstruction of a putative ancestral genome for the family. *BMC Evol. Biol.* **11**, 9 (2011).
53. Guan, Y. Z., Peace, C., Rudell, D., Verma, S. & Evans, K. QTLs detected for individual sugars and soluble solids content in apple. *Mol. Breed.* **35**, 135 (2015).
54. Yamaki, S. Metabolism and accumulation of sugars translocated to fruit and their regulation. *J. Jpn Soc. Hort. Sci.* **79**, 1–15 (2010).