



Mother's Milk: A Purposeful Contribution to the Development of the Infant Microbiota and Immunity

Kirsty Le Doare^{1,2,3,4}, Beth Holder^{1,2}, Aisha Bassett⁵ and Pia S. Pannaraj^{5,6*}

¹Centre for International Child Health, Imperial College London, London, United Kingdom, ²Paediatrics, Imperial College London, London, United Kingdom, ³Paediatric Infectious Diseases Research Group, St. George's, University of London, London, United Kingdom, ⁴Vaccines & Immunity Theme, MRC Unit The Gambia, Fajara, Gambia, ⁵Division of Infectious Diseases, Children's Hospital Los Angeles, Los Angeles, CA, United States, ⁶Department of Pediatrics and Molecular Microbiology and Immunology, University of Southern California, Los Angeles, CA, United States

OPEN ACCESS

Edited by:

Arnaud Marchant,
Université libre de Bruxelles,
Belgium

Reviewed by:

Valerie Verhasselt,
University of Western Australia,
Australia
Sarah Rowland-Jones,
University of Oxford,
United Kingdom

*Correspondence:

Pia S. Pannaraj
ppannaraj@chla.usc.edu

Specialty section:

This article was submitted to
Vaccines and Molecular
Therapeutics,
a section of the journal
Frontiers in Immunology

Received: 28 October 2017

Accepted: 08 February 2018

Published: 28 February 2018

Citation:

Le Doare K, Holder B, Bassett A and
Pannaraj PS (2018) Mother's Milk:
A Purposeful Contribution to the
Development of the Infant
Microbiota and Immunity.
Front. Immunol. 9:361.
doi: 10.3389/fimmu.2018.00361

Breast milk is the perfect nutrition for infants, a result of millions of years of evolution. In addition to providing a source of nutrition, breast milk contains a diverse array of microbiota and myriad biologically active components that are thought to guide the infant's developing mucosal immune system. It is believed that bacteria from the mother's intestine may translocate to breast milk and dynamically transfer to the infant. Such interplay between mother and her infant is a key to establishing a healthy infant intestinal microbiome. These intestinal bacteria protect against many respiratory and diarrheal illnesses, but are subject to environmental stresses such as antibiotic use. Orchestrating the development of the microbiota are the human milk oligosaccharides (HMOs), the synthesis of which are partially determined by the maternal genotype. HMOs are thought to play a role in preventing pathogenic bacterial adhesion through multiple mechanisms, while also providing nutrition for the microbiome. Extracellular vesicles (EVs), including exosomes, carry a diverse cargo, including mRNA, miRNA, and cytosolic and membrane-bound proteins, and are readily detectable in human breast milk. Strongly implicated in cell-cell signaling, EVs could therefore play a further role in the development of the infant microbiome. This review considers the emerging role of breast milk microbiota, bioactive HMOs, and EVs in the establishment of the neonatal microbiome and the consequent potential for modulation of neonatal immune system development.

Keywords: breast milk, microbiota, microbiome, human milk oligosaccharides, exosomes, extracellular vesicles, infant microbiome, breast milk microbiome

INTRODUCTION

Breastfeeding confers protection against respiratory and gastrointestinal infections and is associated with a reduced risk of inflammatory diseases such as asthma, atopy, diabetes, obesity, and inflammatory bowel disease (1–7). Prolonged and exclusively breastfed infants have improved cognitive development (8, 9). Human milk continues the transfer of immunity from mother to child that started *in utero*, providing a nurturing environment that protects against infection and develops the infant intestinal mucosa, microbiota, and their own immunologic defenses. Breast milk is a specialized secretion in which immune response is highly targeted against microorganisms in the mother's gut and airway, providing an important defense against the same pathogens likely encountered by her infant (10). More recent studies suggest that breast milk not only provides passive protection

but also directly modulates the immunological development of the breastfed infant through a variety of personalized microbial and immune factors transmitted from mother to child (11–14). These early imprinting events are crucial for immunologic and metabolic homeostasis.

Breast milk immune factors are at their highest concentrations in the colostrum (15), suggesting an immunologic function of milk when the infant is at highest risk of exposure to new pathogens. However, they continue to be dynamically present throughout the lactation period. Bioactive factors transferred to the infant *via* breastfeeding including immunoglobulins, cytokines, chemokines, growth factors, hormones, and lactoferrin have been reviewed in detail elsewhere (15–17). This review will focus on the roles of breast milk microbiota in the establishment of the infant intestinal microbiota, human milk oligosaccharides (HMOs) in shaping the microbiota, and extracellular vesicles (EVs) in modulation of the host–microbe interactions. Breast milk microbiota, HMOs, and EVs are emerging as areas of potential therapeutic interests due to their implications for infant immune development, health, and scope for therapeutic manipulation.

BREAST MILK MICROBIOTA

Breast milk comprises several hundred bacterial species and harbors bacteria at concentrations of approximately 1,000 colony-forming units (CFUs)/mL (18, 19). It is estimated that breastfed infants ingest up to 800,000 bacteria daily (20). Following a dose of microbes at birth (21), breast milk is the immediate next fundamental source of microbes seeding the infant's gut (22, 23). Many epidemiologic studies have documented differences in the composition of gut microbiota in breastfed and formula-fed infants (24–26). Human milk directly contributes to the establishment of the infant intestinal microbiome (19, 20, 23, 27–29). Multiple studies have documented the sharing of specific microbial strains of *Bifidobacterium*, *Lactobacillus*, *Enterococcus*, and *Staphylococcus* species between breast milk and infant stool (30–32). During the first month of life, infants who primarily breastfeed share 28% of their stool microbes with their mother's milk microbes. The frequency of shared microbes increases with the proportion of daily breast milk intake in a dose-dependent manner (23). These findings strongly suggest the transfer of microbes from breast milk to the infant gut. Although an interindividual variation in the types and abundance of different bacteria in human milk exists, the bacteria found in the infant gut most resemble the bacteria from their own mother (23).


While early studies employed culture-dependent methods, recent development of culture-independent techniques, such as next-generation sequencing, has expanded our understanding of the composition and diversity of the breast milk microbiome (33–35). *Streptococcus* and *Staphylococcus* species are the most commonly identified bacterial families in human milk, followed by *Bifidobacterium*, *Lactobacillus*, *Propionibacteria*, *Enterococcus*, and members of the *Enterobacteriaceae* family (23, 28, 35, 36). Several hundred bacterial species have been identified with higher diversity in colostrum compared to transition and mature milk (18).

The origin of bacteria in breast milk is not well established. Breast tissue itself contains a diverse population of bacteria (37). A dynamic cycling of bacteria between mother and infant with retrograde flow from maternal commensal skin flora to infant mouth flora during breastfeeding (38) likely contributes to the bacterial communities (39). However, commensal contamination does not fully account for the diversity of human milk microbes or the presence of strictly anaerobic species such as *Bifidobacterium*, *Clostridium*, or *Bacteroides* species. Milk microbial community composition has been shown to differ from communities on the surrounding areolar skin and infant mouth (35, 40). Another proposed theory is an enteromammary pathway whereby maternal intestinal bacteria migrate to the mammary glands *via* an endogenous cellular route during pregnancy and lactation (19, 28, 41). It has been hypothesized that bacteria first translocate the maternal gut by internalization in dendritic cells and then circulate to the mammary gland *via* the lymphatic and blood circulation (42). This specialized form of mother–infant communication of transferring microbes from the mother's gut to the infant *via* breastfeeding needs further investigation.

Maternal factors affect milk microbiota composition and diversity (Figure 1). Higher diversity has been reported in milk from mothers who deliver vaginally compared with C-section by some groups (18, 43, 44) but not others (23, 45). Milk bacterial profiles do not significantly differ in relation to maternal age, infant gender, or race/ethnicity within a geographical region but do differ across geographic locations of Europe, Africa, and Asia (23, 45, 46). *Bifidobacterium* species concentration was higher in term deliveries than preterm deliveries (44). Total bacteria concentration using quantitative PCR is lower in colostrum than in transitional and mature milk, with increasing levels of *Bifidobacterium* and *Enterococcus* species over time (18, 44). Maternal health alters milk microbiota composition and diversity as evidenced by comparative studies of healthy mothers to those with obesity, celiac disease, and human immunodeficiency virus (HIV) (18, 47, 48). Immunomodulatory cytokines secreted in breast milk from healthy women such as transforming growth factor beta (TGF β) 1 and TGF β 2 are associated with increased early-life microbial richness, evenness, diversity, and increased abundance of taxa protective against atopic diseases (49). Unsurprisingly, maternal antibiotic use and chemotherapy decrease bacterial diversity in breast milk (50, 51); how this impacts on the infant microbiome and immune system development in the long term is currently unknown. More studies are warranted to understand how maternal genetics, culture, environment, nutritional status, and inflammatory states from acute or chronic diseases affect breast milk microbiota.

Role of Breast Milk Microbiota in the Infant Gut

Breast milk bacteria have both immediate- and long-term roles in reducing the incidence and severity of bacterial infections in breastfed infants by multiple mechanisms. Commensal bacteria can competitively exclude or express antimicrobial properties against pathogenic bacteria. For example, *Lactobacilli* isolated from breast milk have been shown to inhibit adhesion and growth



of gastrointestinal pathogens, including *Escherichia coli*, *Shigella* spp, *Pseudomonas* spp, and *Salmonella* spp strains (52–54). Five breast milk *Lactobacilli* strains increased mucin gene expression by intestinal enterocytes to form an antibacterial barrier (53). Administration of a breast milk *Lactobacilli* strain in a double-blind controlled trial to infants 6–12 months of age reduced the incidence of gastrointestinal, respiratory, and total infections by 46, 27, and 30%, respectively (55). The significant increase in bacterial counts of *Lactobacilli* and bifidobacteria in the experimental group compared with the controls was thought to explain the reduced clinical infection episodes although the pathogenic bacteria counts were not measured. Another study found that 30% of human milk contains nisin-producing bacteria that can survive passage through the intestine (56). Nisin is a bacteriocin used by the dairy industry to prevent spore germination and inhibit *Clostridium botulinum* and *Bacillus cereus*. *Staphylococcus epidermidis* and *Streptococcus salivarius* from expressed breast milk also possesses antimicrobial activity against pathogenic *Staphylococcus aureus* (20). Although there are many studies of antimicrobial peptides and molecules in the intestine, more studies are necessary to understand the specific antimicrobial activities of breast milk bacteria.

Increasing evidence in animals points to the instrumental role of microbiota in the development and instruction of the immune system (57, 58). In the absence of intestinal bacteria, animals have defects in lymphoid tissue development within the spleen, thymus, and lymph nodes. Germ-free intestines have

reduced numbers of lamina propria CD4+ cells, IgA-producing cells, and hypoplastic Peyer's patches (59). Germ-free mice typically are Th2 skewed but achieve a balance of Th1/Th2 cytokine production after the introduction of symbiotic bacteria (60). Breast milk *Lactobacillus* strains have been shown to enhance macrophage production of Th1 cytokines including IL-2, IL-12, and TNF-alpha (61). An early human study has suggested better Th1 responses in breastfed children compared to formula-fed children with immunomodulating effects lasting beyond weaning (62). Another *in vitro* study showed that *Lactobacillus fermentum* and *Lactobacillus salivarius* were potent activators of natural killer cells affecting innate immunity as well as moderate activators of CD4+ and CD8+ T cells and regulator T cells affecting acquired immunity (63). Breastfed rhesus macaque infants develop distinct gut microbiota and robust populations of memory T cells and T helper 17 cells compared to bottle-fed infants (64). Whether these mechanisms also exist in humans is not yet known.

Critical Window of Opportunity for Immune Effects

The World Health Organization recommends exclusive breastfeeding during the first 6 months of life (65). This time period of exclusive milk ingestion is also a critical window for microbial imprinting (23, 66, 67). The infant microbiome comprises a dynamic community of bacteria that transforms throughout infancy and into early childhood, but the community assembly is

non-random and depends on early-life events (57, 66). Dysbiosis during this critical developmental window during a time of exclusive milk ingestion may have long-term health implications (57, 68). Germ-free mice have an overaccumulation of invariant natural killer (iNKT) cells leading to susceptibility to colitis, but colonization with standard microbiota before 2 weeks of life but not after, normalizes iNKT cell numbers and protected against colitis (69). Similarly, germ-free adult mice have elevated serum IgE levels associated with exaggerated allergic responses, but mice colonized with standard microbiota before 4 weeks of age, but not after, have normal IgE levels (70). Oral administration of *Bifidobacterium breve* in mice induces proliferation of FoxP3+ regulatory T cells, but only if administered during the pre-weaning stage (54). Even transient perturbations in the microbiota in early life with penicillin is sufficient to induce sustained metabolic alterations and changes in the expression of immune genes in mice (68). Longitudinal human cohorts have supported the long-term implications of early dysbiosis. Arrieta et al. showed transient gut dysbiosis during the first 100 days of life put infants at higher risk for asthma (71). The relative abundance of *Lachnospira*, *Veillonella*, *Faecalibacterium*, and *Rothia* was significantly lower in children at risk of asthma. These genera are present in breast milk (23, 36). Fujimura et al. found a microbiota conformation that was significantly associated with a higher risk of atopy; the conformation was only detectable in children younger than 6 months. By using fecal water from these infants cultured *ex vivo* with human adult peripheral T cells, the investigators showed enhanced induction of IL4+ CD4+ T cells and decreased abundance of CD4+ CD25+ FOXP3+ cells, suggesting that dysbiosis promotes CD4+ T cell dysfunction associated with atopy (72). The progressive establishment of the infant microbiota is vital for educating their immune system to tolerance and reactivity to maintain health throughout life. A recent study by Bäckhed et al. suggests that cessation of breastfeeding rather than introduction of solid foods is the major driver in the development of an adult microbiota (73). Indeed, Ding and Schloss found that history of breastfeeding as an infant dictated bacterial community composition as adults (74).

Breast Milk Virome

Viruses are also known to be transmitted through breast milk (75) and likely contribute to the gut ecology of the developing infant. The assembly of phage and eukaryotic components of the infant gut virome is affected by health and nutritional status (76). Breitbart et al. did not find similar viral sequences in maternal breast milk and the infant stool in their one infant followed over time (77). However, a recent study of 25 mother–infant pairs identified bifidobacterial communities and bifidophages in maternal milk and infant stool, strongly suggesting vertical transmission through breastfeeding (78). Because the majority of viruses inhabiting the infant and adult gut are bacteriophages (77, 79), they have the ability to kill bacteria or provide them with potentially beneficial gene functions to shape the bacterial community and long-term health. Longitudinal studies to determine the role of breastfeeding in the establishment of the infant gut virome and the viral–bacterial interactions are warranted.


HUMAN MILK OLIGOSACCHARIDES

Human milk oligosaccharides (HMO) may further influence the establishment of a healthy microbiome, by binding potentially harmful bacteria in the intestinal lumen, asserting direct antimicrobial effects, modulating the intestinal epithelial cell immune response, and thereby promoting the growth of “good bacteria” (Figure 2). HMOs are soluble complex carbohydrates that are synthesized in the mammary glands dependent on maternal genotype, including the genes that determine the Lewis blood group antigen.

HMO are indigestible by the infant. Instead, they function as prebiotics, encouraging the growth of certain strains of beneficial bacteria, such as *Bifidobacterium infantis*, within the infant gastrointestinal tract (80), thus preventing infection by allowing the microbiota to outcompete potential pathogenic organisms (81, 82). Once ingested by the infant, HMOs are thought to inhibit the adherence of pathogens to the intestinal epithelium by acting as a decoy receptor for pathogens, which prevents attachment to host cells, thereby preventing pathogen adhesion and invasion (83). HMOs are also thought to have direct antimicrobial effects on certain pathogens (81). Finally, HMOs have been observed to modulate intestinal epithelial cell responses, as well as act as immune modulators. HMOs alter the environment of the intestine, by reducing cell growth, and inducing differentiation and apoptosis (84). They alter immune responses by shifting T cell responses to a balanced Th1/Th2 cytokine production (85).

Genetic differences are responsible for differences in HMO profiles in breast milk (86–89), although HMO abundance changes throughout lactation. Therefore, mothers possessing different genotypes, and thus different HMO profiles, may protect their infants against certain infections to a greater or lesser extent, depending on the presence of specific HMOs. Likewise, the different HMOs produced alter the types of microbiota colonizing infants, as well as the timing of the establishment of the microbiota (90). Because of their complexity, no human milk identical HMOs have been synthesized. However, non-human milk-derived alternatives that may have similar bioactive properties are gaining interest. In a recent placebo-controlled trial of 4,556 infants from India, a plant oligosaccharide, fructooligosaccharide, was given to infants together with *Lactobacillus plantarum* and demonstrated a reduced risk of sepsis and death in those in the treatment arm (RR, 0.6; CI, 0.48–0.74) compared to those in the control arm (91). The results highlight a potential role for HMOs and non-milk oligosaccharides in preventing neonatal infection.

HMO are thought to play an important role in preventing neonatal diarrheal and respiratory tract infections (92, 93). Several HMOs have been implicated in protection against bacterial and viral infections in neonates, including fucosyltransferase enzyme (FUT3), associated with the Lewis–Secretor gene (89) and 2'-fucosyllactose (2'-FL), associated with the Secretor gene (FUT2) (87). High concentrations of 2'-FL are associated with reduced risk of infant *Campylobacter jejuni* (94) and rotavirus infections (95). However, it has been noted that there is a rotavirus strain-specific effect of different HMOs, both alone and in combination (95, 96). Lewis–secretor positive infants in Burkina



Faso and Nicaragua appear to have increased susceptibility to rotavirus infection compared to Lewis-negative infants. As the Lewis antigen is partially responsible for HMO abundance, this finding may explain the reduced efficacy of the live oral rotavirus vaccine in Africa where the majority of women are Lewis–Secretor negative (96). Conversely, an observational study undertaken in the United States found severe rotavirus gastroenteritis to be essentially absent in children who had a genetic polymorphism that inactivates FUT2 expression on the intestinal epithelium, which may indicate further strain-specific adaptations of HMOs (97). Infants who received milk containing a low concentration of lacto-*N*-difucohexaose have an increased incidence of calicivirus diarrhea (98). Other HMO combinations in breast milk have also been associated with reduced risk of HIV transmission in Zambia (99).

It has been suggested that HMOs could be used therapeutically to harness these antibiotic benefits together with standard antibiotics (100, 101). Research to date has primarily focused on developing such adjuncts by investigating antiadhesive properties of HMOs *in vitro*. These include the ability of HMOs to reduce *Streptococcus pneumoniae* adherence to cells of the oropharynx (102) and gastrointestinal adherence with *Escherichia coli* (103–105). Specific HMOs such as FUT3 have been implicated in increased killing of Group B *Streptococcus* (GBS) *in vitro* (106–108). The Bode laboratories have determined that GBS requires specific HMO to proliferate *in vitro* (101). Further *in vitro* investigation revealed that GBS uses a glycosyltransferase, which incorporates HMOs into the cell membrane, preventing bacterial proliferation. The Townsend and Le Doare laboratories have also identified Lewis–Secretor status to be important in

reducing biofilm associated with GBS (106, 107). Further studies have identified that HMO-2'-FL also acts as a decoy receptor for norovirus (109). Animal models also report increased Th1 responses against RSV in mice given a prebiotic containing HMOs (110). HMOs are emerging as a novel potential adjunct to antibiotic therapy, but there is much uncertainty as to individual HMO function and synthesizing individual HMOs in the laboratory for use in clinical trials has proven problematic.

EXTRACELLULAR VESICLES AND THEIR CARGO

One of the most recently identified breast milk components that may alter the intestinal immune response and subsequent establishment of the microbiota are the extracellular vesicles (EV) that contain a rich protein cargo, capable of influencing the local immune response to bacterial challenge (111, 112). Hence, the discovery 10 years ago that human breast milk contains abundant EVs has garnered a lot of attention in the field (113). EVs contain a diverse cargo, including mRNA, miRNA, and cytosolic and membrane proteins and have been demonstrated to be intricately involved in cell–cell signaling. EVs include exosomes, which form through the endosomal pathway, and are released from cells following fusion of multivesicular endosomes with the plasma membrane. The larger (0.1–2 μM), more heterogenous microvesicles are formed through direct blebbing from the cell plasma membrane. It is important to note that many breast milk studies use the term “exosomes,” but do not separate exosomes from other vesicles, neither conceptually nor physically. Unless the isolation procedure takes advantage of exosomes’ known size or flotation density (e.g.,

through sucrose gradients, or size exclusion chromatography) or their known markers (i.e., by immunomagnetic isolation, e.g., anti-tetraspanin beads), isolated vesicles cannot be definitively described as exosomes. Both ultracentrifugation and PEG-based reagents such as Exoquick™, commonly used in breast milk studies to date, will pellet other vesicles as well as non-vesicular proteins, including RNA-binding proteins. Studies that use these methods have still revealed exciting potential for breast milk EVs, in terms of biomarkers, or biological activity *in vivo*. The Nolte-Hoehn group have published a useful study that compares EV isolation methods from breast milk (114).

Breast milk EVs contain RNA (115), miRNA, and long non-coding RNA (116). Several studies that profiled miRNA in breast milk exosomes found enrichment in multiple biological functions, including regulation of actin cytoskeleton, glycolysis/gluconeogenesis, aminoacyl-tRNA biosynthesis, pentose phosphate pathway, galactose metabolism, and fatty acid biosynthesis, as well as a wide range of immunological pathways (117–120). Likewise, proteomic analysis of human breast milk EVs revealed that the majority of proteins mapped to immune cell origin (121). Interestingly, a large number of these proteins had not been previously identified in human breast milk, demonstrating that exploration of EV cargoes may reveal novel biomarkers and functional pathways for further investigation. Exosomes in bovine milk are also enriched in proteins involved in immune response and growth (122).

Exosomes can mediate delivery of novel functional miRNA and mRNA to recipient cells (123). Whether miRNAs in breast milk exosomes are functional in the human digestive system is still relatively unknown; some studies show that exosomes protect miRNAs from digestion (118, 124), while others show that miRNAs are degraded by intestinal contents (125). Certainly, breast milk mRNAs and miRNAs can be taken up by cells and elicit functional effects *in vitro*, suggesting the exciting possibility that they may be able to alter protein expression at the neonatal mucosal surface, impacting on the development of the infant's immune system. These functional effects demonstrated thus far include inhibition of *in vitro* T cell cytokine production and boosting regulatory T cells (113) and inhibition of HIV-1 infection of dendritic cells (126). Liao et al. also recently demonstrated that milk-derived EVs enter human intestinal crypt-like cells, with some localization to the cell nucleus; thus, this is a potential mechanism for delivery of immunoregulatory genetic material from mother-to-infant cells (127). Administration of breast milk exosomes increases intestinal epithelial proliferation in both pigs (128) and rats (129), suggesting that they also have the potential to promote normal intestinal development and function in neonates. In addition to acting in the intestinal tract, EVs could potentially exert effects in the oropharynx and nasopharynx. Thus, breast milk EVs could alter the neonatal immune response to oral vaccines, respiratory pathogens and colonization.

Extracellular vesicles also have the potential to modulate the host-microbe interaction. Epithelial and immune cell responses to gut microbes *Lactobacillus* or *Bifidobacterium* are modulated in the presence of EVs from serum (111). These EV enhance aggregation and phagocytosis of bacteria, as well as modulating

TLR responses. Whether these activities are also performed by breast milk EVs is not known. As well as human milk, EVs also have been detected in porcine (128), bovine (122), and murine (125) milk, enabling the use of animal models to explore this phenomenon, as well as raising the possibility of there being cross-kingdom cell-cell communication *via* unpasteurized milk. Studies in mice have identified that the absence of EVs decreases the diversity of the pup intestinal microbiome (130). Human studies of the role of exosomes and their cargo in modulating infant intestinal microbiome are limited. However, Kosaka et al. identified miRNA associated with immune regulation within exosomes in breast milk that are particularly abundant in the first 6 months of life, when the neonatal mucosal immune system is developing (118). Recent work investigating the role of miRNA in EVs in the ProPACT trial demonstrated an array of miRNA in human milk that differed between mothers given probiotics and those given placebo but no significant differences in atopy outcomes (131).

The few studies of exosomes in breast milk to date have often been cross-sectional (116), and there is only one study of exosomes in human colostrum (113); milk that is delivered at a key stage for early immune priming. A study of bovine exosomes shows that the immunomodulatory protein cargo changes temporally during lactation (122); thus, detailed exploration of human breast milk EV cargo across the course of lactation could yield data that are highly relevant to the development of the neonatal immune system. Isolation of exosomes from breast milk to investigate the miRNA and protein cargo that could be delivered to the infant mucosa would offer novel insight into potential delivery mechanisms for drugs with intestinal immunomodulatory factors (132). Furthermore, improved knowledge of the stability and functionality of EV cargoes *in vivo* is vital for our understanding of how breast milk improves neonatal health and immunity.

FUTURE DIRECTIONS

Breast Milk Microbiota

Many unanswered questions regarding the microbiome need further exploration. We need more studies to define the mechanism by which the microbiota impact immune development and how dysbiosis leads to gut inflammation. Greater comprehension beyond the community profile to elucidate function and metabolites produced by the microbes is integral to utilizing these pathways to improve health or alter disease outcomes. If an enteromammary pathway is confirmed, we could exert a positive influence on infant health by modulating the maternal gut microbiota. Breast milk studies to date have mainly focused on the bacterial component. We also need to further understand how the milk virome and mycobiome influence infant gut health.

Breast Milk HMOs

Further questions surround the HMOs, namely, functions of individual HMO and synthesis of HMO in the laboratory for nutrition supplementation; manipulation of HMO expression; and their delivery to establish a healthy microbiome. It is also possible that early intervention (within the first few days of life) is required for such therapies to be successful.

Breast Milk EVs

For future studies of breast milk EVs, including exosomes, it is key to ensure that the correct nomenclature is utilized, based on the isolation methods used. Utilizing the guidelines of the International Society for Extracellular Vesicles (114) and reporting isolation methods through the new EV-TRACK database (133) will greatly aid the field of breast milk EVs. Apoptotic bodies have been seen as something to deplete in breast milk studies to date. Nothing is known about their cargo nor their function in breast milk, but they could play an important biological function in the neonate, as seen in other fields. We also lack detailed understanding of how breast milk EVs change over the course of lactation in humans. We need to understand better how breast milk EVs survive *in vivo* in the oropharynx, nasopharynx, and the gut, where their delivery would be critical. Finally, a human model of EV interaction with the neonatal microbiome would also give critical insight into possible mechanisms that could be harnessed to protect infants from disease and aid intestinal immune development in term and preterm infants alike.

Summary

Future research studies should aim for enrollment of mother–infant pairs, large sample sizes, and longitudinal sample collections and include a diverse population to further elucidate variability in the breast milk microbiome, HMOs, and EVs on infant health outcomes. Studies should employ metagenomics, metatranscriptomics, and metabolomics approaches to understand the complete taxonomical, functional, and metabolic profile and create a more

accurate picture of the breast milk contribution to infant health. Studies of the breast milk virome and fungome are warranted. Furthermore, ensuring that a repository of maternal and infant samples is kept for future research is useful in determining the long-term health implications of the gut microbiome present during the critical window. A repository can also present the opportunity to study the multigenerational transmission of microbes, HMOs, and EVs, facilitating a comprehensive understanding of the dynamics of the mother's contribution to the infant immune system.

AUTHOR CONTRIBUTIONS

PP conceived and designed the manuscript. KLD, BH, AB, and PP contributed to the drafting and critical revision of this manuscript. All authors approved the final copy of the manuscript.

ACKNOWLEDGMENTS

The authors would like to acknowledge The Royal Society. This review was presented at The Royal Society's scientific meeting "Reducing neonatal infectious morbidity and mortality: joining up our thinking."

FUNDING

BH is supported by MRC The Gambia. PP is supported by NIH K23 HD072774-02. KLD is supported by the Bill and Melinda Gates Foundation OPP115363, the British Research Council and the Thrasher Foundation.

REFERENCES

- American Academy of Pediatrics Section on Breastfeeding. Breastfeeding and the use of human milk. *Pediatrics* (2012) 129(3):e827–41. doi:10.1542/peds.2011-3552
- Klopp A, Vehling L, Becker AB, Subbarao P, Mandhane PJ, Turvey SE, et al. Modes of infant feeding and the risk of childhood asthma: a prospective birth cohort study. *J Pediatr* (2017) 190:192–9.e2. doi:10.1016/j.jpeds.2017.07.012
- Dogaru CM, Nyffenegger D, Pescatore AM, Spycher BD, Kuehni CE. Breastfeeding and childhood asthma: systematic review and meta-analysis. *Am J Epidemiol* (2014) 179(10):1153–67. doi:10.1093/aje/kwu072
- den Dekker HT, Sonnenschein-van der Voort AM, Jaddoe VW, Reiss IK, de Jongste JC, Duijts L. Breastfeeding and asthma outcomes at the age of 6 years: the Generation R Study. *Pediatr Allergy Immunol* (2016) 27(5):486–92. doi:10.1111/pai.12576
- Azad MB, Vehling L, Lu Z, Dai D, Subbarao P, Becker AB, et al. Breastfeeding, maternal asthma and wheezing in the first year of life: a longitudinal birth cohort study. *Eur Respir J* (2017) 49(5). doi:10.1183/13993003.02019-2016
- Horta BL, Loret de Mola C, Victora CG. Long-term consequences of breastfeeding on cholesterol, obesity, systolic blood pressure and type 2 diabetes: a systematic review and meta-analysis. *Acta Paediatr* (2015) 104(467):30–7. doi:10.1111/apa.13133
- Xu L, Lochhead P, Ko Y, Claggett B, Leong RW, Ananthakrishnan AN. Systematic review with meta-analysis: breastfeeding and the risk of Crohn's disease and ulcerative colitis. *Aliment Pharmacol Ther* (2017) 46(9):780–9. doi:10.1111/apt.14291
- Victora CG, Bahl R, Barros AJ, França GV, Horton S, Krasevec J, et al. Breastfeeding in the 21st century: epidemiology, mechanisms, and lifelong effect. *Lancet* (2016) 387(10017):475–90. doi:10.1016/S0140-6736(15)01024-7
- Kramer MS, Aboud F, Mironova E, Vanilovich I, Platt RW, Matush L, et al. Breastfeeding and child cognitive development: new evidence from a large randomized trial. *Arch Gen Psychiatry* (2008) 65(5):578–84. doi:10.1001/archpsyc.65.5.578
- Brandtzaeg P. The mucosal immune system and its integration with the mammary glands. *J Pediatr* (2010) 156(2 Suppl):S8–15. doi:10.1016/j.jpeds.2009.11.014
- Piper KM, Berry CA, Cregan MD. The bioactive nature of human breast-milk. *Breastfeed Rev* (2007) 15(3):5–10.
- Hanson LA, Korotkova M, Lundin S, Håversen L, Silfverdal SA, Mattsby-Baltzer I, et al. The transfer of immunity from mother to child. *Ann N Y Acad Sci* (2003) 987:199–206. doi:10.1111/j.1749-6632.2003.tb06049.x
- Walker WA, Iyengar RS. Breast milk, microbiota, and intestinal immune homeostasis. *Pediatr Res* (2015) 77(1–2):220–8. doi:10.1038/pr.2014.160
- Mueller NT, Bakacs E, Combellick J, Grigoryan Z, Dominguez-Bello MG. The infant microbiome development: mom matters. *Trends Mol Med* (2015) 21(2):109–17. doi:10.1016/j.molmed.2014.12.002
- Ballard O, Morrow AL. Human milk composition: nutrients and bioactive factors. *Pediatr Clin North Am* (2013) 60(1):49–74. doi:10.1016/j.pcl.2012.10.002
- Ruiz L, Espinosa-Martos I, García-Carral C, Manzano S, McGuire MK, Meehan CL, et al. What's normal? Immune profiling of human milk from healthy women living in different geographical and socioeconomic settings. *Front Immunol* (2017) 8:696. doi:10.3389/fimmu.2017.00696
- Andreas NJ, Kampmann B, Mehring Le-Doare K. Human breast milk: a review on its composition and bioactivity. *Early Hum Dev* (2015) 91(11):629–35. doi:10.1016/j.earlhumdev.2015.08.013
- Cabrera-Rubio R, Collado MC, Laitinen K, Salminen S, Isolauri E, Mira A. The human milk microbiome changes over lactation and is shaped by maternal weight and mode of delivery. *Am J Clin Nutr* (2012) 96(3):544–51. doi:10.3945/ajcn.112.037382
- Jeurink PV, van Bergenhenegouwen J, Jiménez E, Knippels LM, Fernández L, Garssen J, et al. Human milk: a source of more life than we imagine. *Benef Microbes* (2013) 4(1):17–30. doi:10.3920/BM2012.0040

20. Heikkilä MP, Saris PE. Inhibition of *Staphylococcus aureus* by the commensal bacteria of human milk. *J Appl Microbiol* (2003) 95(3):471–8. doi:10.1046/j.1365-2672.2003.02002.x
21. Dominguez-Bello MG, Costello EK, Contreras M, Magris M, Hidalgo G, Fierer N, et al. Delivery mode shapes the acquisition and structure of the initial microbiota across multiple body habitats in newborns. *Proc Natl Acad Sci U S A* (2010) 107(26):11971–5. doi:10.1073/pnas.1002601107
22. Gritz EC, Bhandari V. The human neonatal gut microbiome: a brief review. *Front Pediatr* (2015) 3:17. doi:10.3389/fped.2015.00017
23. Pannaraj P, Li F, Cerini C, Bender J, Yang S, Rollie A, et al. Association between breast milk bacterial communities and establishment and development of the infant gut microbiome. *JAMA Pediatr* (2017) 171(7):647–54. doi:10.1001/jamapediatrics.2017.0378
24. Azad MB, Konya T, Maughan H, Guttman DS, Field CJ, Chari RS, et al. Gut microbiota of healthy Canadian infants: profiles by mode of delivery and infant diet at 4 months. *CMAJ* (2013) 185(5):385–94. doi:10.1503/cmaj.121189
25. Bergström A, Skov TH, Bahl MI, Roager HM, Christensen LB, Ejlerskov KT, et al. Establishment of intestinal microbiota during early life: a longitudinal, explorative study of a large cohort of Danish infants. *Appl Environ Microbiol* (2014) 80(9):2889–900. doi:10.1128/AEM.00342-14
26. Gomez-Llorente C, Plaza-Diaz J, Aguilera M, Muñoz-Quezada S, Bermudez-Brito M, Peso-Echarri P, et al. Three main factors define changes in fecal microbiota associated with feeding modality in infants. *J Pediatr Gastroenterol Nutr* (2013) 57(4):461–6. doi:10.1097/MPG.0b013e31829d519a
27. Jost T, Lacroix C, Braegger C, Rochat F, Chassard C. Vertical mother–neonate transfer of maternal gut bacteria via breastfeeding. *Environ Microbiol* (2014) 16(9):2891–904. doi:10.1111/1462-2920.12238
28. Fernández L, Langa S, Martín V, Maldonado A, Jiménez E, Martín R, et al. The human milk microbiota: origin and potential roles in health and disease. *Pharmacol Res* (2013) 69(1):1–10. doi:10.1016/j.phrs.2012.09.001
29. Boix-Amorós A, Collado MC, Mira A. Relationship between milk microbiota, bacterial load, macronutrients, and human cells during lactation. *Front Microbiol* (2016) 7:492. doi:10.3389/fmicb.2016.00492
30. Martín V, Maldonado-Barragán A, Moles L, Rodríguez-Baños M, Campo RD, Fernández L, et al. Sharing of bacterial strains between breast milk and infant feces. *J Hum Lact* (2012) 28(1):36–44. doi:10.1177/0890334411424729
31. Makino H, Kushiro A, Ishikawa E, Muyaert D, Kubota H, Sakai T, et al. Transmission of intestinal *Bifidobacterium longum* subsp. *longum* strains from mother to infant, determined by multilocus sequencing typing and amplified fragment length polymorphism. *Appl Environ Microbiol* (2011) 77(19):6788–93. doi:10.1128/AEM.05346-11
32. Benito D, Lozano C, Jiménez E, Albújar M, Gómez A, Rodríguez JM, et al. Characterization of *Staphylococcus aureus* strains isolated from faeces of healthy neonates and potential mother-to-infant microbial transmission through breastfeeding. *FEMS Microbiol Ecol* (2015) 91(3). doi:10.1093/femsec/fiv007
33. Martín R, Heilig HG, Zoetendal EG, Jiménez E, Fernández L, Smidt H, et al. Cultivation-independent assessment of the bacterial diversity of breast milk among healthy women. *Res Microbiol* (2007) 158(1):31–7. doi:10.1016/j.resmic.2006.11.004
34. Collado MC, Delgado S, Maldonado A, Rodríguez JM. Assessment of the bacterial diversity of breast milk of healthy women by quantitative real-time PCR. *Lett Appl Microbiol* (2009) 48(5):523–8. doi:10.1111/j.1472-765X.2009.02567.x
35. Hunt KM, Foster JA, Forney LJ, Schütte UM, Beck DL, Abdo Z, et al. Characterization of the diversity and temporal stability of bacterial communities in human milk. *PLoS One* (2011) 6(6):e21313. doi:10.1371/journal.pone.0021313
36. Fitzstevens JL, Smith KC, Hagadorn JI, Caimano MJ, Matson AP, Brownell EA. Systematic review of the human milk microbiota. *Nutr Clin Pract* (2017) 32(3):354–64. doi:10.1177/0884533616670150
37. Urbaniak C, Gloor GB, Brackstone M, Scott L, Tangney M, Reid G. The microbiota of breast tissue and its association with breast cancer. *Appl Environ Microbiol* (2016) 82(16):5039–48. doi:10.1128/AEM.01235-16
38. Ramsay DT, Kent JC, Owens RA, Hartmann PE. Ultrasound imaging of milk ejection in the breast of lactating women. *Pediatrics* (2004) 113(2):361–7. doi:10.1542/peds.113.2.361
39. Biagi E, Quercia S, Aceti A, Beghetti I, Rampelli S, Turrone S, et al. The bacterial ecosystem of mother's milk and infant's mouth and gut. *Front Microbiol* (2017) 8:1214. doi:10.3389/fmicb.2017.01214
40. Bender JM, Li F, Martelly S, Byrt E, Rouzier V, Leo M, et al. Maternal HIV infection influences the microbiome of HIV-uninfected infants. *Sci Transl Med* (2016) 8(349):349ra100. doi:10.1126/scitranslmed.aaf5103
41. Rodríguez JM. The origin of human milk bacteria: is there a bacterial enteromammary pathway during late pregnancy and lactation? *Adv Nutr* (2014) 5(6):779–84. doi:10.3945/an.114.007229
42. Perez PF, Doré J, Leclerc M, Levenez F, Benyacoub J, Serrant P, et al. Bacterial imprinting of the neonatal immune system: lessons from maternal cells? *Pediatrics* (2007) 119(3):e724–32. doi:10.1542/peds.2006-1649
43. Cabrera-Rubio R, Mira-Pascual L, Mira A, Collado MC. Impact of mode of delivery on the milk microbiota composition of healthy women. *J Dev Orig Health Dis* (2016) 7(1):54–60. doi:10.1017/S2040174415001397
44. Khodayar-Pardo P, Mira-Pascual L, Collado MC, Martínez-Costa C. Impact of lactation stage, gestational age and mode of delivery on breast milk microbiota. *J Perinatol* (2014) 34(8):599–605. doi:10.1038/jp.2014.47
45. Urbaniak C, Angelini M, Gloor GB, Reid G. Human milk microbiota profiles in relation to birthing method, gestation and infant gender. *Microbiome* (2016) 4(1):1. doi:10.1186/s40168-015-0145-y
46. Kumar H, du Toit E, Kulkarni A, Aakko J, Linderborg KM, Zhang Y, et al. Distinct patterns in human milk microbiota and fatty acid profiles across specific geographic locations. *Front Microbiol* (2016) 7:1619. doi:10.3389/fmicb.2016.01619
47. Olivares M, Albrecht S, De Palma G, Ferrer MD, Castillejo G, Schols HA, et al. Human milk composition differs in healthy mothers and mothers with celiac disease. *Eur J Nutr* (2015) 54(1):119–28. doi:10.1007/s00394-014-0692-1
48. Gonzalez R, Maldonado A, Martín V, Mandomando I, Fumado V, Metzner KJ, et al. Breast milk and gut microbiota in African mothers and infants from an area of high HIV prevalence. *PLoS One* (2013) 8(11):e80299. doi:10.1371/journal.pone.0080299
49. Sitarik AR, Bobbitt KR, Havstad SL, Fujimura KE, Levin AM, Zoratti EM, et al. Breast milk transforming growth factor beta is associated with neonatal gut microbial composition. *J Pediatr Gastroenterol Nutr* (2017) 65(3):e60–7. doi:10.1097/MPG.0000000000001585
50. Soto A, Martín V, Jimenez E, Mader I, Rodríguez JM, Fernandez L. Lactobacilli and bifidobacteria in human breast milk: influence of antibiotic therapy and other host and clinical factors. *J Pediatr Gastroenterol Nutr* (2014) 59(1):78–88. doi:10.1097/MPG.0000000000000347
51. Urbaniak C, McMillan A, Angelini M, Gloor GB, Sumarah M, Burton JP, et al. Effect of chemotherapy on the microbiota and metabolome of human milk, a case report. *Microbiome* (2014) 2:24. doi:10.1186/2049-2618-2-24
52. Jara S, Sanchez M, Vera R, Cofre J, Castro E. The inhibitory activity of *Lactobacillus* spp. isolated from breast milk on gastrointestinal pathogenic bacteria of nosocomial origin. *Anaerobe* (2011) 17(6):474–7. doi:10.1016/j.anaerobe.2011.07.008
53. Olivares M, Diaz-Ropero MP, Martín R, Rodríguez JM, Xaus J. Antimicrobial potential of four *Lactobacillus* strains isolated from breast milk. *J Appl Microbiol* (2006) 101(1):72–9. doi:10.1111/j.1365-2672.2006.02981.x
54. Lyons A, O'Mahony D, O'Brien F, MacSharry J, Sheil B, Ceddia M, et al. Bacterial strain-specific induction of Foxp3+ T regulatory cells is protective in murine allergy models. *Clin Exp Allergy* (2010) 40(5):811–9. doi:10.1111/j.1365-2222.2009.03437.x
55. Maldonado J, Canabate F, Sempere L, Vela F, Sanchez AR, Narbona E, et al. Human milk probiotic *Lactobacillus fermentum* CECT5716 reduces the incidence of gastrointestinal and upper respiratory tract infections in infants. *J Pediatr Gastroenterol Nutr* (2012) 54(1):55–61. doi:10.1097/MPG.0b013e3182333f18
56. Beasley SS, Saris PE. Nisin-producing *Lactococcus lactis* strains isolated from human milk. *Appl Environ Microbiol* (2004) 70(8):5051–3. doi:10.1128/AEM.70.8.5051-5053.2004
57. Arrieta MC, Stiemsma LT, Amenogbe N, Brown EM, Finlay B. The intestinal microbiome in early life: health and disease. *Front Immunol* (2014) 5:427. doi:10.3389/fimmu.2014.00427
58. Gensollen T, Iyer SS, Kasper DL, Blumberg RS. How colonization by microbiota in early life shapes the immune system. *Science* (2016) 352(6285):539–44. doi:10.1126/science.aad9378

59. Macpherson AJ, Harris NL. Interactions between commensal intestinal bacteria and the immune system. *Nat Rev Immunol* (2004) 4(6):478–85. doi:10.1038/nri1373
60. Mazmanian SK, Liu CH, Tzianabos AO, Kasper DL. An immunomodulatory molecule of symbiotic bacteria directs maturation of the host immune system. *Cell* (2005) 122(1):107–18. doi:10.1016/j.cell.2005.05.007
61. Diaz-Ropero MP, Martin R, Sierra S, Lara-Villoslada F, Rodriguez JM, Xaus J, et al. Two *Lactobacillus* strains, isolated from breast milk, differently modulate the immune response. *J Appl Microbiol* (2007) 102(2):337–43. doi:10.1111/j.1365-2672.2006.03102.x
62. Pabst HF, Spady DW, Pilarski LM, Carson MM, Beeler JA, Krezolek MP. Differential modulation of the immune response by breast- or formula-feeding of infants. *Acta Paediatr* (1997) 86(12):1291–7. doi:10.1111/j.1651-2227.1997.tb14900.x
63. Perez-Cano FJ, Dong H, Yaqoob P. In vitro immunomodulatory activity of *Lactobacillus fermentum* CECT5716 and *Lactobacillus salivarius* CECT5713: two probiotic strains isolated from human breast milk. *Immunobiology* (2010) 215(12):996–1004. doi:10.1016/j.imbio.2010.01.004
64. Ardeshir A, Narayan NR, Méndez-Lagares G, Lu D, Rauch M, Huang Y, et al. Breast-fed and bottle-fed infant rhesus macaques develop distinct gut microbiotas and immune systems. *Sci Transl Med* (2014) 6(252):252ra120. doi:10.1126/scitranslmed.3008791
65. World Health Organization. *Exclusive Breastfeeding*. (2018) Available from: http://www.who.int/nutrition/topics/exclusive_breastfeeding/en/
66. Koenig JE, Spor A, Scalfone N, Fricker AD, Stombaugh J, Knight R, et al. Succession of microbial consortia in the developing infant gut microbiome. *Proc Natl Acad Sci U S A* (2011) 108(Suppl 1):4578–85. doi:10.1073/pnas.1000081107
67. Brandtzaeg P. Mucosal immunity: integration between mother and the breast-fed infant. *Vaccine* (2003) 21(24):3382–8. doi:10.1016/S0264-410X(03)00338-4
68. Cox LM, Yamanishi S, Sohn J, Alekseyenko AV, Leung JM, Cho I, et al. Altering the intestinal microbiota during a critical developmental window has lasting metabolic consequences. *Cell* (2014) 158(4):705–21. doi:10.1016/j.cell.2014.05.052
69. Olszak T, An D, Zeissig S, Vera MP, Richter J, Franke A, et al. Microbial exposure during early life has persistent effects on natural killer T cell function. *Science* (2012) 336(6080):489–93. doi:10.1126/science.1219328
70. Cahenzli J, Koller Y, Wyss M, Geuking MB, McCoy KD. Intestinal microbial diversity during early-life colonization shapes long-term IgE levels. *Cell Host Microbe* (2013) 14(5):559–70. doi:10.1016/j.chom.2013.10.004
71. Arrieta MC, Stiemsma LT, Dimitriou PA, Thorson L, Russell S, Yurist-Doutsch S, et al. Early infancy microbial and metabolic alterations affect risk of childhood asthma. *Sci Transl Med* (2015) 7(307):307ra152. doi:10.1126/scitranslmed.aab2271
72. Fujimura KE, Sitarik AR, Havstad S, Lin DL, Levan S, Fadrosh D, et al. Neonatal gut microbiota associates with childhood multisensitized atopy and T cell differentiation. *Nat Med* (2016) 22(10):1187–91. doi:10.1038/nm.4176
73. Bäckhed F, Roswall J, Peng Y, Feng Q, Jia H, Kovatcheva-Datchary P, et al. Dynamics and stabilization of the human gut microbiome during the first year of life. *Cell Host Microbe* (2015) 17(5):690–703. doi:10.1016/j.chom.2015.04.004
74. Ding T, Schloss PD. Dynamics and associations of microbial community types across the human body. *Nature* (2014) 509(7500):357–60. doi:10.1038/nature13178
75. Townsend CL, Peckham CS, Thorne C. Breastfeeding and transmission of viruses other than HIV-1. *Adv Exp Med Biol* (2012) 743:27–38. doi:10.1007/978-1-4614-2251-8_2
76. Reyes A, Blanton LV, Cao S, Zhao G, Manary M, Trehan I, et al. Gut DNA viromes of Malawian twins discordant for severe acute malnutrition. *Proc Natl Acad Sci U S A* (2015) 112(38):11941–6. doi:10.1073/pnas.1514285112
77. Breitbart M, Haynes M, Kelley S, Angly F, Edwards RA, Felts B, et al. Viral diversity and dynamics in an infant gut. *Res Microbiol* (2008) 159(5):367–73. doi:10.1016/j.resmic.2008.04.006
78. Duranti S, Lugli GA, Mancabelli L, Armanini F, Turrone F, James K, et al. Maternal inheritance of bifidobacterial communities and bifidophages in infants through vertical transmission. *Microbiome* (2017) 5(1):66. doi:10.1186/s40168-017-0282-6
79. Lim ES, Zhou Y, Zhao G, Bauer IK, Droit L, Ndao IM, et al. Early life dynamics of the human gut virome and bacterial microbiome in infants. *Nat Med* (2015) 21(10):1228–34. doi:10.1038/nm.3950
80. Ward RE, Ninonuevo M, Mills DA, Lebrilla CB, German JB. In vitro fermentation of breast milk oligosaccharides by *Bifidobacterium infantis* and *Lactobacillus gasseri*. *Appl Environ Microbiol* (2006) 72(6):4497–9. doi:10.1128/AEM.02515-05
81. Bode L. Human milk oligosaccharides: every baby needs a sugar mama. *Glycobiology* (2012) 22(9):1147–62. doi:10.1093/glycob/cws074
82. Ruiz-Palacios GM, Cervantes LE, Ramos P, Chavez-Munguia B, Newburg DS. *Campylobacter jejuni* binds intestinal H(O) antigen (Fuc alpha 1, 2Gal beta 1, 4GlcNAc), and fucosyloligosaccharides of human milk inhibit its binding and infection. *J Biol Chem* (2003) 278(16):14112–20. doi:10.1074/jbc.M207744200
83. Newburg DS, Ruiz-Palacios GM, Morrow AL. Human milk glycans protect infants against enteric pathogens. *Annu Rev Nutr* (2005) 25:37–58. doi:10.1146/annurev.nutr.25.050304.092553
84. Kuntz S, Kunz C, Rudloff S. Oligosaccharides from human milk induce growth arrest via G2/M by influencing growth-related cell cycle genes in intestinal epithelial cells. *Br J Nutr* (2009) 101(9):1306–15. doi:10.1017/S0007114508079622
85. Eiwegger T, Stahl B, Haidl P, Schmitt J, Boehm G, Dehlink E, et al. Prebiotic oligosaccharides: in vitro evidence for gastrointestinal epithelial transfer and immunomodulatory properties. *Pediatr Allergy Immunol* (2010) 21(8):1179–88. doi:10.1111/j.1399-3038.2010.01062.x
86. Thurl S, Munzert M, Henker J, Boehm G, Muller-Werner B, Jelinek J, et al. Variation of human milk oligosaccharides in relation to milk groups and lactational periods. *Br J Nutr* (2010) 104(9):1261–71. doi:10.1017/S0007114510002072
87. Chaturvedi P, Warren CD, Altaye M, Morrow AL, Ruiz-Palacios G, Pickering LK, et al. Fucosylated human milk oligosaccharides vary between individuals and over the course of lactation. *Glycobiology* (2001) 11(5):365–72. doi:10.1093/glycob/11.5.365
88. Zivkovic AM, German JB, Lebrilla CB, Mills DA. Human milk glycomiome and its impact on the infant gastrointestinal microbiota. *Proc Natl Acad Sci U S A* (2011) 108(Suppl 1):4653–8. doi:10.1073/pnas.1000083107
89. Kobata A. Structures and application of oligosaccharides in human milk. *Proc Jpn Acad Ser B Phys Biol Sci* (2010) 86(7):731–47. doi:10.2183/pjab.86.731
90. Lewis ZT, Totten SM, Smilowitz JT, Popovic M, Parker E, Lemay DG, et al. Maternal fucosyltransferase 2 status affects the gut bifidobacterial communities of breastfed infants. *Microbiome* (2015) 3:13. doi:10.1186/s40168-015-0071-z
91. Panigrahi P, Parida S, Nanda NC, Satpathy R, Pradhan L, Chandel DS, et al. A randomized synbiotic trial to prevent sepsis among infants in rural India. *Nature* (2017) 548(7668):407–12. doi:10.1038/nature23480
92. Newburg DS, Walker WA. Protection of the neonate by the innate immune system of developing gut and of human milk. *Pediatr Res* (2007) 61(1):2–8. doi:10.1203/01.pdr.0000250274.68571.18
93. Morrow AL, Ruiz-Palacios GM, Altaye M, Jiang X, Guerrero ML, Meinzen-Derr JK, et al. Human milk oligosaccharide blood group epitopes and innate immune protection against *Campylobacter* and calicivirus diarrhea in breastfed infants. *Adv Exp Med Biol* (2004) 554:443–6. doi:10.1007/978-1-4757-4242-8_61
94. Morrow AL, Ruiz-Palacios GM, Jiang X, Newburg DS. Human-milk glycans that inhibit pathogen binding protect breast-feeding infants against infectious diarrhea. *J Nutr* (2005) 135(5):1304–7. doi:10.1093/jn/135.5.1304
95. Laucirica DR, Triantis V, Schoemaker R, Estes MK, Ramani S. Milk oligosaccharides inhibit human rotavirus infectivity in MA104 cells. *J Nutr* (2017) 147:1709–14. doi:10.3945/jn.116.246090
96. Nordgren J, Sharma S, Bucardo F, Nasir W, Gunaydin G, Ouermi D, et al. Both Lewis and secretor status mediate susceptibility to rotavirus infections in a rotavirus genotype-dependent manner. *Clin Infect Dis* (2014) 59(11):1567–73. doi:10.1093/cid/ciu633
97. Payne DC, Currier RL, Staat MA, Sahni LC, Selvarangan R, Halasa NB, et al. Epidemiologic association between FUT2 secretor status and severe rotavirus gastroenteritis in children in the United States. *JAMA Pediatr* (2015) 169(11):1040–5. doi:10.1001/jamapediatrics.2015.2002
98. Morrow AL, Ruiz-Palacios GM, Altaye M, Jiang X, Guerrero ML, Meinzen-Derr JK, et al. Human milk oligosaccharides are associated with protection

- against diarrhea in breast-fed infants. *J Pediatr* (2004) 145(3):297–303. doi:10.1016/j.jpeds.2004.04.054
99. Bode L, Kuhn L, Kim HY, Hsiao L, Nissan C, Sinkala M, et al. Human milk oligosaccharide concentration and risk of postnatal transmission of HIV through breastfeeding. *Am J Clin Nutr* (2012) 96(4):831–9. doi:10.3945/ajcn.112.039503
100. Barthelson R, Mobasser A, Zopf D, Simon P. Adherence of *Streptococcus pneumoniae* to respiratory epithelial cells is inhibited by sialylated oligosaccharides. *Infect Immun* (1998) 66(4):1439–44.
101. Bode L. The functional biology of human milk oligosaccharides. *Early Hum Dev* (2015) 91(11):619–22. doi:10.1016/j.earlhumdev.2015.09.001
102. Andersson B, Porrás O, Hanson LA, Lagergard T, Svanborg-Eden C. Inhibition of attachment of *Streptococcus pneumoniae* and *Haemophilus influenzae* by human milk and receptor oligosaccharides. *J Infect Dis* (1986) 153(2):232–7. doi:10.1093/infdis/153.2.232
103. Cravioto A, Tello A, Villafan H, Ruiz J, del Vedovo S, Neeser JR. Inhibition of localized adhesion of enteropathogenic *Escherichia coli* to HEp-2 cells by immunoglobulin and oligosaccharide fractions of human colostrum and breast milk. *J Infect Dis* (1991) 163(6):1247–55. doi:10.1093/infdis/163.6.1247
104. He Y, Liu S, Kling DE, Leone S, Lawlor NT, Huang Y, et al. The human milk oligosaccharide 2'-fucosyllactose modulates CD14 expression in human enterocytes, thereby attenuating LPS-induced inflammation. *Gut* (2016) 65(1):33–46. doi:10.1136/gutjnl-2014-307544
105. Idanpaan-Heikkilä I, Simon PM, Zopf D, Vullo T, Cahill P, Sokol K, et al. Oligosaccharides interfere with the establishment and progression of experimental pneumococcal pneumonia. *J Infect Dis* (1997) 176(3):704–12. doi:10.1086/514094
106. Andreas NJ, Al-Khalidi A, Jaiteh M, Clarke E, Hyde MJ, Modi N, et al. Role of human milk oligosaccharides in Group B *Streptococcus* colonisation. *Clin Transl Immunology* (2016) 5(8):e99. doi:10.1038/cti.2016.43
107. Ackerman DL, Doster RS, Weitkamp JH, Aronoff DM, Gaddy JA, Townsend SD. Human milk oligosaccharides exhibit antimicrobial and antibiofilm properties against Group B *Streptococcus*. *ACS Infect Dis* (2017) 3(8):595–605. doi:10.1021/acsinfecdis.7b00064
108. Lin AE, Autran CA, Szyzka A, Escajadillo T, Huang M, Godula K, et al. Human milk oligosaccharides inhibit growth of group B *Streptococcus*. *J Biol Chem* (2017) 292(27):11243–9. doi:10.1074/jbc.M117.789974
109. Koromyslova A, Tripathi S, Morozov V, Schroten H, Hansman GS. Human norovirus inhibition by a human milk oligosaccharide. *Virology* (2017) 508:81–9. doi:10.1016/j.virol.2017.04.032
110. Schijf MA, Kruijssen D, Bastiaans J, Coenjaerts FE, Garssen J, van Bleek GM, et al. Specific dietary oligosaccharides increase Th1 responses in a mouse respiratory syncytial virus infection model. *J Virol* (2012) 86(21):11472–82. doi:10.1128/JVI.06708-11
111. van Bergenhenegouwen J, Kraneveld AD, Rutten L, Kettelarij N, Garssen J, Vos AP. Extracellular vesicles modulate host-microbe responses by altering TLR2 activity and phagocytosis. *PLoS One* (2014) 9(2):e89121. doi:10.1371/journal.pone.0089121
112. Zempleni J, Aguilar-Lozano A, Sadri M, Sukreet S, Manca S, Wu D, et al. Biological activities of extracellular vesicles and their cargos from bovine and human milk in humans and implications for infants. *J Nutr* (2017) 147(1):3–10. doi:10.3945/jn.116.238949
113. Admyre C, Johansson SM, Qazi KR, Filen JJ, Lahesmaa R, Norman M, et al. Exosomes with immune modulatory features are present in human breast milk. *J Immunol* (2007) 179(3):1969–78. doi:10.4049/jimmunol.179.3.1969
114. Zonneveld MI, Brisson AR, van Herwijnen MJ, Tan S, van de Lest CH, Redegeld FA, et al. Recovery of extracellular vesicles from human breast milk is influenced by sample collection and vesicle isolation procedures. *J Extracell Vesicles* (2014) 3. doi:10.3402/jev.v3.24215
115. Lasser C, Alikhani VS, Ekstrom K, Eldh M, Paredes PT, Bossios A, et al. Human saliva, plasma and breast milk exosomes contain RNA: uptake by macrophages. *J Transl Med* (2011) 9:9. doi:10.1186/1479-5876-9-9
116. Karlsson O, Rodosthenous RS, Jara C, Brennan KJ, Wright RO, Baccarelli AA, et al. Detection of long non-coding RNAs in human breastmilk extracellular vesicles: implications for early child development. *Epigenetics* (2016) 11(10):721–29. doi:10.1080/15592294.2016.1216285
117. Yang M, Song D, Cao X, Wu R, Liu B, Ye W, et al. Comparative proteomic analysis of milk-derived exosomes in human and bovine colostrum and mature milk samples by iTRAQ-coupled LC-MS/MS. *Food Res Int* (2017) 92:17–25. doi:10.1016/j.foodres.2016.11.041
118. Kosaka N, Izumi H, Sekine K, Ochiya T. MicroRNA as a new immune-regulatory agent in breast milk. *Silence* (2010) 1(1):7. doi:10.1186/1758-907X-1-7
119. Zhou Q, Li M, Wang X, Li Q, Wang T, Zhu Q, et al. Immune-related microRNAs are abundant in breast milk exosomes. *Int J Biol Sci* (2012) 8(1):118–23. doi:10.7150/ijbs.8.118
120. Henrick BM, Yao XD, Nasser L, Roozrogousheh A, Rosenthal KL. Breastfeeding behaviors and the innate immune system of human milk: working together to protect infants against inflammation, HIV-1, and other infections. *Front Immunol* (2017) 8:1631. doi:10.3389/fimmu.2017.01631
121. van Herwijnen MJ, Zonneveld MI, Goerdal S, Nolte-t Hoen EN, Garssen J, Stahl B, et al. Comprehensive proteomic analysis of human milk-derived extracellular vesicles unveils a novel functional proteome distinct from other milk components. *Mol Cell Proteomics* (2016) 15(11):3412–23. doi:10.1074/mcp.M116.060426
122. Samuel M, Chisanga D, Liem M, Keerthikumar S, Anand S, Ang CS, et al. Bovine milk-derived exosomes from colostrum are enriched with proteins implicated in immune response and growth. *Sci Rep* (2017) 7(1):5933. doi:10.1038/s41598-017-06288-8
123. Valadi H, Ekstrom K, Bossios A, Sjostrand M, Lee JJ, Lotvall JO. Exosome-mediated transfer of mRNAs and microRNAs is a novel mechanism of genetic exchange between cells. *Nat Cell Biol* (2007) 9(6):654–9. doi:10.1038/ncb1596
124. Hata T, Murakami K, Nakatani H, Yamamoto Y, Matsuda T, Aoki N. Isolation of bovine milk-derived microvesicles carrying mRNAs and microRNAs. *Biochem Biophys Res Commun* (2010) 396(2):528–33. doi:10.1016/j.bbrc.2010.04.135
125. Title AC, Denzler R, Stoffel M. Uptake and function studies of maternal milk-derived microRNAs. *J Biol Chem* (2015) 290(39):23680–91. doi:10.1074/jbc.M115.676734
126. Naslund TI, Paquin-Proulx D, Paredes PT, Vallhov H, Sandberg JK, Gabriellsson S. Exosomes from breast milk inhibit HIV-1 infection of dendritic cells and subsequent viral transfer to CD4+ T cells. *AIDS* (2014) 28(2):171–80. doi:10.1097/QAD.0000000000000159
127. Liao Y, Du X, Li J, Lonnerdal B. Human milk exosomes and their microRNAs survive digestion in vitro and are taken up by human intestinal cells. *Mol Nutr Food Res* (2017) 61(11). doi:10.1002/mnfr.201700082
128. Chen T, Xie MY, Sun JJ, Ye RS, Cheng X, Sun RP, et al. Porcine milk-derived exosomes promote proliferation of intestinal epithelial cells. *Sci Rep* (2016) 6:33862. doi:10.1038/srep33862
129. Hock A, Miyake H, Li B, Lee C, Ermini L, Koike Y, et al. Breast milk-derived exosomes promote intestinal epithelial cell growth. *J Pediatr Surg* (2017) 52(5):755–9. doi:10.1016/j.jpedsurg.2017.01.032
130. Zhou F, Paz HA, Sadri M, Fernando SC, Zempleni J. A diet defined by its content of bovine milk exosomes alters the composition of the intestinal microbiome in C57BL/6 mice. *FASEB J* (2017) 31(1 Suppl):965.24.
131. Simpson MR, Brede B, Johansen J, Johnsen R, Storro O, Saetrom P, et al. Human breast milk miRNA, maternal probiotic supplementation and atopic dermatitis in offspring. *PLoS One* (2015) 10(12):e0143496. doi:10.1371/journal.pone.0143496
132. H Rashed M, Bayraktar E, K Helal G, Abd-Ellah MF, Amero P, Chavez-Reyes A, et al. Exosomes: from garbage bins to promising therapeutic targets. *Int J Mol Sci* (2017) 18(3). doi:10.3390/ijms18030538
133. Consortium E-T, Van Deun J, Mestdagh P, Agostinis P, Akay O, Anand S, et al. EV-TRACK: transparent reporting and centralizing knowledge in extracellular vesicle research. *Nat Methods* (2017) 14(3):228–32. doi:10.1038/nmeth.4185

Conflict of Interest Statement: KLD has received funding for investigator led initiatives from GSK and Pfizer. PP receives funding from AstraZeneca and MedImmune for unrelated studies. All other authors declare no conflicts of interest.

Copyright © 2018 Le Doare, Holder, Bassett and Pannaraj. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.