

Cytogenetic aberrations in multiple myeloma are associated with shifts in serum immunoglobulin isotypes distribution and levels

Multiple myeloma (MM) is characterized by secretion of monoclonal (M) protein from clonal plasma cells.¹ In most cases, intact immunoglobulins (Igs) consisting of heavy and light chains are secreted, whereas in about 20% of cases only free light chains (FLCs) are secreted ('light chain only, LCO, MM').¹ MM is also characterized by a number of cytogenetic abnormalities (CAs) categorized into primary (IgH translocations and hyperdiploid trisomies) and secondary (other CAs) events. CAs may be used for MM risk stratification in routine clinical practice.²⁻⁴ One study showed that cases with IgH translocations had higher FLC levels and abnormal FLC ratio.⁵ Recently, prognostic significance was shown for isotype matched Ig ratios (e.g. IgG κ /IgG λ).^{6,8}

So far, the possible impact of CAs on Ig and FLC subtypes and their levels has been poorly studied in MM. The hypothesis is that CAs are associated with the involved Ig isotypes in MM. We carried out a systematic analysis of Ig and FLC subtypes and their serum levels in relation with common CAs using data on 523 patients recruited in the German-Speaking Myeloma Multicenter Group (GMMG)-MM5 trial.^{9,10} For purpose of replication, an independent cohort of 325 cases from the GMMG-HD4 trial was obtained.^{11,12}

Blood samples were collected prior to treatment initiation in the above trials, co-ordinated by University Clinic Heidelberg. Patients' characteristics are summarized in *Online Supplementary Table S1*.^{9,10} Methods used to measure Ig concentrations and subtypes are described elsewhere¹³ and in the *Online Supplementary Appendix*. CAs were identified for available samples using fluorescence *in situ* hybridization (FISH) techniques.¹¹ To assign translocation positivity, 10% or more of the tested affected cells had to demonstrate positive FISH test. All statistical analyses were performed using R software (version 3.2.3). χ^2 test of independence and Fisher exact test were used to test equality of proportions of CA positive cases. Multivariable linear regression models were fitted to test the relationship between clinical variables and CAs. Covariates included in the models were International Staging System (ISS), sex,

light chain type, bone marrow cell count and secondary CAs.

Collection of patients' samples and associated clinical information within both clinical trials was approved by the ethical review board of Heidelberg University, in accordance with the Declaration of Helsinki.

Table 1 depicts the proportion of CA positive cases in three major MM isotypes. In both cohorts, the most frequent CA was hyperdiploidy (57%). Proportions of three CAs varied significantly depending upon MM type. t(4;14) was significantly higher in IgA MM compared with IgG MM (23% vs. 6%; $P=1.1\times 10^{-5}$) and LCO MM (23% vs. 7%; $P=0.02$). Hyperdiploidy occurred more frequently in IgG MM and IgA MM compared with LCO MM (66% vs. 32%, $P=1.3\times 10^{-6}$ and 53% vs. 32%, $P=0.03$, respectively). t(11;14) was most common in LCO MM compared with IgG MM (41% vs. 19%; $P=6.0\times 10^{-4}$) and IgA MM (41% vs. 14%; $P=4.2\times 10^{-4}$). In the GMMG-HD4 cohort, we replicated association of t(4;14) with IgA MM compared to IgG MM (28% vs. 10%; $P=3.3\times 10^{-3}$) and association of hyperdiploidy with IgG MM compared to LCO MM (66% vs. 35%; $P=6.6\times 10^{-4}$). t(14;16) did not influence MM isotype in either cohort. We further analyzed the association between CAs and MM isotypes by considering possible combinations of heavy-light chains. Similar patterns of differences in the proportions were observed across heavy-light chain matched isotypes, as noticed above, for three CAs. Further, isotype-matched pairwise comparisons showed significant differences in proportions of three CAs for the IgG κ and IgG λ pair (*Online Supplementary Table S2*). In the GMMG-HD4 cohort, directions of changes remained but the case numbers were small. No such differences were observed in the isotype-matched pairwise comparisons for IgA and LCO MM (*data not shown*). These results indicate that the above noticed association between CAs and Ig isotypes is not influenced by the light chain type.

Table 2 shows median concentrations of serum M-protein and rFLC (κ/λ ratio) in CA positive and negative patients for IgG MM and IgA MM. M-protein levels were systematically increased in t(4;14) positive patients compared with t(4;14) negative patients for both IgG MM and IgA MM. Statistical significance was reached for IgA MM alone ($P=0.01$; $\beta=14.36$) after adjusting for the covariates (*see above*). This observation was further supported by a sig-

Table 1. Proportion of positivity for cytogenetic abnormalities (CAs) across multiple myeloma (MM) isotypes, IgG, IgA and LCO.

CA	All cases*	Proportion			P for comparison		
		IgG MM	IgA MM	LCO MM	IgG, IgA	IgG, LCO	IgA, LCO
GMMG-MM5	n=523	n=318	n=106	n=90			
t(4;14)	0.09 (44/475)	0.06 (17/294)	0.23 (22/97)	0.07 (5/76)	1.1$\times 10^{-5}$	1.0	0.02
t(11;14)	0.22 (100/461)	0.19 (55/288)	0.14 (13/95)	0.41 (29/71)	0.90	6.0$\times 10^{-4}$	4.2$\times 10^{-4}$
t(14;16)	0.06 (12/197)	0.06 (7/123)	0.04 (2/45)	0.08 (2/25)	1.0	1.0	1.0
Hyperdiploidy	0.57 (263/458)	0.66 (187/285)	0.53 (50/94)	0.32 (23/72)	0.13	1.3$\times 10^{-6}$	0.03
GMMG-HD4	n=325	n=191	n=74	n=55			
t(4;14)	0.14 (41/295)	0.10 (17/173)	0.28 (18/65)	0.11 (6/53)	3.3$\times 10^{-3}$	1.0	0.15
t(11;14)	0.19 (57/296)	0.19 (33/174)	0.12 (8/65)	0.30 (16/53)	0.93	0.36	0.09
t(14;16)	0.02 (5/284)	0.01 (2/166)	0.02 (1/63)	0.04 (2/51)	1.0	0.72	1.0
Hyperdiploidy	0.57 (162/286)	0.66 (111/169)	0.48 (30/62)	0.35 (18/51)	0.08	6.6$\times 10^{-4}$	0.69

Proportion: each entry represents proportion of cytogenetic abnormalities (CA) positive cases. The number of CA positive cases and the number of cases tested for this CA are shown in brackets. Fisher exact test was used when the expected count was less than 5. Significant differences ($P\leq 0.05$) are shown in bold. *All cases included an additional 9 and 5 cases in GMMG-MM5 and GMMG-HD4, respectively, with involved Ig isotypes as IgM or IgD. N: total number of multiple myeloma (MM) cases with this MM isotype; P: Bonferroni corrected P-value from a χ^2 test of pairwise comparisons of proportions in the three groups: IgG, IgA and LCO for each CA. LCO: light chain only.

nificant positive correlation between percentage of t(4;14) harboring plasma cells and M-protein level for IgA MM (Pearson correlation coefficient, $r = 0.46$; $P = 2.6 \times 10^{-4}$). There was a consistent negative relationship, of borderline significance, between M-protein level and hyperdiploidy status for IgG MM and IgA MM ($P = 0.13$; $\beta = -3.86$ and $P = 0.04$; $\beta = -8.23$, respectively). Differences in M-protein level were smaller and insignificant for the remaining five CAs. The rFLC values were outside the reference range (0.26–1.65) for all cases and, with one exception, they were above the reference range, i.e. ratios shifted in favor of κ chains. Significant differences in rFLC were found for gain 1q21 ($P = 0.01$; $\beta = -0.31$) in IgG MM but not in IgA MM ($P = 0.83$; $\beta = 0.05$).

The median M-protein levels were significantly higher in IgG MM cases compared with IgA MM (36.0 g/L vs. 32.3 g/L; $P = 2.2 \times 10^{-3}$) (Online Supplementary Table S3). Consistently, in the GMMG-HD4 cohort, the level in IgG

MM was also higher compared with IgA MM (42.0 g/L vs. 35.8 g/L; $P = 4.0 \times 10^{-4}$). No differences in M-protein levels were observed across the three IgH translocation types ($P > 0.05$). For IgA MM, median M-protein levels in patients with any IgH translocations was more than double the median M-protein level in hyperdiploidy group (42.4 g/L vs. 19.0 g/L; $P = 1.1 \times 10^{-4}$). The difference remained consistent, but smaller and of borderline significant in the GMMG-HD4 cohort (44.9 g/L vs. 39.1 g/L; $P = 0.06$).

The individual FLC κ levels were outside the reference range for 80% of cases, and for 63% cases exceeded the upper limit (>19.4 mg/L) (Online Supplementary Table S4). FLC λ levels were outside the reference range for 63% cases, and for 31% cases surpassed the upper limit (>26.3 mg/L). Online Supplementary Table S4 shows median concentration of uninvolved Igs and FLCs by CA type for three MM isotypes. Practically all uninvolved IgG, IgA and IgM were below the reference values, indicating immunopare-

Table 2. Association between cytogenetic abnormalities (CA) and concentration of serum M-protein and rFLCs (κ/λ ratio) in IgG MM and IgA MM.

CA	Laboratory parameter	Reference range*	Median (m_1, m_2)	Cases (n_1, n_2)	BETA	P
IgG MM (N=318)						
t(4;14)	M-protein (g/L)	–	50.5, 35.3	17, 277	7.68	0.13
	rFLC	0.26–1.65	21.8, 14.8	17, 274	0.04	0.88
t(11;14)	M-protein (g/L)	–	40.2, 35.6	55, 233	2.24	0.48
	rFLC	0.26–1.65	3.2, 17.2	54, 231	0.04	0.81
t(14;16)	M-protein (g/L)	–	34.5, 36.1	7, 116	-3.27	0.74
	rFLC	0.26–1.65	0.01, 30.2	7, 116	-0.49	0.25
Gain 1q21	M-protein (g/L)	–	37.4, 34.5	100, 189	2.49	0.34
	rFLC	0.26–1.65	7.2, 21.2	100, 186	-0.31	0.01
Hyperdiploidy	M-protein (g/L)	–	34.0, 40.2	187, 98	-3.86	0.13
	rFLC	0.26–1.65	19.3, 7.3	186, 96	0.08	0.52
del(13q)	M-protein (g/L)	–	35.4, 35.9	123, 172	-2.97	0.25
	rFLC	0.26–1.65	9.6, 20.3	122, 170	0.04	0.75
del(17p)	M-protein (g/L)	–	29.2, 36.4	30, 266	-4.57	0.25
	rFLC	0.26–1.65	98.2, 13.0	30, 263	0.26	0.16
IgA MM (N=106)						
t(4;14)	M-protein (g/L)	–	45.3, 28.8	22, 75	14.36	0.01
	rFLC	0.26–1.65	23.2, 5.02	22, 74	0.10	0.75
t(11;14)	M-protein (g/L)	–	28.3, 33.8	13, 82	-1.64	0.80
	rFLC	0.26–1.65	9.8, 6.9	13, 81	-0.27	0.45
t(14;16)	M-protein (g/L)	–	3.7, 35.1	2, 43	-22.64	0.16
	rFLC	0.26–1.65	9.9, 7.2	2, 43	-1.17	0.14
Gain 1q21	M-protein (g/L)	–	35.3, 30.4	54, 41	0.45	0.92
	rFLC	0.26–1.65	3.8, 13.5	53, 41	0.05	0.83
Hyperdiploidy	M-protein (g/L)	–	29.2, 36.5	50, 44	-8.23	0.04
	rFLC	0.26–1.65	9.8, 4.6	49, 44	-0.05	0.83
del(13q)	M-protein (g/L)	–	35.6, 27.8	53, 44	4.68	0.27
	rFLC	0.26–1.65	9.7, 5.0	52, 44	0.30	0.21
del(17p)	M-protein (g/L)	–	34.6, 32.3	13, 84	7.44	0.23
	rFLC	0.26–1.65	10.8, 6.9	13, 83	0.08	0.82

m_1, m_2 : median concentration of serum M-protein and rFLC in cytogenetic abnormalities (CA) positive and negative cases, respectively. n_1, n_2 : number of CA positive and negative cases, respectively. BETA, P: estimate of coefficient and P-value for CA in a multiple linear model involving laboratory parameter (e.g. M-protein and rFLC) as dependent variable and CA, International Staging System (ISS), sex, light chain as independent variables. Logarithmic transformation was performed for rFLC. Median concentrations shown in bold for $P \leq 0.05$. *Reference range as reported by the International Myeloma Foundation. - Not applicable.

sis.¹⁴ On the contrary, FLC κ level was above and FLC λ level was within the reference values. For IgG MM, t(11;14) positive cases had suppressed IgA and IgM levels ($P=0.03$; $\beta=-0.14$ and $P=0.02$; $\beta=-0.13$, respectively). Differences in FLC λ levels were observed for gain 1q21 in IgG MM ($P=0.01$; $\beta=0.20$), thus explaining the above noticed differences in rFLC for this CA (Table 2). For IgA MM, IgG levels were significantly suppressed in cases positive for del(13q) or gain 1q21 ($P=0.03$; $\beta=-0.11$ and $P=3.7\times 10^{-3}$; $\beta=-0.15$, respectively), but significantly elevated in hyperdiploidy cases ($P=0.01$; $\beta=0.13$). Cases with gain 1q21 had IgM levels suppressed ($P=0.01$; $\beta=-0.19$) while cases with del(13q) had FLC λ levels suppressed ($P=0.04$; $\beta=-0.34$). For LCO MM, gain 1q21 positive cases had all the Igs levels significantly suppressed. Hyperdiploidy cases had IgA levels elevated ($P=0.04$; $\beta=0.21$). Cases with t(11;14) had FLC κ levels suppressed ($P=0.02$; $\beta=-0.40$) while cases with hyperdiploidy or del(17p) had elevated FLC κ levels ($P=0.05$; $\beta=0.36$ and $P=0.01$; $\beta=0.78$, respectively).

One main finding of this study was that two CAs showed significant associations with the involved Ig isotypes in both cohorts, hyperdiploidy with IgG MM, and t(4;14) with IgA MM. An association was also found for t(11;14) with LCO MM but it did not reach statistical significance in the replication cohort ($P=0.09$), possibly due to a small sample size. In all these cases, CA positivity contributed to higher proportions of the indicated isotypes. What consequences the detected shifts might have remains speculative. t(4;14) positivity was not only related to the increased proportion of IgA isotype but also showed a significant increase in the M-protein level (45.3 g/L vs. 28.8 g/L) which was the highest measured median value for this isotype. Whether such high concentration might contribute to the poor prognosis in t(4;14) still has to be verified. Moreover, future studies should focus on comparing the clinical outcomes of non-IgA and IgA t(4;14) patients.

In a previous study, higher rFLC was found in patients with t(14;16) or del(13).³ We identified a significant suppression of rFLC in cases with gain 1q21 for IgG MM. M-protein levels in IgG MM were higher compared with those in IgA MM cases. This might be expected, as IgG is the principal isotype in the blood and extracellular fluid whereas IgA is the principal isotype in secretions of mucus epithelium of the intestinal and respiratory tracts. The observation that M-protein levels were higher in any IgH translocation group compared with the hyperdiploidy group can possibly be attributed to the active production of M-protein in actively proliferating plasma cells driven by the translocated cyclin D genes.¹⁵ Patients lacking hyperdiploidy had higher M-protein level compared with hyperdiploidy patients, although this difference was only statistically significant for IgA MM.

In summary, our study provides strong evidence for a complex modifying role of at least two CA types on immunophenotype. We also showed that serum Ig and FLC levels were influenced by several CA types. Therefore, our study sheds new light on the role of CA as an important component of MM pathogenesis, influencing immunoglobulin isotypes and the serum Ig and FLC levels.

Pankaj Yadav,¹ Maximilian Merz,² Elias K. Mai,² Asta Försti,^{1,3} Anna Jauch,⁴ Hartmut Goldschmidt,^{2,5,*} Kari Hemminki^{1,3}

*HG and KH share co-senior authorship.

¹Division of Molecular Genetic Epidemiology, German Cancer Research Center (DKFZ), Heidelberg, Germany; ²Department of Internal Medicine V, University Hospital Heidelberg, Germany; ³Center for Primary Health Care Research, Lund University, Malmö,

Sweden; ⁴Institute of Human Genetics, University of Heidelberg, Germany and ⁵National Centre of Tumor Diseases, Heidelberg, Germany

Correspondence: p.yadav@dkfz.de
doi:10.3324/haematol.2017.184226

Funding: this project was supported by the German Cancer Aid, the Harald Huppert Foundations, The German Federal Ministry of Education and Research (eMed, Cliomics 01ZX1309B) and the Multiple Myeloma Research Foundation.

Information on authorship, contributions, and financial & other disclosures was provided by the authors and is available with the online version of this article at www.haematologica.org.

References

- Rajkumar SV, Dimopoulos MA, Palumbo A, et al. International Myeloma Working Group updated criteria for the diagnosis of multiple myeloma. *Lancet Oncol.* 2014;15(12):e538-e548.
- Palumbo A, Avet-Loiseau H, Oliva S, et al. Revised international staging system for multiple myeloma: A report from international myeloma working group. *J Clin Oncol.* 2015;33(26):2863-2869.
- Manier S, Salem KZ, Park J, Landau DA, Getz G, Ghobrial IM. Genomic complexity of multiple myeloma and its clinical implications. *Nat Rev Clin Oncol.* 2016;14(2):100-113.
- Rajan AM, Rajkumar SV. Interpretation of cytogenetic results in multiple myeloma for clinical practice. *Blood Cancer J.* 2015;5(10):e365-e365.
- Kumar S, Zhang L, Dispenzieri A, et al. Relationship between elevated immunoglobulin free light chain and the presence of IgH translocations in multiple myeloma. *Leukemia.* 2010;24(8):1498-1505.
- Ludwig H, Milosavljevic D, Berlanga O, et al. Suppression of the noninvolved pair of the myeloma isotype correlates with poor survival in newly diagnosed and relapsed/refractory patients with myeloma. *Am J Hematol.* 2016;91(3):295-301.
- Bradwell A, Harding S, Fourrier N, et al. Prognostic utility of intact immunoglobulin Ig κ /Ig λ ratios in multiple myeloma patients. *Leukemia.* 2013;27(1):202-207.
- Michallet M, Chapuis-Cellier C, Dejoie T, et al. Heavy-light chain monitoring correlates with clinical outcome in multiple myeloma patients. *Leukemia.* 2017 June 30 [Epub ahead of print].
- Merz M, Salwender H, Haenel M, et al. Subcutaneous versus intravenous bortezomib in two different induction therapies for newly diagnosed multiple myeloma: An interim analysis from the prospective GMMG-MM5 trial. *Haematologica.* 2015;100(7):964-969.
- Mai EK, Bertsch U, Dürig J, et al. Phase III trial of bortezomib, cyclophosphamide and dexamethasone (VCD) versus bortezomib, doxorubicin and dexamethasone (PAD) in newly diagnosed myeloma. *Leukemia.* 2015;29(8):1721-1729.
- Neben K, Lokhorst HM, Jauch A, et al. Administration of bortezomib before and after autologous stem cell transplantation improves outcome in multiple myeloma patients with deletion 17p. *Blood.* 2012;119(4):940-948.
- Goldschmidt H, Lokhorst HM, Mai EK, et al. Bortezomib before and after high-dose therapy in myeloma: long-term results from the phase III HOVON-65/GMMG-HD4 trial. *Leukemia.* 2017 July 4 [Epub ahead of print].
- Bochtler T, Hegenbart U, Heiss C, et al. Evaluation of the serum-free light chain test in untreated patients with AL amyloidosis. *Haematologica.* 2008;93(3):459-462.
- González-Calle V, Cerdá S, Labrador J, et al. Recovery of polyclonal immunoglobulins one year after autologous stem cell transplantation as a long-term predictor marker of progression and survival in multiple myeloma. *Haematologica.* 2017;102(5):922-931.
- Hose D, Rème T, Hielscher T, et al. Proliferation is a central independent prognostic factor and target for personalized and risk-adapted treatment in multiple myeloma. *Haematologica.* 2011;96(1):87-95.