



HHS Public Access

Author manuscript

Obesity (Silver Spring). Author manuscript; available in PMC 2019 April 01.

Published in final edited form as:

Obesity (Silver Spring). 2018 April ; 26(Suppl 2): S35–S44. doi:10.1002/oby.22159.

Accumulating Data to Optimally Predict obesity Treatment (ADOPT) Core Measures: Environmental Domain

Brian E. Saelens, PhD,

University of Washington and Seattle Children's Research Institute, Seattle, WA 98121

S. Sonia Arteaga, PhD,

Division of Cardiovascular Sciences, National Heart, Lung, and Blood Institute, Bethesda MD 20892

David Berrigan, PhD,

Division of Cancer Control and Population Sciences, National Cancer Institute, Bethesda, MD 20892

Rachel M Ballard, MD MPH,

Office of Disease Prevention, Office of the Director, National Institutes of Health, Bethesda, MD 20892

Amy Gorin, PhD,

Department of Psychological Sciences, Institute for Collaboration on Health, Intervention, and Policy, University of Connecticut, Storrs, CT 06269

Tiffany Powell-Wiley, MD, MPH,

Division of Intramural Research of the National Heart, Lung, and Blood Institute of the National Institutes of Health, Bethesda, MD, 20893

Charlotte Pratt, PhD RD,

Division of Cardiovascular Sciences, National Heart, Lung, and Blood Institute, Bethesda MD 20892

Jill Reedy, PhD, and

Division of Cancer Control and Population Sciences, National Cancer Institute, Bethesda, MD 20892

Shannon N. Zenk, PhD

College of Nursing, University of Illinois at Chicago

Abstract

CORRESPONDING AUTHOR: Brian E. Saelens, Seattle Children's Research Institute, P.O. Box 5371, M/S: CW8-6, Seattle, WA 98145-5005; Phone: (206) 884-7800; Fax: (206) 884-7801; bsaelens@u.washington.edu.

DISCLOSURE: All of the authors declared no conflicts of interest that are directly relevant to this work. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health (NIH) or the U.S. Government.

CONTRIBUTIONS: All authors were involved in writing and revising of this manuscript and had final approval of the submitted version.

Background—There is growing interest in how environment is related to adults' weight and activity and eating behaviors. However, little is known about whether environmental factors are related to the individual variability seen in adults' intentional weight loss or maintenance outcomes.

Objectives—The Environment Domain Subgroup of The Accumulating Data to Optimally Predict obesity Treatment (ADOPT) Core Measures Project sought to identify a parsimonious set of objective and perceived neighborhood and social environment constructs and corresponding measures to include in the assessment of response to adult weight loss treatment.

Significance—Starting with home address, the Environment Domain subgroup recommended for inclusion in future weight loss or maintenance studies, constructs and measures related to walkability, perceived land use mix, food outlet accessibility (perceived and objective), perceived food availability, socioeconomics, and crime-related safety (perceived and objective) to characterize the home neighborhood environment. The subgroup also recommended constructs and measures related to social norms (perceived and objective) and perceived support to characterize an individual's social environment. The 12 neighborhood and social environment constructs and corresponding measures provide a succinct and comprehensive set to allow for more systematic examination of the impact of environment on adults' weight loss and maintenance.

Keywords

environment; weight loss; individual variability; neighborhood; social support

Introduction

Physical and social environments likely influence adult's activity and eating behaviors. Environmental factors may also influence behavior change related to weight control, but environmental factors as predictors of adult weight loss success are rarely examined. Perhaps because of limited evidence about environmental factor predictors, published reviews examining factors to explain variability in adult obesity intervention response and treatment adherence focus almost exclusively on behavioral, biological, and psychosocial factors [1–4]. Conceptual models that underpin interventions for adult weight loss also rarely include environmental exposure, despite a growing literature based on cross-sectional and longitudinal observational studies documenting associations between environmental factors and physical activity, diet and weight [5–10].

Environmental exposures are potentially critical components in whether adults initiate and sustain healthy weight-related behaviors for weight loss and maintenance. The importance of the environment is highlighted in many conceptual models of health behaviors [11]. Environmental factors could help explain differential variability in response to adult weight management interventions across sites in multi-center trials, particularly if the physical and social environments vary considerably across study sites. Ongoing exposure to environmental factors in everyday life and the potential for these factors to explain disparate outcomes in meta-analyses of weight loss trials across diverse populations are important reasons to consider them as potential weight loss predictors. Finally, the factors identified to

date as predictors explain only a small amount of the variability in adults' response to weight loss interventions. Perhaps exploring environmental factors or their interaction with biological, psychological, or behavioral factors could improve predictive ability [12, 13].

The Accumulating Data to Optimally Predict obesity Treatment (ADOPT) Core Measures Project aims to better understand the factors influencing individual variability in response to adult obesity treatment by providing a framework for how obesity researchers can generate the evidence base needed to guide the development of tailored treatments of obesity [14]. The identification of a core set of measures across four domains (all included in this issue of *Obesity*)—behavioral, biological, environmental, and psychosocial—to be used in a consistent manner across adult weight loss trials represents a first step in an ongoing process which will be refined and updated dynamically as the science advances and new evidence accumulates.

The goal of the ADOPT Environment domain subgroup was to identify the environmental factors that may predict adult's response to weight loss and weight maintenance. The subgroup defined environmental influences as external factors such as built, natural, economic, or social environments to which individuals are exposed, in contrast to other ADOPT domains focused on cognitions, attitudes, affect, or physiology. The subgroup recommendations focus on environmental variables that could impact weight-related behaviors on a regular basis through on-going exposures. It is recognized that relative to other ADOPT domains there is little existing evidence concerning associations between environmental factors and adult weight loss or weight loss maintenance, but implementing the proposed recommendations would be a first step building systematic evidence.

Identifying high priority constructs and measures for environment

Much of the evidence about environmental factors and weight-related behaviors is observational and cross-sectional, examining differences in weight status and related behaviors at one point in time among individuals living in or otherwise exposed to different physical or social environments. Some more recent studies are longitudinal, examining long-term effects of environmental factors on weight and related behaviors, or alternatively among individuals who move from one environment to another or among those otherwise exposed to environmental changes (e.g., pedestrian infrastructure improvements, opening new food stores) [8, 15, 16]. The ADOPT Environment domain subgroup was only aware of a limited number of published reports that examine the association of environmental factors with weight loss or maintenance among adult weight management trial participants (e.g., [17, 18]).

Narrowing the environmental focus

Given existing evidence and the measures being proposed by other ADOPT subgroups, the Environment Domain subgroup started with a focus on the physical and social aspects of the environments of the home neighborhood that might influence weight loss and maintenance. The physical environment includes the built and food environment, whereas the social environment includes both socioeconomic factors and social factors such as neighborhood crime and safety [19, 20]. The subgroup also decided to include other aspects of the social

environment such as an individual's level of social support for weight loss. The focus on home neighborhood stems from it being the most studied setting in relation to adult weight status and activity and diet behaviors among the various environments an individual encounters routinely [21]. This decision does not diminish the potential importance of other environments encountered on a regular basis (e.g., work environment, chemical/toxin environment) or at higher (e.g., regional environments) or lower levels (e.g., within home activity and food environment) within the socioecological framework. Such other environments are less universal and/or adequate data resources or measurement tools specific to these other environments are lacking. Finally, the focus on the home neighborhood physical and social environments, as well as individual's own social environment as the external contexts in which individuals are routinely embedded, recognizes that these factors are the more proximal influences than more upstream policy or systems influences or broader social contexts.

Criteria for environmental construct and measures selection

Consistent overall ADOPT objectives, the ADOPT Environment domain subgroup identified high priority potential explanatory environmental constructs or factors and the corresponding measures to assess these environmental constructs using the following criteria:

- a.** Environmental factors most directly related to adults' engagement in physical activity (particularly walking behavior given its high prevalence), procurement or consumption of food, or to adherence to other weight loss behaviors;
- b.** Published evidence about the quality of measures (e.g., reliability, validity);
- c.** Feasibility of obtaining the measures within adult weight management trials, including considerations of both participant burden and investigator expertise and burden;
- d.** Attempts to reduce overlap among constructs. For example, residential density and overall walkability are often highly correlated, so only one was recommended.
- e.** Being in an intermediate level within socioecological models - beyond the individual level, but below governmental or organizational policy levels.

Unlike many individual-level constructs (e.g., self-efficacy) which are mostly self-reported or biological measures obtained from biological samples, issues of feasibility for obtaining some objective environmental measures include the availability and cost of local, state, and national geographic information systems (GIS) data, the technical and analytic skills required for creating environmental measures and attributing them to individuals. For example, commonly used objective measures of the home neighborhood food environment require data on the type and location of food stores. Fortunately, many municipalities are improving availability of environmental information by developing publicly available GIS data systems that contain information about activity and food amenities (e.g., parks, food stores) that can be used to characterize local activity and food environments (e.g., <https://octo.dc.gov/service/dc-gis-services>). This includes law enforcement posting publicly

available crime data that are spatially categorized (e.g., linked to census geography or even more precisely to street intersections) and can be used to develop measures of crime-related safety. Other more detailed objective environmental measures (e.g., amenities and quality of parks and recreational facilities, food availability, food prices) that might better discern and characterize more aspects of individuals' environment currently require more intensive measurement efforts (e.g., on-the-ground observations). Given the geographic dispersion of participants in most intervention trials and expense involved in collecting even brief versions of these more detailed measures, these were not deemed currently feasible to recommend.

Considerations Related to the Data Collection in the Environmental Domain

Subjective and Objective Measures

The ADOPT Environment domain subgroup considered it important to include constructs that are directly observed or derived from existing data (e.g., administrative data, road network data), often referred to as “objective,” and constructs based on participant report or perception of environment, often referred to as “subjective” or “perceived.” Objective environment measures are derived from existing or researcher collected (e.g., from on-the-ground or remote audits) geospatial or administrative data. For instance, measures of street connectivity can be derived from data about the connectedness of streets around a location (e.g., an individual's home) using Census Tiger Line Files for roads and intersections. Alternately, perceived or subjective street connectivity can be obtained from querying individuals about the connectedness or pattern of streets in their home neighborhood. Objective and perceived environmental constructs are often intended to capture different constructs or different aspects of a single construct. Subjective measures may also explicitly assess attitudes about environments, e.g., “How concerned are you about the safety in your neighborhood?” versus “Is it safe to walk in your neighborhood?” The validity assessment of subjective and objective environment measures also differs. Obtaining subjective environment measures may capture constructs for which objective data are challenging to obtain such as perceptions of aesthetics. Subjective and objective measures of seemingly similar environmental constructs can have independent associations with physical activity and diet [22–24]. Therefore, resources permitting and as others have recommended [19], the ADOPT Environment domain subgroup advocates collecting a combination of objective and subjective measures of the environment.

The Importance and Use of Home Address

Obtaining objective home neighborhood environmental measures involves multiple steps starting with home address information. Home address, and other key locations such as workplace, can be conceived as the environmental equivalent of a biological specimen – the home address can be stored for later analysis, but if not collected, then measures of objective environment factors will not be available. Indeed, *study teams are strongly encouraged to routinely collect, preserve, and obtain permission to use study participants' home address information, even if the study lacks immediate plans to obtain and analyze objective home neighborhood environmental data.* Ideally, home address is first collected at study enrollment, with prior address information collected if participants have recently moved. Retrospective address collection to determine residential histories is possible, but expensive

and prone to error [25]. Collection of street addresses, rather than Post Office Boxes or ZIP codes alone, is critically important, as is frequent updating of home addresses. Failure to collect new address data to account for moving and to preserve old address data can lead to bias in estimates of participants' environments. Other address information such work address, and even broader activity space locations (i.e., where individuals spend time), might also be beneficial to collect as the effects of environments in and around these locations on weight, weight loss, and weight maintenance are active areas of research [26, 27].

Home address can be used in various ways to develop home neighborhood environment measures. An address can be entered directly into commercially available or free web tools that generate environmental measures specific to that address (e.g. www.walkscore.com). The US Census provides a free high-quality tool for geocoding, with single addresses or batches of up to 1000 addresses able to be entered at a time (<https://www.census.gov/geo/maps-data/data/geocoder.html>). Commercial services are also available that will geocode address files and link the geocode addresses to diverse data resources, including environmental measures from census (e.g. <http://geolytics.com/>) and many others. It is also possible to geocode addresses using commercial or open-sourced computer software. These tools are becoming more accessible and easier to use; however, often personnel with basic GIS training are needed to use such software as programming may be specific to the study question and data resource. Once data are geocoded and linked to census identifiers, measures of various environment features can be calculated such as walkability or neighborhood deprivation.

Confidentiality and privacy protection are critical issues in objective environment measure development, as it requires matching participants' addresses to existing databases containing environmental information; this process represents the sharing of a personal identifier (an address) with the online service provider that does the matching [28]. Bader et al. suggest some alternative approaches such as geographic imputation which involves entering an address from a similar area rather than a participant address. Investigators should work with their IRB to determine how best to proceed when using online or hired commercial geocoding services.

Environment Constructs and Corresponding Measures Recommended by ADOPT

Home Neighborhood Environment

The ADOPT Environment domain subgroup recommends measuring a combination of 5 perceived and objective physical environment constructs that focus on the physical activity and food environments in participants' home neighborhoods (Table 1).

Walkability - Objective—Two approaches for measuring objective walkability are recommended - an individual residence-specific measure of walkability and a neighborhood-wide measure of walkability based on census data. **Walk Score**, a commercially generated index that considers proximities to diverse destinations (e.g., retail, services), provides an estimate of walkability specific to individual addresses and can be obtained for free by

entering addresses into a website or at cost through arrangements with the company (www.walkscore.com). Several studies have demonstrated associations between Walk Score and physical activity obtained by walking, particularly walking for transportation [29–33]. If submission of addresses to a web-based data source is a concern, linkage to pre-calculated area-based measures of walkability is an alternative measures. A number of walkability indices are available for specific areas such as census tracts or blocks [34, 35]. The EPA Office of SmartGrowth recently released a nationwide **EPA Walkability index** at the Census Block group level (<https://www.epa.gov/smartgrowth/smart-location-mapping>). A recent study reported associations between accelerometer measures of physical activity and this walkability index, although it is not known if it is associated with weight trajectories [36]. Nevertheless, it would be straightforward to link census geography for home addresses to this index.

Land use mix - Perceived—Walkability is a multi-faceted construct, with a significant component within it being land use mix, defined for home neighborhood environment purposes as the extent of non-residential destinations nearby to one’s home. The ADOPT Environment domain subgroup recommends obtaining a measure of perceived land use mix. Perceived land use mix has been related to higher rates and amounts of adults’ utilitarian walking behavior [37–39]. Higher land use mix also frequently co-occurs with other aspects of higher neighborhood walkability, including greater residential density and higher street network connectivity. Thus, measuring perceived land use mix allows for a parsimonious proxy assessment of perceived neighborhood walkability. The **land use mix-access subscale of the abbreviated Neighborhood Environment Walkability Survey (NEWS-A)** [40, 41], is a 3-item scale to measure perceived land use mix.

Food Outlet Accessibility and Food Availability—For the food environment, the ADOPT Environment domain subgroup recommends objective and subjective measures of neighborhood food outlet accessibility and a subjective measure of healthy food availability. Food accessibility refers to consumer food supply locations (e.g., food stores, restaurants) and ease of getting to these locations. Food availability refers to the presence or adequacy of healthful and less healthful food products in those locations. The subgroup’s selection of these two dimensions of food access is guided by Caspi’s adaptation of Penchansky and Thomas’s model [42, 43], but also other conceptualizations [44, 45]. The subgroup recognizes that access to both healthful and less healthful food sources and products is relevant, given the evidence that the relative accessibility of healthful to less healthful food outlets and products is associated with diet quality and body weight [46, 47].

The proposed objective and subjective measures of accessibility examine three types of food outlets in the home neighborhood: supermarkets, fast food restaurants, and convenience stores. Supermarkets tend to have the largest selection of healthful foods, while both fast food restaurants and convenience stores tend to predominately sell energy-dense, nutrient-poor foods and beverages with few healthful alternatives [48, 49]. Multiple studies have shown that greater accessibility of supermarkets is associated with healthier dietary intake and lower body weight, while greater accessibility of fast food restaurants and convenience stores is associated with less healthy dietary intake and higher body weight [6, 50]. For

objective measures, for each of the three food outlets, the subgroup suggests measuring the **density of outlets within a given area around the home** (e.g., 1 mile and 3 miles) and distance to the closest outlet. For a subjective measure of food outlet accessibility, the subgroup recommends a 6-item scale on the **perceived presence of food outlets in the neighborhood** developed and tested by Liese and colleagues [51, 52]. For food availability, a 3-item scale is recommended on the **neighborhood availability of fruits and vegetables and low-fat foods** developed for the Multi-Ethnic Study of Atherosclerosis (MESA) study [53, 54], which is the most widely used subjective measure of food access to date. However, there are limitations with this screener and others have updated such screeners to include additional items and other dietary constructs (e.g., whole grains) [55].

Social Environment

In addition to the physical environment around one's home, the social environment in which one lives may impact adult's weight loss and maintenance. The ADOPT Environment domain subgroup recommends collecting measures for 3 neighborhood social environment constructs, focusing on socioeconomics and objective and perceived safety, and 4 individual social environment constructs, focusing on objective and perceived social norms and two aspects of social support.

Neighborhood Socioeconomic Environment—As with other environmental factors, current data are limited on the impact of neighborhood-level socioeconomic characteristics on adult weight loss or weight loss maintenance [17]. However, extensive observational data support inclusion of neighborhood deprivation as an ADOPT Environment domain subgroup construct. Evidence demonstrates that health outcomes differ across neighborhoods of varying socioeconomic status (SES) independent of individuals' socioeconomic level [56, 57]. Studies also show that neighborhood- and individual-level SES discordance, where those with lower individual-level SES live in areas of higher SES and vice versa, exists and has a differential association with mortality when compared to socioeconomic level concordance [58]. Neighborhood deprivation has been associated with weight change over time and weight-related health behaviors [59–61]. Additionally, exposure to increasing deprivation when moving to a lower socioeconomic-level neighborhood has been associated with subsequent weight gain. The Moving To Opportunity (MTO) study, where participants were randomized to live in areas of varying neighborhood socioeconomic status, demonstrated that moving from a high- to low-poverty area was associated with decreased likelihood of obesity among women [8].

Neighborhood deprivation can be measured based on census variables at the census tract- or block group-level. Principal-components factor analyses have been used to identify key variables for inclusion in an index, often comprised of a combination of education, employment and occupation, housing conditions, income and poverty, racial composition, and/or residential stability. Specifically, the variables that make up a neighborhood deprivation index are those that have the highest loading scores in factor analyses; this is also applicable across geographic regions if characterizing neighborhood deprivation for populations in different cities or states. These census variables are then standardized, weighted based on factor loading coefficients, and summed to determine an overall

neighborhood deprivation index. Many potential measures exist to characterize neighborhood deprivation, with the subgroup recommending the **neighborhood deprivation index** developed by Diez Roux and colleagues [59]. Research teams can consider use of other indices, such as the neighborhood socioeconomic position index developed by Krieger [62] or the neighborhood deprivation index developed by Lian and colleagues [63]. Using an additional index may depend on whether the research team has access to a previously calculated index for the geographic location of interest or the team considers specific census-level variables important to include in creating the index (i.e. neighborhood racial composition).

Neighborhood Crime-Related Safety—Another critical aspect of the neighborhood social environment is safety. Thus, the ADOPT Environment domain subgroup recommends including objective and subjective measures of home neighborhood crime-related safety for examining as predictors of adult weight loss and weight loss maintenance. **Police-reported crime** is the recommended objective measure of safety. Higher police-reported crime has been associated with lower walking for physical activity [64]. Police-reported crime can be attributed to individual's home neighborhood environment by linking geocoded address data with publicly available data on the location, date, and type of crime from local or federal government databases. Police-reported crime is then matched to geocoded home address data by calculating the number of crimes within a specified buffer or area surrounding a home address, commonly a 1-mile buffer. Creation of the buffer and calculation of a crime rate can be completed with commercially available or open-source geospatial software. Prior analyses find that increasing perceived safety is associated with increasing walking from cross-sectional data [65] and increasing perceived safety over time is associated with decreasing adiposity in a longitudinal study [66]. For subjective safety, important constructs include perceived safety in walking in one's home neighborhood, perceived neighborhood violence, and perceived safety from crime in the neighborhood. Measurement of perceived safety is often obtained through questions querying respondents about the existence of or their fear of crime in their neighborhood [22, 65]. The subgroup recommends using the two items that Evenson and colleagues used for **perceived neighborhood safety** within the Multi-Ethnic Study of Atherosclerosis [65].

Individual Social Norms and Social Support—Another aspect of the social environment to consider is the influence of friends and family members on health behaviors and weight management. Observational research suggests that weight gain, weight loss, and related behaviors can spread in social networks [9, 10, 67] and clinical trials demonstrate associations between weight loss outcomes and various types of social support and weight-related social norms [68–70]. In the adult weight management literature, much of this research has focused on the interdependence of weight among marital dyads; spouses tend to be of a similar weight status at the start of marriage and follow a similar weight gain trajectory over time [71, 72]. Growing evidence suggests that weight loss tends to have a ripple effect within dyads, with meaningful weight loss observed in the untreated spouses and partners of weight loss participants ([73, 74]. Thus, the ADOPT Environment domain subgroup recommends that the **weights and heights of weight loss treatment participants' spouses or significant others** be directly (i.e., not self-report) measured. There is also

evidence that subjective perceptions of the weight status and eating behaviors of individuals in one's broader social network can influence treatment response and/or intentions to lose weight [69, 75]. To capture these perceptions, the subgroup recommends the use of Leahey and colleagues' measure of **weight-related social norms**. This measure taps social norms for overweight (e.g., "How many of your friends and acquaintances are overweight?"), social norms for weight management (e.g., "How acceptable is it in your social circle to lose weight or control your weight?"), and social norms for weight-related behaviors (e.g., "How acceptable is it in your social circle to eat unhealthy foods / large portions?"). The measure has strong psychometrics and predicts weight loss outcomes in behavioral programs [69, 75].

In addition to social norms, direct support or lack thereof from others around healthy eating and physical activity change has long been recognized as an important determinant of health behaviors and the type of support provided by family and friends is known to influence weight loss response. The **Social Support and Eating Habits Survey and Social Support and Exercise Survey** [76] is recommended by the ADOPT environment domain subgroup because it is a popular measure used to assess several types of support from family and friends (e.g., encouragement and discouragement of healthy eating; participation in exercise; use of rewards and punishment for exercise) and has demonstrated associations with eating and exercise behaviors [77, 78]. A recent update by Kiernan et al. suggests that greater support from family and friends at entry in treatment contribute to the likelihood of weight loss success, although a paradoxical effect was found between the absence of friend support and later weight loss success in group-based lifestyle programs [68]. A more nuanced exploration of the types of support from household partners linked to weight loss outcomes has found that autonomy support (e.g., providing choice and options, listening to the other's perspective) as measured by the **Important Others Questionnaire** [79] to be a more positive and consistent predictor of 6- and 18-month weight loss outcomes than more directive forms of support measured by the Sallis survey [70]. Because these measures assess different aspects of support, the subgroup recommends including both the 23-item Sallis measure and either the 6- or the 12-item version of the Important Others Questionnaire.

Other Environmental Constructs and Measures Considered

Many other aspects of the environment could be explored as potential predictors of adult weight loss and maintenance that the ADOPT Environment domain subgroup considered. Reasons for not recommending other environment constructs and corresponding measures included: insufficient evidence for consistent associations with weight-related behaviors (even within cross-sectional studies), low measurement feasibility, and unknown psychometrics for existing measures. For example, the subgroup considered recommending objective measures of food availability and food prices [80] to more precisely measure healthy versus unhealthy food availability and prices near participants' homes. However, food store and/or restaurant audits were determined to be prohibitively expensive to recommend. When resources are available, there are valid and reliable measures available for such constructs that yield more comprehensive assessments of environments (e.g., [81–84]). Moreover, we considered food price measures derived from secondary data through

companies such as Council for Community and Economic Research (C2ER, formerly ACCRA) and Nielsen, but these are only available for select areas of the country (e.g., metropolitan areas, market areas) and have no variation within an area. These data could be a resource for those clinical trials conducted across a large number of metropolitan areas covered by these data sources. Similarly, more recent attempts to characterize environments for physical activity have included in-person or remote audits (e.g., through Google Street View) [85–88], but such approaches remain expensive and their added utility relative to the already proposed measures is equivocal.

Other noteworthy, but not yet recommended, environmental constructs considered include the within-home food and activity environments, the neighborhood recreation environment, and the worksite environment. Both objective and subjective measures of home food availability are available, but they are typically long (e.g., > ½ hour to complete) and/or have not have undergone more rigorous psychometric testing [89]. Evidence also suggests that adults' home neighborhood recreation environment (e.g., presence of parks, public recreation facilities, gyms near home) is less related to adults' physical activity than other aspects of home neighborhood (e.g., walkability) and that neighborhood recreation environment may influence children's more than adults' physical activity [90, 91]. Considerable evidence exists regarding worksite interventions for adult weight management [92, 93], although many of the interventions have taken an individual-focused rather than environmentally focused approach [94] (for exception see [95]). Less is known about whether and which aspects of one's worksite environment may impact an individual's ability to initiate or sustain adults' healthy behaviors while engaged in active weight loss or maintenance attempts. In addition, the subgroup recognizes that weather, the light environment, the chemical environment, and other environmental factors may be impactful, but require further measurement development and evidence of their potential impact on weight and weight-related behaviors. Food marketing conceptualized as environmental factors could be highly influential on food choices and consumption [80], but corresponding measures are complex. More research is needed on the development of subjective and objective measures of media exposure and exposure to other food marketing, including advertisements within food outlets. Finally, there is growing interest in better characterizing the environments in which individuals actually spend their time and are exposed (i.e., their activity space) rather than relying only on measurements of static environments (e.g., home neighborhood environment) that assume high levels of time spent in the home environments and hence high exposure. Given the realities of contemporary living, particularly for working adults, this assumption may not be valid, but more research is needed to reliably measure activity space.

Limitations and Challenges

The ADOPT Environment domain subgroup recognizes the challenges and potential limitations of the recommended constructs and measures. First, the subgroup sought to balance comprehensiveness and depth of characterizing environment with the feasibility of including environmental measures within adult weight loss trials not designed primarily to examine environmental factors or environmental changes. There exist considerably more detailed and high quality measures of activity environments (e.g., measures of pedestrian

infrastructure) and food (e.g., measures of the healthfulness of different food stores and restaurants); see the National Collaborative on Childhood Obesity Research's Measures Registry (<https://www.nccor.org/nccor-tools/measures/>). Even among the measures proposed herein, feasibility of collecting these measures may differ significantly by investigative team or geographic location. Examination of objective environmental factors has become easier with improvements to the availability of spatial data as well as the tools that help to derive objective environmental measures based on spatial data. Even more user-friendly tools and processes could increase the feasibility of collecting environmental measures and help researchers and practitioners who lack sophisticated spatial methodology expertise. Second, it is also important to note regional and international differences in the availability and detail of administrative and other spatial data. This limits the ability to create the same objective environmental measures across the U.S. or within other countries; a particular challenge for multicenter trials. Therefore, the ADOPT Environment domain subgroup currently recommends objective measures that rely on existing data with at least broader national availability, but this limits the type and perhaps quality of the environmental measures that can be recommended. Third, there are likely cultural, socioeconomic, and other biases to the subjective measures of environment, thus requiring more testing to examine their reliability and validity in diverse populations. Such testing may identify additional constructs that are critical to examine among different sociodemographic populations. Also, only some of the proposed subjective measures have been translated into languages other than English. It is noteworthy that most of the proposed environmental measures were developed for use in urban and suburban setting, so their applicability to rural settings is unknown. Finally, some environmental measures attempt to isolate individual constructs, whereas others are indices that more fully capture various aspects of environments into single overall metrics. For example, indices characterizing aspects of the environment such as overall walkability or socioeconomic deprivation may be additive or may be obtained from multivariate analysis of the selected variables [35, 61]. It is not yet clear how to apply such measures to the smaller sample sizes of many intervention studies and there is a lack of agreed upon and standardized metrics applicable across the country.

These limitations may have discouraged researchers investigating adult weight loss treatment from including environmental measures as potential predictors of success. It could also be that there is little perceived heterogeneity among home neighborhood food or activity environments among individuals in any given weight loss trial, perhaps particularly for small trials. Yet, the ADOPT group now encourages collection of a core set of measures across various domains (biological, behavioral, psychological, and environmental) within all trials to facilitate data pooling and meta-analysis, thus potentially yielding more heterogeneity in participants' environmental data [14]. As most weight loss trials are of relatively short duration, it is unlikely that the set of environmental measures will change in the course of a trial for any individual participant. This may be a larger concern for weight loss trials that last longer or that enroll more low-income participants as this population tends to move residence more frequently. It also may be of greater concern in long-term follow-up for assessment of maintenance of weight loss, although frequent home address verification could mitigate such concerns.

Looking Forward

Individual response to weight loss and weight maintenance treatments is highly variable. To further refine our understanding of environmental influences on adult weight loss and maintenance (e.g., with whom do they have influence? which factors are most influential?, etc.), the ADOPT Environment domain subgroup encourages the use of the proposed set of environmental core measures within adult weight loss trials as the first step in building the empirical evidence base for environmental factors as treatment predictors. More complete details about the proposed ADOPT environmental measures are available in the Grid-Enabled Measures (GEM) database (www.gem-measures.org/Public/wsmeasures.aspx?cat=8&aid=1&wid=35). Environmental factors may explain some of the observed adult weight loss variation, either alone or in combination with behavioral, biological, or psychosocial factors. For example, perhaps individuals with the combination of higher self-efficacy for physical activity and living in a more walkable neighborhood are best able to initiate and sustain higher levels of physical activity. There is growing evidence of environmental factors moderating physical activity interventions [96]. Consistent measures of environmental factors across a variety of intervention types and intervention settings are vital to determine how much of the variation can be explained by the environment. Such evidence could inform the development of tailored or multilevel interventions that address environmental influences on weight-related interventions. The subgroup specifically decided not to establish a prioritization within the proposed constructs and measures. This first set of recommended constructs and measures should be re-evaluated after evidence has been collected on the potential influence of environmental forces. In addition, evidence from research focused on environmental influences on weight-related behaviors should be monitored to consider adding new constructs and measures, including more detailed or comprehensive measures that become more feasible to collect. Whether direct or indirect effects, identifying environments that are supportive or challenging to initiating or sustaining weight loss and maintenance behaviors has the potential to better inform candidates for treatment success, to inform the need to modify individual-based interventions to better fit the contexts in which participants are embedded, and to inform targets for environmental change.

Acknowledgments

FUNDING: The ADOPT Core Measures Working Group was supported by the National Institutes of Health: the National Heart, Lung, and Blood Institute; the National Cancer Institute; and the Office of Disease Prevention. The Grid-Enabled Measures Database (GEM) is supported and administered by the National Cancer Institute.

References

1. Burgess E, Hassmen P, Pumpa KL. Determinants of adherence to lifestyle intervention in adults with obesity: a systematic review. *Clin Obes*. 2017 Jun; 7(3):123–35. [PubMed: 28296261]
2. Lazzarotti L, Rotella F, Pala L, Rotella CM. Assessment of psychological predictors of weight loss: How and what for? *World J Psychiatry*. 2015 Mar 22; 5(1):56–67. [PubMed: 25815255]
3. Stubbs J, Whybrow S, Teixeira P, et al. Problems in identifying predictors and correlates of weight loss and maintenance: implications for weight control therapies based on behaviour change. *Obes Rev*. 2011 Sep; 12(9):688–708. [PubMed: 21535362]

4. Teixeira PJ, Going SB, Sardinha LB, Lohman TG. A review of psychosocial pre-treatment predictors of weight control. *Obes Rev.* 2005 Feb; 6(1):43–65. [PubMed: 15655038]
5. Brownson RC, Hoehner CM, Day K, Forsyth A, Sallis JF. Measuring the built environment for physical activity: state of the science. *Am J Prev Med.* 2009 Apr; 36(4 Suppl):S99–123. e12. [PubMed: 19285216]
6. Cobb LK, Appel LJ, Franco M, Jones-Smith JC, Nur A, Anderson CA. The relationship of the local food environment with obesity: A systematic review of methods, study quality, and results. *Obesity (Silver Spring).* 2015 Jul; 23(7):1331–44. [PubMed: 26096983]
7. French SA, Story M, Jeffery RW. Environmental influences on eating and physical activity. *Annual Review of Public Health.* 2001; 22:309–35.
8. Powell-Wiley TM, Cooper-McCann R, Ayers C, et al. Change in Neighborhood Socioeconomic Status and Weight Gain: Dallas Heart Study. *Am J Prev Med.* 2015 Jul; 49(1):72–9. [PubMed: 25960394]
9. Christakis NA, Fowler JH. The spread of obesity in a large social network over 32 years. *N Engl J Med.* 2007 Jul 26; 357(4):370–9. [PubMed: 17652652]
10. Shakya HB, Christakis NA, Fowler JH. Self-comparisons as motivators for healthy behavior. *Obesity (Silver Spring).* 2015 Dec; 23(12):2477–84. [PubMed: 26465785]
11. Glanz, K.Rimer, B.K., Viswanath, K., editors. *Health Behavior: Theory, Research, and Practice.* 5. San Francisco: Jossey-Bass; 2015.
12. Paquet C, Daniel M, Knauper B, Gauvin L, Kestens Y, Dube L. Interactive effects of reward sensitivity and residential fast-food restaurant exposure on fast-food consumption. *Am J Clin Nutr.* 2010 Mar; 91(3):771–6. [PubMed: 20089726]
13. Paquet C, Dube L, Gauvin L, Kestens Y, Daniel M. Sense of mastery and metabolic risk: moderating role of the local fast-food environment. *Psychosom Med.* 2010 Apr; 72(3):324–31. [PubMed: 20100887]
14. MacLean PS, Rothman AJ, Nicastrò HL, Czajkowski SM, et al. The Accumulating Data to Optimally Predict Obesity Treatment (ADOPT) Core Measures Project: Rationale and Approach. *Obesity (Silver Spring).* in press.
15. Ludwig J, Sanbonmatsu L, Gennetian L, et al. Neighborhoods, obesity, and diabetes--a randomized social experiment. *N Engl J Med.* 2011 Oct 20; 365(16):1509–19. [PubMed: 22010917]
16. Mayne SL, Auchincloss AH, Michael YL. Impact of policy and built environment changes on obesity-related outcomes: a systematic review of naturally occurring experiments. *Obes Rev.* 2015 May; 16(5):362–75. [PubMed: 25753170]
17. Mendez DD, Gary-Webb TL, Goode R, et al. Neighborhood factors and six-month weight change among overweight individuals in a weight loss intervention. *Prev Med Rep.* 2016 Dec.4:569–73. [PubMed: 27818916]
18. Zenk SN, Tarlov E, Wing CM, et al. Long-Term Weight Loss Effects of a Behavioral Weight Management Program: Does the Community Food Environment Matter? *Int J Environ Res Public Health.* 2018 Jan 26.15(2)
19. Gomez SL, Shariff-Marco S, DeRouen M, et al. The impact of neighborhood social and built environment factors across the cancer continuum: Current research, methodological considerations, and future directions. *Cancer.* 2015 Jul 15; 121(14):2314–30. [PubMed: 25847484]
20. Yen IH, Syme SL. The social environment and health: a discussion of the epidemiologic literature. *Annu Rev Public Health.* 1999; 20:287–308. [PubMed: 10352860]
21. Ding D, Gebel K. Built environment, physical activity, and obesity: what have we learned from reviewing the literature? *Health Place.* 2012 Jan; 18(1):100–5. [PubMed: 21983062]
22. McGinn AP, Evenson KR, Herring AH, Huston SL, Rodriguez DA. Exploring associations between physical activity and perceived and objective measures of the built environment. *J Urban Health.* 2007 Mar; 84(2):162–84. [PubMed: 17273926]
23. Orstad SL, McDonough MH, Stapleton S, Altincekic C, Troped PJ. A Systematic Review of Agreement Between Perceived and Objective Neighborhood Environment Measures and Associations With Physical Activity Outcomes. *Environment and Behavior.* 2016; 0(0) 0013916516670982.

24. Moore LV, Diez Roux AV, Nettleton JA, Jacobs DR Jr. Associations of the local food environment with diet quality--a comparison of assessments based on surveys and geographic information systems: the multi-ethnic study of atherosclerosis. *Am J Epidemiol.* 2008 Apr 15; 167(8):917–24. [PubMed: 18304960]
25. Wheeler DC, Wang A. Assessment of Residential History Generation Using a Public-Record Database. *Int J Environ Res Public Health.* 2015 Sep 17; 12(9):11670–82. [PubMed: 26393626]
26. James P, Jankowska M, Marx C, et al. "Spatial Energetics": Integrating Data From GPS, Accelerometry, and GIS to Address Obesity and Inactivity. *Am J Prev Med.* 2016 Nov; 51(5):792–800. [PubMed: 27528538]
27. Zenk SN, Schulz AJ, Matthews SA, et al. Activity space environment and dietary and physical activity behaviors: a pilot study. *Health Place.* 2011 Sep; 17(5):1150–61. [PubMed: 21696995]
28. Bader MD, Mooney SJ, Rundle AG. Protecting Personally Identifiable Information When Using Online Geographic Tools for Public Health Research. *Am J Public Health.* 2016 Feb; 106(2):206–8. [PubMed: 26794375]
29. Duncan DT, Aldstadt J, Whalen J, Melly SJ, Gortmaker SL. Validation of walk score for estimating neighborhood walkability: an analysis of four US metropolitan areas. *Int J Environ Res Public Health.* 2011 Nov; 8(11):4160–79. [PubMed: 22163200]
30. Hirsch JA, Moore KA, Evenson KR, Rodriguez DA, Diez Roux AV. Walk Score(R) and Transit Score(R) and walking in the multi-ethnic study of atherosclerosis. *Am J Prev Med.* 2013 Aug; 45(2):158–66. [PubMed: 23867022]
31. Sriram U, LaCroix AZ, Barrington WE, et al. Neighborhood Walkability and Adiposity in the Women's Health Initiative Cohort. *Am J Prev Med.* 2016 Nov; 51(5):722–30. [PubMed: 27211897]
32. Cole R, Dunn P, Hunter I, Owen N, Sugiyama T. Walk Score and Australian adults' home-based walking for transport. *Health Place.* 2015 Sep; 35:60–5. [PubMed: 26209804]
33. Duncan DT, Meline J, Kestens Y, et al. Walk Score, Transportation Mode Choice, and Walking Among French Adults: A GPS, Accelerometer, and Mobility Survey Study. *Int J Environ Res Public Health.* 2016 Jun 20; 13(6)
34. Frank LD, Sallis JF, Saelens BE, et al. The development of a walkability index: application to the Neighborhood Quality of Life Study. *Br J Sports Med.* 2010 Apr 22; 44:924–33. [PubMed: 19406732]
35. Hoehner CM, Handy SL, Yan Y, Blair SN, Berrigan D. Association between neighborhood walkability, cardiorespiratory fitness and body-mass index. *Soc Sci Med.* 2011 Dec; 73(12):1707–16. [PubMed: 22030212]
36. James P, Hart JE, Hipp JA, et al. GPS-Based Exposure to Greenness and Walkability and Accelerometry-Based Physical Activity. *Cancer Epidemiol Biomarkers Prev.* 2017 Apr; 26(4): 525–32. [PubMed: 28196848]
37. Saelens BE, Handy SL. Built environment correlates of walking: a review. *Med Sci Sports Exerc.* 2008 Jul; 40(7 Suppl):S550–66. [PubMed: 18562973]
38. Sugiyama T, Neuhaus M, Cole R, Giles-Corti B, Owen N. Destination and route attributes associated with adults' walking: a review. *Med Sci Sports Exerc.* 2012 Jul; 44(7):1275–86. [PubMed: 22217568]
39. Van Dyck D, Cerin E, Conway TL, et al. Perceived neighborhood environmental attributes associated with adults' transport-related walking and cycling: Findings from the USA, Australia and Belgium. *Int J Behav Nutr Phys Act.* 2012; 9:70. [PubMed: 22691723]
40. Cerin E, Conway TL, Saelens BE, Frank LD, Sallis JF. Cross-validation of the factorial structure of the Neighborhood Environment Walkability Scale (NEWS) and its abbreviated form (NEWS-A). *Int J Behav Nutr Phys Act.* 2009; 6:32. [PubMed: 19508724]
41. Cerin E, Saelens BE, Sallis JF, Frank LD. Neighborhood Environment Walkability Scale: validity and development of a short form. *Med Sci Sports Exerc.* 2006 Sep; 38(9):1682–91. [PubMed: 16960531]
42. Caspi CE, Sorensen G, Subramanian SV, Kawachi I. The local food environment and diet: a systematic review. *Health Place.* 2012 Sep; 18(5):1172–87. [PubMed: 22717379]

43. Penchansky R, Thomas JW. The concept of access: definition and relationship to consumer satisfaction. *Med Care*. 1981 Feb; 19(2):127–40. [PubMed: 7206846]
44. Apparicio P, Abdelmajid M, Riva M, Shearmur R. Comparing alternative approaches to measuring the geographical accessibility of urban health services: Distance types and aggregation-error issues. *Int J Health Geogr*. 2008 Feb 18;7:7. [PubMed: 18282284]
45. Talen E, Anselin L. Assessing spatial equity: an evaluation of measures of accessibility to public playgrounds. *Environment and Planning A*. 1998; 30:595–613.
46. Clary CM, Ramos Y, Shareck M, Kestens Y. Should we use absolute or relative measures when assessing foodscape exposure in relation to fruit and vegetable intake? Evidence from a wide-scale Canadian study. *Prev Med*. 2015 Feb;71:83–7. [PubMed: 25481095]
47. Spence JC, Cutumisu N, Edwards J, Raine KD, Smoyer-Tomic K. Relation between local food environments and obesity among adults. *BMC Public Health*. 2009 Jun 18;9:192. [PubMed: 19538709]
48. Farley TA, Rice J, Bodor JN, Cohen DA, Bluthenthal RN, Rose D. Measuring the food environment: shelf space of fruits, vegetables, and snack foods in stores. *J Urban Health*. 2009 Sep; 86(5):672–82. [PubMed: 19603271]
49. Zenk SN, Powell LM, Rinkus L, et al. Relative and absolute availability of healthier food and beverage alternatives across communities in the United States. *Am J Public Health*. 2014 Nov; 104(11):2170–8. [PubMed: 25211721]
50. Cobb LK, Appel LJ, Franco M, Jones-Smith JC, Nur A, Anderson CA. Erratum: The relationship of the local food environment with obesity: A systematic review of methods, study quality, and results. *Obesity (Silver Spring)*. 2015 Dec; 23(12):2517–9. [PubMed: 26727119]
51. Barnes TL, Bell BA, Freedman DA, Colabianchi N, Liese AD. Do people really know what food retailers exist in their neighborhood? Examining GIS-based and perceived presence of retail food outlets in an eight-county region of South Carolina. *Spat Spatiotemporal Epidemiol*. 2015 Apr. 13:31–40. [PubMed: 26046635]
52. Ma X, Barnes TL, Freedman DA, Bell BA, Colabianchi N, Liese AD. Test-retest reliability of a questionnaire measuring perceptions of neighborhood food environment. *Health Place*. 2013 May. 21:65–9. [PubMed: 23434497]
53. Moore LV, Diez Roux AV, Franco M. Measuring availability of healthy foods: agreement between directly measured and self-reported data. *Am J Epidemiol*. 2012 May 15; 175(10):1037–44. [PubMed: 22273535]
54. Mujahid MS, Diez Roux AV, Morenoff JD, Raghunathan T. Assessing the measurement properties of neighborhood scales: from psychometrics to ecometrics. *Am J Epidemiol*. 2007 Apr 15; 165(8): 858–67. [PubMed: 17329713]
55. Dubowitz T, Ghosh-Dastidar M, Cohen DA, et al. Diet And Perceptions Change With Supermarket Introduction In A Food Desert, But Not Because Of Supermarket Use. *Health Aff (Millwood)*. 2015 Nov; 34(11):1858–68. [PubMed: 26526243]
56. Diez Roux AV, Merkin SS, Arnett D, et al. Neighborhood of residence and incidence of coronary heart disease. *N Engl J Med*. 2001 Jul 12; 345(2):99–106. [PubMed: 11450679]
57. Diez Roux AV, Mujahid MS, Hirsch JA, Moore K, Moore LV. The Impact of Neighborhoods on CV Risk. *Glob Heart*. 2016 Sep; 11(3):353–63. [PubMed: 27741982]
58. Winkleby M, Cubbin C, Ahn D. Effect of cross-level interaction between individual and neighborhood socioeconomic status on adult mortality rates. *Am J Public Health*. 2006 Dec; 96(12):2145–53. [PubMed: 17077398]
59. Boone-Heinonen J, Diez Roux AV, Kiefe CI, Lewis CE, Guilkey DK, Gordon-Larsen P. Neighborhood socioeconomic status predictors of physical activity through young to middle adulthood: the CARDIA study. *Soc Sci Med*. 2011 Mar; 72(5):641–9. [PubMed: 21316829]
60. Moore K, Diez Roux AV, Auchincloss A, et al. Home and work neighbourhood environments in relation to body mass index: the Multi-Ethnic Study of Atherosclerosis (MESA). *J Epidemiol Community Health*. 2013 Oct; 67(10):846–53. [PubMed: 23868527]
61. Powell-Wiley TM, Ayers C, Agyemang P, et al. Neighborhood-level socioeconomic deprivation predicts weight gain in a multi-ethnic population: longitudinal data from the Dallas Heart Study. *Prev Med*. 2014 Sep;66:22–7. [PubMed: 24875231]

62. Krieger N, Chen JT, Waterman PD, Soobader MJ, Subramanian SV, Carson R. Choosing area based socioeconomic measures to monitor social inequalities in low birth weight and childhood lead poisoning: The Public Health Disparities Geocoding Project (US). *J Epidemiol Community Health*. 2003 Mar; 57(3):186–99. [PubMed: 12594195]
63. Lian M, Struthers J, Liu Y. Statistical Assessment of Neighborhood Socioeconomic Deprivation Environment in Spatial Epidemiologic Studies. *Open J Stat*. 2016 Jun; 6(3):436–42. [PubMed: 27413589]
64. Kerr Z, Evenson KR, Moore K, Block R, Diez Roux AV. Changes in walking associated with perceived neighborhood safety and police-recorded crime: The multi-ethnic study of atherosclerosis. *Prev Med*. 2015 Apr; 73:88–93. [PubMed: 25625690]
65. Evenson KR, Block R, Diez Roux AV, McGinn AP, Wen F, Rodriguez DA. Associations of adult physical activity with perceived safety and police-recorded crime: the Multi-ethnic Study of Atherosclerosis. *Int J Behav Nutr Phys Act*. 2012 Dec 17; 9:146. [PubMed: 23245527]
66. Powell-Wiley TM, Moore K, Allen N, et al. Associations of Neighborhood Crime and Safety and with Changes in Body Mass Index and Waist Circumference: The Multi-Ethnic Study of Atherosclerosis. *Am J Epidemiol*. 2017 May 04.
67. Gorin AA, Wing RR, Fava JL, et al. Weight loss treatment influences untreated spouses and the home environment: evidence of a ripple effect. *Int J Obes (Lond)*. 2008 Nov; 32(11):1678–84. [PubMed: 18762804]
68. Kiernan M, Moore SD, Schoffman DE, et al. Social support for healthy behaviors: scale psychometrics and prediction of weight loss among women in a behavioral program. *Obesity (Silver Spring)*. 2012 Apr; 20(4):756–64. [PubMed: 21996661]
69. Leahey TM, Doyle CY, Xu X, Bihuniak J, Wing RR. Social networks and social norms are associated with obesity treatment outcomes. *Obesity (Silver Spring)*. 2015 Aug; 23(8):1550–4. [PubMed: 26150394]
70. Gorin AA, Powers TA, Koestner R, Wing RR, Raynor HA. Autonomy support, self-regulation, and weight loss. *Health Psychol*. 2014 Apr; 33(4):332–9. [PubMed: 23730718]
71. Jeffery RW, Rick AM. Cross-sectional and longitudinal associations between body mass index and marriage-related factors. *Obes Res*. 2002 Aug; 10(8):809–15. [PubMed: 12181390]
72. The NS, Gordon-Larsen P. Entry into romantic partnership is associated with obesity. *Obesity (Silver Spring)*. 2009 Jul; 17(7):1441–7. [PubMed: 19360012]
73. Cornelius T, Gettens K, Gorin AA. Dyadic Dynamics in a Randomized Weight Loss Intervention. *Ann Behav Med*. 2016 Aug; 50(4):506–15. [PubMed: 26865258]
74. Schierberl Scherr AE, McClure Brenchley KJ, Gorin AA. Examining a ripple effect: do spouses' behavior changes predict each other's weight loss? *J Obes*. 2013; 2013:297268. [PubMed: 24083021]
75. Leahey TM, Gokee LaRose J, Fava JL, Wing RR. Social influences are associated with BMI and weight loss intentions in young adults. *Obesity (Silver Spring)*. 2011 Jun; 19(6):1157–62. [PubMed: 21164501]
76. Sallis JF, Grossman RM, Pinski RB, Patterson TL, Nader PR. The development of scales to measure social support for diet and exercise behaviors. *Prev Med*. 1987 Nov; 16(6):825–36. [PubMed: 3432232]
77. Trost SG, Owen N, Bauman AE, Sallis JF, Brown W. Correlates of adults' participation in physical activity: review and update. *Med Sci Sports Exerc*. 2002 Dec; 34(12):1996–2001. [PubMed: 12471307]
78. Hagler AS, Norman GJ, Zabinski MF, Sallis JF, Calfas KJ, Patrick K. Psychosocial correlates of dietary intake among overweight and obese men. *Am J Health Behav*. 2007 Jan-Feb; 31(1):3–12. [PubMed: 17181457]
79. Williams GC, Lynch MF, McGregor HA, Ryan RM, Sharp D, Deci EL. Validation of the "Important Other" Climate Questionnaire: Assessing autonomy support for health-related change. *Families, Systems, & Health*. 2006; 24:179–94.
80. Powell LM, Chriqui JF, Khan T, Wada R, Chaloupka FJ. Assessing the potential effectiveness of food and beverage taxes and subsidies for improving public health: a systematic review of prices, demand and body weight outcomes. *Obes Rev*. 2013 Feb; 14(2):110–28. [PubMed: 23174017]

81. Glanz K, Sallis JF, Saelens BE, Frank LD. Nutrition Environment Measures Survey in stores (NEMS-S): development and evaluation. *Am J Prev Med.* 2007 Apr; 32(4):282–9. [PubMed: 17383559]
82. Rimkus L, Ohri-Vachaspati P, Powell LM, et al. Development and Reliability Testing of a Fast-Food Restaurant Observation Form. *Am J Health Promot.* 2015 Sep-Oct;30(1):9–18. [PubMed: 24819996]
83. Rimkus L, Powell LM, Zenk SN, et al. Development and reliability testing of a food store observation form. *J Nutr Educ Behav.* 2013 Nov-Dec;45(6):540–8. [PubMed: 23726894]
84. Saelens BE, Glanz K, Sallis JF, Frank LD. Nutrition Environment Measures Study in restaurants (NEMS-R): development and evaluation. *Am J Prev Med.* 2007 Apr; 32(4):273–81. [PubMed: 17383558]
85. Adu-Brimpong J, Coffey N, Ayers C, et al. Optimizing Scoring and Sampling Methods for Assessing Built Neighborhood Environment Quality in Residential Areas. *Int J Environ Res Public Health.* 2017 Mar 08.14(3)
86. Bader MD, Mooney SJ, Lee YJ, et al. Development and deployment of the Computer Assisted Neighborhood Visual Assessment System (CANVAS) to measure health-related neighborhood conditions. *Health Place.* 2015 Jan.31:163–72. [PubMed: 25545769]
87. Wilson JS, Kelly CM, Schootman M, et al. Assessing the built environment using omnidirectional imagery. *Am J Prev Med.* 2012 Feb; 42(2):193–9. [PubMed: 22261217]
88. Rundle AG, Bader MD, Richards CA, Neckerman KM, Teitler JO. Using Google Street View to audit neighborhood environments. *Am J Prev Med.* 2011 Jan; 40(1):94–100. [PubMed: 21146773]
89. Gebremariam MK, Vaque-Crusellas C, Andersen LF, et al. Measurement of availability and accessibility of food among youth: a systematic review of methodological studies. *Int J Behav Nutr Phys Act.* 2017 Feb 14.14(1):22. [PubMed: 28196495]
90. Ding D, Sallis JF, Kerr J, Lee S, Rosenberg DE. Neighborhood environment and physical activity among youth a review. *Am J Prev Med.* 2011 Oct; 41(4):442–55. [PubMed: 21961474]
91. Bancroft C, Joshi S, Rundle A, et al. Association of proximity and density of parks and objectively measured physical activity in the United States: A systematic review. *Soc Sci Med.* 2015 Aug. 138:22–30. [PubMed: 26043433]
92. Lemon SC, Pratt CA. Worksite environmental interventions for obesity control: an overview. *J Occup Environ Med.* 2010 Jan; 52(Suppl 1):S1–3. [PubMed: 20061881]
93. Weerasekara YK, Roberts SB, Kahn MA, LaVertu AE, Hoffman B, Das SK. Effectiveness of Workplace Weight Management Interventions: a Systematic Review. *Curr Obes Rep.* 2016 Jun; 5(2):298–306. [PubMed: 27023071]
94. Thorndike AN, Healey E, Sonnenberg L, Regan S. Participation and cardiovascular risk reduction in a voluntary worksite nutrition and physical activity program. *Prev Med.* 2011 Feb; 52(2):164–6. [PubMed: 21130804]
95. Fernandez ID, Becerra A, Chin NP. Worksite Environmental Interventions for Obesity Prevention and Control: Evidence from Group Randomized Trials. *Curr Obes Rep.* 2014 Jun; 3(2):223–34. [PubMed: 26626604]
96. Schoeny ME, Fogg L, Buchholz SW, Miller A, Wilbur J. Barriers to physical activity as moderators of intervention effects. *Prev Med Rep.* 2017 Mar.5:57–64. [PubMed: 27896045]

What is already known about this subject?

- Evidence for associations between adult energy balance behaviors, neighborhood and social environments suggests environmental factors may be predictors of adult weight-related behavior change.
- Evidence is scant regarding whether or which environmental factors help explain individual variability response to obesity treatment among adults.
- Having a consistent and feasible set of measures to capture aspects of neighborhood and social environments among adults attempting weight loss could start to build on our limited understanding of the influence of environment on adult weight loss.

What does our study add?

- The ADOPT Core Measures Project Environment domain subgroup proposes a succinct set of environmental constructs and corresponding objective and subjective measures (starting with home address collection) to initiate exploration of broadly defined environmental influences on response to weight loss interventions
- ADOPT encourages the consistent use of these measures in adult weight loss trials to aid in the understanding of contextual and environmental factors that promote or hinder weight loss success, and/or have the potential to inform future tailoring of weight loss treatment.
- The proposed environment measures, combined with other ADOPT domain measures, aim to significantly enhance the value of studies by facilitating future meta-analyses to advance our understanding of the individual variability in obesity treatment responses.

Recommended Constructs and Measures from the ADOPT Environment Domain Subgroup of the ADOPT Working Group

Table 1

Construct	Measure	Measure type	Number of Items	Logistics		Source of Evidence ¹
				Administration time (minutes)	Resource Needs	
Home Neighborhood						
Home address	Home address	Self-administered	1	<5	None	
Walkability - Objective	<i>Walk score or EPA walkability</i>	Derived	N/A	N/A		C
Land Use Mix - Perceived	Neighborhood Environment Walkability Scale - Land Use Mix Access Subscale	Self-administered	3	<5	None	C
Food Outlet Accessibility - Objective	<i>Density of (1) Supermarkets, (2) Fast food restaurants, and (3) Convenience stores</i>	Derived	N/A		Basic GIS skills	L, C
Food Outlet Accessibility - Perceived	Perceived presence of (1) Supermarkets, (2) Fast food restaurants, and (3) Convenience stores, plus Others (Liese)	Self-administered	6	<5	None	L, C
Food Availability - Perceived	MESA Neighborhood Healthy Food Availability	Self-administered	8	<5	None	L, C
Socioeconomic Deprivation - Objective	<i>Neighborhood Deprivation Index (Diez Roux et al)</i>	Derived	N/A	N/A	None	L, C
Personal Safety - Objective	<i>Neighborhood Police-Reported Crime</i>	Derived	N/A	N/A	Basic GIS skills	C
Personal Safety - Perceived	MESA Perceived Neighborhood Safety	Self-administered	2	<5	None	C
Individual Social environment						
BMI of Individuals in their Social Network - Objective	<i>BMI of spouse/partner</i>	Interviewer	N/A		None	
Weight status of Individuals in their Social Network - Perceived	Weight-related social norms scale	Self-administered	29 or sub	15–20	None	
Support from Social Network	Modified Sallis Social Support scales	Self-administered	23	10–15	None	
Perceived Autonomy Support	Important Others' Questionnaire	Self-administered	6 or 12	5–10		

Italics= indicates an objective measure

¹L=longitudinal, C=cross-sectional

Evidence included both weight-related outcomes (e.g., weight loss) and behaviors related to weight loss (e.g., physical activity).