



Published in final edited form as:

Semin Musculoskelet Radiol. 2017 November ; 21(5): 648. doi:10.1055/s-0037-1606344.

Erratum: Advanced MRI Techniques for Muscle Imaging

Vivek Kalia, MD, MPH, MS¹, Doris G. Leung, MD, PhD², Darryl B. Sneag, MD¹, Filippo Del Grande, MD, MBA³, and John A. Carrino, MD, MPH¹

¹Department of Radiology and Imaging, Hospital for Special Surgery, New York, New York ²The Center for Genetic Muscle Disorders, Kennedy Krieger Institute, Baltimore, Maryland ³Servizio si Radiologia del Sottoceneri, Ospedale Regionale di Lugano, Lugano, Ticino, Switzerland

It has been brought to our attention that the name Filippo Del Grande was indexed incorrectly in the above article in *Seminars in Musculoskeletal Radiology*, Volume 21, Number 4, 2017 (DOI 10.1055/s-0037-1604007.). The author's name was indexed as "Grande, Filippo Del" but the correct listing should be "Del Grande, Filippo."

In addition, the publisher regrets an error with the placement of a reference citation within the caption for ►**Fig. 10** (p. 466). The corrected reference citation is indicated below.

Fig. 10 T2-weighted MRI and diffusion tensor imaging from a healthy volunteer (left) and a patient with chronic exertional compartment syndrome (CECS) (right) demonstrate right calf muscle before (labeled Pre) and after (labeled Post) treadmill exercise. In the CECS patient, there is clear increased T2 prolongation in the medial and lateral heads of the gastrocnemius that corresponds to areas of higher mean diffusivity (MD) and lower fractional anisotropy (FA). (Reprinted with permission from Sigmund et al.⁹⁷)