

Characterizing the gut microbiome in trauma: significant changes in microbial diversity occur early after severe injury

Benjamin M Howard,¹ Lucy Z Kornblith,¹ Sabrinah A Christie,¹ Amanda S Conroy,¹ Mary F Nelson,¹ Eric M Campion,² Rachael A Callcut,¹ Carolyn S Calfee,³ Brandon J Lamere,⁴ Douglas W Fadrosch,⁴ Susan Lynch,⁴ Mitchell Jay Cohen²

¹Department of Surgery, San Francisco General Hospital, University of California San Francisco, California, USA

²Department of Surgery, Denver Health and Hospital Authority, University of Colorado, Denver, Colorado, USA

³Division of Pulmonary and Critical Care Medicine, Departments of Medicine and Anesthesia, University of California San Francisco, California, USA

⁴Division of Gastroenterology, Department of Medicine, University of California San Francisco, California, USA

Correspondence to

Dr Benjamin M Howard, Department of Surgery, Ward 3A San Francisco General Hospital, 1001 Potrero Avenue, Room 3C-38, San Francisco, CA 94110, USA; benjamin.howard@ucsf.edu

Initial findings presented at the annual meeting of the American Association for the Surgery of Trauma, Las Vegas, Nevada, September 2015.

Received 3 May 2017
Revised 15 June 2017
Accepted 26 June 2017

ABSTRACT

Background Recent studies have demonstrated the vital influence of commensal microbial communities on human health. The central role of the gut in the response to injury is well described; however, no prior studies have used culture-independent profiling techniques to characterize the gut microbiome after severe trauma. We hypothesized that in critically injured patients, the gut microbiome would undergo significant compositional changes in the first 72 hours after injury.

Methods Trauma stool samples were prospectively collected via digital rectal examination at the time of presentation (0 hour). Patients admitted to the intensive care unit (n=12) had additional stool samples collected at 24 hours and/or 72 hours. Uninjured patients served as controls (n=10). DNA was extracted from stool samples and 16S rRNA-targeted PCR amplification was performed; amplicons were sequenced and binned into operational taxonomic units (OTUs; 97% sequence similarity). Diversity was analyzed using principle coordinates analyses, and negative binomial regression was used to determine significantly enriched OTUs.

Results Critically injured patients had a median Injury Severity Score of 27 and suffered polytrauma. At baseline (0 hour), there were no detectable differences in gut microbial community diversity between injured and uninjured patients. Injured patients developed changes in gut microbiome composition within 72 hours, characterized by significant alterations in phylogenetic composition and taxon relative abundance. Members of the bacterial orders Bacteroidales, Fusobacteriales and Verrucomicrobiales were depleted during 72 hours, whereas Clostridiales and Enterococcus members enriched significantly.

Discussion In this initial study of the gut microbiome after trauma, we demonstrate that significant changes in phylogenetic composition and relative abundance occur in the first 72 hours after injury. This rapid change in intestinal microbiota represents a critical phenomenon that may influence outcomes after severe trauma. A better understanding of the nature of these postinjury changes may lead to the ability to intervene in otherwise pathological clinical trajectories.

Level of evidence III

Study type Prognostic/epidemiological

BACKGROUND

While microbial organisms have long held the interest of surgeons in general and critical care specialists in particular, a proliferation of recent research has changed our conception of the relationship between human hosts and their indwelling microbiomes.^{1,2} Improved ability to characterize microbial communities using culture-independent DNA sequencing techniques has led to an increased awareness of the vital influence of such communities on human health and disease states, both acute and chronic.^{3,4} Such investigations have focused on particular demographics and anatomic regions, with resulting findings that have led to paradigm shifts in the way long-studied diseases are understood and treated.⁵⁻¹⁰

The central role of the gut in the response to injury has been well described,¹¹ and major alterations in gut physiology and flora have been associated with critical illness.¹²⁻¹⁴ Though alterations in gut function have been associated with the sterile inflammation that characterizes the critical illness state after injury,^{15,16} no prior studies have used culture-independent profiling techniques to characterize the gut microbiome in a cohort of patient with severe multisystem trauma. Early culture-independent investigations of gut flora in intensive care unit (ICU) patients would suggest that microbial composition at the time of admission may correlate to critical outcomes¹⁷; indeed, investigators have even reported resolution of multiple organ dysfunction syndrome secondary to sepsis by treating with fecal transplant, demonstrating that clinical improvement and reduced inflammation was associated with reconstitution of the gut microbiome.¹⁸ However, to our knowledge, no studies have conducted serial evaluations at multiple time points specifically in a trauma population. As such, it remains difficult to derive conclusions as to the causal implications of such changes, and their importance in a traumatically injured patient.

Given the lack of previous data describing the gut microbiome after severe trauma, we sought to better characterize the changes that occur in critically injured patients using culture-independent DNA sequencing techniques. Our aim was to describe any baseline differences in microbial community in the severely injured, to highlight changes in composition that occur after initial resuscitation and during the early days of ICU stay, and to correlate such changes

To cite: Howard BM, Kornblith LZ, Christie SA, et al. *Trauma Surg Acute Care Open* 2017;**2**:1-6.

**Table 1** Patient demographics, injury characteristics, treatments and outcomes

Age	Sex	Race	ISS	Mechanism	Transfusion			Antibiotics			TBI	ICU	Outcome
					0 hour	24 hours*	72 hours‡	0 hour	24 hours	72 hours			
26	Female	Asian	25	Blunt	No	Yes	No	No	Yes	No	Yes	4	Alive
26	Male	Asian	27	Blunt	Yes	Yes	Yes	No	Yes	Yes	Yes	14	Alive
33	Male	Latino	17	Blunt	No	No	No	No	Yes	Yes	Yes	25	Alive
91	Male	Asian	17	Blunt	No	No	†	No	No	†	Yes	13	Alive
60	Male	White	75	Blunt	No	No	No	No	No	No	Yes	15	Dead
37	Male	Latino	30	Blunt	No	No	†	No	No	†	Yes	19	Alive
42	Male	White	43	Blunt	Yes	Yes	Yes	No	Yes	Yes	No	10	Alive
52	Male	Latino	75	Blunt	Yes	Yes	Yes	No	Yes	Yes	Yes	4	Dead
85	Male	Asian	30	Blunt	No	Yes	†	No	Yes	†	No	25	Alive
70	Male	White	41	Blunt	No	Yes	Yes	No	Yes	Yes	Yes	30	Alive
46	Male	White	14	Penetrating	No	Yes	Yes	No	Yes	No	No	8	Alive
20	Male	Latino	21	Penetrating	No	Yes	No	No	Yes	Yes	No	8	Alive

*Product transfused in the 0 to 24-hour interval.

†No sample collected.

‡Product transfused in the 25 to 72-hour interval.

ICU, days in intensive care unit; ISS, Injury Severity Score; TBI, traumatic brain injury.

to demographic characteristics and clinical interventions. Our long-term goal was to establish a better foundational understanding of microbiome dynamics after trauma, to serve as a basis for future investigation and possible therapeutic intervention.

We hypothesized that in critically injured patients, the gut microbiome would undergo significant compositional changes in the first 72 hours after injury. We further hypothesized that these changes would occur after initial evaluation and resuscitation, and thus that microbial community composition would not differ significantly between severely injured patients and uninjured control patients at time of arrival.

METHODS

Data were prospectively collected from patients with trauma who presented at a single urban level 1 trauma center from 2014 to 2015, under a protocol approved by the University of California, San Francisco Committee on Human Research (CHR). Patients' stool samples were collected via digital rectal examination at the time of presentation in the emergency department (0 hour). Patients who met criteria for highest level critical trauma activation were eligible for inclusion in the study. For injured patients who required admission to the ICU (n=12), additional stool samples were collected at 24 hours and/or 72 hours, with informed consent obtained from the patient or their medical decision maker in accordance with the CHR-approved protocol. All samples had to yield visible amounts of stool to be included; if no visible stool was present at the time of presentation (0 hour), patients were not included in the study. If patients were transferred out of the ICU to lower acuity units prior to 72 hours after admission, they were

excluded from analysis. Thus, patients who were injured but not admitted to the ICU for at least 72 hours were excluded from analysis. Patients with isolated traumatic brain injury were excluded. Patients with activated trauma found to be uninjured, with Injury Severity Score (ISS) of 0 or 1 and length of stay less than 1 day, served as controls (n=10). Comprehensive demographic, injury, clinical and outcomes data were prospectively collected on all included patients.

Initial stool samples were collected by digital rectal examination performed by the examining trauma or emergency room physician at the time of secondary evaluation in the trauma resuscitation room, as part of the standard trauma examination. Subsequent ICU stool samples were collected either by digital rectal examination, or in passed stool if patients had a bowel movement at the time of planned sample collection. Samples were collected on a sterile glove, which was then taken to the lab and placed in a freezer maintained at -80°C within 20 minutes of collection time; the collecting finger of each glove was detached, inverted and placed in a sterile collection tube.

DNA extraction, PCR and DNA sequencing were performed on the Illumina MiSeq gene sequencer (Illumina Inc., San Diego, CA). DNA was extracted from stool samples using the MO BIO PowerSoil DNA Isolation Kit (MO BIO Laboratories Inc., Carlsbad, CA) and Qiagen AllPrep DNA/RNA Mini Kit (Qiagen, Hilden, Germany) according to the manufacturer's recommendations, respectively. Each DNA sample was PCR amplified in triplicate using primer pairs that (1) targeted the V4 hypervariable region of the 16s rRNA gene, (2) contained a unique bar code sequence to enable demultiplexing of pooled samples and (3) contained an adapter sequence enabling the amplicon to bind

Table 2 Measures of alpha-diversity in stool microbiome (p Values)

	Evenness	Richness	Shannon diversity	Simpson diversity	Inverse Simpson diversity
Injured biome over time*	0.279	0.837	0.574	0.268	0.421
Uninjured vs injured biome†	0.148	0.390	0.167	0.261	0.136

*Stool microbiome of injured patients at 0 vs. 24 hours vs. 72 hours.

†Stool microbiome of uninjured vs. injured patients at 0 hour.

Table 3 Measures of beta-diversity in stool microbiome (p Values)

	Canberra	Bray-Curtis	Unweighted UniFrac	Weighted UniFrac
Injured biome over time*	0.627	0.179	0.681	0.023
Uninjured vs injured biome†	0.064	0.177	0.159	0.123

*Stool microbiome of injured patients at 0 vs. 24 hours vs. 72 hours.

†Stool microbiome of uninjured vs. injured patients at 0 hour.

Principle Coordinates Analysis – Weighted UniFrac




Figure 1 Beta-diversity changes during 72 hours. A significant difference between the microbial community composition of critically injured patients from admission (0 hour), 24 or 72 hours after admission was observed when analyzing the weighted UniFrac matrices, as depicted in three-dimensional Principle Coordinates Analysis. This implies that there is both a phylogenetic and relative abundance component to the difference between the microbial communities.

to the MiSeq flow cell. Successful amplicons (50 of 51 samples, 98%) were pooled in equal molar concentrations and sequenced on the Illumina MiSeq.

Paired sequencing reads were quality filtered and demultiplexed using the QIIME software package before being assembled and processed further. Briefly, assembled sequencing read pairs were binned into operational taxonomic units (OTUs) using a 97% similarity to the Greengenes database, and reads that either did not cluster to the Greengenes database or that were chimeric were removed from subsequent analyses. Sample read numbers were rarefied to the read number of the lowest usable sample after processing (44 372), resulting in a rarefied OTU table.

Measures of alpha-diversity, or overall community composition within a tested population (in this case each patient at the specified time point), were derived, with richness (number of different OTUs), evenness (how equal the abundances of the OTUs are) and diversity (Shannon, Simpson and Inverse Simpson) assessed for each comparison group (samples at 0 hour, 24 hours, 72 hours). Beta-diversity, which measures differences in community composition between samples, was calculated and analyzed using principle coordinates analyses, with the derivation of Curtis, Canberra, weighted UniFrac and unweighted UniFrac distance matrices. Negative binomial regression was used to determine significantly enriched OTUs. An alpha of 0.05 was considered significant. All sample processing and subsequent analysis was performed by the authors.

RESULTS

Patients who met inclusion criteria were predominantly male, with an average age of 49 years (table 1). Included patients were severely injured, with median ISS of 27 and mean base deficit

−6.1 mEq/L; all patients suffered polytrauma, or traumatic injuries to multiple systems, and most had blunt mechanisms of injury.

Patients included as controls in analysis all had ISS of 0 or 1, and were either discharged from the emergency department or from the inpatient ward within 24 hours of admission. There were no significant differences in age, gender or ethnicity between these uninjured control patients and the cohort of severely injured patients.

As shown in table 2, there were no significant differences in stool microbiome alpha-diversity between uninjured control patients and severely injured patients at time of admission.

Similarly, there were no differences in beta-diversity indices between these two groups (table 3), indicating that injured and uninjured patients share comparable microbial community composition at the time of admission.

Though overall alpha-diversity measures did not change between time points among injured patients, these patients went on to develop significant changes in gut microbiome composition within 72 hours, as measured by beta-diversity. A significant difference between the microbial community composition of critically injured patients from admission to 72 hours after admission was observed when analyzing the weighted UniFrac matrices, indicating both a phylogenetic composition and taxon relative abundance component to the difference between the microbial communities (weighted UniFrac $p=0.023$, figure 1).

To better understand such differences, changes in significantly enriched OTUs across time points were analyzed. Changes in relative order abundance are depicted in figure 2. This analysis revealed that members of the bacterial orders Bacteroidales, Fusobacteriales and Verrucomicrobiales were depleted during




Figure 2 Microbial changes over time, depicted at order taxonomic level.

72 hours, whereas Clostridiales and Enterococcus members enriched significantly (negative binomial regression $p < 0.05$). Overall, there were 124 significantly enriched OTUs at 0 hours, and a distinct 151 significantly enriched OTUs at 72 hours. These order-level changes are depicted in [figure 3](#).

DISCUSSION

Through novel methods of characterizing microbial community composition, an enhanced understanding of the relationship between commensal organisms and human health is emerging across medical specialties and scientific disciplines. This understanding has led to promising diagnostic and even therapeutic modalities; perhaps the most widely recognized application is that of fecal transplant in colitis due to *Clostridium difficile*.¹⁹ Still, our understanding of such processes remains in its incipient stages, and has yet to be thoroughly examined in trauma populations. Culture-based studies have indicated that critical illness may correlate to significant alterations in microbial populations,¹³ a view that has been supported by increasing numbers of culture-independent investigations.^{14–20} One recent study of burn injury in both animal models and humans suggests that significant restructuring of the microbiome occurs after burn, and may contribute to increased rates of clinical sepsis.²¹ A recent case series demonstrated that in severe burn victims, initially pathogenic changes in the gut microbiota were reversed

in survivors, indicating an association between regeneration of ‘healthy’ microbial communities with survival.²² Another animal investigation demonstrated that altering the gut microbiome led to significant changes in inflammatory responses to remote organ injury (specifically, ischemia reperfusion in lung), thus implicating intestinal microbiota as critical in the response to sterile inflammatory injury.²³ Fecal transplant in murine mouse model has been shown to restore mucosal integrity,²⁴ which has been posited by others as a possible strategy for preventing gut translocation and associated sepsis in surgical patients who are critically ill.²⁵ The emerging understanding of the relationship between the gut microbiome and inflammation^{26–28} represents an area of clear relevance and critical importance to trauma and critical care. To date, however, no dedicated study has prospectively investigated changes in the microbiome in a critically injured trauma population.

In this first prospective study of the gut microbiome after trauma, we demonstrate that significant changes in microbial phylogenetic composition and relative abundance occur in the first 72 hours after injury. Such changes are not apparent at the time of initial assessment, indicating that these patients are no different from controls upon presentation; their injury patterns and subsequent therapeutic interventions are thus correlated to their microbial compositional shifts, via a yet-to-be-determined causal relationship. The short time course in which such



remains an area of active investigation in our lab and others.^{29–31} Though causal relationships remain to be determined, a better understanding of the nature of these postinjury changes may lead to the ability to intervene in otherwise pathological clinical trajectories.

Acknowledgements The authors thank the nurses, physicians and house staff of San Francisco General Hospital for their assistance in making this study possible.

Contributors All authors contributed to study design, sample and data collection, data analysis, or writing and revision, or a combination thereof.

Competing interests None declared.

Ethics approval UCSF Institutional Review Board.

Provenance and peer review Not commissioned; externally peer reviewed.

Open Access This is an Open Access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

© Article author(s) (or their employer(s) unless otherwise stated in the text of the article) 2017. All rights reserved. No commercial use is permitted unless otherwise expressly granted.

REFERENCES

- Turnbaugh PJ, Ley RE, Hamady M, Fraser-Liggett CM, Knight R, Gordon JI. The human microbiome project. *Nature* 2007;449:804–10.
- Ley RE, Knight R, Gordon JI. The human microbiome: eliminating the biomedical/environmental dichotomy in microbial ecology. *Environ Microbiol* 2007;9:3–4.
- Lozupone CA, Stombaugh JI, Gordon JI, Jansson JK, Knight R. Diversity, stability and resilience of the human gut microbiota. *Nature* 2012;489:220–30.
- Gill SR, Pop M, Deboy RT, Eckburg PB, Turnbaugh PJ, Samuel BS, Gordon JI, Relman DA, Fraser-Liggett CM, Nelson KE. Metagenomic analysis of the human distal gut microbiome. *Science* 2006;312:1355–9.
- Turnbaugh PJ, Ley RE, Mahowald MA, Magrini V, Mardis ER, Gordon JI. An obesity-associated gut microbiome with increased capacity for energy harvest. *Nature* 2006;444:1027–131.
- Abreu NA, Nagalingam NA, Song Y, Roediger FC, Pletcher SD, Goldberg AN, Lynch SV. Sinus microbiome diversity depletion and *Corynebacterium tuberculoearicum* enrichment mediates rhinosinusitis. *Sci Transl Med* 2012;4:151ra124.
- van Nood E, Vrieze A, Nieuwdorp M, Fuentes S, Zoetendal EG, de Vos WM, Visser CE, Kuismanen EJ, Bartelsman JF, Tijssen JG, et al. Duodenal infusion of donor feces for recurrent *Clostridium difficile*. *N Engl J Med* 2013;368:407–15.
- Kostic AD, Xavier RJ, Gevers D. The microbiome in inflammatory bowel disease: current status and the future ahead. *Gastroenterology* 2014;146:1489–99.
- Hilty M, Burke C, Pedro H, Cardenas P, Bush A, Bossley C, Davies J, Ervine A, Poulter L, Pachter L, et al. Disordered microbial communities in asthmatic airways. *PLoS One* 2010;5:e8578.
- Forbes JD, Van Domselaar G, Bernstein CN. The gut Microbiota in Immune-Mediated inflammatory diseases. *Front Microbiol* 2016;7:1081.
- Meng M, Klingensmith NJ, Coopersmith CM. New insights into the gut as the driver of critical illness and organ failure. *Curr Opin Crit Care* 2017;23:143–8.
- Alverdy JC, Laughlin RS, Wu L. Influence of the critically ill state on host-pathogen interactions within the intestine: gut-derived sepsis redefined. *Crit Care Med* 2003;31:598–607.
- Shimizu K, Ogura H, Goto M, Asahara T, Nomoto K, Morotomi M, Yoshiya K, Matsushima A, Sumi Y, Kuwagata Y, et al. Altered gut flora and environment in patients with severe SIRS. *J Trauma* 2006;60:126–33.
- McDonald D, Ackermann G, Khailova L, Baird C, Heyland D, Kozar R, Lemieux M, Derenski K, King J, Vis-Kampen C, et al. Extreme dysbiosis of the Microbiome in critical illness. *mSphere* 2016;1:e00199-16.
- Morowitz MJ, Carlisle EM, Alverdy JC. Contributions of intestinal bacteria to nutrition and metabolism in the critically ill. *Surg Clin North Am* 2011;91:771–85.
- Hayakawa M, Asahara T, Henzan N, Murakami H, Yamamoto H, Mukai N, Minami Y, Sugano M, Kubota N, Uegaki S, et al. Dramatic changes of the gut flora immediately after severe and sudden insults. *Dig Dis Sci* 2011;56:2361–5.
- Iapichino G, Callegari ML, Marzorati S, Cigada M, Corbella D, Ferrari S, Morelli L. Impact of antibiotics on the gut microbiota of critically ill patients. *J Med Microbiol* 2008;57:1007–14.
- Wei Y, Yang J, Wang J, Yang Y, Huang J, Gong H, Cui H, Chen D. Successful treatment with fecal microbiota transplantation in patients with multiple organ dysfunction syndrome and diarrhea following severe sepsis. *Crit Care* 2016;20:332.
- Chapman BC, Moore HB, Overbey DM, Morton AP, Harnke B, Gerich ME, Vogel JD. Fecal microbiota transplant in patients with *Clostridium difficile* infection: A systematic review. *J Trauma Acute Care Surg* 2016;81:756–64.
- Alverdy JC, Chang EB. The re-emerging role of the intestinal microflora in critical illness and inflammation: why the gut hypothesis of sepsis syndrome will not go away. *J Leukoc Biol* 2008;83:461–6.
- Earley ZM, Akhtar S, Green SJ, Naqib A, Khan O, Cannon AR, Hammer AM, Morris NL, Li X, Eberhardt JM, et al. Burn Injury alters the intestinal microbiome and increases gut permeability and bacterial translocation. *PLoS One* 2015;10:e0129996.
- Shimizu K, Ogura H, Asahara T, Nomoto K, Matsushima A, Hayakawa K, Ikegawa H, Tasaki O, Kuwagata Y, Shimazu T. Gut microbiota and environment in patients with major burns – a preliminary report. *Burns* 2015;41:e28–33.
- Prakash A, Sundar SV, Zhu YG, Tran A, Lee JW, Lowell C, Hellman J. Lung Ischemia-Reperfusion is a Sterile inflammatory process Influenced by commensal Microbiota in mice. *Shock* 2015;44:272–9.
- Kueth JW, Armocida SM, Midura EF, Rice TC, Hildeman DA, Healy DP, Caldwell CC. Fecal Microbiota transplant restores mucosal integrity in a murine Model of Burn Injury. *Shock* 2016;45:647–52.
- Krezalek MA, DeFazio J, Zaborina O, Zaborin A, Alverdy JC. The shift of an intestinal "Microbiome" to a "Pathobiome" Governs the Course and Outcome of Sepsis Following Surgical Injury. *Shock* 2016;45:475–82.
- Arrieta MC, Finlay BB. The commensal microbiota drives immune homeostasis. *Front Immunol* 2012;3:33.
- Kamada N, Seo SU, Chen GY, Núñez G. Role of the gut microbiota in immunity and inflammatory disease. *Nat Rev Immunol* 2013;13:321–35.
- Yi J, Slaughter A, Kotter CV, Moore EE, Hauser CJ, Itagaki K, Wohlauser M, Frank DN, Silliman C, Banerjee A, et al. A "clean case" of systemic injury: Mesenteric lymph after hemorrhagic shock elicits a sterile inflammatory response. *Shock* 2015;44:336–40.
- Shimizu K, Ogura H, Asahara T, Nomoto K, Morotomi M, Tasaki O, Matsushima A, Kuwagata Y, Shimazu T, Sugimoto H. Probiotic/synbiotic therapy for treating critically ill patients from a gut microbiota perspective. *Dig Dis Sci* 2013;58:23–32.
- Petrof EO, Dhaliwal R, Manzanares W, Johnstone J, Cook D, Heyland DK. Probiotics in the critically ill: a systematic review of the randomized trial evidence. *Crit Care Med* 2012;40:3290–302.
- Manzanares W, Lemieux M, Langlois PL, Wischmeyer PE. Probiotic and synbiotic therapy in critical illness: a systematic review and meta-analysis. *Crit Care* 2016;19:262.