

## CORRIGENDUM

## Arl3 and RP2 regulate the trafficking of ciliary tip kinesins

Nele Schwarz<sup>1</sup>, Amelia Lane<sup>1</sup>, Katarina Jovanovic<sup>1</sup>, David A. Parfitt<sup>1</sup>, Monica Aguila<sup>1</sup>, Clare L. Thompson<sup>2</sup>, Lyndon da Cruz<sup>1,3</sup>, Peter J. Coffey<sup>1</sup>, J. Paul Chapple<sup>4</sup>, Alison J. Hardcastle<sup>1,\*</sup> and Michael E. Cheetham<sup>1,\*</sup>

<sup>1</sup>UCL Institute of Ophthalmology, London EC1V 9EL, UK, <sup>2</sup>Institute of Bioengineering, School of Engineering and Materials Science, Queen Mary University of London, London E1 4NS, UK, <sup>3</sup>Moorfields Eye Hospital, London EC1V 2PD, UK and <sup>4</sup>William Harvey Research Institute, Barts and the London School of Medicine, Queen Mary University of London, London EC1M 6BQ, UK

\*To whom correspondence should be addressed at: UCL Institute of Ophthalmology, 11-43 Bath Street, London EC1V 9EL, UK. Tel: +44 20 7608 6944; Fax: +44 20 7608 6892; Email: michael.cheetham@ucl.ac.uk (M.E.C.); Tel: +44 20 7608 6945; Fax: +44 20 7608 6892; Email: a.hardcastle@ucl.ac.uk (A.J.H.)

*Human Molecular Genetics* 2017, 26, 2480–2492.

doi: 10.1093/hmg/ddx143

The Authors would like to add NC3Rs to the funders acknowledged in their paper.

This has also been corrected online.