

# SCIENTIFIC REPORTS

**OPEN**


## **Author Correction:** Duck plague virus Glycoprotein J is functional but slightly impaired in viral replication and cell-to-cell spread

YuYou<sup>1,2,3</sup>, Tian Liu<sup>1,2,3</sup>, Mingshu Wang<sup>1,2,3</sup>, Anchun Cheng<sup>1,2,3</sup>, Renyong Jia<sup>1,2,3</sup>, Qiao Yang<sup>1,2,3</sup>, Ying Wu<sup>1,2,3</sup>, Dekang Zhu<sup>2,3</sup>, Shun Chen<sup>1,2,3</sup>, Mafeng Liu<sup>1,2,3</sup>, XinXin Zhao<sup>1,2,3</sup>, Shaqiu Zhang<sup>1,2,3</sup>, Yunya Liu<sup>1,2,3</sup>, Yanling Yu<sup>1,2,3</sup> & Ling Zhang<sup>1,2,3</sup>

Correction to: *Scientific Reports* <https://doi.org/10.1038/s41598-018-22447-x>, published online 06 March 2018

In Figure 3B, the upper and the lower immunofluorescence images are the same. The correct Figure 3 appears below as Figure 1.

<sup>1</sup>Institute of Preventive Veterinary Medicine, Sichuan Agricultural University, Wenjiang, Chengdu, Sichuan, 611130, P.R. China. <sup>2</sup>Key Laboratory of Animal Disease and Human Health of Sichuan Province, Sichuan Agricultural University, Wenjiang, Chengdu, Sichuan, 611130, P.R. China. <sup>3</sup>Avian Disease Research Center, College of Veterinary Medicine, Sichuan Agricultural University, Wenjiang, Chengdu, Sichuan, 611130, P.R. China. YuYou, Tian Liu and Mingshu Wang contributed equally to this work. Correspondence and requests for materials should be addressed to A.C. (email: [chenganchun@vip.163.com](mailto:chenganchun@vip.163.com))



**Figure 1.** Rescue mutant viruses and Identification of gJ expression. (A) Purification and enrichment of mutant viruses. Purification and enrichment of mutant viruses were obtained by the three times passage after transfection. (B) Immunofluorescence detection of gJ expression. DEF cells were infected at 1000 TCID<sub>50</sub>, and gJ expression was detected by indirect immunofluorescence at 36 hpi. Rabbit anti-gJ were used as primary antibody, and goat anti-rabbit IgG TRITC were used as secondary antibody. (C) Anti-gJ monoclonal antibody (MAb) was used to detect gJ via western immunoblot analysis. DPV CHv-BAC-ΔgJ infected DEF cells were detected as parental virus.

**Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2018