

Percutaneous Intramedullary Headless Screw Fixation and *Wide-Awake* Anesthesia to Treat Metacarpal Fractures: Early Results in 25 Patients

Andrea Poggetti¹ Anna Maria Nucci¹ Thomas Giesen² Maurizio Calcagni² Stefano Marchetti¹
Michele Lisanti¹

¹Orthopaedic and Trauma Unit, University Hospital of Pisa, Pisa, Italy

²Department of Plastic and Hand Surgery, University Hospital of Zurich, Zurich, Switzerland

Address for correspondence Andrea Poggetti, MD, Orthopaedic and Trauma Unit, University Hospital of Pisa, Via Paradisa 2, Pisa 56100, Italy (e-mail: poggetti.andrea@gmail.com).

J Hand Microsurg 2018;10:16–21

Abstract

Metacarpal fractures constitute 7.8% of the upper extremity fractures. The common treatments remain nonsurgical procedure, but high-demanding patients or unstable fractures require fixation with Kirschner wire (K-wires), plate, and screws. However, these approaches may cause scarring and adhesion with poor functional results. From 2014 to 2015, the authors used an intramedullary headless screw to treat 25 patients (24 men, 1 woman) with metacarpal bones fractures (20 V, 3 IV, 1 III, and 1 II). The fractures patterns were 23 fractures of distal third of metacarpal bone (16 oblique, 5 comminute configurations, and 2 transverse), 1 fracture of the base of the II metacarpal bone. One case presented a multiple metacarpal and phalangeal fracture associated. The authors used *wide-awake* anesthesia (bupivacaine-epinephrine 1:100,000) and intramedullary titanium headless screw fixation percutaneously inserted (CCS Medartis and HCS Synthes 3 mm of diameter). No open reduction was needed. Early active mobilization started with a buddy strapping soon after surgery (0–3 days). The authors followed all patients until satisfactory function was achieved (4–6 weeks) and recorded the time till return to work. All fracture healed with less than 5 degrees of rotational or axial deformities. All patients return to work within 2.38 weeks after surgery (0.5–6 weeks). No cases of complex regional pain syndrome (CRPS), tendon lesions, nerve injuries, infection, hardware protruding, or mobilization were reported. Intramedullary screw fixation with *wide-awake* anesthesia for transverse, oblique, and select comminuted fractures treatment metacarpal fractures represent a reliable option to early active mobilization recovery and a quick return to the work and ordinary activities.

Keywords

- metacarpal fractures
- intramedullary fixation
- headless screws
- *wide-awake* anesthesia

Introduction

The most common treatments of metacarpal fractures remain nonsurgical procedures, but high-demanding patients or unstable fractures require fixation with Kirschner wires (K-wires), plate, and screws. All these procedures may cause scarring and adhesion arising from the implant itself or from the insufficient restoration of skeletal stability.

Studies showed good clinical results with retrograde intramedullary fixation of metacarpal and proximal phalanx

fractures, using cannulated headless screws.^{1,2} This technique may be performed by percutaneous procedure or, if necessary, with minimal exposure and provides primary stability for early active mobilization, minimizing the risk of adhesions.

The aim of this study was to report the outcomes of 25 patients treated with intramedullary headless compression screw and *wide-awake* anesthesia for unstable or secondary displaced fractures of the metacarpals bones.

received
March 12, 2017
accepted after revision
November 6, 2017
published online
March 20, 2018

©2018 Society of Indian Hand & Microsurgeons

DOI <https://doi.org/10.1055/s-0037-1618911>.
ISSN 0974-3227.

Patients and Methods

From March 2014 to December 2015, 25 consecutive extra-articular unstable fractures of the metacarpal in 24 adult patients were treated and reviewed retrospectively (level of evidence IV). Fourteen fractures were initially treated conservatively, but displacement took place after first radiologic control (1 week later). The other 10 fractures were judged to be unstable from the morphology of the fracture. The patients included 24 men and 1 woman with a mean age of 32.8 years (range: 19–59). Nineteen fractures were to the dominant hand and six to the nondominant hand. There were no open or pathologic fractures. A total of 24 patients had an isolated fracture in a single ray. One patient had concomitant fractures in multiple digits (metacarpal and phalanx). There were 20 fractures at the distal third of fifth, 3 of the fourth, 1 of the second, and 1 of the third metacarpals. The mean time from injury to surgical intervention was 5.6 days (range: 1–13). Eight patients (five white-collar and three soldiers) had a health care daily benefit, whereas the others were not covered for medical expenses, as freelancer and student had a high functional demand (cyclists, musician, etc).

The procedure was performed using *wide-awake* anesthesia without tourniquet in all cases except one (polytrauma). The patients were anesthetized using an infiltration with 0.5% bupivacaine and epinephrine (1:100,000) (20 mL), injected 30 minutes before surgery around the dorsal aspect of the ray, from the metacarpal head to metacarpal




Fig. 1 The dorsal metacarpal area involved by anesthesia.

base³ (►**Fig. 1**). In all patients, the diameter of the intramedullary canal, measured on the lateral radiograph, was bigger than 30 mm. Only one patient with the fractures of second metacarpal bone was operated upon with anterograde screw insertion.

Fractures were reduced using Jahss' maneuver under fluoroscopy, and if there was a rotational deformity (using a full active finger flexion after K-wire temporary fixation to the cortical of the metacarpal base), it was corrected by handling of the finger (►**Fig. 2**).

All fingers were operated upon using an intrarticular retrograde technique, inserting the K-guidewire and screw through the dorsoradial/ulnar aspect of the head of the metacarpal, dependent on the configuration of the fractures with the metacarpophalangeal (MCP) joint in approximately 80 to 90 degrees of flexion. Once the guidewire was in the appropriate position and a satisfactory reduction was obtained, a 2-mm skin incision was performed and the screw was inserted over the guide without predrilling (►**Fig. 3**). After, a 3-mm diameter, self-tapping cannulated headless screw was used (CCS Medartis [Basel, Switzerland] in 8 cases and HCS Synthes [Oberdorf, Switzerland] in 17 patients).

In cases of comminute fracture pattern, to decrease the risk collapse of the fragments during screw insertion, the guidewire was put bedside another K-wire. The screw used was as long as possible to avoid excessive forces and secondary displacement of the fractures. The length of the screw was calculated preoperatively and verified under fluoroscopy before insertion. Care was taken to avoid the proximally protrusion of the base of screw. After the fixation, alignment and rotation were controlled asking an immediate patients' active motion. Later, the metacarpal and digit were dressed with buddy strapping to the adjacent fingers. All patients were encouraged to actively mobilize the involved finger immediately after surgery and before starting hand therapy (►**Fig. 4**). All patients were checked within 8 days (range: 7–15) of surgery. The second follow-up was planned at 1 month after surgery (range: 28–45 days) with radiographs. The patients were followed-up for an average of 16.8 weeks (range: 8–24) (►**Fig. 5**).

Patients were addressed to surgical treatment by three hand surgeons. The surgeon who performed the operative fixation and the physician who registered the postoperative data were not the same who enrolled the patients. This was to reduce the possible bias into the medical record.

Manchester-modified or M2 DASH (Disabilities of the Arm, Shoulder and Hand questionnaire) Score Test and Total Active Motion (TAM) of the fingers were measured at each follow-up, and at the final follow-up, grip strength was measured in both the hands using a Jamar dynamometer (Patterson Medical Holding, Warrenville, Illinois, United States).⁴ Hand therapist (included in the study team) performed the tests.

Results

Fractures were radiologically healed within 4.36 weeks (range: 4–6). The functional recovery was checked with M2



Fig. 2 The unstable multifragmentary fracture of the metacarpal neck.



Fig. 3 The percutaneous introduction of cannulated screw, after close reduction.

DASH Score, mean 0.6 (range: 0–5); TAM, mean 250.8 degrees (range: 220–260); and Jamar grip strength, mean 42.24 kg (range: 33–52). At the final follow-up, the average extension lag for the MCP joint was 4 degrees (range: 0–16), for the proximal interphalangeal (PIP) joint it was 7 degrees (range: 0–18), and no extension lag for the distal interphalangeal

(DIP) joint was evaluated (► **Table 1**). No inadequate fixation, intra-articular protruding screws, loss of reduction, malrotation, infection, screw migration, or complex regional pain syndrome (CRPS) was revealed. No implant was removed, and at the final follow-up, no signs of early osteoarthritis was revealed.

Although the results achieved are overall very good, the authors stress the different recovery time between the patients who benefit of a public insurance found (white-collar, soldier, etc.) with a return to work in mean 5 weeks versus the patients without health insurance assistance (freelancer, student, etc.) or with high functional demands (cyclist, musician, etc.) that have a very fast return to daily activities (mean 2.38 weeks).⁴⁻⁸

Discussion

This case series showed that intramedullary headless screws combined with *wide-awake* anesthesia is a safe, quick, and reliable alternative to fix unstable metacarpal fractures. This procedure is even more suitable in patients who require a short period to recovery before return to work or daily life (athletes, white-collar, freelancer, or high demand worker).^{9,10}

To achieve good results and reduce disadvantages, a strict adherence to the surgical technique is mandatory. In practice, it is important to understand the fracture pattern and try to fix it, achieving good primary stability and avoiding shortening, rotation, and kneeling of the metacarpal head. These critical issues can be preventing with a careful preoperative planning, choosing the correct diameter and length of the screw.

Another key point remains the insertion area on the metacarpal head, where the loading area of the articular surface should be respected.^{1,2,7,9,11-13}



Fig. 4 After osteosynthesis, the metacarpal and the adjacent digit were dressed with buddy strapping. The patients were encouraged to actively mobilize finger immediately after surgery.




Fig. 5 The TAM recovered at the end of follow-up.

The technique is even more handy using a *wide-awake* local anesthesia with bupivacaine + epinephrine (1:100,000). In fact, this procedure reduces perioperative time with a concrete cost and time saving.^{3,4}

Furthermore, the *wide-awake* procedure allows to achieve: (1) surgery without tourniquet, (2) reduction in the volume needed to obtain a painless surgery (20 mL rather 40 mL), taking advantage from one single injection dorsal block, (3) immediate malrotation check, (4) a very cost-efficient procedure, reliable in the outpatient setting rather in the main operating room, and (5) prolonged pain control effect. Indeed, bupivacaine is slowly released after injection into the surgical area. This effect is due to a liposomal activity, and it provides prolonged pain analgesia up to 72 hours, clearly longer than bupivacaine alone.^{3,11}

The two main limits of this procedure include the difficult screw management during percutaneous insertion, in particular when the screw base get in the soft tissue, and the

difficult removal that probably implies a more difficult surgery than insertion.^{1,2,9,11}

On the other hand, very good results achieved, especially in term of fast return to work and daily actions, should be a stimulus to return to daily and productive life, primarily for patients who benefit of a public insurance found. These data could be deepen and analyzed with further studies with the aim of improving the public spending review.

Note

The level of evidence is IV.

Conflict of Interest

None.

Acknowledgments

The authors would like to thank Giulia Ghelarducci, Hand Therapist of the University Hospital of Pisa and Fisiolife Upper limb Team Therapy for dedication and professionalism.

Table 1 Functional results at the final follow-up

	No.	Occupation	Age (y)	DASH score pre	DASH Score post	Return to work (w)	Union Rx (w)	Grip Injured hand (Kg)	Grip Uninjured hand (Kg)	Follow-up (w)
TAM (deg)	1	Student	19	35	0	0.5	4	56	58	24
260	2	Engineer	40	49	0	3	5	51	57	18
255	3	White-collar	27	30	0	4	4	58	60	15
245	4	Doctor	54	47	0	1	5	45	44	10
260	5	Musician	28	43	0.7	0.5	5	48	50	24
260	6	Soldier	35	35	5	6	5	53	54	24
245	7	White-collar	37	40	2	3	4	33	28	9
230	8	Cyclist	25	50	0	0.5	4	48	50	8
260	9	Cyclist	22	35	0	0.5	4	40	49	6
260	10	Student	26	43	0	2	4	28	46	12
255	11	White-collar	40	55	0	4	5	33	38	18
250	12	Lawyer	59	40	0	0.5	4	48	45	24
255	13	White-collar	45	35	0.7	4	4	33	36	21
220	14	Soldier	39	49	5	6	4	56	41	24
235	15	White-collar	31	40	1	4	4	41	38	19
250	16	Musician	27	45	0	0.5	4	33	30	17
265	17	Freelancer	22	55	0	1	5	40	41	16
240	18	Cyclist	21	60	0	0.5	6	33	42	24
240	19	Surgeon	34	30	0	0.5	4	41	38	10
245	20	Freelancer	37	50	0.7	2	4	33	45	12
260	21	Soldier	21	49	5	8	4	46	37	11
245	22	Cyclist	19	60	0	0.5	4	38	40	18
260	23	Cyclist	20	45	0	1	4	35	30	16
250	24	White-collar	56	35	0	4	4	41	48	18
260	25	Freelancer	35	40	0	2	5	45	39	24
265		TOTAL	32.8	43.8	0.6	2.38	4.36	42.24	43.36	16.8

Abbreviations: DASH, Disabilities of the Arm, Shoulder and Hand questionnaire; Rx, treatment; TAM, Total Active Motion.

References

- Giesen T, Gazzola R, Poggetti A, Giovanoli P, Calcagni M. Intramedullary headless screw fixation for fractures of the proximal and middle phalanges in the digits of the hand: a review of 31 consecutive fractures. *J Hand Surg Eur Vol* 2016;41(7):688–694
- Borbas P, Dreu M, Poggetti A, Calcagni M, Giesen T. Treatment of proximal phalangeal fractures with an antegrade intramedullary screw: a cadaver study. *J Hand Surg Eur Vol* 2016;41(7):683–687
- Thomson CJ, Lalonde DH. Randomized double-blind comparison of duration of anesthesia among three commonly used agents in digital nerve block. *Plast Reconstr Surg* 2006;118(2):429–432
- Fess EE, Moran CA. Clinical assessment recommendation. American Society of Hand Therapists; 1981
- Ozturk I, Erturer E, Sahin F, et al. Effects of fusion angle on functional results following non-operative treatment for fracture of the neck of the fifth metacarpal. *Injury* 2008;39(12):1464–1466
- Cepni SK, Aykut S, Bekmezci T, Kilic A. A minimally invasive fixation technique for selected patients with fifth metacarpal neck fracture. *Injury* 2016;47(6):1270–1275
- Ruchelsman DE, Puri S, Feinberg-Zadek N, Leibman MI, Belsky MR. Clinical outcomes of limited-open retrograde intramedullary headless screw fixation of metacarpal fractures. *J Hand Surg Am* 2014;39(12):2390–2395
- Page SM, Stern PJ. Complications and range of motion following plate fixation of metacarpal and phalangeal fractures. *J Hand Surg Am* 1998;23(5):827–832
- Malasitt P, Owen JR, Tremblay MA, Wayne JS, Isaacs JE. Fixation for metacarpal neck fracture: a biomechanical study. *Hand (NY)* 2015;10(3):438–443
- Boulton CL, Salzler M, Mudgal CS. Intramedullary cannulated headless screw fixation of a comminuted subcapital metacarpal fracture: case report. *J Hand Surg Am* 2010;35(8):1260–1263
- del Piñal F, Moraleda E, Rúas JS, de Piero GH, Cerezal L. Minimally invasive fixation of fractures of the phalanges and metacarpals with intramedullary cannulated headless compression screws. *J Hand Surg Am* 2015;40(4):692–700

- 12 Doarn MC, Nydick JA, Williams BD, Garcia MJ. Retrograde headless intramedullary screw fixation for displaced fifth metacarpal neck and shaft fractures: short term results. *Hand (NY)* 2015;10(2):314-318
- 13 ten Berg PW, Mudgal CS, Leibman MI, Belsky MR, Ruchelsman DE. Quantitative 3-dimensional CT analyses of intramedullary headless screw fixation for metacarpal neck fractures. *J Hand Surg Am* 2013;38(2):322-330