



# Indole Propionic Acid: a Small Molecule Links between Gut Microbiota and Tuberculosis

Stefan H. E. Kaufmann<sup>a</sup>

<sup>a</sup>Max Planck Institute for Infection Biology, Berlin, Germany

**KEYWORDS** gut microbiota, tuberculosis

Gut bacteria produce numerous metabolites that can circulate systemically and influence host physiology (1). Indole-3-propionic acid (IPA) is the first microbiome-derived metabolite with the potential to impact tuberculosis, now the single leading infectious cause of death in adults, surpassing HIV (2). IPA is found in human serum, has been shown to inhibit  $\beta$ -amyloid fibril formation, and can act as a neuroprotectant against a variety of oxidotoxins (3). It is in early clinical development to treat patients with Friedreich's ataxia (ClinicalTrials.gov identifier NCT01898884). In the human host, IPA is produced by *Clostridium sporogenes* and other gut symbionts from the aromatic amino acid tryptophan, revealing a clear association between circulating IPA and the presence of the producing species in mouse gut (4). Importantly, IPA was shown to impact the intestinal barrier and systemic host immunity in mice (4). The recent findings of Negatu et al., linking IPA and tuberculosis (5), thus constitute exciting new developments in the role of the microbiome in infectious diseases.

In a whole-cell screen of a fragment library designed to identify novel antituberculosis compounds, Negatu et al. discovered that IPA exerts antitubercular activity *in vitro*. Efficacy studies in *M. tuberculosis*-infected mice showed that the compound is also active *in vivo* (5). This suggests for the first time a functional link between gut microbiota and tuberculosis. Whether the effect of IPA is due to direct antibacterial activity, indirect immunomodulatory functions, or a combination of both remains to be determined.

While there is increasing interest in associations between gut microbiome species and disease state, the finding that a metabolite produced by the gut microbiota shows antituberculosis activity *in vitro* and *in vivo* is the first tangible indication for an impact of the microbiome on tuberculosis. This discovery opens up a whole series of questions. Is there a role for IPA in tuberculosis susceptibility or progression to and severity of disease? Does the metabolite support latent tuberculosis infection? Could IPA be useful as an add-on to current antituberculosis drug regimens to accelerate cure? Which bacterial and host factors are involved in the mechanism of action of IPA? Can medicinal chemists design a new antituberculous agent using IPA as a lead compound?

Negatu's results highlight the promise of IPA as a potential "low-hanging fruit" to treat tuberculosis, since it is in clinical development for another disease indication and has thus met stringent tolerability criteria in preclinical studies. Not only could IPA be investigated as an add-on to current antituberculosis drug regimens, it also constitutes—owing to its small size and attractive physicochemical properties—an ideal starting point for target-based drug optimization.

**Citation** Kaufmann SHE. 2018. Indole propionic acid: a small molecule links between gut microbiota and tuberculosis. *Antimicrob Agents Chemother* 62:e00389-18. <https://doi.org/10.1128/AAC.00389-18>.

**Copyright** © 2018 American Society for Microbiology. All Rights Reserved.

Address correspondence to kaufmann@mpiib-berlin.mpg.de.

Ed. Note: The authors of the published article did not feel that a response was necessary.

**REFERENCES**

1. Wikoff WR, Anfora AT, Liu J, Schultz PG, Lesley SA, Peters EC, Siuzdak G. 2009. Metabolomics analysis reveals large effects of gut microflora on mammalian blood metabolites. *Proc Natl Acad Sci U S A* 106: 3698–3703. <https://doi.org/10.1073/pnas.0812874106>.
2. WHO. 2017. Global tuberculosis report. World Health Organization, Geneva, Switzerland.
3. Bendheim PE, Poeggeler B, Neria E, Ziv V, Pappolla MA, Chain DG. 2002. Development of indole-3-propionic acid (OXIGON) for Alzheimer's disease. *J Mol Neurosci* 19:213–217. <https://doi.org/10.1007/s12031-002-0036-0>.
4. Dodd D, Spitzer MH, Van Treuren W, Merrill BD, Hryckowian AJ, Higginbottom SK, Le A, Cowan TM, Nolan GP, Fischbach MA, Sonnenburg JL. 2017. A gut bacterial pathway metabolizes aromatic amino acids into nine circulating metabolites. *Nature* 551:648–652. <https://doi.org/10.1038/nature24661>.
5. Negatu DA, Liu JJJ, Zimmerman M, Kaya F, Dartois V, Aldrich CC, Gengenbacher M, Dick T. 2018. Whole-cell screen of fragment library identifies gut microbiota metabolite indole propionic acid as antitubercular. *Antimicrob Agents Chemother* 62:e01571-17. <https://doi.org/10.1128/AAC.01571-17>.