



REVIEW

Beginning at the ends: telomeres and human disease [version 1; referees: 4 approved]

Sharon A. Savage

Clinical Genetics Branch, Division of Cancer Epidemiology and Genetics, National Cancer Institute, Bethesda, Maryland, USA

v1 First published: 01 May 2018, 7(F1000 Faculty Rev):524 (doi: 10.12688/f1000research.14068.1)

Latest published: 01 May 2018, 7(F1000 Faculty Rev):524 (doi: 10.12688/f1000research.14068.1)

Abstract

Studies of rare and common illnesses have led to remarkable progress in the understanding of the role of telomeres (nucleoprotein complexes at chromosome ends essential for chromosomal integrity) in human disease. Telomere biology disorders encompass a growing spectrum of conditions caused by rare pathogenic germline variants in genes encoding essential aspects of telomere function. Dyskeratosis congenita, a disorder at the severe end of this spectrum, typically presents in childhood with the classic triad of abnormal skin pigmentation, nail dystrophy, and oral leukoplakia, accompanied by a very high risk of bone marrow failure, cancer, pulmonary fibrosis, and other medical problems. In contrast, the less severe end of the telomere biology disorder spectrum consists of middle-age or older adults with just one feature typically seen in dyskeratosis congenita, such as pulmonary fibrosis or bone marrow failure. In the common disease realm, large-scale molecular epidemiology studies have discovered novel associations between illnesses, such as cancer, heart disease, and mental health, and both telomere length and common genetic variants in telomere biology genes. This review highlights recent findings of telomere biology in human disease from both the rare and common disease perspectives. Multi-disciplinary collaborations between clinicians, basic scientists, and epidemiologist are essential as we seek to incorporate new telomere biology discoveries to improve health outcomes.

Keywords

telomere, dyskeratosis congenita, telomere biology disorder, cancer, Coats plus, Hoyeraal Hreidsarsson syndrome, epidemiology

Open Peer Review

Referee Status:

	Invited Referees			
	1	2	3	4
version 1 published 01 May 2018				

F1000 Faculty Reviews are commissioned from members of the prestigious F1000 Faculty. In order to make these reviews as comprehensive and accessible as possible, peer review takes place before publication; the referees are listed below, but their reports are not formally published.

- 1 **Dirk Hockemeyer**, University of California, Berkeley, USA
- 2 **Tracy Bryan**, University of Sydney, Australia
- 3 **Steven E. Artandi**, Stanford University School of Medicine, USA
Stanford University School of Medicine, USA
Stanford University School of Medicine, USA
- 4 **Fadi Charchar**, Federation University, Australia

Discuss this article

Comments (0)

Corresponding author: Sharon A. Savage (savagesh@mail.nih.gov)

Author roles: Savage SA: Conceptualization, Data Curation, Funding Acquisition, Investigation, Project Administration, Writing – Original Draft Preparation, Writing – Review & Editing

Competing interests: No competing interests were disclosed.

How to cite this article: Savage SA. **Beginning at the ends: telomeres and human disease [version 1; referees: 4 approved]**
F1000Research 2018, 7(F1000 Faculty Rev):524 (doi: [10.12688/f1000research.14068.1](https://doi.org/10.12688/f1000research.14068.1))

Copyright: © 2018 Savage SA. This is an open access article distributed under the terms of the [Creative Commons Attribution Licence](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. The author(s) is/are employees of the US Government and therefore domestic copyright protection in USA does not apply to this work. The work may be protected under the copyright laws of other jurisdictions when used in those jurisdictions.

Grant information: This work was supported by the intramural research program of the Division of Cancer Epidemiology and Genetics, National Cancer Institute, National Institutes of Health.

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

First published: 01 May 2018, 7(F1000 Faculty Rev):524 (doi: [10.12688/f1000research.14068.1](https://doi.org/10.12688/f1000research.14068.1))

Introduction

Telomeres are the nucleoprotein complex at chromosome ends with essential roles in maintaining chromosomal integrity. They shorten with each cell division because of incomplete replication of the 3' ends of DNA and thus are markers of cellular aging. Over the last decade, there has been remarkable growth in the breadth and depth of understanding the multiple roles of telomere biology in human disease. At one end of the spectrum, very rare pathogenic germline genetic variants in telomere biology genes cause exceedingly short telomeres, resulting in dyskeratosis congenita (DC) and its related telomere biology disorders. The other end of the spectrum consists of large-scale population-based studies seeking to determine associations between telomere length human disease, environmental exposures, or common genetic variants as well as the interactions between these factors.

The complexity of these interactions requires an integrated understanding of telomere basic science, clinical medicine, and epidemiology (Figure 1). Each of these topics is worthy

of an in-depth critical review beyond the scope of this article. Instead, I will highlight some key findings and methodologic considerations and discuss where additional research is needed to aid in understanding the contribution of telomere biology to both rare and common human diseases.

Dyskeratosis congenita – a direct connection between germline telomere biology and human disease

DC was first described in a 1906 case report of males with the mucocutaneous triad of abnormal skin pigmentation, nail dystrophy, and oral leukoplakia (Figure 2)¹. Additional similar cases were reported, including the first female case in 1963²⁻⁵. Patients with DC also have very high rates of bone marrow failure; stenosis of the esophagus, urethra, or lacrimal ducts (or a combination of these); head and neck squamous cell carcinoma (HNSCC); myelodysplastic syndrome (MDS); acute myeloid leukemia (AML); pulmonary fibrosis; liver disease; avascular necrosis of the hips; and other medical problems (Table 1)⁶. DC is inherited in X-linked recessive, autosomal dominant, or autosomal recessive patterns.




Figure 1. Factors associated with human disease are integrally connected to telomere biology. This schematic illustrates the complex relationships between telomere biology, disease, aging, genetics, and environmental exposures, all of which should be considered in studies of telomeres and human disease.



Figure 2. Mucocutaneous features of dyskeratosis congenita of an adult male whose disease is due to a *DKC1* mutation. (A) Dystrophic and ridged fingernails. (B) Hyper- and hypo-pigmented skin of neck and upper chest. (C) Irregular leukoplakia of the tongue.

Table 1. Clinical features of telomere biology disorders.

Disorder	Key clinical features*
Dyskeratosis congenita (DC)	Mucocutaneous triad of nail dysplasia, abnormal skin pigmentation (hyper/hypopigmented, lacy, reticular pigmentation), and oral leukoplakia. BMF, PF, PAVM, liver disease, avascular necrosis of hips or shoulders (or both), urethral stenosis, lacrimal duct stenosis, esophageal stenosis, HNSCC, MDS, AML, and/or developmental delay. Traditional diagnosis of DC: classic triad or one of the triad, BMF, and two other findings ⁷ .
Revesz syndrome	Features of DC plus bilateral exudative retinopathy. Intracranial calcifications and developmental delay also reported.
Hoyeraal Hreidarsson syndrome	Features of DC plus cerebellar hypoplasia. Immunodeficiency has been reported as presenting problem.
Coats plus	Bilateral retinopathy, intracranial calcifications, leukodystrophy, anemia, osteopenia, and poor bone healing
DC-like	BMF, AA, MDS, or PF occurring in presence of at least one other DC-associated feature or family history suspicious of DC
Aplastic anemia	Progressive multi-lineage cytopenias, non-immune mediated. May occur in the absence of DC-associated features.
Myelodysplastic syndrome	Cytopenias with cellular dysplasia or clonal chromosomal translocations or both. May occur in the absence of DC-associated features.
Acute myeloid leukemia	May progress from MDS or aplastic anemia. May occur in the absence of DC-associated features.
Pulmonary fibrosis	May occur in the absence of DC-associated features.
Liver fibrosis	Non-alcoholic, non-infectious liver disease. May occur in the absence of DC-associated features.
Familial melanoma	Multiple family members with melanoma, often early age at onset
Familial lymphoproliferative disease	Multiple-affected family members with chronic lymphocytic leukemia, or non-Hodgkin lymphoma
Li-Fraumeni-like syndrome	Cancer family history notable for angiosarcoma and other cancers

*Key references are noted in Table 2. AA, aplastic anemia; AML, acute myeloid leukemia; BMF, bone marrow failure; DC, dyskeratosis congenita; HNSCC, head and neck squamous cell carcinoma; MDS, myelodysplastic syndrome; PAVM, pulmonary arteriovenous malformation; PF, pulmonary fibrosis.

The first DC genetic locus was mapped to Xq28 in 1996 and specifically to mutations in dyskerin (encoded by *DKC1*) in 1999⁸⁻¹⁰. The seminal work by Mitchell and Collins was the first to show a connection between telomere biology and human disease through aberrant dyskerin function and the resultant very short telomeres now well known in patients with DC¹¹. Currently, *DKC1* mutations account for about 25% of classic DC cases. A combination of candidate gene sequencing, genetic linkage studies, and whole exome sequencing occurring over the last 15 years has since identified at least 14 telomere biology genes associated with DC or DC-like phenotypes: telomerase holoenzyme complex (*DKC1*, *TERC*, *TERT*, *NOP10*, and *NHP2*), shelterin telomere protection complex (*ACD*, *TINF2*, and *POT1*), telomere capping proteins (*CTCI* and *STN1*), and other proteins that directly or indirectly interact with these key cellular processes (*RTEL1*, *NAFI*, *WRAP53*, and *PARN*) (Figure 3 and Table 2) (reviewed in 6).

The germline mutations in DC-associated telomere biology genes result in very short telomere lengths for age (Figure 4). This knowledge made it possible to develop a diagnostic test for DC, flow cytometry with fluorescent *in situ* hybridization (flow FISH), in leukocyte subsets¹². Lymphocyte telomeres less than the first percentile for age are more than 95% sensitive and

specific for differentiating patients with DC from their unaffected relatives or patients with other inherited bone marrow failure syndromes^{13,14}. In addition to aiding diagnosis, using telomeres less than the first percentile for age has greatly added in discovering the genetic causes of DC¹⁵.

Telomere biology disorders – many names connected by pathophysiology

The discovery of the multiple genetic causes and modes of inheritance has led to a growing appreciation that there is a wide range of clinical phenotypes associated with mutations in telomere biology genes. This started with the identification of *TERT* and *TERC* mutations in patients with apparently isolated aplastic anemia or pulmonary fibrosis¹⁶⁻¹⁸. As defined above, classic DC is a complex multi-system disorder, but variable penetrance and expressivity of the clinical manifestations have identified a growing number of patients with one or a few features of DC as well as germline mutations in telomere biology genes and short telomeres (Table 1 and Table 2). This spectrum of illnesses has been termed telomeropathy, short telomere syndromes, or telomere biology disorders (TBDs)^{6,19-22}. The last of these, TBD, was proposed and is favored because it is more descriptive and reflective of the underlying biology that unites these disorders^{6,22}.




Figure 3. Proteins encoding key components of telomere biology associated with disease. Protein names are noted in the figure. Associated disease and mode(s) of inheritance are shown in Table 1. The asterisk indicates proteins encoded by genes with single-nucleotide polymorphisms associated with cancer. The pound sign indicates proteins encoded by genes with single-nucleotide polymorphisms associated with telomere length.

Table 2. Germline genetics of telomere biology disorders.

Gene	Protein name(s)	Disorder(s)	Mode of inheritance	Year gene first associated with any disease, relevant reference(s)
<i>DKC1</i>	DKC1, dyskerin	DC, HH	XLR	1998 ²³
<i>TERC</i>	hTr, telomerase RNA component (encodes an RNA)	DC, AA, PF	AD	2001 ^{16,18,24}
<i>TERT</i>	TERT, telomerase	DC, AA, MDS, AML, PF, LD, FM	AD	2005 ^{16,17,25-27}
		HH	AR	
<i>NOP10</i>	NOP10, NOLA3, nucleolar protein family A, member 3	DC	AR	2007 ²⁸
<i>NHP2</i>	NHP2, NOLA2 nucleolar protein family A, member 2	DC	AR	2008 ²⁹
<i>TINF2</i>	TIN2, TERF1 (TRF1)-interacting nuclear factor 2	DC, HH, RS	AD	2008 ¹⁵
<i>WRAP53</i>	TCAB1, telomere cajal body associated protein 1	DC, HH	AR	2011 ^{30,31}
<i>CTC1</i>	CTC1, conserved telomere maintenance component 1	CP, DC	AR	2012 ³²⁻³⁴
<i>RTEL1</i>	RTEL1, regulator of telomere elongation helicase 1	PF, AA	AD	2013 ³⁵⁻⁴⁰
		DC, HH	AR	
<i>TERF2IP</i>	RAP1, TERF2-interacting protein	FM	AD	2015 ⁴¹
<i>PARN</i>	PARN, poly(A)-specific ribonuclease	PF	AD	2015 ⁴²⁻⁴⁵
		DC, HH	AR	
<i>ACD</i>	TPP1, telomere protection protein 1	AA, FM, FLPD	AD	2014 ^{41,46,47}
		HH	AR	
<i>STN1</i>	STN1, CST-complex subunit	CP	AR	2016 ⁴⁹
<i>POT1</i>	POT1, protection of telomeres 1	FM, FLPD, LFL	AD	2014 ^{47,50-52}
		CP	AR	
<i>NAF1</i>	NAF1, nuclear assembly factor 1 ribonucleoprotein	PF	AD	2016 ⁵⁴

AA, aplastic anemia; AD, autosomal dominant; AML, acute myeloid leukemia; AR, autosomal recessive; CP, Coats plus; DC, dyskeratosis congenita; FLPD, familial lymphoproliferative disease; FM, familial melanoma; HH, Hoyeraal-Hreidarsson syndrome; LD, liver disease; LFL, Li-Fraumeni-like syndrome; MDS, myelodysplastic syndrome; PF, pulmonary fibrosis; RS, Revesz syndrome; XLR, X-linked recessive.




Figure 4. Schematic representation of the connections between age, telomere length, and human disease. Clinically significant telomeres associated with telomere biology disorders are generally at or below the first percentile for age (blue shape). Many association studies of telomere length and human phenotypes, including cancer, have identified statistically significant, but perhaps not clinically significant, differences in telomere length between cases and controls (represented by yellow and red). Some studies have identified rare families with germline mutations in components of the shelterin telomere protection complex as associated with longer telomeres (green shape). AA, aplastic anemia; CLL, chronic lymphocytic leukemia; CP, Coats plus; DC, dyskeratosis congenita; FM, familial melanoma; HH, Hoyeraal Hreidarsson syndrome; LD, liver disease; LFL, Li-Fraumeni-like syndrome; MDS, myelodysplastic syndrome; PF, pulmonary fibrosis; RS, Revesz syndrome.

The most complex TBDs are those disorders presenting very early in childhood, namely Hoyeraal Hreidarsson syndrome (HH), Revesz syndrome (RS), and Coats plus^{6,55-60}. In addition to having features of DC, patients with HH have cerebellar hypoplasia and immunodeficiency, whereas those with RS also have bilateral exudative retinopathy. Coats plus, a disorder characterized by retinal and gastrointestinal vascular abnormalities, poor bone healing, leukodystrophy, and cerebellar calcifications, joined the TBD spectrum when its primary cause was identified as autosomal recessive *CTCI* mutations^{32,33,59}.

The other end of the clinical spectrum includes patients with middle or later age at onset of pulmonary fibrosis, liver disease, or bone marrow failure and heterozygous germline mutations in *NAF1*, *TERT*, *TERC*, *PARN*, or *RTEL1*^{17,18,42,61}. Additionally, it is important to note that most patients do not have all of the DC-associated medical complications. The mucocutaneous triad is diagnostic but varies with the age of onset and is usually progressive over time. Many patients, even members

of the same family, may present with just one feature but develop more over time because of variable penetrance and expressivity of germline telomere biology defects.

There is also a growing role of mutations in the shelterin complex and cancer-prone families without DC-related clinical manifestations. Heterozygous rare, pathogenic variants in *POT1* resulting in longer telomeres have been reported in familial melanoma, familial chronic lymphocytic leukemia (CLL), and a Li-Fraumeni-like syndrome family⁵⁰⁻⁵². *POT1* somatic mutations in CLL have also been associated with CLL outcomes⁶²⁻⁶⁵. Familial melanoma has also been associated with germline mutations in *ACD* (TPP1) and *TERF2IP* (RAP1)⁴¹. These studies suggest an interesting dichotomy in clinical phenotypes resulting from long versus short telomeres.

Telomeres and cancer

Telomeres are closely connected to cancer biology because of the role they play in chromosomal stability. There is a

detailed body of work in this realm and only a few of the key features will be highlighted herein as they have been reviewed in detail elsewhere^{66–73}.

The primary hypotheses connecting telomeres and cancer are based on the fact that telomeres shorten with each cell division⁷⁴. In general, cellular senescence or apoptosis is triggered when telomeres reach a critically short state. It likely takes just one critically short telomere on one chromosome arm to trigger these events as suggested in a *TERC* mouse model by Hemann *et al.*⁷⁵. A cellular survival advantage is gained through bypassing apoptosis or senescence through the upregulation of telomerase, inactivation of TP53 or RB or both, initiation of alternative lengthening of telomeres (ALT), and other key biological pathways^{76–80}. The continued division of cells originally destined for death is hypothesized to lead to continued accumulation of mutations, and sticky chromosome ends due to abnormal telomeres can contribute to chromosomal aneuploidy. Unchecked cellular growth can occur if these genetic aberrations result in a growth advantage.

Activating somatic mutations in the *TERT* promoter have been described in melanoma, bladder, thyroid, and some central nervous system cancers^{81–83}. These somatic mutations in the *TERT* promoter result in increased telomerase expression and suggest that this activation could convey a growth advantage as cancer cells continue to divide despite the presence of aberrant telomeres.

Patients with DC/TBD have significantly increased risks of MDS, AML, HNSCC, and other malignancies^{84–86}. The 2017 update of cancer in the National Cancer Institute Inherited Bone Marrow Failure Syndrome cohort reported that cancer in patients with DC occurs at an approximately four fold higher incidence and a younger age than the general population⁸⁷. MDS and AML occurred at 578- and 24-fold greater incidence, respectively, than the general population. There was also an excess of solid cancers in patients with DC with observed/expected ratios of 74 for any HNSCC and 216 for tongue HNSCC⁸⁷. The mechanisms by which cancer develops in patients with DC/TBDs is unknown and represents an important research opportunity.

Cells of patients with TBD already have a “first hit” in a key component of telomere biology. Studies of the next steps in carcinogenesis in patient-derived cells could lead to important insights into the carcinogenic process.

The advent of telomere molecular epidemiology

Telomere molecular epidemiology has emerged with the development of high-throughput telomere length measurement methods, genome-wide genotyping platforms, and keen interest in the role of telomere biology in human disease^{88,89}. These large, often population-based studies seek to determine (1) whether telomere length is associated with disease, (2) whether common genetic variants (that is, single-nucleotide polymorphisms, or SNPs) are associated with telomere length, (3) the degree to which SNPs contribute to telomere biology, and (4) interactions between telomere length, SNPs, and disease or phenotypes (Figure 1 and Table 3). Although a great deal of excitement has been generated by these studies, it is important to point out that differences in telomere length between cases and controls in large population-based studies may be statistically significant but not clinically relevant. “Short telomeres” in a large case-control or cohort study are still within the clinically “normal” range and not nearly as short as telomeres of patients with TBDs (Figure 4).

Robust and accurate telomere length measurement is at the crux of all telomere length association studies. Blood or buccal cell DNA telomere length has been evaluated in a wide array of association studies, including cancer, cardiovascular disease, mental health, inflammatory diseases, environmental exposures, and many other settings. There are numerous methods to determine telomere length in tissues, single cells, and DNA preparations, each appropriate for different applications and reviewed extensively^{90,91}. Quantitative polymerase chain reaction (qPCR) is amenable to large studies because it uses very small amounts of DNA and can be scaled up rapidly^{92,93}. However, qPCR telomere assays generate a relative telomere length and are very sensitive to DNA extraction methods and storage⁹⁴. These challenges have led to significant challenges in reproducing data in case-control or cohort studies of qPCR relative telomere length and phenotypes^{95,96}. The telomere restriction fragment

Table 3. Features of robust telomere length association studies.

- Strong *a priori* hypothesis of why telomere biology might be important in disease of interest
- Comprehensive clinical phenotyping
- Accurately measured exposure of interest
- Large sample size with power calculations reported
- Collection of samples prior to disease onset
- Detailed information on how samples were collected, processed, and stored
- Telomere length measurement methods described in detail, especially if any adaptations to published methods
- Accurate and reproducible telomere length measurement
- Strong statistical justification of association findings

method uses restriction enzymes to cut the subtelomeric ends of chromosomes in a DNA preparation and is most widely used in basic science laboratories, although a few groups use it in population-based studies^{90,91}.

Blood or buccal cell telomere length association studies

This section highlights just a few key topics within the growing literature of telomere length association studies. For example, individuals of African ancestry have longer telomeres than those of European ancestry and thus ancestry should be accounted for in analyses^{97,98}. Since self-reported ancestry can be quite variable, genomic approaches may be helpful in classifying cases and controls in order to appropriately adjust for ancestry.

There is also a growing understanding of associations between environmental exposures and telomere length. Smokers have shorter telomeres than non-smokers and thus it is important to adjust for smoking in association analyses⁹⁹. Associations between prenatal exposures to smoking and air pollution as well as exposure to certain occupational chemicals have also been explored but with varying results^{100–105}. In each of these studies, it is essential to precisely quantify the exposure of interest in addition to using a robust and reproducible telomere length measurement.

Early cancer-telomere length association studies suggested shorter telomeres as a cancer risk factor^{106,107} but studies of other cancers were not consistent^{108–112}. Meta-analyses found that most studies with blood or buccal cell DNA collected prior to cancer diagnosis were null but that case-control studies were more likely to find associations^{113,114}. Similarly, a direct comparison of prospectively and retrospectively collected DNA samples from patients with breast or colorectal cancer reported that telomere shortening occurred primarily after cancer diagnosis¹¹⁵. Many of these inconsistencies have been attributed to reverse causation bias due to the presence of cancer, underlying inflammation, or prior therapy at the time of sample collection^{113,116}. Currently, the most consistent studies are those of longer telomeres in pre-diagnostic samples of patients with lung cancer and melanoma^{117–119}. Interestingly, shorter leukocyte telomeres were associated with overall cancer mortality but not with cancer in a large prospective study of 64,637 individuals who developed 2,420 cancers¹²⁰.

There is a growing body of telomere length association studies and different aspects of mental health, including measures of perceived stress in caregivers, exposure to early life adversity, and in patients with schizophrenia, bipolar disorder, and depression^{121–126}. The biological mechanisms underlying these findings are unknown but current hypotheses include stress responses inducing oxidative stress, resulting in DNA damage and telomere shortening. Notably, abnormalities in brain development are present in patients with HH (cerebellar hypoplasia), RS (intracranial calcifications), and Coats plus (leukodystrophy and intracranial calcifications)^{6,55}. The only study to date of neuropsychiatric complications in DC found higher-than-expected occurrence of developmental delay and psychiatric disorders¹²⁷.

Studies of DC/TBD patients by psychiatrists and neurobiologists constitute an unstudied area highly likely to generate important insights into telomere biology and brain development.

There is also a great deal of interest in using telomere length as a measure of biological age and even in modulating telomere length through lifestyle interventions. Numerous studies suggest associations between lifestyle, exercise, and telomere lengths^{128–133}. However, a recent review suggests that telomere length in and of itself is not sufficient as a specific aging biomarker¹³⁴.

Current data consistently report shorter leukocyte telomeres in individuals with atherosclerotic cardiovascular disease than in unaffected controls^{135–137}. The biology underlying this association is thought to be related to chronic inflammation and oxidative stress coupled with aging of the vasculature. This prompted Aviv *et al.* to propose a model whereby age-dependent telomere shortening varies on the basis of the replicative needs of the specific tissue^{138,139}. They hypothesize that skeletal muscle (a minimally replicative cell type) may represent telomere length closer to the time of birth and that the gap between skeletal muscle and leukocyte (a rapidly dividing cell type) telomere length attrition could serve to aid understanding of the associations between telomere length and human disease with each patient, in effect, serving as their own control. The first such study testing this hypothesis showed that increased attrition of telomeres in leukocytes was associated with atherosclerotic cardiovascular disease¹⁴⁰.

Single-nucleotide polymorphisms, telomere biology genes, and disease

The advent of genome-wide association studies (GWAS) opened the door to understanding associations between common genetic variants (that is, greater than 1% minor allele frequency, SNPs) and human disease or phenotypes (Figure 5) (reviewed in 141). GWAS genotype hundreds of thousands of SNPs in thousands of cases and controls and use methods to fine-tune risk estimates through large-scale replication studies and polygenic risk score computation¹⁴².

Numerous GWAS of cancer etiology have identified variants in telomere biology genes as being associated with cancer risk or outcomes. SNPs in the *TERT-CLPTMIL* locus on chromosome 5p15.33 are associated with multiple cancer types, including lung, pancreatic, breast, bladder, ovarian, prostate, and testicular germ cell cancers as well as glioma, melanoma, and non-melanoma skin cancers^{143–146}. There are specific regions of this locus associated with different cancers, but these variants do not specifically encode deleterious coding alleles in *TERT*. They do, however, appear to be connected to telomere length through long-range regulation of this locus¹⁴⁷.

SNPs in *RTEL1* are associated with glioma in large GWAS of this rare brain cancer. The glioma-associated *RTEL1* SNPs are intronic, but functional studies have not yet been completed to understand their potential functions^{148–152}. These findings are intriguing because patients with DC or HH due to autosomal recessive *RTEL1* mutations often have abnormal brain development




Figure 5. Relationship between telomere length, variant allele frequency, and human disease. The majority of genetic variants associated with common disease have a minor allele frequency (MAF) greater than 1% and telomeres in the “normal” range (that is, between the 1st and 99th percentiles for age). In contrast, genetic variants associated with rare and more highly penetrant disease are rare with MAF often much less than 1% and the extremes of telomere length. AA, aplastic anemia; CLL, chronic lymphocytic leukemia; CP, Coats plus; DC, dyskeratosis congenita; FM, familial melanoma; HH, Hoyeraal Hreidarsson syndrome; LD, liver disease; LFL, Li-Fraumeni-like syndrome; MDS, myelodysplastic syndrome; PF, pulmonary fibrosis; RS, Revesz syndrome.

in the form of cerebellar hypoplasia^{35–37,153}. Although the specific genetic loci are different, it is intriguing to speculate that there could be an important biological connection between these findings.

In addition to GWAS of cancer or other illnesses, several GWAS have been conducted to identify novel loci associated with telomere length. SNPs in known telomere biology genes, including *TERT*, *OBFC1* (encodes STN1), *CTCI*, *TERC*, *NAFI*, and *RTEL1*, as well as genes not previously known to be associated with telomere biology have been discovered^{154–157}. These studies illustrate the complexities of telomere length regulation by showing that even common genetic variants, and especially combinations of common genetic variants, are associated with telomere length in the general population.

The existence of telomere length GWAS in various populations set the stage for even larger studies using Mendelian randomization methods in which telomere length-associated SNPs serve as surrogates for telomere length¹⁵⁸. One such study used nine telomere length-associated SNPs to create a telomere length surrogate score and found longer telomere length scores associated with lung adenocarcinoma but not the other cancers¹⁵⁹. Renal cell carcinoma, one of the cancers with reproducible

telomere length association data, was also studied using nine telomere length surrogate SNPs and it was found that genetically longer telomeres were associated with renal cell carcinoma¹⁶⁰.

In 2017, a Mendelian randomization study of 16 telomere length-associated SNPs from 103 GWAS with summary data on 35 cancers and 48 non-neoplastic diseases found that genetically longer telomeres associated with elevated risk of many cancers, including glioma, ovarian cancer, lung cancer, neuroblastoma, bladder, skin, testicular germ cell cancer and kidney cancer, and endometrial cancer¹⁶¹. That study also found an association between genetically shorter telomeres and the risk of interstitial lung disease, celiac disease, abdominal aortic aneurysm, and coronary heart disease but not of other inflammatory or psychiatric diseases¹⁶¹.

Although several studies suggest that telomere length is associated with depression, one study using Mendelian randomization and three SNPs—one each in *TERT*, *TERC*, and *OBFC1*—as surrogates for telomere length in 67,000 individuals did not find an association between depression and genetically shorter telomeres¹⁶². These investigators used the same three SNPs to investigate genetically predicted telomere length and risk of

ischemic heart disease¹⁶³. They found small but statistically significant associations in a dataset of 60,837 ischemic heart disease cases compared with controls.

The studies briefly described above have generated a great deal of enthusiasm but are not without limitations. In many instances, qPCR was used to measure the telomere lengths in GWAS and this assay can be variable between studies. The sensitivity of the assay telomere length measurement and relatively small contributions of SNPs to telomere length should be considered in interpreting large-scale telomere length Mendelian randomization studies.

The way forward

The connections between telomere biology and human disease are complex and myriad and require a multi-disciplinary approach to truly understand the clinically relevant data, important basic science questions, and implications of epidemiologic analyses (Figure 1). As protectors of chromosome ends, telomeres are clearly integral to all aspects of cell biology. They are markers of biological aging and are regulated by a wide range of proteins.

Both very rare and very common germline genetic variants in telomere biology genes are associated with human disease, although the specific clinical phenotypes comprise a wide-ranging disease spectrum. Inheritance of telomere length inheritance and epigenetic regulation are also important aspects of telomere biology and should be incorporated into collaborative studies of rare and common telomere phenotypes. Additionally, optimization of telomere length measurement methods and improved understanding of environmental factors contributing to telomere biology will be essential in order to thoroughly understand these complexities. It is of the utmost importance

for clinicians, epidemiologists, and basic scientists, all of whom study telomeres for a wide variety of different yet important reasons, to work together to build upon the expertise they each possess and incorporate that into improved understanding of telomere biology in human disease. This multi-disciplinary approach will enable the discovery of therapeutics and disease prevention modalities effective for patients with TBDs and for the general population.

Abbreviations

AML, acute myeloid leukemia; CLL, chronic lymphocytic leukemia; DC, dyskeratosis congenita; GWAS, genome-wide association study; HH, Hoyeraal Hreidarsson syndrome; HNSCC, head and neck squamous cell carcinoma; MDS, myelodysplastic syndrome; qPCR, quantitative polymerase chain reaction; RS, Revesz syndrome; SNP, single-nucleotide polymorphism; TBD, telomere biology disorder.

Competing interests

The author declares that she has no competing interests.

Grant information

This work was supported by the intramural research program of the Division of Cancer Epidemiology and Genetics, National Cancer Institute, National Institutes of Health.

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Acknowledgments

The author acknowledges the invaluable contributions of all patients and their families who have taught us so much about the complexities of telomere biology disorders in all their forms.

References



- Zinsser F: **Atrophia cutis reticularis cum pigmentatione, dystrophia unguimet leukoplakia oris.** *Ikongr Dermatol (Hyoto)*. 1906; 5: 219–223.
[Reference Source](#)
- Engman MF: **A unique case of reticular pigmentation of the skin with atrophy.** *Arch Derm Syph.* 1926; 13: 685–687.
[Reference Source](#)
- COLE HN, RAUSCHKOLB JE, TOOMEY J: **DYSKERATOSIS CONGENITA WITH PIGMENTATION, DYSTROPHIA UNGUIS AND LEUKOKERATOSIS ORIS.** *Arch Dermatol.* 1930; 21(1): 71–95.
[Publisher Full Text](#)
- COSTELLO MJ, BUNCKE CM: **Dyskeratosis congenita.** *AMA Arch Derm.* 1956; 73(2): 123–32.
[PubMed Abstract](#) | [Publisher Full Text](#)
- SORROW JM Jr, HITCH JM: **DYSKERATOSIS CONGENITA. FIRST REPORT OF ITS OCCURRENCE IN A FEMALE AND A REVIEW OF THE LITERATURE.** *Arch Dermatol.* 1963; 88(3): 340–7.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Bertuch AA: **The molecular genetics of the telomere biology disorders.** *FNA Biol.* 2016; 13(8): 696–706.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Vulliamy TJ, Marrone A, Knight SW, *et al.*: **Mutations in dyskeratosis congenita: their impact on telomere length and the diversity of clinical presentation.** *Blood.* 2006; 107(7): 2680–5.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Knight SW, Vulliamy T, Forni GL, *et al.*: **Fine mapping of the dyskeratosis congenita locus in Xq28.** *J Med Genet.* 1996; 33(12): 993–5.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Knight SW, Vulliamy TJ, Heiss NS, *et al.*: **1.4 Mb candidate gene region for X linked dyskeratosis congenita defined by combined haplotype and X chromosome inactivation analysis.** *J Med Genet.* 1998; 35(12): 993–6.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Knight SW, Heiss NS, Vulliamy TJ, *et al.*: **X-linked dyskeratosis congenita is predominantly caused by missense mutations in the DKC1 gene.** *Am J Hum Genet.* 1999; 65(1): 50–8.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Mitchell JR, Wood E, Collins K: **A telomerase component is defective in the human disease dyskeratosis congenita.** *Nature.* 1999; 402(6761): 551–5.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
- Baerlocher GM, Mak J, Tien T, *et al.*: **Telomere length measurement by fluorescence *in situ* hybridization and flow cytometry: tips and pitfalls.**

- Cytometry*. 2002; **47**(2): 89–99.
[PubMed Abstract](#) | [Publisher Full Text](#)
13. **F** Alter BP, Baerlocher GM, Savage SA, *et al.*: Very short telomere length by flow fluorescence *in situ* hybridization identifies patients with dyskeratosis congenita. *Blood*. 2007; **110**(5): 1439–47.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 14. Alter BP, Rosenberg PS, Giri N, *et al.*: Telomere length is associated with disease severity and declines with age in dyskeratosis congenita. *Haematologica*. 2012; **97**(3): 353–9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 15. **F** Savage SA, Giri N, Baerlocher GM, *et al.*: *TINF2*, a component of the shelterin telomere protection complex, is mutated in dyskeratosis congenita. *Am J Hum Genet*. 2008; **82**(2): 501–9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 16. **F** Tsakiri KD, Cronkhite JT, Kuan PJ, *et al.*: Adult-onset pulmonary fibrosis caused by mutations in telomerase. *Proc Natl Acad Sci U S A*. 2007; **104**(18): 7552–7.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 17. **F** Armanios MY, Chen JJ, Cogan JD, *et al.*: Telomerase mutations in families with idiopathic pulmonary fibrosis. *N Engl J Med*. 2007; **356**(13): 1317–26.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 18. Yamaguchi H, Baerlocher GM, Lansdorp PM, *et al.*: Mutations of the human telomerase RNA gene (*TERC*) in aplastic anemia and myelodysplastic syndrome. *Blood*. 2003; **102**(3): 916–8.
[PubMed Abstract](#) | [Publisher Full Text](#)
 19. Aalbers AM, Kajigaya S, van den Heuvel-Eibrink MM, *et al.*: Human telomere disease due to disruption of the CCAAT box of the *TERC* promoter. *Blood*. 2012; **119**(13): 3060–3.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 20. **F** Diaz de Leon A, Cronkhite JT, Katzenstein AL, *et al.*: Telomere lengths, pulmonary fibrosis and telomerase (*TERT*) mutations. *PLoS One*. 2010; **5**(5): e10680.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 21. **F** Barbaro PM, Ziegler DS, Reddel RR: The wide-ranging clinical implications of the short telomere syndromes. *Intern Med J*. 2016; **46**(4): 393–403.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 22. Savage SA, Bertuch AA: The genetics and clinical manifestations of telomere biology disorders. *Genet Med*. 2010; **12**(12): 753–64.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 23. **F** Heiss NS, Knight SW, Vulliamy TJ, *et al.*: X-linked dyskeratosis congenita is caused by mutations in a highly conserved gene with putative nucleolar functions. *Nat Genet*. 1998; **19**(1): 32–8.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 24. **F** Vulliamy T, Marrone A, Goldman F, *et al.*: The RNA component of telomerase is mutated in autosomal dominant dyskeratosis congenita. *Nature*. 2001; **413**(6854): 432–5.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 25. Marrone A, Walne A, Tamary H, *et al.*: Telomerase reverse-transcriptase homozygous mutations in autosomal recessive dyskeratosis congenita and Hoyeraal-Hreidarsson syndrome. *Blood*. 2007; **110**(13): 4198–205.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 26. **F** Horn S, Figl A, Rachakonda PS, *et al.*: *TERT* promoter mutations in familial and sporadic melanoma. *Science*. 2013; **339**(6122): 959–61.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 27. Vulliamy TJ, Walne A, Baskaradas A, *et al.*: Mutations in the reverse transcriptase component of telomerase (*TERT*) in patients with bone marrow failure. *Blood Cells Mol Dis*. 2005; **34**(3): 257–63.
[PubMed Abstract](#) | [Publisher Full Text](#)
 28. Walne AJ, Vulliamy T, Marrone A, *et al.*: Genetic heterogeneity in autosomal recessive dyskeratosis congenita with one subtype due to mutations in the telomerase-associated protein NOP10. *Hum Mol Genet*. 2007; **16**(13): 1619–29.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 29. Vulliamy T, Beswick R, Kirwan M, *et al.*: Mutations in the telomerase component NHP2 cause the premature ageing syndrome dyskeratosis congenita. *Proc Natl Acad Sci U S A*. 2008; **105**(23): 8073–8.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 30. **F** Zhong F, Savage SA, Shkreli M, *et al.*: Disruption of telomerase trafficking by TCAB1 mutation causes dyskeratosis congenita. *Genes Dev*. 2011; **25**(1): 11–6.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 31. **F** Venteicher AS, Abreu EB, Meng Z, *et al.*: A human telomerase holoenzyme protein required for Cajal body localization and telomere synthesis. *Science*. 2009; **323**(5914): 644–8.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 32. **F** Anderson BH, Kasher PR, Mayer J, *et al.*: Mutations in *CTC1*, encoding conserved telomere maintenance component 1, cause Coats plus. *Nat Genet*. 2012; **44**(3): 338–42.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 33. Polvi A, Linnankivi T, Kivelä T, *et al.*: Mutations in *CTC1*, encoding the CTS telomere maintenance complex component 1, cause cerebroretinal microangiopathy with calcifications and cysts. *Am J Hum Genet*. 2012; **90**(3): 540–9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 34. Keller RB, Gagne KE, Usmani GN, *et al.*: *CTC1* Mutations in a patient with dyskeratosis congenita. *Pediatr Blood Cancer*. 2012; **59**(2): 311–4.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 35. **F** Ballew BJ, Yeager M, Jacobs K, *et al.*: Germline mutations of regulator of telomere elongation helicase 1, *RTEL1*, in Dyskeratosis congenita. *Hum Genet*. 2013; **132**(4): 473–80.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 36. Deng Z, Glousker G, Molczan A, *et al.*: Inherited mutations in the helicase *RTEL1* cause telomere dysfunction and Hoyeraal-Hreidarsson syndrome. *Proc Natl Acad Sci U S A*. 2013; **110**(36): E3408–16.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 37. **F** Walne AJ, Vulliamy T, Kirwan M, *et al.*: Constitutional mutations in *RTEL1* cause severe dyskeratosis congenita. *Am J Hum Genet*. 2013; **92**(3): 448–53.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 38. Kannengiesser C, Borie R, Ménard C, *et al.*: Heterozygous *RTEL1* mutations are associated with familial pulmonary fibrosis. *Eur Respir J*. 2015; **46**(2): 474–85.
[PubMed Abstract](#) | [Publisher Full Text](#)
 39. Cogan JD, Kropski JA, Zhao M, *et al.*: Rare variants in *RTEL1* are associated with familial interstitial pneumonia. *Am J Respir Crit Care Med*. 2015; **191**(6): 646–55.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 40. Le Guen T, Jullien L, Touzot F, *et al.*: Human *RTEL1* deficiency causes Hoyeraal-Hreidarsson syndrome with short telomeres and genome instability. *Hum Mol Genet*. 2013; **22**(16): 3239–49.
[PubMed Abstract](#) | [Publisher Full Text](#)
 41. Aoude LG, Pritchard AL, Robles-Espinoza CD, *et al.*: Nonsense mutations in the shelterin complex genes *ACD* and *TERF2IP* in familial melanoma. *J Natl Cancer Inst*. 2015; **107**(2): pii: djv408
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 42. **F** Stuart BD, Choi J, Zaidi S, *et al.*: Exome sequencing links mutations in *PARN* and *RTEL1* with familial pulmonary fibrosis and telomere shortening. *Nat Genet*. 2015; **47**(5): 512–7.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 43. Tummala H, Walne A, Collopy L, *et al.*: Poly(A)-specific ribonuclease deficiency impacts telomere biology and causes dyskeratosis congenita. *J Clin Invest*. 2015; **125**(5): 2151–60.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 44. Moon DH, Segal M, Boyraz B, *et al.*: Poly(A)-specific ribonuclease (*PARN*) mediates 3'-end maturation of the telomerase RNA component. *Nat Genet*. 2015; **47**(12): 1482–8.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 45. Dhanraj S, Gunja SM, Deveau AP, *et al.*: Bone marrow failure and developmental delay caused by mutations in poly(A)-specific ribonuclease (*PARN*). *J Med Genet*. 2015; **52**(11): 738–48.
[PubMed Abstract](#) | [Publisher Full Text](#)
 46. Guo Y, Kartawinata M, Li J, *et al.*: Inherited bone marrow failure associated with germline mutation of *ACD*, the gene encoding telomere protein TPP1. *Blood*. 2014; **124**(18): 2767–74.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 47. Speedy HE, Kinnerley B, Chubb D, *et al.*: Germ line mutations in shelterin complex genes are associated with familial chronic lymphocytic leukemia. *Blood*. 2016; **128**(19): 2319–26.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 48. Kocak H, Ballew BJ, Bisht K, *et al.*: Hoyeraal-Hreidarsson syndrome caused by a germline mutation in the TEL patch of the telomere protein TPP1. *Genes Dev*. 2014; **28**(19): 2090–102.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 49. Simon AJ, Lev A, Zhang Y, *et al.*: Mutations in *STN1* cause Coats plus syndrome and are associated with genomic and telomere defects. *J Exp Med*. 2016; **213**(8): 1429–40.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 50. Shi J, Yang XR, Ballew B, *et al.*: Rare missense variants in *POT1* predispose to familial cutaneous malignant melanoma. *Nat Genet*. 2014; **46**(5): 482–6.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 51. Robles-Espinoza CD, Harland M, Ramsay AJ, *et al.*: *POT1* loss-of-function variants predispose to familial melanoma. *Nat Genet*. 2014; **46**(5): 478–81.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 52. Calvete O, Martinez P, Garcia-Pavia P, *et al.*: A mutation in the *POT1* gene is responsible for cardiac angiosarcoma in TP53-negative Li-Fraumeni-like families. *Nat Commun*. 2015; **6**: 8383.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 53. Takai H, Jenkinson E, Kabir S, *et al.*: A *POT1* mutation implicates defective telomere end fill-in and telomere truncations in Coats plus. *Genes Dev*. 2016; **30**(7): 812–26.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 54. Stanley SE, Gable DL, Wagner CL, *et al.*: Loss-of-function mutations in the RNA biogenesis factor *NAF1* predispose to pulmonary fibrosis-empyema. *Sci*


- Transl Med.* 2016; **8**(351): 351ra107.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
55. Savage SA: **Dyskeratosis Congenita**. in GeneReviews(R). R.A. Pagon, et al., Editors. 1993; Seattle (WA).
[PubMed Abstract](#)
56. Yaghmai R, Kimyai-Asadi A, Rostamiani K, et al.: **Overlap of dyskeratosis congenita with the Hoyeraal-Hreidarsson syndrome**. *J Pediatr.* 2000; **136**(3): 390–3.
[PubMed Abstract](#) | [Publisher Full Text](#)
57. Kirwan M, Dokal I: **Dyskeratosis congenita: a genetic disorder of many faces**. *Clin Genet.* 2008; **73**(2): 103–12.
[PubMed Abstract](#) | [Publisher Full Text](#)
58. Revesz T, Fletcher S, al-Gazzali LI, et al.: **Bilateral retinopathy, aplastic anaemia, and central nervous system abnormalities: a new syndrome?** *J Med Genet.* 1992; **29**(9): 673–5.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
59. Savage SA: **Connecting complex disorders through biology**. *Nat Genet.* 2012; **44**(3): 238–40.
[PubMed Abstract](#) | [Publisher Full Text](#)
60. Crow YJ, McMenamin J, Haenggeli CA, et al.: **Coats' plus: a progressive familial syndrome of bilateral Coats' disease, characteristic cerebral calcification, leukoencephalopathy, slow pre- and post-natal linear growth and defects of bone marrow and integument**. *Neuropediatrics.* 2004; **35**(1): 10–9.
[PubMed Abstract](#) | [Publisher Full Text](#)
61. Yamaguchi H, Calado RT, Ly H, et al.: **Mutations in TERT, the gene for telomerase reverse transcriptase, in aplastic anemia**. *N Engl J Med.* 2005; **352**(14): 1413–24.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
62. Ramsay AJ, Quesada V, Foronda M, et al.: **POT1 mutations cause telomere dysfunction in chronic lymphocytic leukemia**. *Nat Genet.* 2013; **45**(5): 526–30.
[PubMed Abstract](#) | [Publisher Full Text](#)
63. Véronèse L, Tourmilhac O, Callanan M, et al.: **Telomeres and chromosomal instability in chronic lymphocytic leukemia**. *Leukemia.* 2013; **27**(2): 490–3.
[PubMed Abstract](#) | [Publisher Full Text](#)
64. Lin TT, Norris K, Heppel NH, et al.: **Telomere dysfunction accurately predicts clinical outcome in chronic lymphocytic leukaemia, even in patients with early stage disease**. *Br J Haematol.* 2014; **167**(2): 214–23.
[PubMed Abstract](#) | [Publisher Full Text](#)
65. Pepper C, Baird D, Fegan C: **Telomere analysis to predict chronic lymphocytic leukemia outcome: a STELA test to change clinical practice?** *Expert Rev Hematol.* 2014; **7**(6): 701–3.
[PubMed Abstract](#) | [Publisher Full Text](#)
66. Aviv A, Anderson JJ, Shay JW: **Mutations, Cancer and the Telomere Length Paradox**. *Trends Cancer.* 2017; **3**(4): 253–8.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
67. Feldser DM, Greider CW: **Short telomeres limit tumor progression in vivo by inducing senescence**. *Cancer Cell.* 2007; **11**(5): 461–9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
68. Dagg RA, Pickett HA, Neumann AA, et al.: **Extensive Proliferation of Human Cancer Cells with Ever-Shorter Telomeres**. *Cell Rep.* 2017; **19**(12): 2544–56.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
69. Reddel RR: **Telomere maintenance mechanisms in cancer: clinical implications**. *Curr Pharm Des.* 2014; **20**(41): 6361–74.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
70. Tian X, Doerig K, Park R, et al.: **Evolution of telomere maintenance and tumour suppressor mechanisms across mammals**. *Philos Trans R Soc Lond B Biol Sci.* 2018; **373**(1741): pii: 20160443.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
71. Robinson NJ, Schiemann WP: **Means to the ends: The role of telomeres and telomere processing machinery in metastasis**. *Biochim Biophys Acta.* 2016; **1866**(2): 320–9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
72. Rousseau P, Autexier C: **Telomere biology: Rationale for diagnostics and therapeutics in cancer**. *RNA Biol.* 2015; **12**(10): 1078–82.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
73. Maciejowski J, de Lange T: **Telomeres in cancer: tumour suppression and genome instability**. *Nat Rev Mol Cell Biol.* 2017; **18**(3): 175–86.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
74. Harley CB, Futcher AB, Greider CW: **Telomeres shorten during ageing of human fibroblasts**. *Nature.* 1990; **345**(6274): 458–60.
[PubMed Abstract](#) | [Publisher Full Text](#)
75. Hemann MT, Strong MA, Hao LY, et al.: **The shortest telomere, not average telomere length, is critical for cell viability and chromosome stability**. *Cell.* 2001; **107**(1): 67–77.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
76. Dilley RL, Greenberg RA: **ALTernative Telomere Maintenance and Cancer**. *Trends Cancer.* 2015; **1**(2): 145–56.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
77. Londoño-Vallejo JA: **Telomere instability and cancer**. *Biochimie.* 2008; **90**(1): 73–82.
[PubMed Abstract](#) | [Publisher Full Text](#)
78. Chiodi I, Mondello C: **Telomere and telomerase stability in human diseases and cancer**. *Front Biosci (Landmark Ed).* 2016; **21**: 203–24.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
79. García-Cao M, Gonzalo S, Dean D, et al.: **A role for the Rb family of proteins in controlling telomere length**. *Nat Genet.* 2002; **32**(3): 415–9.
[PubMed Abstract](#) | [Publisher Full Text](#)
80. Pantic M, Zimmermann S, El Daly H, et al.: **Telomere dysfunction and loss of p53 cooperate in defective mitotic segregation of chromosomes in cancer cells**. *Oncogene.* 2006; **25**(32): 4413–20.
[PubMed Abstract](#) | [Publisher Full Text](#)
81. Pópulo H, Boaventura P, Vinagre J, et al.: **TERT promoter mutations in skin cancer: the effects of sun exposure and X-irradiation**. *J Invest Dermatol.* 2014; **134**(8): 2251–7.
[PubMed Abstract](#) | [Publisher Full Text](#)
82. Vinagre J, Almeida A, Pópulo H, et al.: **Frequency of TERT promoter mutations in human cancers**. *Nat Commun.* 2013; **4**: 2185.
[PubMed Abstract](#) | [Publisher Full Text](#)
83. Huang FW, Hodis E, Xu MJ, et al.: **Highly recurrent TERT promoter mutations in human melanoma**. *Science.* 2013; **339**(6122): 957–9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
84. Alter BP, Giri N, Savage SA, et al.: **Malignancies and survival patterns in the National Cancer Institute inherited bone marrow failure syndromes cohort study**. *Br J Haematol.* 2010; **150**(2): 179–88.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
85. Alter BP, Giri N, Savage SA, et al.: **Cancer in dyskeratosis congenita**. *Blood.* 2009; **113**(26): 6549–57.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
86. Walne AJ, Dokal I: **Dyskeratosis Congenita: a historical perspective**. *Mech Ageing Dev.* 2008; **129**(1–2): 48–59.
[PubMed Abstract](#) | [Publisher Full Text](#)
87. Alter BP, Giri N, Savage SA, et al.: **Cancer in the National Cancer Institute inherited bone marrow failure syndrome cohort after fifteen years of follow-up**. *Haematologica.* 2018; **103**(1): 30–9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
88. Bodelon C, Savage SA, Gadalla SM: **Telomeres in molecular epidemiology studies**. *Prog Mol Biol Transl Sci.* 2014; **125**: 113–31.
[PubMed Abstract](#) | [Publisher Full Text](#)
89. Aviv A, Valdes AM, Spector TD: **Human telomere biology: pitfalls of moving from the laboratory to epidemiology**. *Int J Epidemiol.* 2006; **35**(6): 1424–9.
[PubMed Abstract](#) | [Publisher Full Text](#)
90. Lai TP, Wright WE, Shay JW: **Comparison of telomere length measurement methods**. *Philos Trans R Soc Lond B Biol Sci.* 2018; **373**(1741): pii: 20160451.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
91. Aubert G, Hills M, Lansdorp PM: **Telomere length measurement-caveats and a critical assessment of the available technologies and tools**. *Mutat Res.* 2012; **730**(1–2): 59–67.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
92. Cawthon RM: **Telomere length measurement by a novel monochrome multiplex quantitative PCR method**. *Nucleic Acids Res.* 2009; **37**(3): e21.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
93. Cawthon RM: **Telomere measurement by quantitative PCR**. *Nucleic Acids Res.* 2002; **30**(10): e47.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
94. Dagnall CL, Hicks B, Teshome K, et al.: **Effect of pre-analytic variables on the reproducibility of qPCR relative telomere length measurement**. *PLoS One.* 2017; **12**(9): e0184098.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
95. Martin-Ruiz CM, Baird D, Roger L, et al.: **Reproducibility of telomere length assessment: an international collaborative study**. *Int J Epidemiol.* 2015; **44**(5): 1673–83.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
96. Gadalla SM, Khincha PP, Katki HA, et al.: **The limitations of qPCR telomere length measurement in diagnosing dyskeratosis congenita**. *Mol Genet Genomic Med.* 2016; **4**(4): 475–9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
97. Hansen ME, Hunt SC, Stone RC, et al.: **Shorter telomere length in Europeans than in Africans due to polygenetic adaptation**. *Hum Mol Genet.* 2016; **25**(11): 2324–30.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
98. Hunt SC, Chen W, Gardner JP, et al.: **Leukocyte telomeres are longer in African Americans than in whites: the National Heart, Lung, and Blood Institute Family Heart Study and the Bogalusa Heart Study**. *Aging Cell.* 2008; **7**(4): 451–8.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
99. Astuti Y, Wardhana A, Watkins J, et al.: **Cigarette smoking and telomere length: A systematic review of 84 studies and meta-analysis**. *Environ Res.* 2017; **158**: 480–9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)

100. Salihi HM, Pradhan A, King L, *et al.*: **Impact of intrauterine tobacco exposure on fetal telomere length.** *Am J Obstet Gynecol.* 2015; 212(2): 205.e1–8.
[PubMed Abstract](#) | [Publisher Full Text](#)
101. **F** Ip P, Chung BH, Ho FK, *et al.*: **Prenatal Tobacco Exposure Shortens Telomere Length in Children.** *Nicotine Tob Res.* 2017; 19(1): 111–8.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
102. **F** Martens DS, Cox B, Janssen BG, *et al.*: **Prenatal Air Pollution and Newborns' Predisposition to Accelerated Biological Aging.** *JAMA Pediatr.* 2017; 171(12): 1160–7.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
103. **F** Lu L, Johnman C, McGlynn L, *et al.*: **Association between exposure to second-hand smoke and telomere length: cross-sectional study of 1303 non-smokers.** *Int J Epidemiol.* 2017; 46(6): 1978–84.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
104. Andreotti G, Hoppin JA, Hou L, *et al.*: **Pesticide Use and Relative Leukocyte Telomere Length in the Agricultural Health Study.** *PLoS One.* 2015; 10(7): e0133382.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
105. Hou L, Andreotti G, Baccarelli AA, *et al.*: **Lifetime pesticide use and telomere shortening among male pesticide applicators in the Agricultural Health Study.** *Environ Health Perspect.* 2013; 121(8): 919–24.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
106. Wu X, Amos CI, Zhu Y, *et al.*: **Telomere dysfunction: a potential cancer predisposition factor.** *J Natl Cancer Inst.* 2003; 95(16): 1211–8.
[PubMed Abstract](#) | [Publisher Full Text](#)
107. Jang JS, Choi YY, Lee WK, *et al.*: **Telomere length and the risk of lung cancer.** *Cancer Sci.* 2008; 99(7): 1385–9.
[PubMed Abstract](#) | [Publisher Full Text](#)
108. Mirabello L, Garcia-Closas M, Cawthon R, *et al.*: **Leukocyte telomere length in a population-based case-control study of ovarian cancer: a pilot study.** *Cancer Causes Control.* 2010; 21(1): 77–82.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
109. Mirabello L, Huang WY, Wong JY, *et al.*: **The association between leukocyte telomere length and cigarette smoking, dietary and physical variables, and risk of prostate cancer.** *Aging Cell.* 2009; 8(4): 405–13.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
110. Gramatges MM, Telli ML, Balise R, *et al.*: **Longer relative telomere length in blood from women with sporadic and familial breast cancer compared with healthy controls.** *Cancer Epidemiol Biomarkers Prev.* 2010; 19(4): 605–13.
[PubMed Abstract](#) | [Publisher Full Text](#)
111. de Vivo I, Prescott J, Wong JY, *et al.*: **A prospective study of relative telomere length and postmenopausal breast cancer risk.** *Cancer Epidemiol Biomarkers Prev.* 2009; 18(4): 1152–6.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
112. Shen J, Terry MB, Gurvich I, *et al.*: **Short telomere length and breast cancer risk: a study in sister sets.** *Cancer Res.* 2007; 67(11): 5538–44.
[PubMed Abstract](#) | [Publisher Full Text](#)
113. Wentzensen IM, Mirabello L, Pfeiffer RM, *et al.*: **The association of telomere length and cancer: a meta-analysis.** *Cancer Epidemiol Biomarkers Prev.* 2011; 20(6): 1238–50.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
114. Ma H, Zhou Z, Wei S, *et al.*: **Shortened telomere length is associated with increased risk of cancer: a meta-analysis.** *PLoS One.* 2011; 6(6): e20466.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
115. Pooley KA, Sandhu MS, Tyrer J, *et al.*: **Telomere length in prospective and retrospective cancer case-control studies.** *Cancer Res.* 2010; 70(8): 3170–6.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
116. Zhu X, Han W, Xue W, *et al.*: **The association between telomere length and cancer risk in population studies.** *Sci Rep.* 2016; 6: 22243.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
117. Seow WJ, Cawthon RM, Purdue MP, *et al.*: **Telomere length in white blood cell DNA and lung cancer: a pooled analysis of three prospective cohorts.** *Cancer Res.* 2014; 74(15): 4090–8.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
118. Sanchez-Espiridon B, Chen M, Chang JY, *et al.*: **Telomere length in peripheral blood leukocytes and lung cancer risk: a large case-control study in Caucasians.** *Cancer Res.* 2014; 74(9): 2476–86.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
119. Rode L, Nordestgaard BG, Bojesen SE: **Long telomeres and cancer risk among 95 568 individuals from the general population.** *Int J Epidemiol.* 2016; 45(5): 1634–43.
[PubMed Abstract](#) | [Publisher Full Text](#)
120. Weischer M, Nordestgaard BG, Cawthon RM, *et al.*: **Short telomere length, cancer survival, and cancer risk in 47102 individuals.** *J Natl Cancer Inst.* 2013; 105(7): 459–68.
[PubMed Abstract](#) | [Publisher Full Text](#)
121. **F** Osler M, Bendix L, Rask L, *et al.*: **Stressful life events and leukocyte telomere length: Do lifestyle factors, somatic and mental health, or low grade inflammation mediate this relationship? Results from a cohort of Danish men born in 1953.** *Brain Behav Immun.* 2016; 58: 248–53.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
122. Lindqvist D, Epel ES, Mellon SH, *et al.*: **Psychiatric disorders and leukocyte telomere length: Underlying mechanisms linking mental illness with cellular aging.** *Neurosci Biobehav Rev.* 2015; 55: 333–64.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
123. Shalev I, Entringer S, Wadhwa PD, *et al.*: **Stress and telomere biology: a lifespan perspective.** *Psychoneuroendocrinology.* 2013; 38(9): 1835–42.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
124. **F** Mathur MB, Epel E, Kind S, *et al.*: **Perceived stress and telomere length: A systematic review, meta-analysis, and methodologic considerations for advancing the field.** *Brain Behav Immun.* 2016; 54: 158–69.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
125. Hoen PW, de Jonge P, Na BY, *et al.*: **Depression and leukocyte telomere length in patients with coronary heart disease: data from the Heart and Soul Study.** *Psychosom Med.* 2011; 73(7): 541–7.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
126. **F** Révész D, Verhoeven JE, Milaneschi Y, *et al.*: **Depressive and anxiety disorders and short leukocyte telomere length: mediating effects of metabolic stress and lifestyle factors.** *Psychol Med.* 2016; 46(11): 2337–49.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
127. Rackley S, Pao M, Seratti GF, *et al.*: **Neuropsychiatric conditions among patients with dyskeratosis congenita: a link with telomere biology?** *Psychosomatics.* 2012; 53(3): 230–5.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
128. Mundstock E, Zatti H, Louzada FM, *et al.*: **Effects of physical activity in telomere length: Systematic review and meta-analysis.** *Ageing Res Rev.* 2015; 22: 72–80.
[PubMed Abstract](#) | [Publisher Full Text](#)
129. Ludlow AT, Ludlow LW, Roth SM: **Do telomeres adapt to physiological stress? Exploring the effect of exercise on telomere length and telomere-related proteins.** *Biomed Res Int.* 2013; 2013: 601368.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
130. Mathur S, Ardestani A, Parker B, *et al.*: **Telomere length and cardiorespiratory fitness in marathon runners.** *J Investig Med.* 2013; 61(3): 613–5.
[PubMed Abstract](#) | [Publisher Full Text](#)
131. Østhus IB, Sgura A, Berardinelli F, *et al.*: **Telomere length and long-term endurance exercise: does exercise training affect biological age? A pilot study.** *PLoS One.* 2012; 7(12): e52769.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
132. **F** García-Calzón S, Martínez-González MA, Razquin C, *et al.*: **Mediterranean diet and telomere length in high cardiovascular risk subjects from the PREDIMED-NAVARRA study.** *Clin Nutr.* 2016; 35(6): 1399–405.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
133. Sun Q, Shi L, Prescott J, *et al.*: **Healthy lifestyle and leukocyte telomere length in U.S. women.** *PLoS One.* 2012; 7(5): e38374.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
134. **F** Chilton W, O'Brien B, Charchar F: **Telomeres, Aging and Exercise: Guilty by Association?** *Int J Mol Sci.* 2017; 18(12): pii: E2573.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
135. **F** Haycock PC, Heydon EE, Kaptoge S, *et al.*: **Leukocyte telomere length and risk of cardiovascular disease: systematic review and meta-analysis.** *BMJ.* 2014; 349: g4227.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
136. D'Mello MJ, Ross SA, Briel M, *et al.*: **Association between shortened leukocyte telomere length and cardiometabolic outcomes: systematic review and meta-analysis.** *Circ Cardiovasc Genet.* 2015; 8(1): 82–90.
[PubMed Abstract](#) | [Publisher Full Text](#)
137. **F** Fitzpatrick AL, Kronmal RA, Gardner JP, *et al.*: **Leukocyte telomere length and cardiovascular disease in the cardiovascular health study.** *Am J Epidemiol.* 2007; 165(1): 14–21.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
138. **F** Sabharwal S, Verhulst S, Guirguis G, *et al.*: **Telomere length dynamics in early life: the blood-and-muscle model.** *FASEB J.* 2018; 32(1): 529–34.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
139. **F** Stone RC, Horvath K, Kark JD, *et al.*: **Telomere Length and the Cancer-Atherosclerosis Trade-Off.** *PLoS Genet.* 2016; 12(7): e1006144.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
140. **F** Benetos A, Toupance S, Gautier S, *et al.*: **Short Leukocyte Telomere Length Precedes Clinical Expression of Atherosclerosis: The Blood-and-Muscle Model.** *Circ Res.* 2018; 122(4): 616–23.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
141. **F** Sud A, Kinnerley B, Houlston RS: **Genome-wide association studies of cancer: current insights and future perspectives.** *Nat Rev Cancer.* 2017; 17(11): 692–704.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
142. **F** Shi J, Park JH, Duan J, *et al.*: **Winner's Curse Correction and Variable Thresholding Improve Performance of Polygenic Risk Modeling Based on Genome-Wide Association Study Summary-Level Data.** *PLoS Genet.* 2016; 12(12): e1006493.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
143. **F** Zhang M, Wang Z, Obazee O, *et al.*: **Three new pancreatic cancer susceptibility signals identified on chromosomes 1q32.1, 5p15.33 and 8q24.21.**

- Oncotarget*. 2016; 7(41): 66328–43.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
144. Wang Z, Zhu B, Zhang M, *et al.*: **Imputation and subset-based association analysis across different cancer types identifies multiple independent risk loci in the *TERT-CLPTM1L* region on chromosome 5p15.33.** *Hum Mol Genet*. 2014; 23(24): 6616–33.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
145. Kote-Jarai Z, Saunders EJ, Leongamornlert DA, *et al.*: **Fine-mapping identifies multiple prostate cancer risk loci at 5p15, one of which associates with *TERT* expression.** *Hum Mol Genet*. 2013; 22(12): 2520–8.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
146. Bojesen SE, Pooley KA, Johnatty SE, *et al.*: **Multiple independent variants at the *TERT* locus are associated with telomere length and risks of breast and ovarian cancer.** *Nat Genet*. 2013; 45(4): 371–84, 384e1–2.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
147. **F** Fang J, Jia J, Makowski M, *et al.*: **Functional characterization of a multi-cancer risk locus on chr5p15.33 reveals regulation of *TERT* by ZNF148.** *Nat Commun*. 2017; 8: 15034.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
148. Melin B, Dahlin AM, Andersson U, *et al.*: **Known glioma risk loci are associated with glioma with a family history of brain tumours -- a case-control gene association study.** *Int J Cancer*. 2013; 132(10): 2464–8.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
149. Jin TB, Zhang JY, Li G, *et al.*: ***RTEL1* and *TERT* polymorphisms are associated with astrocytoma risk in the Chinese Han population.** *Tumour Biol*. 2013; 34(6): 3659–66.
[PubMed Abstract](#) | [Publisher Full Text](#)
150. **F** Shete S, Hosking FJ, Robertson LB, *et al.*: **Genome-wide association study identifies five susceptibility loci for glioma.** *Nat Genet*. 2009; 41(8): 899–904.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
151. **F** Wrensch M, Jenkins RB, Chang JS, *et al.*: **Variants in the *CDKN2B* and *RTEL1* regions are associated with high-grade glioma susceptibility.** *Nat Genet*. 2009; 41(8): 905–8.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
152. **F** Delgado DA, Zhang C, Chen LS, *et al.*: **Genome-wide association study of telomere length among South Asians identifies a second *RTEL1* association signal.** *J Med Genet*. 2018; 55(1): 64–71.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
153. Ballew BJ, Joseph V, De S, *et al.*: **A recessive founder mutation in regulator of telomere elongation helicase 1, *RTEL1*, underlies severe immunodeficiency and features of Hoyeraal Hreidarsson syndrome.** *PLoS Genet*. 2013; 9(8): e1003695.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
154. Pooley KA, Bojesen SE, Weischer M, *et al.*: **A genome-wide association scan (GWAS) for mean telomere length within the COGS project: identified loci show little association with hormone-related cancer risk.** *Hum Mol Genet*. 2013; 22(24): 5056–64.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
155. Mangino M, Hwang SJ, Spector TD, *et al.*: **Genome-wide meta-analysis points to *CTC1* and *ZNF676* as genes regulating telomere homeostasis in humans.** *Hum Mol Genet*. 2012; 21(24): 5385–94.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
156. **F** Mirabello L, Yu K, Kraft P, *et al.*: **The association of telomere length and genetic variation in telomere biology genes.** *Hum Mutat*. 2010; 31(9): 1050–8.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
157. Prescott J, Kraft P, Chasman DI, *et al.*: **Genome-wide association study of relative telomere length.** *PLoS One*. 2011; 6(5): e19635.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
158. Emdin CA, Khera AV, Kathiresan S: **Mendelian Randomization.** *JAMA*. 2017; 318(19): 1925–6.
[PubMed Abstract](#) | [Publisher Full Text](#)
159. Zhang C, Doherty JA, Burgess S, *et al.*: **Genetic determinants of telomere length and risk of common cancers: a Mendelian randomization study.** *Hum Mol Genet*. 2015; 24(18): 5356–66.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
160. Machiela MJ, Hofmann JN, Carreras-Torres R, *et al.*: **Genetic Variants Related to Longer Telomere Length are Associated with Increased Risk of Renal Cell Carcinoma.** *Eur Urol*. 2017; 72(5): 747–54.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
161. **F** Telomeres Mendelian Randomization Collaboration, Haycock PC, Burgess S, *et al.*: **Association Between Telomere Length and Risk of Cancer and Non-Neoplastic Diseases: a Mendelian Randomization Study.** *JAMA Oncol*. 2017; 3(5): 636–651.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
162. Wium-Andersen MK, Ørsted DD, Rode L, *et al.*: **Telomere length and depression: prospective cohort study and Mendelian randomisation study in 67 306 individuals.** *Br J Psychiatry*. 2017; 210(1): 31–8.
[PubMed Abstract](#) | [Publisher Full Text](#)
163. **F** Scheller Madrid A, Rode L, Nordestgaard BG, *et al.*: **Short Telomere Length and Ischemic Heart Disease: Observational and Genetic Studies in 290 022 Individuals.** *Clin Chem*. 2016; 62(8): 1140–9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)

Open Peer Review

Current Referee Status:



Editorial Note on the Review Process

F1000 Faculty Reviews are commissioned from members of the prestigious F1000 Faculty and are edited as a service to readers. In order to make these reviews as comprehensive and accessible as possible, the referees provide input before publication and only the final, revised version is published. The referees who approved the final version are listed with their names and affiliations but without their reports on earlier versions (any comments will already have been addressed in the published version).

The referees who approved this article are:

Version 1

- 1 **Fadi Charchar** School of Applied and Biomedical Sciences, Faculty of Science and Technology, Federation University, Ballarat, Victoria, Australia
Competing Interests: No competing interests were disclosed.
- 1 **Steven E. Artandi** ^{1,2,3} 1 Department of Medicine, Stanford University School of Medicine, Stanford, California, USA
² Cancer Biology Program, Stanford University School of Medicine, Stanford, California, USA
³ Department of Biochemistry, Stanford University School of Medicine, Stanford, California, USA
Competing Interests: No competing interests were disclosed.
- 1 **Tracy Bryan** Children's Medical Research Institute, University of Sydney, Sydney, New South Wales, Australia
Competing Interests: No competing interests were disclosed.
- 1 **Dirk Hockemeyer** Department of Molecular and Cell Biology, University of California, Berkeley, Berkeley, California, USA
Competing Interests: No competing interests were disclosed.

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research