

Cite this: *Chem. Sci.*, 2017, 8, 6290

# Triple bonds of niobium with silicon, germanium and tin: the tetrylidyne complexes $[(\kappa^3\text{-tmps})(\text{CO})_2\text{Nb}\equiv\text{E}-\text{R}]$ (E = Si, Ge, Sn; $\text{tmps} = \text{MeSi}(\text{CH}_2\text{PMe}_2)_3$ ; R = aryl)<sup>†</sup>

Alexander C. Filippou,<sup>ID\*</sup> David Hoffmann<sup>ID</sup> and Gregor Schnakenburg<sup>ID</sup>

A systematic, efficient approach to first complexes containing a triple bond between niobium and the elements silicon, germanium or tin is reported. The approach involves a metathetical exchange of the niobium-centered nucleophile  $(\text{NMe}_4)[\text{Nb}(\text{CO})_4(\kappa^2\text{-tmps})]$  (**1**) ( $\text{tmps} = \text{MeSi}(\text{CH}_2\text{PMe}_2)_3$ ) with a suitable organotetrel(II)halide. Compound **1** was obtained from  $(\text{NMe}_4)[\text{Nb}(\text{CO})_6]$  and the triphosphane  $\text{tmps}$  by photodecarbonylation. Reaction of **1** with the disilene  $E\text{-Tbb}(\text{Br})\text{Si}=\text{Si}(\text{Br})\text{Tbb}$  in the presence of 4-dimethylaminopyridine afforded selectively the red-brown silylidyne complex  $[(\kappa^3\text{-tmps})(\text{CO})_2\text{Nb}\equiv\text{Si}-\text{Tbb}]$  (**2-Si**, Tbb = 4-*tert*-butyl-2,6-bis(bis(trimethylsilyl)methyl)phenyl). Similarly, treatment of **1** with  $\text{E}(\text{Ar}^{\text{Mes}})\text{Cl}$  (E = Ge, Sn;  $\text{Ar}^{\text{Mes}} = 2,6\text{-mesitylphenyl}$ ) afforded after elimination of  $(\text{NMe}_4)\text{Cl}$  and two CO ligands the deep magenta colored germylidyne complex  $[(\kappa^3\text{-tmps})(\text{CO})_2\text{Nb}\equiv\text{Ge}-\text{Ar}^{\text{Mes}}]$  (**3-Ge**), and the deep violet, light-sensitive stannylidyne complex  $[(\kappa^3\text{-tmps})(\text{CO})_2\text{Nb}\equiv\text{Sn}-\text{Ar}^{\text{Mes}}]$  (**3-Sn**), respectively. Formation of **3-Sn** proceeds *via* the niobiastannylene  $[(\kappa^3\text{-tmps})(\text{CO})_3\text{Nb}-\text{SnAr}^{\text{Mes}}]$  (**4-Sn**), which was detected by IR and NMR spectroscopy. The niobium tetrylidyne complexes **2-Si**, **3-Ge** and **3-Sn** were fully characterized and their solid-state structures determined by single-crystal X-ray diffraction studies. All complexes feature an almost linear tetrel coordination and the shortest Nb–E bond lengths ( $d(\text{Nb}-\text{Si}) = 232.7(2)$  pm;  $d(\text{Nb}-\text{Ge}) = 235.79(4)$  pm;  $d(\text{Nb}-\text{Sn}) = 253.3(1)$  pm) reported to date. Reaction of **3-Ge** with a large excess of  $\text{H}_2\text{O}$  afforded upon cleavage of the Nb–Ge triple bond the hydridogermanediol  $\text{Ge}(\text{Ar}^{\text{Mes}})\text{H}(\text{OH})_2$ . Photodecarbonylation of  $[\text{CpNb}(\text{CO})_4]$  (Cp =  $\eta^5\text{-C}_5\text{H}_5$ ) in the presence of  $\text{Ge}(\text{Ar}^{\text{Mes}})\text{Cl}$  afforded the red-orange chlorogermlydene complex  $[\text{Cp}(\text{CO})_3\text{Nb}=\text{Ge}(\text{Ar}^{\text{Mes}})\text{Cl}]$  (**5-Ge**). The molecular structure of **5-Ge** features an upright conformation of the germlydene ligand, a trigonal-planar coordinated Ge atom, and a Nb–Ge double bond length of 251.78(6) pm, which lies in-between the Nb–Ge triple bond length of **3-Ge** (235.79(4) pm) and a Nb–Ge single bond length (267.3 pm). Cyclic voltammetric studies of **2-Si**, **3-Ge**, and **3-Sn** reveal several electron-transfer steps. One-electron oxidation and reduction of the germlydene complex of **3-Ge** in THF are electrochemically reversible suggesting that both the radical cation and radical anion of **3-Ge** are accessible species in solution.

Received 18th June 2017

Accepted 2nd July 2017

DOI: 10.1039/c7sc02708g

rsc.li/chemical-science

## Introduction

Complexes of the general formula  $[\text{L}_n\text{M}\equiv\text{E}-\text{R}]$  (M = d-block metal; E = Si–Pb; R = singly bonded group (*e.g.* alkyl, aryl);  $\text{L}_n$  = ligand sphere) featuring a triple bond between a d-block metal and the tetrels Si/Ge/Sn/Pb are an intriguing class of

compounds with an auspicious synthetic potential originating from the highly reactive, polar  $\text{M}\equiv\text{E}$  bond.<sup>1–4</sup> Isolation of these compounds is very challenging and requires specific stereo-electronic properties of the metal fragment  $\text{L}_n\text{M}$  as well as a steric protection of the electrophilic tetrel center by a tailor-made, bulky substituent R to circumvent a head-to-tail cyclo-dimerisation or unintentional intra- or intermolecular  $\sigma$ -bond activations destroying the  $\text{M}\equiv\text{E}-\text{R}$  functionality. Whereas earlier work concentrated exclusively on group 6 metals, recent studies have shown that also group 7,<sup>2l,3d,4d</sup> group 8 (ref. 1c and 5) and even group 10 metals<sup>6</sup> can be incorporated into triple bonding with the tetrels Si–Pb. Extension of this chemistry to the group 5 elements V–Ta seemed attractive to investigate whether the lower electronegativity and larger metallic radii of these elements compared to Cr–W would have an effect on the

*Institut für Anorganische Chemie, Rheinische Friedrich-Wilhelms-Universität Bonn, Gerhard-Domagk-Straße 1, 53121 Bonn, Germany. E-mail: filippou@uni-bonn.de*

<sup>†</sup> Electronic supplementary information (ESI) available: Syntheses and analytical data of **1**, **2-Si**, **3-Ge**, **3-Sn** and **5-Ge**, illustrations of the IR and heteronuclear magnetic resonance spectra of **1**, **2-Si**, **3-Ge**, **3-Sn** and **5-Ge**, details of the cyclic voltammetric studies of **2-Si**, **3-Ge** and **3-Sn**, and crystal structure determination of **2-Si**, **3-Ge** (THF), **3-Sn** (toluene) and **5-Ge**. CCDC 1553387–1553389 and 1555671. For ESI and crystallographic data in CIF or other electronic format see DOI: 10.1039/c7sc02708g




$M\equiv E$  functionality. Group 5 metal complexes featuring a triple bond to the heavier tetrrels ( $E = \text{Si-Pb}$ ) are presently not known, and even compounds with a  $M=E$  double bond are very scarce and poorly characterized illustrating the challenge to make such compounds.<sup>7</sup> We decided to address this issue, and present herein a systematic, efficient approach to the first complexes containing  $\text{Nb}\equiv E$  ( $E = \text{Si-Sn}$ ) triple bonds.

## Results and discussion


Two methods have been employed so far for the formation of transition metal-tetrrel ( $\text{Si-Pb}$ ) triple bonds. The first method, abbreviated as the “salt elimination method”, involves a substitution reaction of a suitable anionic 18 VE metal complex with an organotetrrel(II) halide, as exemplified by the synthesis of Cp-substituted group 6 metal tetrylidyne complexes (Scheme 1).<sup>2a,2b,2i,2j,2m</sup>

The second method, commonly termed “ $\text{N}_2/\text{PMe}_3$  elimination method”, takes advantage of the exchange of labile ligands (mostly  $\text{N}_2$  or  $\text{PMe}_3$ ) in neutral 18 VE metal complexes by suitable organotetrrel(II) halides. This approach may afford directly neutral ylidyne complexes, as demonstrated by the syntheses of phosphane-substituted group 6 and 7 metal tetrylidyne complexes (Scheme 2).<sup>2f,2g,2l,3a,4a,4b</sup>



Alternatively, haloylidene complexes are initially obtained by this method, which are subsequently converted to cationic ylidyne complexes by halide abstraction. Examples demonstrating this reaction path include the preparation of group 8 and 10 ylidyne complexes (Scheme 3).<sup>5,6</sup>



Scheme 1 Preparation of half-sandwich group 6 metal germlylidyne complexes by the salt elimination method.



Scheme 2 Preparation of neutral group 6 and 7 metal tetrylidyne complexes by the  $\text{N}_2/\text{PMe}_3$  elimination method.



$M = \text{Ni, Pt}$ ;  $\text{Tbb} = \text{C}_6\text{H}_2-2,6-(\text{CH}(\text{SiMe}_3)_2)_2-4-\text{tBu}$ ;  $\text{Ar}^F = \text{C}_6\text{H}_3-3,5-(\text{CF}_3)_2$

Scheme 3 Preparation of group 8 and 10 metal tetrylidyne complexes via haloylidene complexes using the  $\text{N}_2/\text{PMe}_3$  elimination method.

We decided to apply the first method, given the availability of anionic niobium carbonyl complexes.<sup>8</sup> At first, the homoleptic carbonyl niobate  $[\text{Nb}(\text{CO})_6]^-$  was chosen. For this purpose the canary yellow salts  $(\text{NR}_4)[\text{Nb}(\text{CO})_6]$  ( $R = \text{Me, Et}$ ) were prepared, following the method developed by J. E. Ellis *et al.*<sup>9</sup> However, these compounds proved to be unreactive towards the *m*-terphenyltetrrel(II)halides  $\text{E}(\text{Ar}^{\text{Mes}})\text{Cl}$  ( $E = \text{Ge, Sn}$ ;  $\text{Ar}^{\text{Mes}} = 2,6$ -mesitylphenyl; mesityl ( $\text{Mes}$ ) = 2,4,6-trimethylphenyl).<sup>10</sup> For example, IR monitoring of the reaction of  $(\text{NET}_4)[\text{Nb}(\text{CO})_6]$  with  $\text{Ge}(\text{Ar}^{\text{Mes}})\text{Cl}$  in refluxing toluene did not provide any evidence for a conversion of the niobate even after prolonged heating, probably due to the poor nucleophilicity of  $[\text{Nb}(\text{CO})_6]^-$ . Therefore, as next we turned our attention to niobates containing ligands with a higher  $\sigma$ -donor/ $\pi$ -acceptor ratio than CO, such as trialkyl- or triarylphosphanes. Various carbonyl(phosphane) niobates of the general formula  $[\text{Nb}(\text{CO})_4\text{L}_2]^-$  ( $\text{L}_2 =$  bidentate di- or oligo-arylphosphane ligand) have been accessed from  $[\text{Nb}(\text{CO})_6]^-$  upon photolytic CO substitution.<sup>11</sup> In order to increase the electron density at the metal centre, we decided to use the highly basic, albeit, very oxygen-sensitive, tripodal alkylphosphane  $\text{MeSi}(\text{CH}_2\text{PMe}_2)_3$  (*tmps*).<sup>12</sup>

Photolysis of  $(\text{NMe}_4)[\text{Nb}(\text{CO})_6]$  was carried out in the presence of one equivalent of *tmps* in THF at room temperature. A high-power blue light LED ( $\lambda = 465 \text{ nm}$ ) was used instead of a high-pressure mercury UV-lamp. The use of a nearly monochromatic source with an exciting wavelength close to the longest-wavelength absorption maximum of  $[\text{Nb}(\text{CO})_6]^-$  ( $\lambda_{\text{max}} = 440 \text{ nm}$  in  $\text{CH}_2\text{Cl}_2$ )<sup>13</sup> was conceived to be advantageous preventing the formation of insoluble brown decomposition



products formed during the photolysis using a high-pressure mercury-lamp.<sup>11a</sup>

In fact, IR-monitoring of the reaction revealed a slow, but very selective conversion into the tetracarbonyl niobate  $[\text{Nb}(\text{CO})_4(\kappa^2\text{-tmps})]^-$  proceeding *via* the pentacarbonyl intermediate  $[\text{Nb}(\text{CO})_5(\kappa^1\text{-tmps})]^-$  ( $\nu(\text{CO})$  in THF: 1966 (m), 1821 (vs)  $\text{cm}^{-1}$ ). After work-up the salt  $(\text{NMe}_4)[\text{Nb}(\text{CO})_4(\kappa^2\text{-tmps})]$  (**1**) was isolated in nearly quantitative yield (97%) as an orange, analytically pure, very air-sensitive powder, which decolorizes immediately upon exposure to air. The salt decomposes upon heating at 142 °C to a dark brown mass, and is well soluble in acetonitrile and tetrahydrofuran (THF), but only moderately soluble in benzene, toluene, and diethyl ether. Attempts to grow suitable single crystals of **1** for an X-ray diffraction study failed, however unambiguous proof for the composition and structure of **1** was provided by elemental analysis, IR spectroscopy and  $^1\text{H}$ ,  $^{13}\text{C}\{^1\text{H}\}$ ,  $^{31}\text{P}\{^1\text{H}\}$  and  $^{29}\text{Si}\{^1\text{H}\}$  NMR spectroscopy. The IR spectrum of **1** in THF displays four  $\nu(\text{CO})$  absorption bands at 1900, 1787, 1764 and 1732  $\text{cm}^{-1}$  (Fig. 1a), the band pattern being typical for octahedral *cis*-disubstituted metal tetracarbonyl complexes with a local  $C_{2v}$  symmetry of the  $\text{M}(\text{CO})_4$  fragment.<sup>14</sup> All  $\nu(\text{CO})$  bands of **1** are shifted to lower frequencies than those of  $[\text{Nb}(\text{CO})_4(\text{Ph}_2\text{PCH}_2\text{CH}_2\text{PPh}_2)]^-$  ( $\nu(\text{CO})$  in THF = 1908, 1806, 1782 and 1746  $\text{cm}^{-1}$ ) or related disubstituted arylphosphane-carbonyl niobates.<sup>11b</sup> This shift to lower

frequencies evinces the stronger +I effect of the P-bonded alkyl substituents in **1**, which enhances the electron density at the metal center and leads to a stronger  $\text{Nb}(\text{d}\pi) \rightarrow \text{CO}(\pi^*)$  back-bonding and softening of the CO bonds in **1**. The NMR spectra of **1** corroborate the presence of an overall  $C_s$  symmetric complex, in which one of the arms of the tripodal ligand tmps is pendant and the other two arms are bonded to the niobium center. For example, the  $^{31}\text{P}\{^1\text{H}\}$  NMR spectrum of **1** displays a sharp singlet for the  $^{31}\text{P}$  nucleus of the pendant  $\text{CH}_2\text{PMe}_2$  arm, which appears at almost the same position ( $\delta(\text{P}_A) = -55.8$  ppm in benzene- $d_6$ ) as that of the non-coordinated ("free") tmps ( $\delta(\text{P}) = -55.1$  ppm in benzene- $d_6$ ), and a very broad signal for the two symmetry-equivalent Nb-bonded  $^{31}\text{P}$  nuclei at considerably lower field ( $\delta(\text{P}_B) = -11.6$  ppm in benzene- $d_6$ ) (Fig. 1b). The broadness of the second signal ( $\Delta\nu_{1/2}$  (full width at half maximum) = 696 Hz) is caused by the quadrupole moment of the  $^{93}\text{Nb}$  nucleus ( $Q = -0.32 \times 10^{-28} \text{ m}^2$ ;  $I = 9/2$ , 100% natural abundance) and its effect on the relaxation time.<sup>15</sup> Further structural information was provided by the  $^{29}\text{Si}\{^1\text{H}\}$  NMR spectrum of **1**, which shows a sharp signal for the bridgehead Si atom, that is split to a doublet of triplets (Fig. 1c) due to coupling with the two chemically different types of  $^{31}\text{P}$  nuclei in the integral ratio 1 : 2 ( $^2J(\text{Si}, \text{P}_A) = 14.7 \text{ Hz}$ ,  $^2J(\text{Si}, \text{P}_B) = 8.2 \text{ Hz}$ ). A positional exchange of the pendant and the Nb-bonded arms of the tmps ligand in **1** was not observed in solution at 298 K.

Complex **1** was found to be a very suitable nucleophile for the formation of  $\text{Nb}\equiv\text{E}$  triple bonds ( $\text{E} = \text{Si}-\text{Sn}$ ). Thus addition of a freshly prepared, orange-colored solution of a mixture of the 1,2-dibromodisilene *E*-Tbb(Br)Si=Si(Br)Tbb<sup>16</sup> and 4-dimethylamino pyridine (4-DMAP) (molar ratio 1 : 4), to a solution of one equiv. of **1** in toluene at ambient temperature was accompanied by an immediate color change to red-brown, and precipitation of a white solid  $(\text{NMe}_4)\text{Br}$ . IR monitoring revealed a complete and selective conversion to the silyldiynide complex  $[(\kappa^3\text{-tmps})(\text{CO})_2\text{Nb}\equiv\text{Si}-\text{Tbb}]$  (**2-Si**, Scheme 4). After work-up, complex **2-Si** was isolated in 59% yield as a red-brown, extremely air-sensitive, microcrystalline solid, which





Fig. 1 (a) FT-IR spectrum of **1** in THF in the range of 2000–1600  $\text{cm}^{-1}$ . (b)  $^{31}\text{P}\{^1\text{H}\}$  NMR spectrum of **1** in benzene- $d_6$ ; an enlarged excerpt with the broad signal at  $\delta = -11.6$  ppm is shown in the inset. (c)  $^{29}\text{Si}\{^1\text{H}\}$  NMR spectrum of **1** in THF- $d_6$ ; an enlarged excerpt with the signal at  $\delta = -0.44$  ppm is depicted in the inset.



Scheme 4 Synthesis of the niobium silyldiynide complex **2-Si** and the germyldiynide complex **3-Ge**.



decolorizes immediately upon exposure to air. Compound **2-Si** is remarkably thermostable, and decomposes to a dark brown mass at 258 °C. It is moderately soluble in *n*-pentane, but readily soluble in benzene, toluene and THF.

Similarly, treatment of complex **1** with the *m*-terphenylgermanium(II) chloride Ge(Ar<sup>Mes</sup>)Cl in toluene at -40 °C followed by warming to room temperature afforded rapidly and selectively the germylydine complex [(κ<sup>3</sup>-tmps)(CO)<sub>2</sub>Nb≡Ge-Ar<sup>Mes</sup>] (**3-Ge**) (Scheme 4). Compound **3-Ge** was isolated as a deep-magenta, very air-sensitive, thermally stable powder (dec. at 284 °C), that is moderately soluble in benzene and toluene, and well soluble in THF. No evidence for the formation of the putative metallogermylene intermediate [(κ<sup>3</sup>-tmps)(CO)<sub>3</sub>Nb-GeAr<sup>Mes</sup>] could be obtained during IR monitoring of the reaction of **1** with Ge(Ar<sup>Mes</sup>)Cl in toluene, the reaction starting at -35 °C and proceeding rapidly with CO evolution below 0 °C.

In comparison, reaction of the analogous *m*-terphenyltin(II) chloride Sn(Ar<sup>Mes</sup>)Cl with **1** in toluene afforded after stirring at ambient temperature the brick-red metallostanlylene [(κ<sup>3</sup>-tmps)(CO)<sub>3</sub>Nb-SnAr<sup>Mes</sup>] (**4-Sn**) with a small amount of the stannylydine complex [(κ<sup>3</sup>-tmps)(CO)<sub>2</sub>Nb≡SnAr<sup>Mes</sup>] (**3-Sn**) (Scheme 5). Prolonged heating at 80 °C and periodic evacuation of the reaction tube was necessary to remove the released CO and to convert **4-Sn** almost quantitatively into the stannylydine complex **3-Sn**, which after work-up was isolated as a dark violet, very air-sensitive powder in 70% yield. Complex **3-Sn** is as **3-Ge** thermally stable and decomposes upon heating at 266 °C. However, unlike **3-Ge**, complex **3-Sn** was found to be extremely light sensitive. Thus exposure of the deep-violet solutions of **3-Sn** to fluorescent, ambient light or sun light lead to deposition of a tin mirror and formation of tmps and 1,3-dimesitylbenzene as evidenced by <sup>1</sup>H NMR spectroscopy. Therefore, all operations during the synthesis, isolation and characterization of **3-Sn** had to be carried out under exclusion of light.

Decarbonylation of **4-Sn** to afford **3-Sn** is a remarkable, new type of reaction in the chemistry of metallostanlylenes. In fact previous attempts to transform the metallostanlylenes [Cp(CO)<sub>3</sub>M-SnR] (M = Cr, Mo, W; R = Ar<sup>Mes</sup>, Ar<sup>Trip</sup>; Ar<sup>Trip</sup> = C<sub>6</sub>H<sub>3</sub>-2,6-Trip<sub>2</sub>, Trip = C<sub>6</sub>H<sub>2</sub>-2,4,6-iPr<sub>3</sub>),<sup>17</sup> [Cp(CO)<sub>2</sub>Fe-SnR] (R = Ar<sup>Dipp</sup>, Ar<sup>Trip</sup>; Ar<sup>Dipp</sup> = C<sub>6</sub>H<sub>3</sub>-2,6-Dipp<sub>2</sub>, Dipp = C<sub>6</sub>H<sub>3</sub>-2,6-iPr<sub>2</sub>)<sup>18</sup> or [Cp\*(CO)<sub>3</sub>W-Sn(IDipp)]<sup>+</sup> (IDipp = C[N(Dipp)CH]<sub>2</sub>, Dipp = C<sub>6</sub>H<sub>3</sub>-2,6-iPr<sub>2</sub>)<sup>20</sup> into terminal stannylydine complexes failed. We assume, that the increased steric pressure imposed by the

tripodal ligand at the metal center weakens the Nb-CO bonds in the seven-coordinate complex **4-Sn** and decreases thereby the barrier for a CO dissociation. In addition, formation of a strong Nb≡Sn triple bond resulting from the higher energy and larger radial extension of the d orbitals, which are engaged in the Nb(dπ) → SnR(π\*) back bonding, may be also a driving force for the reaction.

The tetrylydine complexes **2-Si**, **3-Ge** and **3-Sn** were characterized by elemental analyses, IR spectroscopy and <sup>1</sup>H, <sup>13</sup>C{<sup>1</sup>H}, <sup>31</sup>P{<sup>1</sup>H}, <sup>29</sup>Si{<sup>1</sup>H} and <sup>119</sup>Sn{<sup>1</sup>H} NMR spectroscopy. In addition their molecular structures were determined by single-crystal X-ray crystallography (Fig. 2 and 3). All complexes are distorted octahedral and feature a tridentate (κ<sup>3</sup>-bonded) tmps ligand, which spans three facial coordination sites with the P-Nb-P bite angles varying in a small range (85.3–87.9°). A view along the Si...Nb vector reveals that the CH<sub>2</sub> groups connecting the bridgehead Si atom with the P donors are twisted out creating a local C<sub>3</sub> symmetric, right or left-handed conformation, which reduces the bite of the chelating triphosphane ligand and optimizes the bonding with the niobium center (Fig. 3b). In solution, however, a rapid interchange of the two conformational enantiomers occurs according to NMR spectroscopy leading to averaged C<sub>s</sub> symmetric structures.

The tetrylydine complexes **2-Si**, **3-Ge** and **3-Sn** feature the shortest Nb-Si, Nb-Ge and Nb-Sn bonds reported to date. In practice, the Nb-Si bond of **2-Si** (232.7(2) pm) is ca. 28 pm shorter than the Nb-Si single bonds of silyl complexes (*d*(Nb-Si)<sub>mean</sub> of 28 structurally characterized complexes = 261.3 pm),<sup>19</sup> and the Nb-Ge bond of **3-Ge** (235.79(4) pm) ca. 31 pm shorter than a Nb-Ge single bond (*d*(Nb-Ge)<sub>mean</sub> = 267.3 pm).<sup>20</sup> Similarly, the Nb-Sn bond of **3-Sn** (253.3(1) pm) is ca. 30 pm shorter than a Nb-Sn single bond (*d*(Nb-Sn)<sub>mean</sub> 282.9 pm).<sup>21</sup>





Fig. 2 DIAMOND plot of the molecular structure of the silylydine complex **2-Si** in the solid state. Thermal ellipsoids were set at 30% electronic probability at 100 K. Hydrogen atoms and the methyl groups of the C<sup>2,6</sup>-CH(SiMe<sub>3</sub>)<sub>2</sub> substituents were omitted for clarity. Selected bond lengths [pm] and angles [°]: Nb-Si1 232.7(2), Nb-P1 259.9(2), Nb-P2 258.4(2), Nb-P3 259.3(2), Nb-C35 206.8(9), Nb-C36 206.3(7), Si1-C1 189.0(7), C35-O1 117.6(8), C36-O2 117.9(7); Nb-Si1-C1 159.2(2), P1-Nb-P2 85.61(7), P1-Nb-P3 85.31(6), P2-Nb-P3 87.92(6), C35-Nb-C36 93.3(3).



Scheme 5 Synthesis of the niobium stannylydine complex **3-Sn** via the niobiastannylene **4-Sn**.






Fig. 3 (a) DIAMOND plot of the molecular structure of the stannylydne complex **3-Sn** (toluene) in the solid state. Thermal ellipsoids were set at 30% electronic probability at 100 K. Hydrogen atoms and methyl groups of the  $\text{Ar}^{\text{Mes}}$  substituent were omitted for clarity. Selected bond lengths [pm] and angles [ $^\circ$ ] of **3-Sn** (toluene) (bond lengths and angles for **3-Ge** (THF) are given in brackets): Nb–Sn 253.3(1) [235.79(4)], Nb–P1 260.6(4) [259.5(1)], Nb–P2 255.1(3) [258.0(1)], Nb–P3 258.6(4) [261.2(1)], Nb–C35 205.7(14) [206.0(5)], Nb–C36 207.1(16) [206.5(5)], Sn–C1 214.2(1) [196.3(4)], C35–O1 116.8(18) [116.8(6)], C36–O2 115.4(18) [115.5(6)]; Nb–Sn–C1 160.9(3) [164.0(1)], P1–Nb–P2 87.7(1) [86.30(4)], P1–Nb–P3 85.9(1) [86.08(4)], P2–Nb–P3 87.4(1) [87.82(4)], C35–Nb–C36 90.9(5) [92.5(2)]. (b) Top view of **3-Sn** along the Si...Nb vector illustrating the  $C_3$ -symmetric twist of the tmps ligand.

Notably, a comparison of the Nb–E triple bond lengths of **2-Si**, **3-Ge** and **3-Sn** with those of related molybdenum tetrylydne complexes (e.g.  $d(\text{Mo}\equiv\text{Si})$  in  $[\text{Cp}(\text{CO})_2\text{Mo}\equiv\text{Si}-\text{Ar}^{\text{Trip}}] = 222.41(7)$  pm;<sup>1a</sup>  $d(\text{Mo}\equiv\text{Ge})$  in  $[\text{Cp}(\text{CO})_2\text{Mo}\equiv\text{Ge}-\text{R}]$  (R =  $\text{C}(\text{SiMe}_3)_3$ ,  $\text{Ar}^{\text{Mes}}$ ,  $\text{Ar}^{\text{Trip}} = 227$ – $228$  pm;<sup>2a,2b,2i,2m</sup>  $d(\text{Mo}\equiv\text{Sn})$  in  $\text{trans}-[\text{X}(\text{PMe}_4)\text{Mo}\equiv\text{Sn}-\text{Ar}^{\text{Mes}}]$  (X = Cl, Br, I) =  $248$ – $249$  pm)<sup>22</sup>) reveals that the differences in the M $\equiv$ E triple bond lengths (E = Si: 10 pm; E = Ge: 8–9 pm; E = Sn: 5–6 pm) compare reasonably well with the difference (7 pm) of the metallic radii of the two elements ( $r_{\text{Nb}} = 147$  pm,  $r_{\text{Mo}} = 140$  pm; radii for a coordination number of 12).<sup>23</sup> A series of additive triple bond radii for most elements of the periodic table have been predicted by P. Pyykkö *et al.*<sup>24</sup> The experimental Nb–E triple bond lengths **2-Si**, **3-Ge** and **3-Sn**, are however, longer than the sum of the theoretically predicted triple bond radii ( $d(\text{Nb}\equiv\text{E})_{\text{calc}} = \text{Si}: 218$  pm, Ge: 230 pm, Sn: 248 pm).

In all complexes the tetrylydne ligand is slightly bent at the tetrel center as evidenced by the bonding angle Nb–E–C1 (**2-Si**:  $159.2(2)^\circ$ , **3-Ge**:  $164.0(1)^\circ$ , **3-Sn**:  $160.9(3)^\circ$ ). Bending occurs in all cases towards the CO ligands. It is presently unclear, whether this phenomenon, which is also observed in a series of group 6 metal dicarbonyl ylydne complexes, is of steric or electronic origin or both. No clear evidence for steric congestion is at least provided by the molecular structures of **2-Si**, **3-Ge** and **3-Sn**. For example, the closest van der Waals contacts were found in **2-Si** between the methyl groups of the tmps ligand and the  $\text{SiMe}_3$  methyl groups of the Tbb substituent ( $d(\text{H}\cdots\text{H}) = 244$  pm). These contacts are longer than twice the van der Waals radius of hydrogen ( $r_{\text{vdW}}(\text{H}) = 110$  pm).<sup>25</sup> It should be also taken into consideration, that deviation of the M $\equiv$ E–R atom sequence from linearity does not require a lot of energy, indicating that subtle electronic effects may cause such a bending.<sup>26</sup>

Further structural information was obtained from the IR and NMR spectra of the tetrylydne complexes. The IR spectra of

**2-Si**, **3-Ge** and **3-Sn** display two  $\nu(\text{CO})$  bands of almost equal intensity, which are typical for *cis*-dicarbonyl complexes and can be assigned to the in-phase ( $A'$  symmetric) and out-of-phase ( $A''$  symmetric) CO stretching modes assuming local  $C_s$  symmetry of the  $\text{M}(\text{CO})_2$  fragment (Fig. 4a). The  $\nu(\text{CO})$  bands of **3-Sn** appear at lower frequencies ( $1851$  and  $1791$   $\text{cm}^{-1}$  in toluene) than those of **3-Ge** ( $1868$  and  $1805$   $\text{cm}^{-1}$  in toluene), which suggests that the stannylydne ligand  $\text{SnAr}^{\text{Mes}}$  has a higher  $\sigma$ -donor/ $\pi$ -acceptor ratio than the germylydne ligand  $\text{GeAr}^{\text{Mes}}$ . Notably, the  $\nu(\text{CO})$  bands of **2-Si** appear also at lower wavenumbers ( $1855$  and  $1790$   $\text{cm}^{-1}$  in toluene) than those of **3-Ge**. This shift can be




Fig. 4 (a) IR  $\nu(\text{CO})$  absorption bands of the germylydne complex **3-Ge** in toluene. (b)  $^{31}\text{P}\{^1\text{H}\}$  NMR signal of the germylydne complex **3-Ge** in  $\text{THF}-d_8$  at 283 K. (c)  $^{31}\text{P}\{^1\text{H}\}$  NMR signals of the germylydne complex **3-Ge** in  $\text{THF}-d_8$  at 193 K.



rationalized with the stronger +I effect of the Tbb substituent, leading to a higher  $\sigma$ -donor/ $\pi$ -acceptor ratio of the silyldiylidene ligand SiTbb than that of the germyldiylidene ligand GeAr<sup>Mes</sup>. The low-frequency position of the  $\nu(\text{CO})$  bands of **2-Si**, **3-Ge** and **3-Sn** suggests the presence of an electron-rich Nb center that is engaged in strong Nb( $d\pi$ )  $\rightarrow$  CO( $\pi^*$ ) backbonding. Additional evidence for a strong Nb( $d\pi$ )  $\rightarrow$  CO( $\pi^*$ ) backbonding is provided by the  $^{13}\text{C}\{^1\text{H}\}$  NMR spectra, which all display a broad CO signal at even lower field ( $\delta_{\text{CO}} = 238.7$  ppm (**2-Si**), 239.2 ppm (**3-Ge**), 238.9 ppm (**3-Sn**)) than that of **1** ( $\delta_{\text{CO}} = 226.5$  ppm).<sup>27</sup> The number and relative intensity of the NMR signals indicate an averaged  $C_s$  symmetry of the tetryldiylidene complexes in solution and a rapid rotation of the tetrel-bonded aryl group about the E-C<sub>aryl</sub> bond. The signals of all nuclei directly attached to the quadrupolar  $^{93}\text{Nb}$  nucleus are significantly broadened due to fast relaxation (*vide supra*). For example, the  $^{29}\text{Si}\{^1\text{H}\}$  NMR spectrum of **2-Si** displays at 298 K a very broad signal ( $\Delta\nu_{1/2} = 130$  Hz) for the Nb $\equiv$ Si nucleus at  $\delta = 267.8$  ppm, for which the  $^2J(^{29}\text{Si}, ^{31}\text{P})$  coupling could not be resolved. In comparison, the remote positioned bridgehead Si atom of the tmps ligand and the SiMe<sub>3</sub> groups of the Tbb substituent give rise to sharp signals at  $\delta = -0.7$  ppm and +1.5 ppm, respectively, with the first of these signals being split into a quartet due to coupling to the three  $^{31}\text{P}$  nuclei ( $^2J(^{29}\text{Si}, ^{31}\text{P}) = 9.7$  Hz) (Fig. S16 and S17 (ESI<sup>†</sup>)). Similarly, the  $^{31}\text{P}\{^1\text{H}\}$  NMR spectra of **2-Si** and **3-Ge** show only one broad signal at  $\delta = -13.0$  ppm ( $\Delta\nu_{1/2} \approx 182$  Hz at 298 K) and  $-10.7$  ppm ( $\Delta\nu_{1/2} \approx 187$  Hz at 283 K), respectively, instead of two  $^{31}\text{P}$  NMR signals expected for an AX<sub>2</sub> spin system (Fig. 4b). Broadness of the signals can be influenced by the temperature given the well known relationship between the quadrupole-coupled nuclear relaxation time and the temperature dependent molecular correlation time.<sup>28</sup> In fact, lowering of the temperature lead to a “decoupling” of the Nb nucleus and allowed to resolve the two  $^{31}\text{P}$  NMR signals and their  $^2J(\text{P}, \text{P})$  coupling of 20.9 Hz as illustrated by the  $^{31}\text{P}$  NMR spectrum of **3-Ge** at 193 K (Fig. 4c). Taking advantage of the same effect, also the  $^{119}\text{Sn}$  resonance of **3-Sn**, that was not observable in THF-*d*<sub>8</sub> at room temperature, could be detected at 243 K as a very broad signal ( $\Delta\nu_{1/2} \approx 1297$  Hz) at  $\delta = 829.7$  ppm (Fig. S36 (ESI<sup>†</sup>)).


First studies reveal a marked difference in the reactivity of the niobium germyldiylidene complex **3-Ge** and the related molybdenum germyldiylidene complexes [Cp(CO)<sub>2</sub>Mo $\equiv$ Ge-R] (R = C(SiMe<sub>3</sub>)<sub>3</sub>, Ar<sup>Mes</sup>, Ar<sup>Triip</sup>). Thus treatment of [Cp(CO)<sub>2</sub>Mo $\equiv$ Ge-R] with H<sub>2</sub>O or MeOH (one equiv.) in diethyl ether at 0 °C followed by warming to ambient temperature afforded within one hour selectively the brown hydroxy/methoxygermylidene complexes [Cp(CO)<sub>2</sub>(H)Mo=Ge(O'R)] (R' = H, Me), which were fully characterized.<sup>2m</sup> In contrast, no reaction of **3-Ge** with H<sub>2</sub>O (one equiv.) was observed in THF even at 60 °C. The inertness of **3-Ge** can be rationalized with the stronger metal-germyldiylidene Nb( $d\pi$ )  $\rightarrow$  GeR( $\pi^*$ ) back bonding, which reduces the electrophilicity of the Ge center in **3-Ge**, and increases in combination with the steric protection of the metal center by the tridentate tmps ligand the activation barrier for the H<sub>2</sub>O addition at the Nb $\equiv$ Ge bond. In fact, a large excess of water (925 equiv.) and prolonged heating (3 h) was necessary to effectuate a full conversion of **3-Ge** accompanied by a color change of the reaction solution

from magenta to orange. IR monitoring of the reaction did not provide any evidence for the formation of the anticipated H<sub>2</sub>O addition products. Instead, a continuous decrease in intensity of the two  $\nu(\text{CO})$  bands of **3-Ge** was observed suggesting the formation of mainly CO-free products. Benzene extraction of the orange-brown solid obtained after solvent evaporation afforded a benzene soluble, pale-orange part containing mainly the germanediol Ge(Ar<sup>Mes</sup>)H(OH)<sub>2</sub>, as well as a benzene-insoluble brownish part. The unprecedented hydridogermanediol<sup>29</sup> was isolated as a pale yellow solid and characterized by IR and  $^1\text{H}$  NMR spectroscopy. Its IR spectrum displays two  $\nu(\text{OH})$  bands at 3600 and 3398 cm<sup>-1</sup> and a characteristic  $\nu(\text{Ge-H})$  band at 2104 cm<sup>-1</sup>, the latter one appearing at a close position to that of GeBr<sub>2</sub>HMe<sub>3</sub> ( $\nu(\text{Ge-H}) = 2105$  cm<sup>-1</sup>).<sup>30</sup> In the  $^1\text{H}$  NMR spectrum a distinctive doublet signal is observed for the Ge(OH)<sub>2</sub> protons at  $\delta = 0.91$  ppm and a triplet signal for the Ge-H functionality at  $\delta = 5.61$  ppm ( $^2J(\text{H}, \text{H}) = 3.5$  Hz) in the integral ratio of 2 : 1. Notably, the Ge-OH protons of the germanetriol Ge(Ar<sup>Triip</sup>)(OH)<sub>3</sub> have a similar chemical shift ( $\delta = 0.77$  ppm in CDCl<sub>3</sub>).<sup>29k</sup>

Attempts were also undertaken to access cationic tetryldiylidene complexes. For this purpose, [CpNb(CO)<sub>4</sub>]<sup>31</sup> was prepared using a slightly modified procedure<sup>32</sup> and irradiated in THF with a high-power blue light LED ( $\lambda = 465$  nm) in the presence of one equivalent of Ge(Ar<sup>Mes</sup>)Cl. IR monitoring of the reaction revealed a quite selective decarbonylation leading to the chlorogermlylidene complex **5-Ge**, which after work-up was isolated as red-orange, air-sensitive crystals in 25% yield (Scheme 6). Remarkably attempts to abstract the chloride from **5-Ge** and to form the putative germyldiylidene complex cation [Cp(CO)<sub>3</sub>Nb $\equiv$ GeAr<sup>Mes</sup>]<sup>+</sup> were not successful so far. For example, no reaction of **5-Ge** with Na[B(Ar<sup>F</sup>)<sub>4</sub>] (Ar<sup>F</sup> = C<sub>6</sub>H<sub>3</sub>-3,5-(CF<sub>3</sub>)<sub>2</sub>) was observed in C<sub>6</sub>H<sub>5</sub>F at room temperature.

Complex **5-Ge** is the first niobium germylidene complex to be reported. Its solid-state molecular structure was determined by single-crystal X-ray crystallography (Fig. 5). The four-legged piano stool complex is  $C_s$  symmetric and features a trigonal-planar coordinated Ge-atom (sum of angles at the Ge atom = 360.0°). The symmetry plane passes through the atoms Nb, Ge, C1 and Cl, and bisects the CpNb(CO)<sub>3</sub> fragment and the central ring of the *m*-terphenyl substituent.

The germylidene ligand adopts an upright conformation, with the Ar<sup>Mes</sup> substituent pointing towards the cyclopentadienyl ring. The Nb-Ge distance (251.78(6) pm) of **5-Ge** lies in-between that found for the Nb-Ge triple bond of **3-Ge** (235.79(4) pm) (*vide supra*) and that of a Nb-Ge single bond



Scheme 6 Synthesis of the niobium chlorogermlylidene complex **5-Ge**.






Fig. 5 DIAMOND plot of the molecular structure of **5-Ge** in the solid state. Thermal ellipsoids were set at 30% electronic probability at 100 K, and hydrogen atoms were omitted for clarity. Selected bond lengths [pm] and angles [°]: Nb–Ge 251.78(6), Ge–C1 196.2(4), Ge–Cl 219.1(1), Nb–C17 207.4(3), Nb–C18 206.1(4), C17–O1 115.2(4), C18–O2 114.5(5); Nb–Ge–C1 141.4(1), Nb–Ge–Cl 118.75(4), Cl–Ge–C1 99.8(1).

( $d(\text{Nb-Ge}) = 267.3 \text{ pm}$ )<sup>20</sup> indicating the presence of a Nb–Ge double bond in **5-Ge**. The angles at the Ge atom differ markedly with the Nb–Ge–C<sub>aryl</sub> angle (141.4(1)°) being much larger than the C<sub>aryl</sub>–Ge–Cl angle (99.8(1)°). This distortion can be attributed to the large steric demand of the Ar<sup>Mes</sup> substituent and the low tendency of germanium for isovalent hybridization.<sup>1a,1b,2f,3d</sup> The Ge–Cl bond of **5-Ge** (219.1(1) pm) compares well with that of Ge(Ar<sup>TRIP</sup>)Cl (220.3(2) pm),<sup>33</sup> but is considerably shorter than those of chlorogermylidene complexes containing electron-rich metal centers, such as [(dmpe)<sub>2</sub>Fe=Ge(Ar<sup>Mes</sup>)Cl] ( $d(\text{Ge-Cl}) = 232.2(1) \text{ pm}$ ),<sup>5</sup> [(PMe<sub>3</sub>)<sub>3</sub>Ni=Ge(Ar<sup>Mes</sup>)Cl] ( $d(\text{Ge-Cl}) = 230.03(8) \text{ pm}$ )<sup>6</sup> or [(PMe<sub>3</sub>)<sub>3</sub>Pd=Ge(Ar<sup>Mes</sup>)Cl] ( $d(\text{Ge-Cl}) = 227.3(1) \text{ pm}$ ),<sup>6</sup> in which a strong M(dπ) → Ge(pπ) back bonding is presumed to cause a strong polarization of Ge–Cl bond leading to a facile chloride abstraction by Lewis acids. The reduced polarization of the Ge–Cl bond of **5-Ge** provides a rationale for its inertness towards mild chloride abstraction reagents.

The solution IR and NMR spectra of **5-Ge** are fully consistent with its solid-state molecular structure. Thus, the IR spectrum of **5-Ge** in THF displays three intense ν(CO) absorption bands at 1980, 1910 and 1899 cm<sup>-1</sup>, as expected for a Nb(CO)<sub>3</sub> fragment with local C<sub>s</sub> symmetry, which are assigned to the A' (all three CO modes in phase), A' (two CO<sub>lat</sub> modes in phase; CO<sub>diag</sub> mode out-of-phase) and A'' symmetric (two CO<sub>lat</sub> modes out-of-phase) CO stretching modes, respectively. The ν(CO) absorption bands of **5-Ge** are high-frequency shifted compared to those of [CpNb(CO)<sub>3</sub>THF] (ν(CO) in THF = 1961, 1840 cm<sup>-1</sup>)<sup>31</sup> or [CpNb(CO)<sub>3</sub>PET<sub>3</sub>] (ν(CO) in THF = 1953, 1850 cm<sup>-1</sup>),<sup>28b</sup> but appear at roughly the same position as those of [CpNb(CO)<sub>3</sub>N<sub>2</sub>] (ν(CO) in *n*-heptane = 1991, 1905 cm<sup>-1</sup>)<sup>34</sup> suggesting a similar σ-donor/π-acceptor ratio of the germylidene GeAr<sup>Mes</sup>Cl and the N<sub>2</sub> ligand. The <sup>1</sup>H and <sup>13</sup>C{<sup>1</sup>H} NMR spectra also confirm the C<sub>s</sub> symmetry of **5-Ge** in solution. Rotation of the *m*-terphenyl substituent about the Ge–C<sub>aryl</sub> bond occurs fast on the NMR




Fig. 6 Single-scan cyclic voltammograms of the reversible one-electron oxidation of **3-Ge** at different scan rates in THF at –11 °C (supporting electrolyte: [NBu<sub>4</sub>][PF<sub>6</sub>] (0.1 M); reference electrode: 0.004 M [Fe(C<sub>5</sub>Me<sub>5</sub>)<sub>2</sub>]<sup>+1/0</sup>/0.1 M [NBu<sub>4</sub>][PF<sub>6</sub>]/THF).

time-scale at ambient temperature leading to an exchange of the two diastereotopic *ortho* (C<sup>2,6</sup>) and *meta* (C<sup>3,5</sup>) positions of the enantiotopic mesityl substituents. Therefore, only one singlet signal is observed in the <sup>1</sup>H NMR spectrum of **5-Ge** for the C<sup>2,6</sup>-bonded methyl groups and C<sup>3,5</sup>-bonded protons of the mesityl substituents, respectively.

### Electrochemical studies

Electrochemical studies of the tetrylidyne complexes **2-Si**, **3-Ge** and **3-Sn** were carried out using cyclic voltammetry to elucidate the redox properties of these compounds. All complexes display a rich electrochemistry involving several electron-transfer steps (see ESI, chapter 3†). Remarkably, both the one-electron reduction and oxidation of the germylidene complex **3-Ge** are electrochemically reversible occurring at a half wave potential ( $E_{1/2}$ ) of –2.612 mV and –405 mV vs. the dmfc<sup>+1/0</sup> redox couple (dmfc = decamethylferrocene), respectively (Fig. 6).<sup>35</sup>

In comparison, the corresponding redox steps of **2-Si** and **3-Sn** are irreversible (ESI, chapter 3†), but one-electron oxidation **2-Si** and **3-Sn** occurs at similar potentials as that of **3-Ge** ( $E_{\text{pa}} + E_{\text{pc}}/2 = -468 \text{ mV}$ , **3-Sn**:  $E_{\text{pa}} + E_{\text{pc}}/2 = -435 \text{ mV}$  (scan rate = 100 mV s<sup>-1</sup>)). Evidence that the redox process at  $E_{1/2} = -405 \text{ mV}$  involves a one electron oxidation of **3-Ge** was provided by chemical means. Thus, no reaction of **3-Ge** with the one-electron reducing agent cobaltocene ( $E_{1/2}$  of CoCp<sub>2</sub> in DME = –740 mV) was observed in fluorobenzene even at 70 °C, whereas an instantaneous oxidation of **3-Ge** occurred upon treatment with one equivalent of [Fe(η<sup>5</sup>-C<sub>5</sub>Me<sub>5</sub>)<sub>2</sub>][B(Ar<sup>F</sup>)<sub>4</sub>] in fluorobenzene solution at –30 °C. Unfortunately, attempts to isolate the putative germylidene complex radical cation [(κ<sup>3</sup>-tmps)(CO)<sub>2</sub>-Nb(GeAr<sup>Mes</sup>)<sup>+</sup>] failed so far.<sup>36</sup> Notably, the redox potential for the one-electron oxidation of **3-Ge** is slightly lower than that of the molybdenum tetrylidyne complexes *trans*-[ClMo(PMe<sub>3</sub>)<sub>4</sub>≡E-Ar<sup>Mes</sup>] (E = Ge:  $E_{1/2}$  in C<sub>6</sub>H<sub>5</sub>F = –340 mV; E = Sn:  $E_{1/2}$  in THF = –350 mV; E = Pb:  $E_{1/2}$  in THF = –358 mV) verifying the presence of an electron-rich Nb center in **3-Ge**.

## Conclusion

The synthesis of the tailor-made carbonyl-niobate (NMe<sub>4</sub>)[Nb(CO)<sub>4</sub>(κ<sup>2</sup>-tmps)] allowed to explore its reactivity towards



a series of organotetrel(II) halides, which lead to the isolation of the first niobium complexes featuring triple bonds with the elements Si, Ge and Sn. Photochemical CO substitution in  $[\text{CpNb}(\text{CO})_4]$  ( $\text{Cp} = \eta^5\text{-C}_5\text{H}_5$ ) by  $\text{Ge}(\text{Ar}^{\text{Mes}})\text{Cl}$  afforded also the novel chlorogermlydene complex  $[\text{Cp}(\text{CO})_3\text{Nb}=\text{Ge}(\text{Ar}^{\text{Mes}})\text{Cl}]$ . The structural, spectroscopic and electrochemical data of the tetrylidyne complexes  $[(\kappa^3\text{-tmps})(\text{CO})_2\text{Nb}\equiv\text{Si-Tbb}]$  (**2-Si**),  $[(\kappa^3\text{-tmps})(\text{CO})_2\text{Nb}\equiv\text{Ge-Ar}^{\text{Mes}}]$  (**3-Ge**) and  $[(\kappa^3\text{-tmps})(\text{CO})_2\text{Nb}\equiv\text{Sn-Ar}^{\text{Mes}}]$  (**3-Sn**) suggest the presence of an electron-rich metal center that is engaged into strong metal ( $d\pi$ )  $\rightarrow$  ER( $\pi^*$ ) and metal ( $d\pi$ )  $\rightarrow$  CO( $\pi^*$ ) back bonding. Remarkably, one-electron oxidation and reduction of the germlydene complex **3-Ge** are electrochemically reversible.

## Acknowledgements

We thank the Jürgen Manchot Stiftung and the University of Bonn for the financial support of this work. We also thank Mrs Charlotte Rödde, Dr Senada Nozinovic, Dipl.-Ing. Karin Prochnicki, Dr Burhanshah Lewall, Mrs Kerstin Kühnel-Lysek and Mrs Anna Martens for their contributions to the single-crystal X-ray diffraction studies, the NMR measurements, the cyclic voltammetric studies and the elemental analyses.

## References

- For experimental studies on compounds with  $\text{M}\equiv\text{Si}$  bonds (silylidyne complexes) see: (a) A. C. Filippou, O. Chernov, K. W. Stumpf and G. Schnakenburg, *Angew. Chem. Int. Ed.*, 2010, **49**, 3296; *Angew. Chem.*, 2010, **122**, 3368; (b) A. C. Filippou, O. Chernov and G. Schnakenburg, *Angew. Chem. Int. Ed.*, 2011, **50**, 1122; *Angew. Chem.*, 2011, **123**, 1154; (c) P. G. Hayes, Z. Xu, C. Beddie, J. M. Keith, M. B. Hall and T. D. Tilley, *J. Am. Chem. Soc.*, 2013, **135**, 11780; (d) A. C. Filippou, B. Baars, O. Chernov, Y. N. Lebedev and G. Schnakenburg, *Angew. Chem. Int. Ed.*, 2014, **53**, 565; *Angew. Chem.*, 2014, **126**, 576; (e) T. Fukuda, T. Yoshimoto, H. Hashimoto and H. Tobita, *Organometallics*, 2016, **35**, 921; (f) T. Yoshimoto, H. Hashimoto, N. Hayakawa, T. Matsuo and H. Tobita, *Organometallics*, 2016, **35**, 3444.
- For experimental studies on germlydene complexes see: (a) R. S. Simons and P. P. Power, *J. Am. Chem. Soc.*, 1996, **118**, 11966; (b) L. Pu, B. Twamley, S. T. Haubrich, M. M. Olmstead, B. V. Mork, R. S. Simons and P. P. Power, *J. Am. Chem. Soc.*, 2000, **122**, 650; (c) A. C. Filippou, A. I. Philippopoulos, P. Portius and D. U. Neumann, *Angew. Chem. Int. Ed.*, 2000, **39**, 2778; *Angew. Chem.*, 2000, **112**, 2881; (d) A. C. Filippou, A. I. Philippopoulos, P. Portius and G. Schnakenburg, *Organometallics*, 2004, **23**, 4503; (e) A. C. Filippou, G. Schnakenburg, A. I. Philippopoulos and N. Weidemann, *Angew. Chem. Int. Ed.*, 2005, **44**, 5979; *Angew. Chem.*, 2005, **117**, 6133; (f) A. C. Filippou, N. Weidemann, A. I. Philippopoulos and G. Schnakenburg, *Angew. Chem. Int. Ed.*, 2006, **45**, 5987; *Angew. Chem.*, 2006, **118**, 6133; (g) A. C. Filippou, A. Barandov, G. Schnakenburg, B. Lewall, M. van Gastel and A. Marchanka, *Angew. Chem.*, 2012, **124**, 813; *Angew. Chem. Int. Ed.*, 2012, **51**, 789; (h) H. Hashimoto, T. Fukuda, H. Tobita, M. Ray and S. Sakaki, *Angew. Chem., Int. Ed.*, 2012, **51**, 2930; *Angew. Chem.*, 2012, **124**, 2984; (i) A. C. Filippou, K. W. Stumpf, O. Chernov and G. Schnakenburg, *Organometallics*, 2012, **31**, 748; (j) J. Hicks, T. J. Hadlington, C. Schenk, J. Li and C. Jones, *Organometallics*, 2013, **32**, 323; (k) T. Fukuda, H. Hashimoto and H. Tobita, *Chem. Commun.*, 2013, **49**, 4232; (l) A. C. Filippou, U. Chakraborty and G. Schnakenburg, *Chem.-Eur. J.*, 2013, **19**, 5676; (m) K. W. Stumpf, Germlydene and Germlydene Complexes of Molybdenum, Dissertation, University of Bonn, Verlag Dr Hut, 2014, ISBN 978-3-8439-1849-7; (n) Y. Lebedev, U. Das, G. Schnakenburg and A. C. Filippou, *Organometallics*, 2017, **36**, 1530.
- For experimental studies on stannylidyne complexes see: (a) A. C. Filippou, P. Portius, A. I. Philippopoulos and H. Rohde, *Angew. Chem. Int. Ed.*, 2003, **42**, 445; *Angew. Chem.*, 2003, **115**, 461; (b) A. C. Filippou, A. I. Philippopoulos and G. Schnakenburg, *Organometallics*, 2003, **22**, 3339; (c) H. Rohde, M. Menzel, F. Renz and A. C. Filippou, *Hyperfine Interact.*, 2008, **185**, 129; (d) A. C. Filippou, P. Ghana, U. Chakraborty and G. Schnakenburg, *J. Am. Chem. Soc.*, 2013, **135**, 11525; (e) C. Lindlahr, Novel Tetrylidyne Complexes of Group 6 Metals with Trialkylphosphane Ligands, Dissertation, University of Bonn, Verlag Dr Hut, 2015, ISBN 978-3-8439-2126-8.
- For experimental studies on plumbylidyne complexes see: (a) A. C. Filippou, H. Rohde and G. Schnakenburg, *Angew. Chem. Int. Ed.*, 2004, **43**, 2243; *Angew. Chem.*, 2004, **116**, 2293; (b) A. C. Filippou, N. Weidemann, G. Schnakenburg, H. Rohde and A. I. Philippopoulos, *Angew. Chem. Int. Ed.*, 2004, **43**, 6512; *Angew. Chem.*, 2004, **116**, 6674; (c) A. C. Filippou, N. Weidemann and G. Schnakenburg, *Angew. Chem. Int. Ed.*, 2008, **47**, 5799; *Angew. Chem.*, 2008, **120**, 5883; (d) U. Chakraborty, Multiple Bonds between Group 7 Transition Metals and Heavier Tetrel Elements (Ge-Pb), Dissertation, University of Bonn, Dr Hut Verlag, 2013, ISBN 978-3-8439-1248-8.
- B. Blom, Reactivity of Ylenes at Late Transition Metal Centers, Dissertation, University of Bonn, Cuvillier Verlag, 2011, ISBN 978-3-86955-907-0.
- I. Papazoglou, Unprecedented Tetrylidyne Complexes of Group 6 and Group 10 Metals, Dissertation, University of Bonn, Dr Hut Verlag, 2017, ISBN 978-3-8439-3155-7.
- (a) A. V. Lalov, M. P. Egorov, O. M. Nefedov, V. K. Cherkasov, N. L. Ermolaev and A. V. Piskunov, *Russ. Chem. Bull.*, 2005, **54**, 807; (b) A. Shinohara, J. McBee, R. Waterman and T. D. Tilley, *Organometallics*, 2008, **27**, 5717; (c) H. Arp, J. Baumgartner, C. Marschner, P. Zark and T. Müller, *J. Am. Chem. Soc.*, 2012, **134**, 10864.
- (a) A. Davison and J. E. Ellis, *J. Organomet. Chem.*, 1972, **36**, 113; (b) F. Calderazzo, G. Pampaloni and G. Pelizzi, *J. Organomet. Chem.*, 1982, **233**, C41; (c) K. M. Pfahl and J. E. Ellis, *Organometallics*, 1984, **3**, 230; (d) F. Calderazzo, M. Castellani, G. Pampaloni and P. F. Zanazzi, *J. Chem.*





- Soc.*, *Dalton Trans.*, 1985, 1989; (e) E. M. Carnahan and S. J. Lippard, *J. Am. Chem. Soc.*, 1992, **114**, 4166; (f) M. V. Barybin, J. E. Ellis, M. K. Pomije, M. L. Tinkham and G. F. Warnock, *Inorg. Chem.*, 1998, **37**, 6518; (g) P. J. Daff, P. Legzdins and S. J. Rettig, *J. Am. Chem. Soc.*, 1998, **120**, 2688.
- 9 (a) C. G. Dewey, J. E. Ellis, K. L. Fjare, K. M. Pfahl and G. F. P. Warnock, *Organometallics*, 1983, **2**, 388; (b) F. Calderazzo, U. Englert, G. Pampaloni, G. Pelizzi and R. Zamboni, *Inorg. Chem.*, 1983, **22**, 1865; (c) F. Calderazzo, G. Pampaloni and J. E. Ellis, *Inorganic Syntheses*, John Wiley & Sons, Hoboken, NJ, USA, 1990, p. 192.
- 10 R. S. Simons, L. Pu, M. M. Olmstead and P. P. Power, *Organometallics*, 1997, **16**, 1920.
- 11 (a) A. Davison and J. E. Ellis, *J. Organomet. Chem.*, 1971, **31**, 239; (b) H.-C. Bechthold and D. Rehder, *J. Organomet. Chem.*, 1982, **233**, 215; (c) K. Bachmann and D. Rehder, *J. Organomet. Chem.*, 1984, **276**, 177; (d) F. Calderazzo, G. Pampaloni, U. Englert and J. Strähle, *J. Organomet. Chem.*, 1990, **383**, 45.
- 12 H. H. Karsch and A. Appelt, *Z. Naturforsch., B: Anorg. Chem., Org. Chem.*, 1983, **38**, 1399.
- 13 M. S. Wrighton, D. I. Handeli and D. L. Morse, *Inorg. Chem.*, 1976, **15**, 434.
- 14 K. Nakamoto, *Infrared and Raman Spectra of Inorganic and Coordination Compounds*, Wiley, Hoboken, 2009.
- 15 (a) P. Raghavan, *At. Data Nucl. Data Tables*, 1989, **42**, 189; (b) J. H. Nelson, *Nuclear Magnetic Resonance Spectroscopy*, Pearson Education, 2003.
- 16 T. Agou, N. Hayakawa, T. Sasamori, T. Matsuo, D. Hashizume and N. Tokitoh, *Chem.-Eur. J.*, 2014, **20**, 9246.
- 17 B. E. Eichler, A. D. Phillips, S. T. Haubrich, B. V. Mork and P. P. Power, *Organometallics*, 2002, **21**, 5622.
- 18 H. Lei, J.-D. Guo, J. C. Fettinger, S. Nagase and P. P. Power, *Organometallics*, 2011, **30**, 6316.
- 19 According to a CSD survey (13.06.17) the Nb–Si bond lengths of silyl complexes range from 254.1 to 267.9 pm leading to a mean value of 261.3 pm and a median of 261.2 pm for the Nb–Si single bond length: (a) J. Arnold, T. D. Tilley, A. L. Rheingold and S. J. Geib, *Organometallics*, 1987, **6**, 473; (b) A. Antiñolo, F. Carrillo, M. Fajardo, A. Otero, M. Lanfranchi and M. A. Pellinghelli, *Organometallics*, 1995, **14**, 1518; (c) G. I. Nikonov, L. G. Kuzmina, D. A. Lemenovskii and V. V. Kotov, *J. Am. Chem. Soc.*, 1995, **117**, 10133; (d) G. I. Nikonov, L. G. Kuzmina, S. F. Vyboishchikov, D. A. Lemenovskii and J. A. K. Howard, *Chem.-Eur. J.*, 1999, **5**, 2947; (e) V. I. Bakhmutov, J. A. K. Howard, D. A. Keen, L. G. Kuzmina, M. A. Leech, G. I. Nikonov, E. V. Vorontsov and C. C. Wilson, *J. Chem. Soc., Dalton Trans.*, 2000, 1631; (f) G. I. Nikonov, P. Mountford, J. C. Green, P. A. Cooke, M. A. Leech, A. J. Blake, J. A. K. Howard and D. A. Lemenovskii, *Eur. J. Inorg. Chem.*, 2000, 1917; (g) G. I. Nikonov, L. G. Kuzmina and J. A. K. Howard, *J. Chem. Soc., Dalton Trans.*, 2002, 3037; (h) G. I. Nikonov, P. Mountford and S. R. Dubberley, *Inorg. Chem.*, 2003, **42**, 258; (i) K. Y. Dorogov, E. Dumont, N.-N. Ho, A. V. Churakov, L. G. Kuzmina, J.-M. Poblet, A. J. Schultz, J. A. K. Howard, R. Bau, A. Lledos and G. I. Nikonov, *Organometallics*, 2004, **23**, 2845; (j) K. Y. Dorogov, A. V. Churakov, L. G. Kuzmina, J. A. K. Howard and G. I. Nikonov, *Eur. J. Inorg. Chem.*, 2004, 771; (k) K. Y. Dorogov, M. Yousufuddin, N.-N. Ho, A. V. Churakov, L. G. Kuzmina, A. J. Schultz, S. A. Mason, J. A. K. Howard, D. A. Lemenovskii, R. Bau and G. I. Nikonov, *Inorg. Chem.*, 2007, **46**, 147; (l) S. K. Ignatov, N. H. Rees, A. A. Merkoulov, S. R. Dubberley, A. G. Razuvaev, P. Mountford and G. I. Nikonov, *Chem.-Eur. J.*, 2008, **14**, 296.
- 20 A CSD survey (13.06.17) revealed that only two niobium germyl complexes have been structurally characterized. The Nb–Ge distances range from 260.8 to 271.0 pm, leading to a mean value of 267.3 pm and a median of 270.2 pm for the Nb–Ge single bond length: (a) A. Antiñolo, F. Carrillo-Hermosilla, A. Castel, M. Fajardo, J. Fernández-Baeza, M. Lanfranchi, A. Otero, M. A. Pellinghelli, G. Rima, J. Satgé and E. Villaseñor, *Organometallics*, 1998, **17**, 1523; (b) G. I. Nikonov, A. V. Churakov and M. Y. Antipin, *Organometallics*, 2003, **22**, 2178.
- 21 A CSD survey (13.06.17) revealed that 9 niobium stannyl complexes have been structurally characterized. The Nb–Sn bond lengths range from 276.3 to 286.2 pm, giving a mean value of 282.9 pm and a median of 283.0 pm for the Nb–Sn single bond length. The respective complexes can be found in [ref. 20a] and in the following references: (a) F. Näumann, J. Kopf and D. Rehder, *J. Organomet. Chem.*, 1984, **267**, 249; (b) Y. V. Skripkin, O. G. Volkov, A. A. Pasynskii, A. S. Antsyshkina, L. M. Dikareva, V. N. Ostrikova, M. A. Porai-Koshits, S. L. Davydova and S. G. Sakharov, *J. Organomet. Chem.*, 1984, **263**, 345; (c) Y. V. Skripkin, O. G. Volkov, A. A. Pasynskii, M. A. Porai-Koshits, A. S. Antsyshkina, L. M. Dikareva and V. N. Ostrikova, *Koord. Khim.*, 1985, **11**, 1136; (d) M. L. H. Green, A. K. Hughes and P. Mountford, *J. Chem. Soc., Dalton Trans.*, 1991, 1407; (e) M. L. H. Green, A. K. Hughes and P. Mountford, *J. Chem. Soc., Dalton Trans.*, 1991, 1699; (f) G. I. Nikonov, L. G. Kuzmina, J. Lorberth and J. A. K. Howard, *Eur. J. Inorg. Chem.*, 1999, 825.
- 22 H. Rohde, *Synthesis and Reactivity Studies of Stannyliidyne Complexes*, Dissertation, University of Bonn, 2007.
- 23 A. F. Holleman, E. Wiberg and N. Wiberg, *Inorganic Chemistry*, Academic Press, San Diego, 2001.
- 24 (a) P. Pykkö, S. Riedel and M. Patzschke, *Chem.-Eur. J.*, 2005, **11**, 3511; (b) P. Pykkö, *J. Phys. Chem. A*, 2015, **119**, 2326.
- 25 M. Mantina, A. C. Chamberlin, R. Valero, C. J. Cramer and D. G. Truhlar, *J. Phys. Chem. A*, 2009, **113**, 5806.
- 26 For example quantum chemical studies on the model compounds *trans*-[Cl(PH<sub>3</sub>)<sub>4</sub>W≡EPh] (E = Si–Pb) at the B3LYP/TZVPP(W,E)/6-311G\*(C,H,P,Cl) level of theory reveal that the energy required for bending the tetryliidyne ligand up to 20° is rather small (max. 20 kJ mol<sup>-1</sup>) and decreases with increasing atomic number of E.



- 27 L. J. Todd and J. R. Wilkinson, *J. Organomet. Chem.*, 1974, **77**, 1.
- 28 (a) P. S. Ireland, C. A. Deckert and T. L. Brown, *J. Magn. Reson.*, 1976, **23**, 485; (b) D. Rehder and H.-C. Bechthold, *Z. Naturforsch., B: Anorg. Chem., Org. Chem.*, 1984, **39**, 323 and references therein.
- 29 Only few four-coordinated germanediols and two germanetriols ( $\text{Ge}(\text{Ar}^{\text{TriP/DiP}})(\text{OH})_3$ ) were reported so far: (a) H. Puff, S. Franken, W. Schuh and W. Schwab, *J. Organomet. Chem.*, 1983, **254**, 33; (b) L. Fajari, J. Carilla, L. Juliá, J. Riera, A. Párraga, M. Coll and X. Solans, *J. Organomet. Chem.*, 1994, **474**, 89; (c) K. Kishikawa, N. Tokitoh and R. Okazaki, *Chem. Lett.*, 1996, **25**, 695; (d) L. Pu, N. J. Hardman and P. P. Power, *Organometallics*, 2001, **20**, 5105; (e) T. Matsumoto, Y. Nakaya and K. Tatsumi, *Organometallics*, 2006, **25**, 4835; (f) Y. Sugiyama, T. Sasamori, Y. Hosoi, Y. Furukawa, N. Takagi, S. Nagase and N. Tokitoh, *J. Am. Chem. Soc.*, 2006, **128**, 1023; (g) E. Bonnefille, S. Mazières, C. Bibal, N. Saffon, H. Gornitzka and C. Couret, *Eur. J. Inorg. Chem.*, 2008, 4242; (h) E. Bonnefille, S. Mazières, N. Saffon and C. Couret, *J. Organomet. Chem.*, 2009, **694**, 2246; (i) M. Saito, M. Nakamura and T. Tajima, *Heterocycles*, 2009, **78**, 657; (j) L. Li, T. Fukawa, T. Matsuo, D. Hashizume, H. Fueno, K. Tanaka and K. Tamao, *Nat. Chem.*, 2012, **4**, 361; (k) L.-C. Pop, N. Kurokawa, H. Ebata, K. Tomizawa, T. Tajima, M. Ikeda, M. Yoshioka, M. Biesemans, R. Willem, M. Minoura and M. Saito, *Can. J. Chem.*, 2014, **92**, 542.
- 30 P. Rivière, M. Rivière-Baudet, A. Castel, J. Satgé and E. A. Lavabre, *Synth. React. Inorg. Met.-Org. Chem.*, 1987, **17**, 539.
- 31 W. A. Herrmann and H. Biersack, *J. Organomet. Chem.*, 1980, **191**, 397 and refs. therein.
- 32 T. E. Bitterwolf, S. Gallagher, J. T. Bays, B. Scallorn, A. L. Rheingold, I. A. Guzei, L. Liable-Sands and J. C. Linehan, *J. Organomet. Chem.*, 1998, **557**, 77.
- 33 L. Pu, M. M. Olmstead, P. P. Power and B. Schiemenz, *Organometallics*, 1998, **17**, 5602.
- 34 G. I. Childs, D. C. Grills, S. Gallagher, T. E. Bitterwolf and M. W. George, *J. Chem. Soc., Dalton Trans.*, 2001, 1711.
- 35 All potentials in the text are given vs. the  $[\text{Fe}(\eta^5\text{-C}_5\text{Me}_5)_2]^{1+/0}$  ( $\text{dmfc}^{1+/0}$ ) redox couple, which was chosen as the reference standard for the CV experiments due to its favorable properties vs. the  $[\text{Fe}(\eta^5\text{-C}_5\text{H}_5)_2]^{1+/0}$  redox couple: (a) I. Noviandri, K. N. Brown, D. S. Fleming, P. T. Gulyas, P. A. Lay, A. F. Masters and L. Phillips, *J. Phys. Chem. B*, 1999, **103**, 6713; (b) J. R. Aranzaes, M.-C. Daniel and D. Astruc, *Can. J. Chem.*, 2006, **84**, 288. For comparison reasons the half-wave potential of the  $[\text{Fe}(\eta^5\text{-C}_5\text{H}_5)_2]^{1+/0}$  redox couple was also determined under the same conditions (THF,  $-11^\circ\text{C}$ ) and found to be +440 mV vs. the  $[\text{Fe}(\eta^5\text{-C}_5\text{Me}_5)_2]^{1+/0}$  redox couple.
- 36 Rapid oxidation of **3-Ge** was evidenced by the immediate disappearance of the green color of the fluorobenzene solution of  $[\text{Fe}(\eta^5\text{-C}_5\text{Me}_5)_2][\text{B}(\text{Ar}^{\text{F}})_4]$  at the dropping site and the change of the color of the reaction solution to brown-red. After evaporation of the solvent and washing of the residue with *n*-pentane to remove  $[\text{Fe}(\eta^5\text{-C}_5\text{Me}_5)_2]$ , a brown oil was obtained, which displayed in the IR spectrum in THF mainly two intense, broad  $\nu(\text{CO})$  bands at 1954 and 1889  $\text{cm}^{-1}$ . The brown oil contained according to  $^{31}\text{P}$  NMR spectroscopy diamagnetic impurities and could not be crystallized.

