

CORRECTION

Correction: Oral anticoagulant re-initiation following intracerebral hemorrhage in non-valvular atrial fibrillation: Global survey of the practices of neurologists, neurosurgeons and thrombosis experts

Yan Xu, Ashkan Shoamanesh, Sam Schulman, Dar Dowlatshahi, Rustam Al-Shahi Salman, Ioana Doina Moldovan, Philip Stephen Wells, Fahad AlKherayf

The fifth author is listed incorrectly in the citation. The correct citation is: Xu Y, Shoamanesh A, Schulman S, Dowlatshahi D, Al-Shahi Salman R, Moldovan ID, et al. (2018) Oral anticoagulant re-initiation following intracerebral hemorrhage in non-valvular atrial fibrillation: Global survey of the practices of neurologists, neurosurgeons and thrombosis experts. PLoS ONE 13(1): e0191137. <https://doi.org/10.1371/journal.pone.0191137>.

An incorrect file was used for Fig 1. Please see the complete, correct Fig 1 here.

The images for Figs 2 and 3 are incorrectly switched. The image that appears as Fig 2 should be Fig 3, and the image that appears as Fig 3 should be Fig 2. The figure captions appear in the correct order.



OPEN ACCESS

Citation: Xu Y, Shoamanesh A, Schulman S, Dowlatshahi D, Al-Shahi Salman R, Moldovan ID, et al. (2018) Correction: Oral anticoagulant re-initiation following intracerebral hemorrhage in non-valvular atrial fibrillation: Global survey of the practices of neurologists, neurosurgeons and thrombosis experts. PLoS ONE 13(5): e0198031. <https://doi.org/10.1371/journal.pone.0198031>

Published: May 24, 2018

Copyright: © 2018 Xu et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.




Fig 1. Overall response from survey participants on timing of OAC re-initiation across 11 clinical scenarios. ICH, intracerebral haemorrhage; DOAC, direct oral anticoagulants; IPH, intraparenchymal haemorrhage; HTN, hypertension; CHADS₂, Congestive heart failure, hypertension, age (≥75), diabetes, stroke/TIA score.

<https://doi.org/10.1371/journal.pone.0198031.g001>




Fig 2. Choice of anticoagulant for re-initiation across thrombosis experts, stroke neurologists and neurosurgeons.

<https://doi.org/10.1371/journal.pone.0198031.g002>




Fig 3. Rates of neuro-imaging utilization for risk stratification among patients with anticoagulant-associated ICH across specialties. Intracranial vessel imaging includes CT angiography, MR angiography or Digital Subtraction Angiography.

<https://doi.org/10.1371/journal.pone.0198031.g003>

Reference

1. Xu Y, Shoamanesh A, Schulman S, Dowlatshahi D, Salman RA-S, Moldovan ID, et al. (2018) Oral anti-coagulant re-initiation following intracerebral hemorrhage in non-valvular atrial fibrillation: Global survey of the practices of neurologists, neurosurgeons and thrombosis experts. PLoS ONE 13(1): e0191137. <https://doi.org/10.1371/journal.pone.0191137>. PMID: 29370183