

## Research Article

# Single-Word Speech Intelligibility in Children and Adults With Down Syndrome

Alyssa Wild,<sup>a</sup> Houri K. Vorperian,<sup>a</sup> Ray D. Kent,<sup>a</sup> Daniel M. Bolt,<sup>b</sup> and Diane Austin<sup>a</sup>

**Purpose:** A single-word identification test was used to study speech production in children and adults with Down syndrome (DS) to determine the developmental pattern of speech intelligibility with an emphasis on vowels.

**Method:** Speech recordings were collected from 62 participants with DS aged 4–40 years and 25 typically developing participants aged 4–7 years. Panels of 5 adult lay listeners transcribed the speech recordings orthographically, and their responses were scored in comparison with the speakers' target words.

**Results:** Speech intelligibility in persons with DS improved with age, especially between the ages of 4 and 16 years.

Whereas consonants contribute to intelligibility, vowels also played an important role in reduced intelligibility with an apparent developmental difference in low versus high vowels, where the vowels /æ/ and /ɑ/ developed at a later age than /i/ and /u/. Interspeaker variability was large, with male individuals being generally less intelligible than female individuals and some adult men having very low intelligibility.

**Conclusion:** Results show age-related patterns in speech intelligibility in persons with DS and identify the contribution of dimensions of vowel production to intelligibility. The methods used clarify the phonetic basis of reduced intelligibility, with implications for assessment and treatment.

Reduced speech intelligibility in persons with Down syndrome (DS) frequently results in communication difficulties and frustration for both speakers and their communication partners (Kent & Vorperian, 2013; Kumin, 1994; Roberts, Price, & Malkin, 2007). It is a challenge to study speech intelligibility comprehensively, because it is affected by many factors related to the speaker, listener, and context (Hustad, 2012; Kent, Miolo, & Bloedel, 1994). In persons with DS, factors likely to influence intelligibility are voice, articulation, resonance, fluency, and/or prosody (Kent & Vorperian, 2013). Because many factors contribute to intelligibility, no single test can capture all aspects of intelligibility in DS. Studies of intelligibility in DS have used a variety of approaches that can be classified into two general categories: global and analytical. Hustad (2012) made a similar distinction using the respective terms “subjective” and “objective.” Global approaches entail overall ratings made by listeners (such as percentage of words understood or

values on an interval scale) serving as a gauge of the severity of the communication problem, whereas analytical approaches involve scoring a unit of analysis, such as a word, phoneme, or phonetic feature, as correct or incorrect and serve to identify the reason for reduced speech intelligibility in speakers with DS. Our use of the term *analytical* compares with the term *diagnostic intelligibility testing*, which “aims to reach beyond mere quantification of degree of overall intelligibility and gain insights into specifically why someone is difficult to understand” (Miller, 2013, p. 604).

## Global Approaches

Reduced speech intelligibility by both familiar and unfamiliar listeners has been documented in several studies. Questionnaires administered to parents of persons with DS make it clear that family members recognize reduced intelligibility as an obstacle to communication and that the problem can be lifelong. Kumin (1994) reported that across their life spans, 95% of persons with DS experience some difficulty being understood. Kumin (2006) also reported that persons with DS aged 1–21 years had a mean speech intelligibility rating from their parents of 4.97 on a 10-point scale (1 being the least and 10 being the most), with boys with DS being less intelligible than girls with DS in the same age group. Toğram (2015) reported similar results

<sup>a</sup>433 Waisman Center, University of Wisconsin-Madison

<sup>b</sup>Department of Educational Psychology, University of Wisconsin-Madison

Correspondence to Houri K. Vorperian:  
vorperian@waisman.wisc.edu

Editor-in-Chief: Krista Wilkinson

Editor: Rebecca McCauley

Received January 6, 2017

Revision received September 29, 2017

Accepted October 11, 2017

[https://doi.org/10.1044/2017\\_AJSLP-17-0002](https://doi.org/10.1044/2017_AJSLP-17-0002)

**Disclosure:** The authors have declared that no competing interests existed at the time of publication.

from parents of children with DS aged 1–19 years. Reduced speech intelligibility persists into adulthood, but there are few published reports that describe the problem in detail or show the general pattern of change with age (Coppens-Hofman, Maassen, van Schrojenstein Lantman-de Valk, & Snik, 2012).

Similarly, overall ratings at the word level made by unfamiliar listeners, such as percentage of words correct, have often been used as an outcome measure in studies of communication in persons with DS. For example, ratings of intelligibility reported for connected speech are 81.2% for children with DS aged 4–16 years (Barnes et al., 2009), 87% for conversation for persons with DS aged 5–20 years (Chapman, Seung, Schwartz, & Kay-Raining Bird, 1998), and 66.4% for male individuals with DS aged 10–17 years (Rosin, Swift, Bless, & Vetter, 1988). Differences in methodology may account for part of the variability across these studies. These word-level analyses show that persons with DS often have difficulties producing speech that is easily understood.

### ***Analytical Approaches***

Studies using analytical approaches provide insights into the phonetic components that affect intelligibility in persons with DS, which could help in guiding treatment plans to enhance intelligibility. Higher error rates for consonant than vowel production among children and young adults with DS were reported by Dodd and Thompson (2001) for ages 5–15 years and Borsel (1996) for ages 15–28 years. However, both consonant and vowel errors in all articulatory positions except for the high-front vowel /i/ have been reported in a single-word phonetic intelligibility test of five men with DS (Bunton, Leddy, & Miller, 2007). Although Bunton et al. (2007) did not report total error rates for consonants and vowels, more errors in the high–low dimension of vowel production than the front–back dimension were noted. The impact of phonological processes on intelligibility in persons with DS has not been studied systematically. However, persons with DS reportedly have both delayed and disordered phonological processes, which likely affect intelligibility (Kent & Vorperian, 2013).

### ***Converging Conclusions From Global and Analytic Studies***

The two approaches converge on some general conclusions. The main conclusion, which motivates this study, is that reduced speech intelligibility is a common problem in persons with DS and can be a lifelong issue for some individuals. The available data address overall intelligibility but reveal little about developmental changes in factors underlying intelligibility, for example, patterns of phonetic features for vowels and consonants. Another conclusion is that speech intelligibility is highly variable within the population of individuals with DS, but little is known about the factors underlying this variability. A related conclusion,

albeit somewhat tentative because of the limited data, is that boys with DS have poorer articulation and intelligibility than girls with DS (Kumin, 2006; Martin, Klusek, Estigarribia, & Roberts, 2009; Roberts et al., 2005). What is not clear is whether this sex difference persists into adulthood.

### ***Research Strategy for a Life Span Perspective on Speech Intelligibility***

An understanding of the reduced speech intelligibility in DS, across the life span, requires a multifaceted program of research that considers the nature of the deficit in speech intelligibility and the atypical phonological patterns observed at different stages of development, while taking into account the articulatory–motor difficulties suspected to occur in persons with DS as well as the anatomic features commonly associated with the syndrome.

Research on DS is further complicated because the syndrome has a complex phenotype consisting of features that are not fully penetrant (Antonarakis & Epstein, 2006; Vilardell et al., 2011). To address this issue of heterogeneity within the population of individuals with DS, recent research studies have placed emphasis to the study of developmental trajectories (Fidler, 2005) and individual profiles in development (Karmiloff-Smith et al., 2016; Tsao & Kindelberger, 2009). Such a strategy is well-suited for research on a life span perspective of communication skills and can encompass issues such as sex differences, time of emergence of critical skills or deviation from the norm (i.e., emergence of differences), and age intervals of greatest change.

This study is one step in the research process and focuses on the life span study of speech intelligibility deficit using a single-word intelligibility test designed specifically to assess vowel production. The rationale for the methodology used is presented next.

### ***Methods to Study Life Span Speech Intelligibility***

#### ***Tool to Study Life Span Speech Intelligibility***

A tool that is ideal for the study of the developmental trajectory of speech intelligibility is one that is suited to individuals of different ages and different levels of capability. A single-word identification test is advantageous for quantifying changes in speech intelligibility over development in persons with DS for several reasons: First, many persons with DS can participate in a single-word repetition task regardless of their verbal ability and/or age. Single word production is viewed as a general strength in children with DS (Grieco, Pulsifer, Seligsohn, Skotko, & Schwartz, 2015). Second, this approach provides an opportunity for systematic phonetic analysis to characterize the phonetic basis of intelligibility reduction in persons with DS of all ages. Third, the recordings can be used to determine the acoustic correlates of phonetic errors, for example, the voice-onset-time data for voiced–voiceless contrasts in consonants or the formant–frequency patterns for vowel productions

offering a unique opportunity to establish correlations among acoustic, perceptual, articulatory, and even anatomic data. Fourth, a narrow phonetic focus can serve as a touchpoint for studies of acoustic and articulatory correlates (e.g., using vowel space area as an index of articulatory working space for vowels [more on this in the following section]). Thus, keeping in mind the need for a life span perspective, the rationale for an initial focus on vowels is that (a) vowels typically are acquired early in speech development and can be followed from babbling onward, (b) vowels serve as syllable nuclei and therefore have a central role in emerging phonologies, and (c) vowels can be described in terms of articulatory-acoustic correlates such as the F1–F2 vowel plot interpreted as articulatory features.

### Corner Vowels Depicting Articulatory Working Space

The syndrome-specific craniofacial characteristics in DS include a smaller mid and lower facial skeleton but no differences in tongue size (Uong et al., 2001). A typically sized tongue inside a small oral cavity may limit articulatory performance, specifically restricting the articulatory working space for vowels (Kent & Vorperian, 2013). This idea is supported by research on acoustic vowel space area and/or vowel overlap (Abe, 1973; Bunton & Leddy, 2011; Whitworth & Bray, 2015) and by a report of reduced acoustic contrast in the F2 values of the high vowels /i/ and /u/ (Moura et al., 2008). In fact, Moura et al. (2008) suggested that the smaller ratio of the F2 frequencies for vowels /i/ and /u/ is the consequence of anatomic features in DS and accordingly named the relationship between these vowels the *DS vocalic anatomical functional ratio*. Aside from acoustic studies, difficulties with vowel production in DS are evidenced by parent reports (Toğram, 2015), listener judgments of prolonged vowels (Moran, 1986), and speech intelligibility tests (Bunton et al., 2007). Despite these findings indicating that vowel production may impact intelligibility, the relationship between the dimensions of vowel production (front, back, high, low) and intelligibility reduction in persons with DS has not yet been reported. Such knowledge could be invaluable in guiding speech therapy and/or assessing progress in speech treatment. In addition, such information can be used in shaping future anatomic studies and/or evaluating surgical intervention strategies used in upper airway treatment.

### Purpose

The purpose of this perceptual study was to use a single-word identification test with lay listeners to identify phonetic components that contribute to reduced intelligibility in female and male speakers with DS aged 4–40 years. Specific research questions are listed below:

1. What does an open-set word intelligibility test reveal about the pattern of speech intelligibility (words correct scores) in male and female speakers with DS aged 4–40 years? We hypothesized that the results would show (a) improving intelligibility with age, reaching asymptote at adulthood, and (b) substantial

interspeaker variability, owing in part to a sex difference with male individuals having lower intelligibility scores than female individuals.

2. Do vowels differ in their contribution to speech intelligibility? We hypothesized that vowel errors would occur in DS and that the front–back dimension of vowel production would have more vowel identification errors than the high–low dimension because of the reduced mid and lower face skeleton associated with DS.

## Methods

### Part I: Speech Sample Production

#### Participants

All study participants were native speakers of English from the Midwest and followed the same procedure as approved by the University of Wisconsin-Madison Health Sciences Institutional Review Board. Participants (speakers) included 62 persons with DS (29 female and 33 male participants, aged 4–40 years) and a control group of 25 typically developing (TD) participants (11 female and 14 male participants, aged 4–7 years). They were recruited via community flyers and the Developmental Disabilities Registry at the Waisman Center. The participants with DS had Trisomy 21 with no other concomitant diagnosis such as autism or dementia. Elective intake information regarding speech and hearing concerns, history, and/or treatment was provided by 92% of the caregivers of participants with DS and all TD participants. Findings reflected a history of speech therapy services, between the ages 4 and 21 years, for most of the participants with DS (92%), with 35% of those participants receiving services at the time of their speech recording, and hearing concerns were reported for 38% of the participants, with 17% of those indicating it being a former concern. A more formal hearing assessment (hearing screening or audiogram), available for 60% of the participants with DS, reflected hearing status to have a wide range from no hearing loss, to mild, to moderate, and even to severe high-frequency hearing loss in two cases (who wore hearing aids). No developmental concerns and no speech and hearing concerns or treatment were reported for any of the TD participants. Observations of the participants' orofacial structures and function revealed participants with DS to have open mouth posture at rest (37%), underbite (32%), tongue thrust with swallow (21%), and drooling (3%). No similar observations were made for any of the TD participants. The participants with DS were studied in a combined cross-sectional and longitudinal (multiple-session) design where some of them made repeat visits for speech recordings as a venue to explore intraspeaker variability during the course of development. Given that the hearing status was highly variable across participants with DS and within participants with DS who came in for multiple visits, speech recording was carried out after confirming that the participant could follow simple commands and repeat words. Each participant was involved in at least one speech recording session,

with a minimum span of 9 months between recording dates for participants who had multiple sessions. Of the 62 participants with DS, 24 speakers participated in one session and 38 participated in two to seven sessions, yielding 143 speech recordings. As for the 25 TD participants, 23 speakers participated in one session and three participated in two sessions, yielding 29 total speech recordings. The limited age range of TD participants was related to the purpose of determining the age at which children are fully intelligible. TD participants of the following ages were included: 4 years (two girls and two boys), 5 years (three girls and six boys), 6 years (three girls and eight boys), and 7 years (three girls and two boys).

### Procedure

Speech stimuli, drawn from a larger speech task, consisted of 20 words, with each word composed of one of the four corner vowels within a consonant–vowel, vowel–consonant, or consonant–vowel–consonant syllable. Words listed by the vowel they contain are as follows: /i/ = *bead, bee, eat, sheep, and feet*; /u/ = *boo, boot, zoo, hoot, and shoe*; /æ/ = *bath, bat, cat, hat, and sad*; and /a/ = *dot, hop, pot, top, and hot*. Production of corner vowels requires that the tongue be in the most extreme articulatory positions, so these words were chosen to facilitate analysis of the four dimensions of vowel production (high, low, front, and back). Stimuli were also selected to be developmentally appropriate for children and to have a high phonological neighborhood density, because this has been shown to maximize acoustic vowel space (Munson & Solomon, 2004).

Participants were seated in a quiet room and, when possible, completed a pure-tone hearing screening in accordance with American Speech-Language-Hearing Association (2016) standards. Then, speech stimuli were randomly presented on a 14-in. laptop computer or an 11-in. tablet using the TOCS+ Platform (Hodge & Daniels, 2007). The printed word for each stimulus was displayed below its corresponding image, as shown in Figure 1, and a previous recording of the word, produced by a man judged to be highly intelligible and with good voice quality, was presented aurally through loudspeakers at the same time. Participants were instructed to say each word after hearing it. An imitation task allowed for later comparison of listeners' responses to known target stimuli. Participants spoke into a unidirectional Shure SM48 microphone (Shure Inc.) that was placed approximately 15 cm from the speaker's mouth and was stabilized using a floor stand. The microphone was directly connected to a Marantz PMD660 digital audio recorder (Marantz Professional) that digitized speech at 48 kHz with 16-bit resolution. Targeted recording level on the audio recorder's volume unit meter was between 6 and 12 dB below the maximum level. Before producing the selected stimuli, participants counted from 1 to 5 and/or repeated two practice stimuli while adjustments were made to ensure adequate mouth-to-microphone distance and/or recording level. If additional adjustments were needed during the recording session, repeat productions were elicited.

**Figure 1.** An example of stimuli presented to speakers to elicit the production of a target word, which consisted of an image of the target word and the printed word displayed below it, along with a played audio recording of the target word.



## Part II: Intelligibility (Perceptual) Study

### Participants

All 54 adult lay listening participants (17 men and 37 women) were native speakers of English from the Midwest and determined by self-report to be free of communication disorder. They ranged in age from 19 to 58 years, with a mean age of 30 years ( $SD = 11.5$  years).

### Procedure

The stimuli for the listening task were prepared as follows. Each speech recording (as described above) was segmented at the word level, and words were saved as separate sound files using Praat (Boersma & Weenink, 2010). Each word was normalized for amplitude to an average root mean square intensity of  $-10$  dB SPL using Sound Forge (Version 8.0; Sony Creative Software, Inc., 2006) to ensure that fluctuations in speakers' sound levels would not affect intelligibility ratings. Speakers' recordings were assembled into 32 groups for presentation to the listeners. Each group contained speech recordings from five to seven speakers. Twenty-seven groups included speakers with DS, and five groups included TD speakers. Age and sex of DS speakers were balanced to ensure that recordings from children, adults, and male and female individuals were included in each group. Similarly, TD speakers' sex was balanced to ensure that recordings from both boys and girls were presented in each group. Groups with TD speakers included either 4- to 5- or 6- to 7-year-olds. TD speakers were separated by age given inconsistent reports on the age at which children are fully intelligible to unfamiliar listeners (Coplan & Gleason, 1988; Hustad, Oakes, & Allison, 2015).

Panels of five listeners heard one group of recordings per session and participated in two to three sessions on different days. Thus, each group of recordings was presented to five different listeners, because variability in listener

performance has been reported among studies of intelligibility (DePaul & Kent, 2000; Hustad et al., 2015). Listeners were seated in a quiet room to complete the study, which began with a pure-tone hearing screening in accordance with American Speech-Language-Hearing Association (2016) standards. If listeners did not pass the hearing screening, they completed only the first of three listening sessions and their data were not used. After the hearing screening, listeners read instructions for the listening task. They also viewed a list of the speakers' target words to reduce learning effects, because listeners can learn as a task progresses, even with no instruction or feedback (Hustad et al., 2015). In addition, listeners were verbally instructed to type what they heard and not what they thought the speaker was attempting to say. The listening task was completed using a 15-in. laptop with a wireless mouse. Before beginning the task, listeners chose a comfortable listening level by manually adjusting the laptop volume while a prerecorded voice counted from 1 to 10. Then, speech stimuli were presented to listeners through headphones via an in-house computer program written using *Python 3.2* (Python Software Foundation). After each word was presented, listeners typed what they heard. Words were presented only once, and listeners controlled the rate of stimulus presentation. After typing a response, listeners clicked a "next" button, and another word was presented after a 2-s time delay. All words from each speaker were presented before listeners heard another speaker. The order of words within each speaker's recording and the order of speakers' recordings within each listening group were randomized during each session to prevent potential order effects.

### Data Analysis

A software program with two analysis features was written in-house using *Python 3.2* to analyze listeners' responses. The first analysis option scored listeners' identification of each target phoneme and word as correct or incorrect. The program searched for the phonetic representation of each listener's response within the Carnegie Mellon University (CMU) Pronouncing Dictionary (Version 0.7a; Carnegie Mellon University, 2008), compared it with the phonetic representation of the speaker's target word, and then scored each phoneme and word in the listener's response.

Because the CMU dictionary contains only English words, phonetic representations of some listeners' responses were not available for automatic scoring, and manual scoring was required for these responses. Manual editing was also completed as needed to verify scoring accuracy. An example of a response that would require manual scoring is *deeb* for the target word *bead*, because the phonetic representation of *deeb* is not in the CMU dictionary. The word, initial consonant, and final consonant would be manually scored as incorrect, but the vowel would be correct. Thus, speakers were given credit for all phonemes produced accurately in the correct word position, regardless of whether the word was correct. In addition, listeners' responses that were blank or consisted of one letter were considered incomplete and were not included in the analysis. This prevented bias due to listeners' mistakes, such as accidentally

clicking the "next" button before typing their entire response.

After manual editing, the software's second analysis feature calculated the speaker's average percentages correct across all five listeners' responses for words and each of the following phonetic components: total consonants, initial consonants, final consonants, total vowels, and individual vowels /i/, /u/, /æ/, and /a/. These percentages correct, averaged across five listeners for each speaker, are referred to as words correct or phonetic components correct and used as the primary measure of speech intelligibility for each speaker.

### Statistical Analysis

To determine the developmental pattern of speech intelligibility in male and female individuals with DS aged 4–40 years (first research question), we used words correct as the primary measure of intelligibility, with total consonants, initial consonants, final consonants, total vowels, and individual vowels /i/, /u/, /æ/, and /a/ correct as the phonetic components of words correct. Because words correct is a percentage, we applied a generalized linear mixed model (GLMM) with a logit link and binomial errors using PROC GLIMMIX, Statistical Analysis System software (Version 9.4 by SAS Institute, Inc., 2013). Because most participants with DS had multiple sessions, a random intercept for each participant was included in this model to account for correlation across multiple recordings. The model included predictors related to both age and sex (male/female) of the speaker. Because scatter plots of words correct suggested a nonlinear relationship between words correct and age, a fractional polynomial technique was used to select the best-fitting transformation of age (Royston & Altman, 1994). The best transformation was found to be the inverse (1/age). There was no significant interaction between 1/age and sex.

To examine differences in the frequency of errors across vowel types (second research question), we applied a different form of analysis to account for the multivariate outcome. In contrast to the words correct analysis, we now have two sources of statistical dependence in our data: (a) dependence across vowel types due to observations being collected at the same recording occasion and (b) dependence due to the same participants providing recordings over time. We thus specified a multivariate multilevel model in which the error frequencies of each vowel type are the outcomes and vowel-type binary indicators are entered as speaking occasion predictor variables. Because of the use of a multivariate outcome, we adopted a slightly different approach to handling proportional outcomes; specifically, we applied an arcsine transformation of the square root of proportion correct and entered binary indicator predictors for all but the fourth vowel type /u/, which served as the reference category. The choice of /u/ as a reference category was arbitrary, and statistical equivalence involving alternative reference categories was confirmed. To apply this model, we used HLM 7.01 (Scientific Software International; Raudenbush, Bryk, & Congdon, 2010) and

applied follow-up contrasts within the software that conduct statistical tests of differences between all possible pairs of vowel types. Then, to compare patterns of speech intelligibility (words/vowels correct scores) between male and female individuals with DS, in a separate analysis, we entered a binary indicator predictor for speaker sex (entered at the speaker level of the model) as well as interactions between the speaker sex and vowel-type predictors. Tests of statistical significance are based on corresponding statistical contrasts of the relevant coefficient estimates in the model. Using HLM, these statistical tests are single degree-of-freedom chi-square tests.

Listener reliability for words correct scores was assessed using the intraclass correlation coefficient (ICC; Shrout & Fleiss, 1979). Because words correct scores for each speaker were averaged across five listeners, the ICC for a mean of five random raters (1,5) was calculated for a random sample of 25 speakers.

## Results

### *Listener Reliability of Words Correct*

The average of five listeners per speaker was included in the study design to ameliorate effects related to potential differences among listeners. As noted above, the ICC was calculated to assess listener reliability using the mean of five random raters for 25 randomly selected speakers. The ICC was .95, with a 95% confidence interval of .91–.97, confirming high reliability for words correct.

### *Words Correct as a Function of Age and Speaker Sex*

Figures 2a and 2b provide an overall view of the data for words correct for individual speakers with DS compared with data for the TD speaker groups. Data for persons with DS are shown by scatter plots for cross-sectional data and scatter plots connected with lines for longitudinal data. Data for the TD groups are shown by box and whisker plots in the top left corner of each graph. TD girls aged 4–5 years had a mean percent correct of 76 ( $SD = 9$ ), whereas 6- to 7-year-olds had a mean of 92 ( $SD = 7$ ). Similarly, the mean percent words correct for TD boys aged 4–5 years was 81 ( $SD = 11$ ), whereas 6- to 7-year-olds had a mean of 93 ( $SD = 5$ ). As expected, words correct improved with age for all speakers, both TD speakers and those with DS. The 4- to 5- and 6- to 7-year-old TD children were more intelligible than children with DS of the same age and sex. Children with DS also had greater variability than TD speakers of the same age and sex, with the exception of 4- to 5-year-old TD boys. Furthermore, TD 6- to 7-year-olds were more intelligible than most children and adults with DS of the same sex, indicating markedly reduced speech intelligibility in many persons with DS. For children with DS younger than 14 years old, 100% of boys and 95% of girls were less intelligible than TD 6- to 7-year-olds. For persons with DS

aged 14 years and older, 74% of male individuals and 49% of female individuals remained less intelligible than TD 6- to 7-year-olds.

Despite the overall trend of improved speech intelligibility with age in persons with DS, there was considerable intraspeaker and interspeaker variability. As shown by the longitudinal data (points for the same speaker are connected) in Figures 2a and 2b, most persons with DS showed improved intelligibility across sessions between ages 4 and 16 years. However, after 16 years old, intelligibility patterns across sessions were more variable. Thus, intelligibility did not always improve across sessions, and reduced intelligibility persisted into adulthood for some persons with DS. Considerable interspeaker variability among persons with DS can be seen in Figures 2a and 2b for any age–sex combination.


The possibility of sex differences is also evident in Figures 2a and 2b, where visual examination of the data shows female individuals with DS being more intelligible at the word level than male individuals, particularly for persons aged 9 years and older. Words correct for female individuals with DS was greater than 40% by the age of 9 years, but most male individuals did not reach this level of intelligibility until the age of 16 years. In addition, all female individuals with DS aged 8 years and older had greater than 40% words correct, whereas some adult men with DS did not exceed 40% words correct.

Statistical testing of the developmental trajectories was conducted using the GLMM described earlier. The coefficient for the inverse of age ( $1/\text{age}$ ) was  $-15.64$ , which was highly significant,  $F(1, 80) = 43.36, p < .0001$ . A significant main effect coefficient of 0.65 was also observed for sex,  $F(1, 61) = 5.39, p = .024$ , with female speakers yielding higher proportion correct. As noted earlier, there was no detectable interaction between sex and inverse of age. The resulting model-based trajectories by sex are shown in Figure 3, demonstrating that word intelligibility increased significantly with age among both male and female individuals with DS at seemingly similar rates and with a larger rate of increase during the developmental ages (especially 4–16 years).

### *Words Correct as a Function of Phonetic Component*

As described earlier, our scoring permitted a finer assessment of how both consonant and vowel errors contributed to diminished speech intelligibility. Figures 4 and 5 illustrate that both consonant and vowel errors contribute to intelligibility as measured by percent words correct. Figure 4 highlights the relationship between words correct and total consonants correct (initial and final consonants combined; Figure 4a), and total vowels correct (Figure 4b). Figure 5 displays the finer detail of the relationships between words correct and initial consonants correct (Figure 5a) and final consonants correct (Figure 5b) as well as words correct and each of the individual vowels correct: /i/ (Figure 5c), /æ/ (Figure 5d), /u/ (Figure 5e), and /a/

**Figure 2.** (a–b) Words correct for all speakers, with female individuals shown in red and male individuals in blue. Data from TD speakers are represented by box and whisker plots. The box plots show the 25th and 75th percentiles, with a dotted line representing the mean and a solid line representing the median. The whiskers show the 10th and 90th percentiles, and outlying data are shown as dots. Data from persons with DS are shown by scatter plots. Speakers with only one speech recording are represented by triangles, and speakers with multiple speech recordings are represented by different symbols (squares, circles, and cross) connected by dotted or solid lines. DS = Down syndrome; TD = typically developing.




(Figure 5f). Closer examination of Figure 5 reveals that the relationship between percent words correct and each of the vowels correct was nonuniform, with particular vowel types associated with a greater occurrence of errors. We therefore carried out statistical analyses to test for differences among vowels (second research question), as described in the following section.

### *Vowel Error Frequency as a Function of Vowel Type*

Statistical analysis contrasting vowel types (based on the multivariate multilevel analysis described earlier) indicated statistically significant differences in error frequencies between vowel types /a/ and /u/ ( $\chi^2 = 131.63$ ,  $df = 1$ ,  $p < .001$ ),

**Figure 3.** Words correct for persons with DS, with data from male individuals shown by blue triangles and data from female individuals shown by red circles. Colored lines indicate the equation for generalized linear mixed models for male and female individuals, and colored bands indicate 95% confidence intervals for the models. DS = Down syndrome.




*/æ/* and */u/* ( $\chi^2 = 53.79$ ,  $df = 1$ ,  $p < .001$ ), */a/* and */æ/* ( $\chi^2 = 24.82$ ,  $df = 1$ ,  $p < .001$ ), */a/* and */i/* ( $\chi^2 = 175.63$ ,  $df = 1$ ,  $p < .001$ ), and */æ/* and */i/* ( $\chi^2 = 65.94$ ,  $df = 1$ ,  $p < .001$ ). The only pairwise comparison that did not yield statistical significance was between */i/* and */u/* ( $\chi^2 = 0.79$ ,  $df = 1$ ,  $p > .500$ ). This can be seen in Table 1, where the error frequencies are compiled by vowel type and support these results with the highest frequency of errors occurring for */a/*, followed

by */æ/*, and with */i/* and */u/* having the lowest error frequencies.

### *Vowel Errors as a Function of Speaker Sex*


Given that the GLMM analysis found a significant main effect related to sex for overall intelligibility, we also proceeded to examine potential sex differences related to

**Figure 4.** Relationship between percent words correct for persons with DS and total consonants correct (a) and total vowels correct (b). The percentage of words perceived as correct was averaged across five listeners' responses, similar to the calculations used to create Figures 2 and 3. DS = Down syndrome.





**Figure 5.** Relationship between percent words correct for persons with DS and initial consonants correct (a), and final consonants correct (b), and individual vowels correct /i/, /u/, /æ/, and /a/ (c-f). DS = Down syndrome.



**Table 1.** Descriptive statistics for the proportion of vowels correct across speech recordings from persons with Down syndrome ( $N$  = total number of speech recordings, generally several recordings per speaker).


| Speaker Group | Vowel type | $N$ | Minimum | Maximum | Mean | $SD$ |
|---------------|------------|-----|---------|---------|------|------|
| Overall       | /i/        | 143 | 0.42    | 1.00    | 0.94 | 0.11 |
|               | /u/        | 143 | 0.17    | 1.00    | 0.93 | 0.15 |
|               | /æ/        | 143 | 0.12    | 1.00    | 0.82 | 0.24 |
|               | /a/        | 143 | 0.09    | 1.00    | 0.73 | 0.25 |
| Male          | /i/        | 75  | 0.42    | 1.00    | 0.91 | 0.14 |
|               | /u/        | 75  | 0.17    | 1.00    | 0.90 | 0.19 |
|               | /æ/        | 75  | 0.17    | 1.00    | 0.74 | 0.26 |
|               | /a/        | 75  | 0.09    | 1.00    | 0.69 | 0.26 |
| Female        | /i/        | 68  | 0.80    | 1.00    | 0.98 | 0.05 |
|               | /u/        | 68  | 0.60    | 1.00    | 0.97 | 0.07 |
|               | /æ/        | 68  | 0.12    | 1.00    | 0.89 | 0.17 |
|               | /a/        | 68  | 0.24    | 1.00    | 0.78 | 0.23 |

*Note.* The table displays the statistics broken down by vowel type for all (top), male individuals only (center), and female individuals only (bottom).  $N$  = speaking occasions.

vowels specifically. Our second analysis used speaker sex as an additional predictor of vowel proportion correct, as described earlier. Findings revealed a significant main effect related to sex ( $\chi^2 = 5.87$ ,  $df = 1$ ,  $p = .015$ ) such that male individuals tended to have more vowel errors. An omnibus test for interactions between sex and vowel type based on the coefficients of product variables between sex and vowel type was not significant ( $\chi^2 = 1.18$ ,  $df = 3$ ,  $p = .276$ ), suggesting a similar profile of vowel errors across sex. Table 1 displays the descriptive statistics of error frequencies by vowel type for each sex.

To examine the possibility of a developmental trend for vowel intelligibility in persons with DS, percent correct (averaged across five listeners) was plotted for each of the four corner vowels, as shown in Figure 6. Individual vowels correct for persons with DS are shown by triangles for cross-sectional data and circles for longitudinal data. As shown in Figure 6, high vowels /i/ and /u/ developed at an earlier age and were more intelligible than low vowels /æ/ and /a/ for persons with DS. High vowels were developed by 4 years old for most persons with DS. Female

**Figure 6.** Percent correct for vowels /i/, /u/, /a/, and /æ/ for persons with DS as a function of age, with female individuals shown in red and male individuals in blue. Speakers with a single session are represented by triangles, whereas speakers with repeat sessions are represented by circles. Refer to Table 2 for percent vowels correct data for TD speakers. DS = Down syndrome; TD = typically developing.



individuals with DS of all ages had a mean percent correct of 97 ( $SD = 4$ ) for high vowels, whereas male individuals had a mean percent correct of 92 or greater ( $SD = 15$ ). However, percent correct continued to increase through 12 years old for low vowel /æ/ and 16 years old for low vowel /a/ for persons with DS. Furthermore, persons with DS had fewer vowels correct and greater variability for low vowels /æ/ and /a/ at all ages. Female individuals with DS aged 13 years and older had a mean percent correct of 93 ( $SD = 10$ ) for vowel /æ/, whereas male individuals had a mean percent correct of 86 ( $SD = 19$ ). For vowel /a/, female individuals with DS aged 17 years and older had a mean percent correct of 84 ( $SD = 20$ ), whereas male individuals had a mean percent correct of 80 ( $SD = 23$ ). Although high vowels were mastered at younger ages than low vowels in female and male individuals with DS, male individuals had fewer vowels correct and more variability than female individuals for each individual vowel.

TD speaker groups had a mean percent correct of 95 or greater ( $SD = 7$ ) for every vowel, except 4- to 5-year-old TD girls for vowel /a/. Table 2 displays the mean percent correct and standard deviation of each vowel for TD

speaker groups. Intelligibility and age of mastery for high vowels were similar for TD speaker groups and persons with DS, but TD speakers had greater intelligibility and earlier age of mastery for low vowels.

## Discussion

This study supports and extends previous research showing reduced intelligibility in children and adults with DS. To our knowledge, this is the first study to analyze speech intelligibility in children and adults with DS by using a uniform methodology focusing on patterns of intelligibility at the word and phonetic levels, specifically vowels. Our results confirm increased speech intelligibility with age in persons with DS, especially between the ages of 4 and 16 years. This 12-year window of improved intelligibility holds implications for intervention focused on speech communication. The data also reveal considerable interspeaker variability. Male individuals with DS were less intelligible than female individuals, particularly during early childhood. As expected, both initial and final consonants contributed to intelligibility, but vowels also contribute to

**Table 2.** Mean and SD for percent vowels correct for typically developing boys and girls.

| Vowel | Sex | 4- to 5-year-olds:        |       | 6- to 7-year-olds: |                           |     |
|-------|-----|---------------------------|-------|--------------------|---------------------------|-----|
|       |     | percent correct mean (SD) |       | Sex                | percent correct mean (SD) |     |
| /i/   | F   | 100                       | (0)   | F                  | 100                       | (0) |
|       | M   | 100                       | (0)   | M                  | 100                       | (0) |
| /u/   | F   | 99                        | (2)   | F                  | 99                        | (2) |
|       | M   | 98                        | (4)   | M                  | 98                        | (5) |
| /æ/   | F   | 97                        | (5)   | F                  | 98                        | (5) |
|       | M   | 98                        | (4)   | M                  | 98                        | (4) |
| /ɑ/   | F   | 89                        | (10)* | F                  | 97                        | (5) |
|       | M   | 95                        | (7)   | M                  | 98                        | (3) |

*Note.* The only speaker group with a mean of less than 95% and a standard deviation greater than 7% is marked with an asterisk (\*). F = female; M = male.

intelligibility and should not be neglected, as they also provide valuable information about the phonetic bases of reduced intelligibility. High vowels developed at an earlier age than low vowels in DS.

### Variability

The considerable intraspeaker and interspeaker variability observed in this study agrees with previous findings documenting individual variability in error patterns among men with DS (Bunton et al., 2007) and variability in speech intelligibility among children with DS (Kumin, 2006). In this study, intelligibility patterns were the most variable across sessions for persons with DS over 16 years old, as seen in Figures 2a and 2b. Such intraspeaker variability could be due to a variety of factors, such as the time of day speech was recorded, physiological or psychological state, and variations in medical condition and/or speech and language abilities. Informal assessment of the time of day of speech recording did not show any consistent patterns related to speech intelligibility.

Interspeaker variability could be rooted in factors intrinsic to the speaker, such as craniofacial dysmorphology, speech motor impairment (including hypotonia and articulatory placement and sequencing disorders), hearing loss, auditory processing deficiencies, language impairment, and/or cognitive impairment (Kent & Vorperian, 2013). Informal assessment of hearing screening results, hearing status, and/or hearing history (when available) revealed no consistent pattern between hearing loss and speech intelligibility. Variability could also be due to differences in speech therapy services received. Of the 52 participants with DS with information available, 48 speakers had a history of speech therapy. No consistent pattern of intelligibility was observed among the four participants who did not receive speech therapy. However, detailed information about type and frequency of speech therapy was not collected, and these variables likely influence intelligibility. Further research is needed to examine these factors, and their likely interactions, as they relate to intelligibility. As emphasized by Karmiloff-Smith et al. (2016), individual differences in DS are notable across genetic, cellular, neural, cognitive,

behavioral, and environmental observations. The authors suggested that examination of these differences is one of the best approaches to understand genotype/phenotype relations in DS. The striking heterogeneity in persons with DS makes it difficult to classify research participants in groups according to such potentially influential factors as hearing status, type and severity of craniofacial anomaly, degree of hypotonia in the oral musculature, and dosage and type of speech therapy. Determining the effects and interactions of these variables would require a large sample.

### Speaker Sex and Age

Analysis of speaker sex showed that female individuals with DS were more intelligible than male individuals, in agreement with results reported by Kumin (2006), Martin et al. (2009), and Roberts et al. (2005). A possible explanation is that speech intelligibility of girls with DS develops at a faster rate than that of boys with DS through the age of 15 years, as shown in Figures 2a, 2b, and 3. This developmental trend is similar to that in TD children, as most TD girls aged 6 years and younger master phonemes earlier than TD boys (Smit, Hand, Freilinger, Bernthal, & Bird, 1990). For adults with DS, greater variability was more evident among men than women. The influence of speaker sex on speech intelligibility in DS needs further investigation, especially with respect to causal factors that may include possible anatomic differences of the vocal tract.

A brief comment should be made on the TD data. On the basis of reports in the literature (Coplan & Gleason, 1988), we expected that TD 4- to 5-year olds would achieve nearly complete intelligibility for a single-word test. However, the mean words correct for TD 4- to 5-year-olds was 76% for girls and 81% for boys, whereas TD 6- to 7-year-old boys and girls had greater than 90% words correct. Our results support the conclusion of Hustad et al. (2015) that TD children are not fully intelligible until about 6 years old. It should be emphasized that our results pertain to a single-word test based on highly familiar lexical items and with a limited repertoire of phonetic elements. It is likely that

speech intelligibility would be even lower for more complex speech materials. Despite the use of stimuli that were developmentally appropriate for speakers of all ages, TD 4- to 5- and 6- to 7-year-olds were more intelligible than most persons with DS of all ages, highlighting the speech intelligibility difficulties faced by many persons with DS.

### ***Contribution of Consonants and Vowels to Intelligibility***

As expected, total consonants correct contributed to word intelligibility in DS. Higher accuracy for vowel production than consonant production has been reported for persons with DS (Dodd & Thompson, 2001; van Bysterveldt, Gillon, & Foster-Cohen, 2010). In this study, comparisons of vowel and consonant contributions to intelligibility are limited because of the structure of the speech material. The relative distribution of consonants and vowels in the stimuli words was such that 15 of 20 words contained two consonants and one vowel.

This study was designed to examine aspects of vowel production in relation to word intelligibility, and we were interested to see if and how vowel errors can be used as a gauge of speech intelligibility in DS. Vowel errors are likely to occur because of several factors that also contribute to interspeaker variability, including anatomic, acoustic, and/or physiological differences in persons with DS.

### ***Intelligibility of Individual Vowels***

High vowels were more intelligible and developed earlier than low vowels for both male and female individuals with DS (see Figure 6). For most children with DS, high vowels developed before 4 years old, whereas intelligibility of low vowels continued to increase through 12 years old for vowel /æ/ and through 16 years old for vowel /a/ (see Figure 6). In addition, persons with DS of all ages had similar intelligibility to TD children aged 4–7 years for high vowels, although persons with DS had more variability. Furthermore, mean intelligibility of low vowels was greatly reduced for persons with DS as compared with high vowels, and vowel /a/ had the greatest variability (see Figure 6). Male individuals with DS had a lower mean intelligibility for each individual vowel and greater variability as compared with female individuals in this study. Bunton et al. (2007) studied five men with DS, and their study is the only other study that has used an analytical approach to study vowel intelligibility. The present finding of reduced intelligibility for low vowels agrees with Bunton et al.'s report that men with DS had more errors in the high–low dimension of vowel production than in the front–back dimension. The present results suggest that this trend may be true for female and male individuals with DS of all ages. Bunton et al. also reported no errors in vowel /i/ for men with DS, and in this study, vowel /i/ had the fewest errors of the corner vowels.

The difficulty in the production of low vowels in DS may be explained as follows. Precise tongue–jaw coordination is needed to produce low vowels, but little tactile feedback is provided between the tongue and the maxilla. Low and high vowels may differ not only in sensory feedback but also in mechanical support from peripheral structures such as the molars, palate, and pharynx. The term *bracing* has been used to describe a lingual posture in speech wherein the tongue is in contact with a rigid vocal-tract surface, such as the teeth or palate (Gick, Allen, Roewer-Després, & Stavness, 2017). The nature of the bracing differs with the articulatory features of vowels. For example, the high vowels are likely to be braced by the palate and the maxillary dentition, whereas the low–back vowels are more likely to be braced by the pharynx. Another factor to be considered is that the articulatory distance between low vowels can be smaller than that between high vowels. Possibly, the tongue movement between low vowels may be especially restricted because of anatomic limitations in persons with DS. The result that the low dimension of vowel production is more related to intelligibility than the high dimension is in line with previous reports of a smaller mid and lower facial skeleton. This anatomic feature may contribute to a reduced articulatory working space and a consequent reduction of the vowel acoustic space in DS.

### ***Limitations and Future Directions***

To our knowledge, this is the first study to use an open-set, single-word intelligibility test to determine the contribution of specific phonetic factors to speech intelligibility in children and adults with DS. Although the design of this study facilitated phonetic analysis across a wide age range, intelligibility data at the word level may not be representative of speakers' intelligibility during connected speech tasks such as conversation. Further research analyzing intelligibility at the phrase or sentence level is needed to obtain a more comprehensive picture of intelligibility reduction in DS, and research should account for the considerable intraspeaker and interspeaker variability within this population. An important caveat is that children with DS appear to be capable of single-word production as compared with more complex utterances (Grieco et al., 2015), which makes these materials more suitable than complex utterances for testing basic articulation. The speech sample used in this study was specifically designed to examine the corner vowels. It would be informative to use a speech sample that incorporates consonant contrasts and additional vowels.

Longitudinal intelligibility data from persons with DS are needed to examine individual developmental patterns using both global and analytic methods. In addition, data from young children with DS (younger than 4 years old) should provide a comprehensive view of intelligibility during development. Other variables that warrant further study are (a) the effects of listener familiarity and the context of the utterance; (b) speaker-related factors, such

as voice and prosody; and (c) the relationship between acoustic and intelligibility measures. As recommended by Miller (2013), combining “diagnostic” or analytic intelligibility testing with global measures of intelligibility is likely necessary for a full understanding of speech intelligibility in clinical populations. There are no specific data on underlying mechanisms that negatively impact speech, and there is no systematic assessment procedure available for evaluating the speech of adults with DS.

### ***Clinical Implications***

The word intelligibility test used in the present research provides both an overall measure of intelligibility and an analysis of some phonetic factors that account for variations in overall intelligibility. Bunton et al. (2007) used a single-word test to determine factors underlying reduced intelligibility in five men with DS. The most disrupted features involved (a) word-initial and word-final cluster simplification and (b) contrasts of tongue posture, control, and timing (e.g., high–low vowel, front–back vowel; place of articulation for stops and fricatives). These features may be particularly important for assessing speech intelligibility in persons with DS and planning treatment that is focused on factors contributing to intelligibility.

Although this study was not designed to examine consonant production in detail, it is clear that consonants carry a significant load in speech intelligibility. The difficulties with consonants noted in this study are especially remarkable given that stimuli words were composed mostly of consonants that are acquired early in typical development (e.g., /b p h t d k/). Given the developmental and life span perspective of this general project, the selection of words was guided by the criteria of familiarity, high phonological neighborhood density, and relative ease of production. More demanding versions of the test could be developed to explore additional phonetic aspects, but even the current version appears to be applicable and informative for studies of speech intelligibility in children and adults with DS.

The protracted improvement in speech intelligibility observed in this study provides justification for treatment targeting intelligibility. It is noteworthy that many individuals with DS achieve relatively high levels of intelligibility by the age of 16 years. Although some adults with DS have persisting low levels of intelligibility, the prospects appear encouraging for many individuals to achieve speech that is reasonably intelligible to unfamiliar listeners. In view of the continued improvement in intelligibility in DS from ages 4 to 16 years, it is possible that intervention efforts during this period would accelerate intelligibility gains. This is not to say that treatment at earlier or later ages is inappropriate but only to suggest that individuals between the ages of 4 and 16 years appear to be good candidates for intervention focused on improving intelligibility. The protracted intelligibility improvements in DS, compared with that of TD children who are intelligible by the

age of 7 years, may be evidence for self-adaptations and maturational processes that could be reinforced or even expanded with focused treatments.

One novel result of this study is that reduced contrast between low–front and low–back vowels is a factor accounting for reduced intelligibility in DS. A likely explanation is that these vowels have a slower development than high–front and high–back vowels and therefore account for changes in intelligibility with maturation. The low vowels may offer insight into the developmental processes underlying intelligibility and therefore may be sensitive for purposes of assessment. They also could serve as suitable targets for early phases of treatment, not only for more correct vowel production per se but also to leverage better control of lingual articulation that could lead to a more accurate consonant production. The production of consonants is predicated in part on the mastery of vowels, which requires control of tongue body articulation. Strategies for the treatment of vowels in children are discussed by Gibbon (2013) and Speake, Stackhouse, and Pascoe (2012). Mahler and Jones (2012) reported that treatment of two adults with DS using the Lee Silverman Voice Treatment had positive outcomes, leading to improved vowel production in one individual. More generally, there appears to be promise in the use of principles of motor learning to improve motor skills in persons with DS (Horvat, Croce, & Fallaize, 2016), and intervention for speech could incorporate the same approach, especially if due consideration is given to the structural, motor, sensory, and cognitive issues pertaining to speech production.

It is also possible that combinations of behavioral and other kinds of treatment can facilitate gains in speech intelligibility in persons with DS. For example, interventions that directly alter the anatomy of the orofacial complex, such as a stimulating palatal plate (Matthews-Brzozowska, Cudzilo, Walasz, & Kawala, 2014), may contribute to improved intelligibility by helping speakers to overcome anatomic limitations. Concurrent behavioral intervention in concert with such anatomically directed treatment may achieve desired functional outcomes. However, it cannot be assumed that all interventions affecting the tongue or oral cavity will lead to improved speech. Partial glossectomy does not necessarily result in improved speech intelligibility in either DS or Beckwith–Wiedemann syndrome (Margar-Bacal, Witzel, & Munro, 1987; Van Lierde et al., 2012). A variety of interventions have been introduced to improve appearance, reduce drooling, and facilitate swallowing in DS, but it is important to ensure that such treatments do not have deleterious effects on speech production. Clinical speech assessment during treatment regimens is highly advisable to ensure that speech is not negatively affected. It is also imperative to keep in mind the individual differences notable in the population with DS.

Bittles, Bower, Hussain, and Glasson (2007) commented on an extraordinary point of progress in improving the lives of individuals with DS: “A highly significant change in the survival of people with DS has occurred during

the last two generations, with life expectancy estimates increasing from 12 to nearly 60 years of age” (p. 224). This increased life expectancy would be accompanied by improved quality of life for many individuals through enhanced communication skills.

## Conclusion

This study makes a novel contribution by showing that an open-set, single-word intelligibility test can be used to determine the developmental pattern of word-level speech intelligibility and to examine phonetic factors relating to intelligibility scores. The results reveal that intelligibility gains are evident in speakers with DS, especially in the age range of 4–16 years, that female individuals have higher intelligibility scores than male individuals, and that the corner vowels make different contributions to speech intelligibility. Results of this perceptual study add to the understanding of the pervasive intelligibility difficulties experienced by children and adults with DS. Clarifying the basis of these speech intelligibility challenges is a critical step to direct future research efforts and design interventions that will enhance communication and, subsequently, quality of life.

## Acknowledgments

This work was supported by Research Grant R01 DC006282 to Houri K. Vorperian from the National Institute on Deafness and Other Communication Disorders. This study was also supported in part by core grants P30 HD03352 and U54 HD090256 to the Waisman Center from the National Institute of Child Health and Human Development. Special thanks to Allison Carolan, Ekaterini Derdemezis, Allison Petska, Daniel Reilly, Peggy Rosin, and Emily Reinicke for assistance with experimental design, task setup, and data collection; Robert Olson for software programming; Carlyn Burris, Erin Douglas, Julie Eichhorn, Ellie Fisher, Katie Lester, Erin Nelson, and Katelyn K. Tillman for data collection; Elaine Romenesko and Lela Murchison for data analysis and data collection; Gabriel Jardim for assistance with figure edits; and Abigail Lamers and Courtney Wagner for assistance with references. Portions of this research were presented in 2015 at the American Speech-Language-Hearing Association Convention in Denver, CO.

## References

- Abe, M. (1973). A developmental study of vowels in Down’s syndrome: An acoustic study [Japanese]. *The Japan Journal of Logopedics and Phoniatrics*, 14, 2–11.
- American Speech-Language-Hearing Association. (2016). *Adult hearing screening*. Retrieved from [http://www.asha.org/PRPSpecificTopic.aspx?folderid=8589942721&section=Key\\_Issues](http://www.asha.org/PRPSpecificTopic.aspx?folderid=8589942721&section=Key_Issues)
- Antonarakis, S. E., & Epstein, C. J. (2006). The challenge of Down syndrome. *Trends in Molecular Medicine*, 12(10), 473–479.
- Barnes, E., Roberts, J., Long, S. H., Martin, G. E., Berni, M. C., Mandulak, K. C., & Sideris, J. (2009). Phonological accuracy and intelligibility in connected speech of boys with fragile X syndrome or Down syndrome. *Journal of Speech, Language, and Hearing Research*, 52(4), 1048–1061. [https://doi.org/10.1044/1092-4388\(2009/08-0001\)](https://doi.org/10.1044/1092-4388(2009/08-0001))
- Bittles, A. H., Bower, C., Hussain, R., & Glasson, E. J. (2007). The four ages of Down syndrome. *European Journal of Public Health*, 17(2), 221–225.
- Boersma, P., & Weenink, D. (2010). Praat (version 5.1.31) [software]. Amsterdam, The Netherlands: Phonetic Sciences, University of Amsterdam. Retrieved from <http://www.fon.hum.uva.nl/praat>
- Borsel, J. V. (1996). Articulation in Down’s syndrome adolescents and adults. *European Journal of Disorders of Communication*, 31(4), 415–444.
- Bunton, K., & Leddy, M. (2011). An evaluation of articulatory working space area in vowel production of adults with Down syndrome. *Clinical Linguistics & Phonetics*, 25(4), 321–334. <https://doi.org/10.3109/02699206.2010.535647>
- Bunton, K., Leddy, M., & Miller, J. (2007). Phonetic intelligibility testing in adults with Down syndrome. *Down’s Syndrome, Research and Practice*, 12(1), 1–24. <https://doi.org/10.3104/reports.2027>
- Carnegie Mellon University. (2008). *Carnegie Mellon University Pronouncing Dictionary* (Version 0.7a). Retrieved from <http://svn.code.sf.net/p/cmuspshinx/code/trunk/cmudict>
- Chapman, R. S., Seung, H. K., Schwartz, S. E., & Kay-Raining Bird, E. (1998). Language skills of children and adolescents with Down syndrome: II. Production deficits. *Journal of Speech, Language, and Hearing Research*, 41(4), 861–873.
- Coplan, J., & Gleason, J. R. (1988). Unclear speech: Recognition and significance of unintelligible speech in preschool children. *Pediatrics*, 82(3), 447–452.
- Coppens-Hofman, M. C., Maassen, B. A., van Schroyen Lantman-de Valk, H., & Snik, A. F. (2012). Speech difficulties and poor speech intelligibility in adults with Down syndrome: A review of the literature. *Journal of Hearing Science*, 2(1), 9–16.
- De Moura, C. P., Andrade, D., Cunha, L. M., Tavares, M. J., Cunha, M. J., Vaz, P., . . . Clemente, M. P. (2008). Down syndrome: Otolaryngological effects of rapid maxillary expansion. *The Journal of Laryngology and Otology*, 122(12), 1318–1324.
- DePaul, R., & Kent, R. D. (2000). A longitudinal case study of ALS: Effects of listener familiarity and proficiency on intelligibility judgments. *American Journal of Speech-Language Pathology*, 9(3), 230–240.
- Dodd, B., & Thompson, L. (2001). Speech disorder in children with Down’s syndrome [comparative study]. *Journal of Intellectual Disability Research*, 45(Pt. 4), 308–316.
- Fidler, D. J. (2005). The emerging Down syndrome behavioral phenotype in early childhood: Implications for practice. *Infants & Young Children*, 18(2), 86–103.
- Gibbon, F. E. (2013). Therapy for children with abnormal vowels. In M. J. Ball & F. E. Gibbon (Eds.), *Handbook of vowels and vowel disorders* (pp. 429–446). New York, NY: Psychology Press.
- Gick, B., Allen, B., Roewer-Després, F., & Stavness, I. (2017). Speaking tongues are actively braced. *Journal of Speech, Language, and Hearing Research*, 60(3), 494–506.
- Grieco, J., Pulsifer, M., Seligsohn, K., Skotko, B., & Schwartz, A. (2015). Down syndrome: Cognitive and behavioral functioning across the lifespan. *American Journal of Medical Genetics Part C: Seminars in Medical Genetics*, 169(2), 135–149.
- Hodge, M., & Daniels, J. (2007). TOCS+ software platform for stimulus presentation and audio recording [Computer software]. Edmonton, Canada: University of Alberta.
- Horvat, M., Croce, R., & Fallaize, A. (2016). Information processing and motor control in Down syndrome. *Journal of Down Syndrome & Chromosome Abnormalities*, 2, 1. <https://doi.org/10.4172/jdsca.1000107>

- Hustad, K. C. (2012). Speech intelligibility in children with speech disorders. *Perspectives on Language Learning and Education*, 19(1), 7–11.
- Hustad, K. C., Oakes, A., & Allison, K. (2015). Variability and diagnostic accuracy of speech intelligibility scores in children. *Journal of Speech, Language, and Hearing Research*, 58(6), 1695–1707. [https://doi.org/10.1044/2015\\_JSLHR-S-14-0365](https://doi.org/10.1044/2015_JSLHR-S-14-0365)
- Karmiloff-Smith, A., Al-Janabi, T., D'Souza, H., Groet, J., Massand, E., Mok, K., ... Strydom, A. (2016). The importance of understanding individual differences in Down syndrome. *F1000Research*, 5, 389. <https://doi.org/10.12688/f1000research.7506.1>
- Kent, R., & Vorperian, H. (2013). Speech Impairment in Down syndrome: A Review. *Journal of Speech, Language, and Hearing Research*, 56(1), 178–210. [https://doi.org/10.1044/1092-4388\(2012\)12-0148](https://doi.org/10.1044/1092-4388(2012)12-0148)
- Kent, R. D., Miolo, G., & Bloedel, S. (1994). The intelligibility of children's speech: A review of evaluation procedures. *American Journal of Speech-Language Pathology*, 3(2), 81–95.
- Kumin, L. (1994). Intelligibility of speech in children with Down syndrome in natural settings: Parents' perspective. *Perceptual and Motor Skills*, 78(1), 307–313.
- Kumin, L. (2006). Speech intelligibility and childhood verbal apraxia in children with Down syndrome. *Down's Syndrome, Research and Practice*, 10(1), 10–22.
- Mahler, L. A., & Jones, H. N. (2012). Intensive treatment of dysarthria in two adults with Down syndrome. *Developmental Neurorehabilitation*, 15(1), 44–53. <https://doi.org/10.3109/17518423.2011.632784>
- Margar-Bacal, F., Witzel, M. A., & Munro, I. R. (1987). Speech intelligibility after partial glossectomy in children with Down's syndrome. *Plastic and Reconstructive Surgery*, 79(1), 44–47.
- Martin, G. E., Klusek, J., Estigarribia, B., & Roberts, J. E. (2009). Language characteristics of individuals with Down syndrome. *Topics in Language Disorders*, 29(2), 112–132.
- Matthews-Brzozowska, T., Cudziło, D., Walasz, J., & Kawala, B. (2014). Rehabilitation of the orofacial complex by means of a stimulating plate in children with Down syndrome. *Advances in Clinical and Experimental Medicine*, 24(2), 301–305. <https://doi.org/10.17219/acem/40451>
- Miller, N. (2013). Measuring up to speech intelligibility. *International Journal of Language & Communication Disorders*, 48(6), 601–612. <https://doi.org/10.1111/1460-6984.12061>
- Moran, M. J. (1986). Identification of Down's syndrome adults from prolonged vowel samples. *Journal of Communication Disorders*, 19(5), 387–394.
- Moura, C. P., Cunha, L. M., Vilarinho, H., Cunha, M. J., Freitas, D., Palha, M., ... Pais-Clemente, M. (2008). Voice parameters in children with Down syndrome. *Journal of Voice*, 22(1), 34–42.
- Munson, B., & Solomon, N. P. (2004). The effect of phonological neighborhood density on vowel articulation. *Journal of Speech, Language, and Hearing Research*, 47(5), 1048–1058. [https://doi.org/10.1044/1092-4388\(2004\)078](https://doi.org/10.1044/1092-4388(2004)078)
- Raudenbush, S. W., Bryk, A. S., & Congdon, R. (2010). HLM 7.00 for Windows [Computer software]. Lincolnwood, IL: Scientific Software International, Inc.
- Roberts, J., Long, S. H., Malkin, C., Barnes, E., Skinner, M., Hennon, E. A., & Anderson, K. (2005). A comparison of phonological skills of boys with fragile X syndrome and Down syndrome. *Journal of Speech, Language, and Hearing Research*, 48(5), 980–995.
- Roberts, J. E., Price, J., & Malkin, C. (2007). Language and communication development in Down syndrome. *Developmental Disabilities Research Reviews*, 13(1), 26–35.
- Rosin, M. M., Swift, E., Bless, D., & Vetter, D. K. (1988). Communication profiles of adolescents with Down syndrome. *Journal of Childhood Communication Disorders*, 12, 49–64.
- Royston, P., & Altman, D. G. (1994). Regression using fractional polynomials of continuous covariates: Parsimonious parametric modelling. *Applied Statistics*, 43, 429–467. <https://doi.org/10.2307/2986270>
- SAS Institute, Inc. (2013). Statistical Analysis System (Version 9.4) [Computer software]. Cary, NC: Author.
- Shrout, P. E., & Fleiss, J. L. (1979). Intra-class correlations: Uses in assessing rater reliability. *Psychological Bulletin*, 86(2), 420–428.
- Smit, A. B., Hand, L., Freilinger, J. J., Bernthal, J. E., & Bird, A. (1990). The Iowa articulation norms project and its Nebraska replication. *Journal of Speech and Hearing Disorders*, 55(4), 779–798.
- Sony Creative Software, Inc. (2006). Sound Forge (version 8.0) [Computer software]. Middleton, WI: Author.
- Speake, J., Stackhouse, J., & Pascoe, M. (2012). Vowel targeted intervention for children with persisting speech difficulties: Impact on intelligibility. *Child Language Teaching and Therapy*, 28(3), 277–295. <https://doi.org/10.1177/0265659012453463>
- Togram, B. (2015). How do families of children with Down syndrome perceive speech intelligibility in Turkey? *BioMed Research International*, 2015, 707134. <https://doi.org/10.1155/2015/707134>
- Tsao, R., & Kindelberger, C. (2009). Variability of cognitive development in children with Down syndrome: Relevance of good reasons for using the cluster procedure. *Research in Developmental Disabilities*, 30(3), 426–432.
- Uong, E. C., McDonough, M., Tayag-Kier, C. E., Zhao, H., Mahboubi, S., Schwab, R. J., ... Arens, R. (2001). Magnetic resonance imaging of the upper airway in children with Down syndrome. *American Journal of Respiratory & Critical Care Medicine*, 163(3), 731–736.
- van Bysterveldt, A. K., Gillon, G., & Foster-Cohen, S. (2010). Integrated speech and phonological awareness intervention for pre-school children with Down syndrome. *International Journal of Language & Communication Disorders*, 45(3), 320–335. <https://doi.org/10.3109/13682820903003514>
- Van Lierde, K., Galiwango, G., Hodges, A., Bettens, K., Luyten, A., & Vermeersch, H. (2012). Impact of tongue reduction on overall speech intelligibility, articulation and oromyofunctional behavior in 4 children with Beckwith-Wiedemann syndrome. *Folia Phoniatrica et Logopaedica*, 64, 55–63.
- Vilardell, M., Rasche, A., Thormann, A., Maschke-Dutz, E., Pérez-Jurado, L. A., Lehrach, H., & Herwig, R. (2011). Meta-analysis of heterogeneous Down syndrome data reveals consistent genome-wide dosage effects related to neurological processes. *BMC Genomics*, 12, 229.
- Whitworth, N., & Bray, M. (2015). Emergence of the vowel space in very young children with Down syndrome: An exploratory case study. In *ICPhS2015: Proceedings of the 18th International Congress of Phonetic Sciences*, August 10–14, 2015, SECC, Glasgow, Scotland, United Kingdom. Retrieved from <http://www.internationalphoneticassociation.org/icphs-proceedings/ICPhS2015/Papers/>