



Published in final edited form as:

J Am Chem Soc. 2018 May 09; 140(18): 5895–5898. doi:10.1021/jacs.8b02751.

Formation of Chiral Allylic Ethers via an Enantioselective Palladium-Catalyzed Alkenylation of Acyclic Enol Ethers


Harshkumar H. Patel, Matthew B. Prater, Scott O. Squire Jr., and Matthew S. Sigman*

Department of Chemistry, University of Utah, 315 South 1400 East, Salt Lake City, Utah 84112, United States

Abstract

This report details a palladium-catalyzed process to access highly functionalized optically active allylic aryl ethers. A number of electron-deficient alkenyl triflates underwent enantio- and site-selective coupling with acyclic aryl enol ethers in the presence of a chiral palladium catalyst. This transform provides chiral allylic ether products in high yields and excellent enantiomeric ratios furnishing a unique disconnection to incorporate heteroatoms at a stereocenter. Finally, the applicability of the products to target synthesis was demonstrated through the formation of a chiral allylic alcohol and the generation of a flavone-inspired product.

Graphical Abstract



Access to enantiomerically enriched aryl allyl ethers is of high synthetic utility as they are represented in numerous ether containing natural products and pharmaceuticals.¹ Substantial progress has been reported for their synthesis using transition-metal catalysis.² Most methods strategically rely on the reaction of symmetric π -allyl intermediates of Pd,³ Rh,⁴ and Ir⁵ using phenoxides or excess phenol as the trapping exogenous nucleophile. Alternatively, branched allylic ethers can be accessed in high enantioselectivity using a distinct Pd(II)-catalyzed S_N2' reaction of allylic trichloroacetamidates.⁶ In spite of these advances, the alkenyl components installed using these methods are relatively simple inspiring us to consider new reactions to obtain such products. Herein we describe a palladium-catalyzed redox-relay enantioselective Heck reaction of a new class of substrates,

Corresponding Author: sigman@chem.utah.edu.

Notes

The authors declare no competing financial interests.

Supporting Information

Experimental procedures and characterization data for new compounds. This material is available free of charge via the Internet at <http://pubs.acs.org>.

namely acyclic aryl enol ethers. As illustrated below, this reaction provides facile access to a range of diversely functionalized aryl allyl ethers, especially in terms of the alkenyl component, in high enantioselectivity.

In previous reports, the redox-relay Heck strategy has led to the successful enantioselective coupling of acyclic disubstituted alkenols with a range of aryl, alkenyl, and alkynyl groups (Figure 1a).⁷ The site-selectivity observed in the migratory insertion step is sensitive to the polarization of the alkene carbons in the transition state, which is proposed to be stabilized by the CO dipole of the alcohol moiety.⁸ Although the measured site-selectivity for allylic alcohols is generally high, it decreases substantially with the increasing chain-length between the alkene and the alcohol.⁹ To presumably further bias the migratory insertion improving site selectivity of more remote functionalization reactions, we envisioned the coupling of alkenyl electrophiles with acyclic enol ether substrates (Figure 1b). This is proposed to be due to increased nucleophilicity of carbon distal to the oxygen in the enol ether substrate (**2**, Figure 1c).¹⁰ Furthermore, high site-selectivity was anticipated irrespective of the number of carbons between the alkene and alcohol in the enol ether substrate. Successful implementation incorporates a heteroatom at the stereocenter formed in the migratory insertion process to access enantioenriched aryl allyl ether products (**3**).

At the outset, we were concerned that the migratory insertion event would be followed by β -alkoxy elimination instead of β -hydride elimination towards the alcohol (Figure 1d) as these processes could be competitive on the basis of previous efforts of our group.¹¹ Additionally, the β -alkoxy aldehyde products **3** ($n=0$) may not survive the relatively acidic conditions and undergo an E1cB to form the α,β -unsaturated carbonyl product.¹² Consistent with this, early experiments of enol ethers **2a** and **2b**, easily prepared through a two-step sequence from the propiolate ester,¹³ with the electron deficient vinylogous lactone triflate **1a** in the presence of a Pd⁽⁰⁾ catalyst and a chiral pyridine-oxazoline ligand **L1** (Figure 2a), exclusively led to the undesired anticipated E1cB product **4** in 70% yield. Considering the relatively Brønsted/Lewis acidic conditions of this process, it was hypothesized that lowering the basicity of the oxygen on the enol ether may slow the presumed E1cB process.

To test this hypothesis, an enol ether substrate **2c** containing an aryl substituent, instead of an alkyl group, was prepared and reacted with alkenyl triflate **1a** under the standard redox-relay Heck conditions (Figure 2b). The evaluation of the crude reaction mixture by ¹H NMR indicated the formation of the desired β -alkenylated aldehyde product **3c** in high yield (> 90% NMR yield). However, an attempt to purify the β -phenoxy aldehyde product **3c** via silica gel chromatography resulted in partial decomposition of the product and formation of product **4**, previously observed. Thus, for characterization purposes and ease of handling, the obtained aldehydes were transformed to the corresponding alcohols upon reduction of the crude mixture with sodium borohydride. This allowed the resultant alcohol **5a** to be isolated in 90% yield and an excellent enantiomeric ratio (er) of 99:1 (Table 1, 5a).

With the optimal conditions determined, we began the assessment of a variety of electron poor alkenyl triflates (Table 1). The cyclic five and seven-membered alkenyl triflates performed well in the redox-relay Heck reaction to generate **5b** and **5c** in high yields and high enantiomeric ratio of 98:2 and 94:6, respectively (see Supporting Information for the

determination of absolute configuration of compounds in Table 1, Table 2 and Table 3). The β -alkenylated product **5d** containing a protected amine was also formed in good yield and high enantioselectivity, although sodium tris(1,1,1,3,3,3-hexafluoroisopropoxy)borohydride¹⁴ was used instead of sodium borohydride during the reduction step. The relay Heck reaction also tolerated a hindered tetrasubstituted cyclohexenone triflate. Unfortunately, in this case, a selective reduction of the aldehyde in the presence of the enone proved to be challenging. Thus, over reduction afforded a diastereomeric mixture of diols (see Supporting Information), which was treated with MnO₂ to ultimately afford the chiral allylic ether **5e** in good yield and 93:7 er.¹⁵ Next, a variety of acyclic alkenyl triflates, conveniently prepared from β -ketoesters in a stereodefined manner, were subjected to the redox-relay Heck reaction.¹⁶ Both (*E*) and (*Z*)-alkenyl triflates were added to enol ether **2c**, forming **5f** and **5g** in high yield and enantiomeric ratio of 93:7 and 92:8, respectively. Additionally, acyclic alkenyl triflates having a benzyl substitution, a remotely positioned phthalimide, and an alkyl silane were successfully integrated to yield **5h–5j** in modest to good yields and generally good enantioselectivity. Importantly, the olefins incorporated in **5f–5j** did not undergo isomerization, thereby providing a means to generate stereodefined tetrasubstituted alkenes adjacent to a stereocenter. Lastly, a dihydrobenzofuran derived alkenyl triflate was also incorporated in **5k**, albeit in low yield but high enantioselectivity. It is noteworthy that di- and tri-substituted alkenyl triflates performed poorly in this reaction with low conversion of **2c** (see Supporting Information).¹⁷ Although, the precise cause for this difference in reactivity is unknown, it is assumed that the tetrasubstituted alkenyl triflates may slow β -hydride elimination to generate the diene.^{7b}

Next, we evaluated a range of electronically diverse aryl enol ethers. Electron-rich substituents on the phenol provided the desired allylic ether products **5l–5o** in good yields and high enantioselectivity (Table 2). The hindered ortho-substitution in **5m** showed no deleterious effects on either the yield or the enantioselectivity of the reaction. Interestingly, the use of an electron-poor group (for instance a *p*-Br substitution) on the phenyl ring of the acyclic enol ether did not undergo reaction under these conditions. Consequently, to tackle this issue, new conditions were identified empirically using THF as solvent, and ligand **L2** (-CF₃ substituted) instead of **L1**. Thereby, using the newly optimized reaction conditions, a number of enol ethers having an electron poor substituent on the phenyl ring, underwent the Heck reaction, and resulted in the formation of alcohols **5r–5t**, upon reduction of the aldehydes, in consistently high yield and enantioselectivity. The reason for such disparity in the reactivity of electronically different arenes is not currently understood.

As the redox-relay strategy is unique in establishing remote stereocenters, a range of aryl enol ether substrates having a different number of methylenes between the alkene and the alcohol were investigated (Table 3).¹³ Significantly, the use of a homoallylic alcohol may allow the isolation of aldehyde products directly, as the degradation via an E1cB process is not possible. Thus, the reaction of the cyclic vinylogous lactone triflate **1a** with a *p*-OMe-substituted aryl enol ether substrate **2u**, using slightly higher catalyst loadings provided the γ -alkenylated aldehyde product **3u** in 87% yield and 97:3 enantiomeric ratio. Similarly, a sesamol or phenol derived enol ether also provided the remotely alkenylated aldehyde products **3v** and **3w** in consistently high yield and enantioselectivity. The alkenyl triflate was

further installed at a more remote position δ from the aldehyde to form allylic ethers **3x** and **3y** in good yield and high enantioselectivity.

The use of a cleavable group in the acyclic enol ether substrate has the potential to deliver enantioenriched chiral allylic alcohols. To this end, a one electron oxidative deprotection of the *p*-methoxyphenyl (PMP) ether was accomplished by treating compound **6** with CAN (Scheme 1a).¹⁸ The desired allylic alcohol **6a** was obtained in 80% isolated yield with no loss of the stereochemical integrity (Scheme 1). In order to further demonstrate the synthetic applicability of such chiral allylic ethers, we envisioned accessing flavone derivatives containing an alkenyl group at the 2-position. To accomplish this, the β -alkenylated aldehyde **3c** generated after the Heck reaction was subjected to a sequential Pinnick oxidation/Friedel-Crafts cyclization¹⁹ affording the desired alkenylated flavone product **7** in 78% overall yield and 94:6 enantiomeric ratio, in single purification (Scheme 1b). A slight erosion of the stereochemical integrity was observed and found to be associated with the Pinnick oxidation step (see Supporting Information for details). Such a strategy allows effective incorporation of highly functionalized alkenyl groups into flavonoids, thereby expanding the chemical space, as most reported flavonoids contain an aryl group.²⁰

In summary, we have developed a unique method for the construction of highly functionalized chiral allylic ethers using a redox-relay Heck strategy. The process utilizes conveniently synthesized aryl enol ether substrates and readily accessible alkenyl triflates to generate chiral allylic ether products in good yield and enantioselectivity. The simple deprotection of the PMP ether provides direct access to chiral allylic alcohols in high enantiomeric ratio. Future work is focused on expanding this approach to other heteroatom containing alkenyl substrates.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

The work was supported by National Institute of Health (NIGMS R01GM063540) and the National Science Foundation REU Program for S.O.S. (CHE-1358740).

References

- (a) Trost BM, Crawley ML. *Chem Rev.* 2003; 103:2921. [PubMed: 12914486] (b) Cao B, Park H, Joullié MM. *J Am Chem Soc.* 2002; 124:520. [PubMed: 11804473] (c) Tanaka H, Sawayama AM, Wandless TJ. *J Am Chem Soc.* 2003; 125:6864. [PubMed: 12783528] (d) Mizuguchi E, Achiwa K. *Chem Pharm Bull.* 1997; 45:1209. (e) Pochetti G, Mitro N, Lavecchia A, Gilardi F, Besker N, Scotti E, Aschi M, Re N, Fracchiolla G, Laghezza A, Tortorella P, Montanari R, Novellino E, Mazza F, Crestani M, Loiodice F. *J Med Chem.* 2010; 53:4354. [PubMed: 20462215] (f) Nicolaou KC, Pfefferkorn JA, Roecker AJ, Cao GQ, Barluenga S, Mitchell HJ. *J Am Chem Soc.* 2000; 122:9939. (g) Ishibashi H, Ishihara K, Yamamoto H. *J Am Chem Soc.* 2004; 126:11122. [PubMed: 15355072]
- Zhang J, Wei YF, Luo RS. *J Org Chem Res.* 2017; 5:51.
- (a) Trost BM, Tsuji HC, Toste FD. *J Am Chem Soc.* 2000; 122:3534. (b) Trost BM, Toste FD. *J Am Chem Soc.* 1999; 121:4545. (c) Trost BM, Toste FD. *J Am Chem Soc.* 1998; 120:815. (d) Uozumi Y, Kimura M. *Tetrahedron: Asymmetry.* 2006; 17:161. (e) Trost BM, Machacek MR, Tsui HC. *J Am Chem Soc.* 2005; 127:7014. [PubMed: 15884945] (f) Trost BM, Crawley ML. *Chem Eur J.* 2004;

- 10:2237. [PubMed: 15112213] (g) Trost BM, Guzner JL, Dirat O, Rhee YH. *J Am Chem Soc.* 2002; 124:10396. [PubMed: 12197742]
4. (a) Li C, Breit B. *Chem Eur J.* 2016; 22:14655. [PubMed: 27620970] (b) Lysek R, Borsuk K, Furman B, Kaluza Z, Kazimierski A, Chmielewski M. *Curr Med Chem.* 2004; 11:1813. [PubMed: 15279570] (c) Evans PA, Leahy DK. *J Am Chem Soc.* 2000; 122:5012.
5. (a) Stanley LM, Bai C, Ueda M, Hartwig JF. *J Am Chem Soc.* 2010; 132:8918. [PubMed: 20552969] (b) Shekhar S, Trantow B, Leitner A, Hartwig JF. *J Am Chem Soc.* 2006; 128:11770. [PubMed: 16953612] (c) Leitner A, Shu C, Hartwig JF. *Org Lett.* 2005; 7:1093. [PubMed: 15760147] (d) Kiener CA, Incarvito C, Hartwig JF. *J Am Chem Soc.* 2003; 125:14272. [PubMed: 14624564] (e) Lopez F, Ohmura T, Hartwig JF. *J Am Chem Soc.* 2003; 125:3426. [PubMed: 12643693] (f) Kim D, Reddy S, Singh OV, Lee JS, Kong SB, Han H. *Org Lett.* 2013; 15:512. [PubMed: 23317334]
6. (a) Kirsch SF, Overman LE. *J Am Chem Soc.* 2005; 127:2866. [PubMed: 15740118] (b) Kirsch SF, Overman LE, White NS. *Org Lett.* 2007; 9:911. [PubMed: 17274626] (c) Cannon JS, Overman LE. *Acc Chem Res.* 2016; 49:2220. [PubMed: 27689745] (d) Cannon JS, Kirsch SF, Overman L, Sneddon HF. *J Am Chem Soc.* 2010; 132:15192. [PubMed: 20942424]
7. (a) Werner EW, Mei TS, Burckle AJ, Sigman MS. *Science.* 2012; 338:1455. [PubMed: 23239733] (b) Patel HH, Sigman MS. *J Am Chem Soc.* 2015; 137:3462. [PubMed: 25738548] (c) Patel HH, Sigman MS. *J Am Chem Soc.* 2016; 138:14226. [PubMed: 27768842] (d) Chen ZM, Nervig CS, Deluca RJ, Sigman MS. *Angew Chem, Int Ed.* 2017; 56:6651. (e) Zhang C, Santiago CB, Crawford JM, Sigman MS. *J Am Chem Soc.* 2015; 137:15668. [PubMed: 26624236]
8. (a) Xu L, Hilton MJ, Zhang X, Norrby PO, Wu YD, Sigman MS, Wiest O. *J Am Chem Soc.* 2014; 136:1960. [PubMed: 24410393] (b) Hilton MJ, Xu LP, Norrby PO, Wu YD, Wiest O, Sigman MS. *J Org Chem.* 2014; 79:11841. [PubMed: 25186804]
9. Mei TS, Werner EW, Burckle AJ, Sigman MS. *J Am Chem Soc.* 2013; 135:6830. [PubMed: 23607624]
10. (a) Ozawa F, Kobatake Y, Hayashi T. *Tetrahedron Lett.* 1993; 34:2505. (b) Kilroy TC, Cozzi PG, Nicole E, Patrick J. *Synthesis.* 2004; 11:1879. (c) Loiseleur O, Hayashi M, Schmees N, Pfaltz A. *Synthesis.* 1997; 11:1338. (d) Loiseleur O, Meier P, Pfaltz A. *Angew Chem, Int Ed Engl.* 1996; 35:200. (e) Gilbertson SR, Fu Z. *Org Lett.* 2001; 3:161. [PubMed: 11430024] (f) Gilbertson SR, Denov DG, Rheingold AL. *Org Lett.* 2000; 2:2885. [PubMed: 10964390] (g) Kaukoranta P, Kallstrom K, Andersson PG. *Adv Syn Catal.* 2007; 349:2595. (h) Tietze LF, Thede K, Sannicola F. *Chem Comm.* 1999:1811. (i) Hennessy AJ, Malone YM, Guiry PJ. *Tetrahedron Lett.* 2000; 41:2261.
11. Race NJ, Schwalm CS, Nakamuro T, Sigman MS. *J Am Chem Soc.* 2016; 138:15881. [PubMed: 27960316]
12. (a) Bertelsen S, Dinér P, Johansen RL, Jørgensen KA. *J Am Chem Soc.* 2007; 129:1536. [PubMed: 17284000] (b) Jiang H, Holub N, Jørgensen KA. *Proc Natl Acad Sci USA.* 2010; 107:20630. [PubMed: 20547884] (c) Sundén H, Ibrahim I, Zhao GL, Eriksson L, Córdova A. *Chem Eur J.* 2007; 13:574. [PubMed: 17039558]
13. Tellam JP, Kociok-Köhn G, Carbery DR. *Org Lett.* 2008; 10:5199. [PubMed: 18928292] . For more preparations of other enol ether substrates see: Ma D, Cai Q, Xie X. *Synlett.* 2005:1767. Kundu D, Maity P, Ranu BC. *Org Lett.* 2014; 16:1040. [PubMed: 24502267] Wan C, Jones CD, Koenig TM, Pu J, Mitchell D. *Tetrahedron Lett.* 2003; 44:8257. Willis MC, Taylor D, Gillmore AT. *Chem Comm.* 2003:2222. [PubMed: 13678211] Nordman G, Buchwald SL. *J Am Chem Soc.* 2003; 125:4978. [PubMed: 12708838]
14. Kuroiwa Y, Matsumura S, Toshima K. *Synlett.* 2008:2523.
15. Gritter RJ, Wallace TJ. *J Org Chem.* 1959; 24:1051.
16. Babinski D, Soltani O, Frantz DE. *Org Lett.* 2008; 10:2901. [PubMed: 18540630] . For alternative procedures see: Specklin S, Bertus P, Weibel JM, Pale P. *J Org Chem.* 2008; 73:7845. [PubMed: 18767806] Ritter K. *Synthesis.* 1993:735. Crisp GT, Meyer AGJ. *Org Chem.* 1992; 57:6972.
17. (a) Crouch IT, Neff RK, Frantz DE. *J Am Chem Soc.* 2013; 35:4970. (b) Crouch IT, Dreier T, Frantz DE. *Angew Chem, Int Ed.* 2011; 50:6128.
18. Qin D, Byun HS, Bittman R. *J Am Chem Soc.* 1999; 121:662.

19. (a) Zhao D, Beiring B, Glorius F. *Angew Chem, Int Ed.* 2013; 52:8454. (b) Vila C, Hornillos V, Fañanás-Mastral M, Feringa BL. *Chem Commun.* 2013; 49:5933.
20. Chahar MK, Sharma N, Dobhal MP, Joshi YC. *Pharmacog Rev.* 2011; 5:1.


Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript


a) Previously reported enantioselective redox-relay Heck reaction




R = aryl, alkenyl, alkynyl

♦ arylboronic acids ♦ alkenyl triflates ♦ alkynyl iodonium salts

b) Proposed enantioselective alkenylation of acyclic enol ethers



c) Mechanistic hypothesis



d) Anticipated issues





Figure 1.

(a) Previously reported enantioselective redox-relay Heck reaction. (b) Proposed enantioselective alkenylation of acyclic enol ether. (c) Mechanistic hypothesis. (d) Anticipated issues.

a) Palladium-catalyzed enantioselective alkenylation of alkyl enol ether



b) Palladium-catalyzed enantioselective alkenylation of aryl enol ether






Figure 2.

(a) Palladium-catalyzed enantioselective alkenylation of alkyl enol ether (b) Palladium-catalyzed enantioselective alkenylation of acyclic aryl enol ether.

a) PMP deprotection



b) Access to enantioenriched alkenylated flavone:




i. $\text{Pd}_2\text{dba}_3 \cdot \text{CHCl}_3$ (5 mol%), **L1** (12 mol%), 3 Å MS, DMF, N_2 , rt. ii. NaClO_2 , NaH_2PO_4 , 2-methyl-2-butene, $t\text{-BuOH}/\text{H}_2\text{O}$. iii. TFA/TFAA, CH_2Cl_2 .

Scheme 1.

Derivatization of alkenylated products.

Table 1

Scope of alkenyl triflates in the redox-relay Heck reaction of acyclic aryl enol ether.




Yields are reported as an average of two parallel experiment. Enantioselectivity determined by SFC equipped with a chiral column. Reaction performed on 0.5 mmol scale. Absolute configuration determined to be (*S*) for a synthetic derivative of **5z** (see Supporting Information) arising from the (*E*)-alkene. All other products were assigned by analogy.

^aNaBH(HFIP)₃ in HFIP was used for the reduction of aldehyde to alcohol.

^bMnO₂ oxidation of the diol yields the ketone **5e** (overall yield reported).

Table 2

Scope of allylic aryl enol ethers.




Yields are reported as an average of two parallel experiments. Enantioselectivity determined by SFC equipped with a chiral column. Reaction performed on 0.25 mmol scale. Absolute configuration determined to be (*S*) for a synthetic derivative of **5z** (see Supporting Information) arising from the (*E*)-alkene. All other products were assigned by analogy.

^aTHF and **L2** ligand used instead of DMF and **L1** ligand.

Table 3

Evaluation of aryl enol ether with varying chain lengths.



Yields are reported as an average of two parallel experiments. Enantioselectivity determined by SFC equipped with a chiral column. Reaction performed on 0.25 mmol scale. Absolute configuration determined to be (*S*) for a synthetic derivative of **5z** (see Supporting Information) arising from the (*E*)-alkene. All other products were assigned by analogy.