

Research article

In what extent anemia coexists with cognitive impairment in elderly: a cross-sectional study in Greece

Stella Argyriadou^{1,2}, Ioannis Vlachonikolis³, Haritini Melisopoulou¹, Kostas Katachanakis³ and Christos Lionis*²

Address: ¹Health Centre of Chrisoupolis (HCC), Chrissoupolis 64 200, Macedonia, Greece, ²Clinic of Social and Family, School of Medicine, University of Crete, PO Box 1393, Heraklion, Crete, Greece and ³Laboratory of Biostatistics, School of Medicine, University of Crete, PO Box 1393, Heraklion, Crete, Greece

E-mail: Stella Argyriadou - starge@otenet.gr; Ioannis Vlachonikolis - I.Vlachonikolis@surrey.ac.uk; Haritini Melisopoulou - starge@otenet.gr; Kostas Katachanakis - drknk@hotmail.com; Christos Lionis* - lionis@med.uoc.gr

*Corresponding author

Published: 25 October 2001

Received: 30 August 2001

BMC Family Practice 2001, 2:5

Accepted: 25 October 2001

This article is available from: <http://www.biomedcentral.com/1471-2296/2/5>

© 2001 Argyriadou et al; licensee BioMed Central Ltd. Verbatim copying and redistribution of this article are permitted in any medium for any non-commercial purpose, provided this notice is preserved along with the article's original URL. For commercial use, contact info@biomedcentral.com

Abstract

Background: A project aimed at studying the frequency of dementia and depression in the catchment area of the Health Centre of Chrissoupolis (HCCh), Northern Greece, was carried out. This paper reports the association between AD and anemia among the elderly participants in this Greek study.

Methods: Eligible participants were people 65 years or over who were (a) living in the Elderly People's Home (all 48 subjects included); (b) visiting the Open Center for Elderly People during a 20 workday period (75 subjects) and (c) visiting the HCCh for routine medical care. The Mini Mental State Examination (MMSE) was used in assessing the cognitive capacity of the participants. Blood was drawn for serum hematocrit, vitamin B₁₂ and folate determination.

Results: The prevalence proportions of possible cognitive impairment among anemic and non-anemic males were 55.6% and 34.4%, respectively ($\chi^2 = 5.8$, d.f. = 1, $p = 0.016$). The corresponding proportions in females were 47.5% and 40.1 % ($\chi^2 = 1.1$, d.f. = 1, $p = 0.305$). Using logistic regression analysis, age-group (≥ 80 yrs), type of Institute, vitamin B₁₂ and anemia had significant independent associations with possible cognitive impairment.

Conclusions: Anemia is a frequent finding in elderly and it may be a risk factor for dementia, but the extent of the associated deterioration of cognitive impairment or the relation with AD is not known. GPs should be aware of this coexistence and recommend for screening, assaying and treating elderly people.

Background

Alzheimer's disease (AD) and other neurodegenerative diseases affect large parts of the population worldwide and they constitute a serious public health problem for the aging European population [1–4]. General Practitioners seem to be able to diagnose early cognitive im-

pairments, after training, among their elderly patients [5,6]. Identifying reversible and avoidable cognitive impairments seems to be also a challenge for future General Practitioners (GPs). There are many controversies regarding various associated factors, which can affect the incidence and progress of Alzheimer's disease (AD) [7,8].

Among these factors, vitamin B12 deficiency and folate status deterioration have been extensively discussed. Although some studies reported that anemia might be a risk factor for AD in the elderly [9–12] this subject is still debated in the literature.

A project with the aim of studying the frequency of dementia and depression among the elderly people in the catchment area of the Chrisoupolis Health Center (HCCh) in Northern Greece has been carried out during the past few years. A high prevalence of cognitive impairment has recently been reported in this area [13]. Thus, it was interesting to study and discuss the laboratory findings of the eligible subjects together with their mental status ability. This paper reports the association between cognitive impairment and anemia among the elderly participants in this Greek study.

Patients and Methods

Setting

This cross-sectional study was carried out at the catchment area of the (HCCh) in Northern Greece, which comprises 22,000 inhabitants.

Participants

Eligible participants were people 65 years or over who were (a) living in the Elderly People's Home (all 48 subjects included); (b) visiting the Open Center for Elderly People during a 20 workday period (75 subjects) and (c) visiting the HCCh for routine medical care. The latter

group comprised of 413 subjects, which were randomly selected among the elderly people visiting one of the two outpatient clinics of the health center for the same period of time. Thus the total number of participants was 536, 245 males and 291 females. Baseline data of the studied population were presented in Table 1.

The study has been approved by the Scientific and Ethical Committee of the General Hospital of Kavala, Greece.

Tests

The Greek validated version of Mini Mental State Examination test (MMSE) was used [14]. For diagnostic classification of cognitive and mood disorders the criteria of ICD-10 were applied [15]. Any subject with an MMSE score of 24 or less was included in the possible cognitive impairment group for analyses.

Blood (10 cc) was drawn for mean hematocrit (HCT), vitamin B12 (Vit.B12) and serum folate concentrations and was assayed in the lab of the HCCh. HCT was defined with microHt method. Normal values were considered when HCT was ≥ 38% for males and ≥ 36% for females. Vit.B12 was defined with the Elisa method in the micro Merieux analyzer. Normal values for Vit.B12 were considered those higher or equal to 145 pg/mL. Folate was determined with ICIA method (Ion Capture Imune Assay), in Imx automatic analyzer. Normal values for folate were considered those higher or equal to 1.8 ng/mL. Quality control is applied at this lab on a regular basis.

Table 1: Demographic and backgrounds characteristics of the studied population

Variable	Elderly	Open	Chrissoupolis	Total
	People's Home	Centre for Elderly People	Health Centre	
	N (%)	N (%)	N (%)	N
Age				
65–74 years	-	73 (27.1)	196 (72.9)	269
75–84	32 (17.1)	2 (1.1)	153 (81.8)	187
85+	16 (20.0)	-	64 (80.0)	80
Total	48 (9.0)	75 (14.0)	413 (77.1)	536
Sex				
Males	19 (7.75)	72 (29.4)	154 (62.9)	245
Females	29 (10.0)	28 (9.6)	234 (80.5)	291
Marital status				
Married	-	71 (18.0)	324 (82.0)	395
Widowed	48 (34.04)	4 (2.8)	89 (63.1)	141
Education level				
<6 years education	17 (13.4)	9 (7.1)	101 (79.5)	127
6 years education	11 (3.5)	32 (10.2)	270 (86.3)	313
6–12 years education	20 (20.8)	34 (35.4)	42 (43.7)	96

Table 2: HCT, Vit.B₁₂ and Folate and cognitive capacity (number of people and prevalence proportions)

	MMSE > 24 N (%)	MMSE ≤ 24 N (%)	Total N (%)
HCT			
Normal*			
Male	137 (65.6)	72 (34.4)	209 (100)
Female	139 (59.9)	93 (40.1)	232 (100)
Total	276 (62.6)	165 (37.4)	441 (100)
Abnormal			
Male	16 (44.4)	20 (55.6)	36 (100)
Female	31 (52.5)	28 (47.5)	59 (100)
Total	47 (49.5)	48 (50.5)	95 (100)
Vit.B₁₂			
Normal			
Male	144 (65.2)	77 (34.8)	221 (100)
Female	154 (59.5)	105 (40.5)	259 (100)
Total	298 (62.1)	182 (37.9)	480 (100)
Abnormal			
Male	9 (37.5)	15 (62.5)	24 (100)
Female	16 (50)	16 (50)	32 (100)
Total	25 (44.6)	31 (55.4)	56 (100)
Folate			
Normal			
Male	149 (62.3)	90 (37.7)	239 (100)
Female	166 (58.9)	116 (41.1)	282 (100)
Total	315 (60.3)	206 (39.5)	521 (100)
Abnormal			
Male	4 (66.7)	2 (33.3)	6 (100)
Female	4 (44.4)	5 (55.6)	9 (100)
Total	8 (53.3)	7 (46.7)	15 (100)

* Normal value of HCT ≥ 38% in male and >36 in females

Statistical analysis

All statistical analyses were carried out using SPSS for Windows (Release 10.0). The associations between categorical variables, such as different subject groups and prevalence of disorders, were tested by means of Pearson's chi square-test or Fisher's test where was appropriate. To show the strength of the association, p-values are shown in the text [16]. Comparisons of mean values between groups were made by Student's t-test or analysis of variance. The independent influence of all potential factors in identifying the presence of cognitive impairment was studied by means of unconditional logistic regression analysis [17].

Results

Anemia and cognitive impairment

Ninety-five (17.7%) out of the 536 participants were found to be anemic. Of those, 47 scored higher than 24 in the MMSE and 48 lower or equal to 24 (possibly cognitive impairment) (Table 2). The association between

anemia and possible cognitive impairment was statistically significant ($X^2 = 5.6$, d.f. = 1, $p = 0.018$). A separate analysis by gender revealed that this association was true only for males and not for females. In fact, in males the prevalence proportions of possible cognitive impairment among anemic and non-anemic subjects were 55.6% and 34.4%, respectively ($X^2 = 5.8$, d.f. = 1, $p = 0.016$). The corresponding proportions in females were 47.5% and 40.1% ($X^2 = 1.1$, d.f. = 1, $p = 0.305$).

Vitamin B₁₂ and cognitive impairment

The analysis of blood sample of the participants indicated an abnormal value of Vit.B₁₂ in 56 (10.4%) subjects. Of those, 25 subjects scored as normal in the cognitive test and 31 were assessed as possibly cognitive impaired. The prevalence proportions of possible cognitive impairment among male participants with normal and abnormal value of Vit.B₁₂ were significantly different, 34.8% and 62.5%, respectively ($X^2 = 7.1$, d.f. = 1, $p = 0.008$). The corresponding proportions in females were not significantly different, 40.5% and 50.0% ($X^2 = 1.0$, d.f. = 1, $p = 0.306$).

Folate and cognitive impairment

Fifteen (2.7%) subjects (7 with a MMSE lower or equal to 24) were found to have an abnormal value of folic acid. The prevalence proportions of possible cognitive impairment among participants with normal or abnormal value of folic acid were 39.5% and 46.7%, respectively. The association between folic acid and possible cognitive impairment was not significant ($X^2 = 0.3$, d.f. = 1, $p = 0.578$).

Differences among the three institutions

A two-way analysis of variance, by MMSE classification and institution, revealed significant differences between the three institutions in HCT and folate ($F = 4.2$, d.f. = 2 and 530, $p = 0.016$ and $F = 4.9$, d.f. = 2 and 530, $p = 0.008$, respectively), and marked, but not significant differences in Vit.B₁₂ ($F = 2.7$, d.f. = 2 and 530, $p = 0.068$). The results depicted in Table 3, show that subjects living in the Old People's Home (OPH), had the lowest value of HCT and Vit.B₁₂, while those participating in the Open Centre for the Elderly people (OCE) had the lowest value in folate. There were similar results with respect to differences between the two MMSE categories: participants with possible cognitive impairment had lower values in HCT, Vit.B₁₂ and folate, than those with normal MMSE score. The results in Table 3, show that the above results for the MMSE categories were not consistent across all three institutions, in particular those for Vit.B₁₂ and folate, where a reverse difference was observed in subjects from the OPH. This MMSE classification by institution interaction was significant only for folate ($F = 9.1$, d.f. = 2 and 530, $p < 0.001$).

Table 3: Mean values (and standard deviations) of HCT, Vit.B₁₂ and Folate in different places

Variable	HCCh M.A.(SD)	OCE M.A.(SD)	OPH M.A.(SD)	Total M.A.(SD)
Hct (%)				
MMSE > 24	40.1 (3.3)	40.1 (3.6)	39.5(3.7)	40.1 (3.3)
MMSE ≤ 24	39.6 (4.7)	36.5 (6.9)	37.9 (4.3)	39.1 (4.8)
Total	39.9(3.8)	39.5 (4.5)	38.2 (4.2)	39.7 (4)
Vit.B₁₂ (pg/mL)				
MMSE > 24	463.5 (197.6)	465.0(261.9)	377.2 (207.6)	460.9(211.6)
MMSE ≤ 24	438.3 (203.7)	330.9 (242.2)	434.3 (294.9)	431.1 (224.6)
Total	453.6 (200.2)	441.7(262.1)	421.2(276.3)	449 (217.1)
Folate (ng/mL)				
MMSE > 24	9.3 (3.3)	11.3(5.6)	8.5 (7.8)	9.6(4.1)
MMSE ≤ 24	8.0 (3.7)	7.9 (5.9)	13.4(8.5)	8.9 (5.4)
Total	8.8 (3.5)	10.7(5.8)	12.3(8.5)	9.4 (4.7)

Table 4: Results of the unconditional regression analysis

Variable	B	SE	P-value	Odds ratio	95% CI
Age ^a					
Age ^b	0.360	0.250	0.150	1.4	0.9–2.3
Age ^c	1.123	0.256	0.001	3.1	1.9–5.1
Place					
Group					
Group ^d	-0.854	0.349	0.014	0.4	0.2–0.8
Group ^f	1.513	0.407	0.001	4.5	2.0–10.1
Bl ₂ ^g	0.714	0.338	0.034	2.0	1.1–4.0
Anemia ^g	0.535	0.262	0.041	1.7	1.02–2.9
Folate ^g	1.323	0.714	0.064	3.8	0.9–15.2

a: 64–70 yrs b: 71–79 c: ≥ 80 d: Health Centre e: Open Centre for Elderly f: Old Peoples Home g: Abnormal value

Logistic regression analysis

The results of the logistic regression analysis (backward, unconditional maximum likelihood) are illustrated in Table 4. Age-group (≥ 80 yrs), type of institution, anemia, Vit.B₁₂ and folate concentrations had marked or significant independent influences on the presence of cognitive impairment, as indicated by MMSE score less than the cutoff point of ≤ 24. Table 4, shows also the odds – ratios and 95% CI for each of the above factors; it can be seen that the risk of cognitive impairment (as measured by the odds-ratio) for age greater than 80 years, OPH residence and abnormal value of HCT, Vit.B₁₂ is significantly higher than one.

Discussion

It is widely accepted that anemia is a common health problem among elderly people, and cobalamin deficiency increases with advancing age. Although there are several studies which report an association between cobalamin deficiency and cognitive impairment [18–21], more studies are required to determine whether there is a specific profile of cognitive impairments typical of Vit.B₁₂ deficiency. It is also widely accepted that cobalamin deficiency may result in a variety of atypical symptoms and hematological changes typical of megaloblastic anemia are frequently absent in the majority of patients with mental disorders [22].

Our study indicated an association between cognitive impairment and anemia, but causation cannot be as-

sessed on the basis of the results of this cross-sectional study. Why this association exists needs further discussion. The impact which the changes on olfactory function and taste associated with AD may have on dietary intake, nutritional status and the haematinic factors is not examined in this study. The possible contribution of the concomitant diseases on the causation of anemia should not be ignored, when the results are interpreted. The present study is based on people who have visited the Health Centre and other older people who live at the community setting. This situation reflects the current figure of the real context in which the GPs are invited to work. Anemia was also based on the HCT findings, but both HCT and the blood Hb concentration remain the best available indirect indicators [23,24]. The validity and reliability of the used laboratory methods is an important issue as it can affect the study results. However the lab of the HCCh was using quality controls and all the blood tests were administered by the same assistant. The use of MMSE as a screening tool has also been discussed widely in the literature [25,26]. Even the MMSE's specificity and sensitivity can be affected by factors such as sex, education level, age [27] this tool remains to be a simple screening tool at GP's disposal for cognitive impairment and cannot be used to make even tentative psychiatric diagnoses [26,27].

Despite these limitations, our study adds to the existing literature. This study also reveals that people, who are living in Old People's Homes, were recognized at a specific risk and it is also in accordance with other reports [28,29]. The greater risk of male subjects having cognitive impairment when they are anemic in comparison with females has been reported in another study [9].

This Greek study may have some serious implications on the GPs attitude and practices when they identify elderly cognitive impaired subjects; specifically, they could recommend them a hematological investigation including HCT, CBC, Vit.B₁₂ and folic acid levels. The initiation of a adequate treatment with Vit.B₁₂ and/or folate when laboratory indications exist, seems to be an effective strategy in General Practice and this treatment is recommended, even in cases with a normal score with MMSE. Some authorities recommend that serum B₁₂ screening should be initially performed to all elderly patients, with further testing in those with a Vit.B₁₂ level of less than 350 pg/ mL (260 pmol / L) [20]. Seeking the effectiveness of different screening strategies, clinicians need to use their own judgment to decide how to identify Vit.B₁₂ deficiency before it leads to anemia or neurologic disease.

Conclusions

Anemia seems to be a risk factor for cognitive impairment. There are also many questions whether Vit.B₁₂ and folate deficiency work as risk factors on the onset of dementia and whether the appearance of dementia can be avoided by a supplemented treatment. Until definitive clarifying of those questions GPs should be warranted to screening, assaying, and treating elderly people.

Competing interests

None declared

References

- Jorm AF, Korten AE, Henderson AS: **The prevalence of dementia: a quantitative integration of the literature. Application of a new method for making projections.** *Acta Psychiatr Scand* 1987, **78**:493-500
- Olafsdottir M, Marcusson J, Skoog I: **Mental disorders among elderly people in primary care: The Linköping study.** *Acta Psychiatr Scand* 2001, **104**:12-18
- Kalaria RN, Ballard C: **Stroke and cognition.** *Curr Atheroscler Rep* 2001, **3**:334-339
- Honig LS, Mayeux R: **Natural history of Alzheimer's disease.** *Ageing* 2001, **13**:171-182
- Mant A, Eylan EA, Pond DC, Saunders NA, Chancellor AH: **Recognition of dementia in general practice: comparison of general practitioners' opinions with assessments using the minimal state examination and the Blessed dementia rating scale.** *Fam Pract* 1998, **5**:184-188
- Lionis C, Tzagournissakis M, Iatraki E, Kozyraki M, Antonakis N, Platakis A: **Are Primary Care Physicians Able to Assess Dementia? An estimation of their capacity after a short-term training program in rural Crete.** *Am J Geriatr Psychiatry* 2001, **9**:315
- Fylkesnes SI, Nygaard HA: **Dementia and hypothyroidism.** *Tidsskr Nor Lægeforen* 2000, **120**:905-907
- Burke D, Sengoz A, Schwartz R: **Potentially reversible cognitive impairment in patients presenting to a memory disorders clinic.** *J Clin Neurosci* 2000, **7**:120-123
- Beard CM, Kokmen E, O'Brien PC, Ania BJ, Melton LJ 3rd: **Risk of Alzheimer's disease among elderly patients with anemia: population - based investigations in Olmsted County, Minnesota.** *Ann Epidemiol* 1997, **7**:219-224
- Rabins P: **Pernicious anemia and reversible dementia: Strachan and Henderson 30 years later.** *Int J Geriatr Psychiatry* 1998, **13**:139-140
- Jacques PF, Selhub J, Bostom AG, Wilson PW, Rosenberg IH: **The effect of folic acid fortification on plasma folate and total homocysteine concentrations.** *N Engl J Med* 1999, **340**:1449-1454
- Healton EB, Savage DG, Brust JC, Garrett TJ, Lindenbaum J: **Neurologic aspects of cobalamin deficiency.** *Medicine* 1991, **70**:229-245
- Argyriadou S, Melissopoulou H, Krania E, Karagiannidou A, Vlachonicolis J, Lionis C: **Dementia and depression: two frequent disorders of the aged in primary health care in Greece.** *Fam Practice* 2001, **18**:87-91
- Fountoulakis C, Tsolaki M, Chazi H, Kazis , Mini Mental State Examination (MMSE): **A validation study in demented patients from the elderly Greek population.** *Encephalos (abstract in English)* 1994, **31**:93-102
- World Health Organization: **The ICD-10 classification of mental and behavioral disorders: clinical descriptions and diagnostic guidelines.** Translated in Greek by Stefanis K, Soldatos K, Mavreas V (BETA Publications) Athens 1993
- Armitage P, Berry G: **Statistical Methods in Medical Research.** 2nd ed. Oxford: Blackwell Scientific Publications 1987
- Collett D: **Modeling Binary Data.** London: Chapman & Hall 1991
- Milward EA, Grayson DA, Creasey H, Janu MR, Brooks WS, Broe GA: **Evidence for association of anemia with vascular dementia.** *Neuroreport* 1999, **10**:2377-2381

19. Carmel R, Gott PS, Waters CH, Cairo K, Green R, Bondareff W, et al: **The frequently low cobalamin levels in dementia usually signify treatable metabolic, neurologic and electrophysiologic abnormalities.** *Eur J Haematol* 1995, **54**:245-253
20. Douglas LS: **Anemia in the elderly.** *Am Fam Physician* 2000, **62**:1565-1572
21. Bopp-Kistler I, Ruegger-Frey B, Grob D, Six P: **Vitamin B12 deficiency in geriatrics.** *Schwetz Med Prax* 1999, **88**:1867-1875
22. Larner AJ, Janssen JC, Cipolotti L, Rossor MN: **Cognitive profile in dementia associated with vitamin B12 deficiency due to pernicious anemia.** *J Neurol* 1999, **246**:317-319
23. Walsh JR: **Hematologic problems.** In: *Geriatric Medicine (Edited by Cassel CK, et al).* New York. N.Y.: Springer 1997:627-636
24. Mock DM, Bell EF, Lankford GL, Widness JA: **Hematocrit correlates well with circulating red blood cell volume in very low birth weight infants.** *Pediatr Res* 2001, **50**:525-531
25. Weissgarten J, Modai D, Oz D, Chen Levy Z, Cohn M, Marcus O, et al: **Vitamin B (6) therapy does not improve hematocrit in hemodialysis patients supplemented with iron and erythropoietin.** *Nephron* 2001, **87**:328-332
26. Wind AW, Schellevis FG, Van Staveren G, Scholten RP, Jonker C, et al: **Limitations of the Mini - Mental State Examination in diagnosing dementia in general practice.** *Int J Geriatr Psychiatry* 1997, **12**:101-108
27. O'Connor DW, Pollitt PA, Hyde JB, Fellows JL, Miller ND, et al: **The reliability and validity of the Mini-Mental State in a British community survey.** *J Psychiatr Res* 1989, **23**:87-96
28. O'Connor DW, Pollitt PA, Treasure FP, Brook CPB, Reiss BB: **The influence of education, social class and sex on Mini-Mental State scores.** *Psychol* 1989, **19**:771-776
29. Essama-Tatjani JC, Guillard JC, Potier de Courcy G, Fuchs F, Richard D: **Folate status worsens in recently institutionalized elderly people without evidence of functional deterioration.** *J Am Coll Nutr* 2000, **19**:392-404
30. Weyerer S, Mann AH, Ames D: **Prevalence of depression and dementia in residents of old age homes in Mannheim and Camden.** *Z Gerontol Geriatr* 1995, **28**:169-178

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMedcentral will be the most significant development for disseminating the results of biomedical research in our lifetime."

Paul Nurse, Director-General, Imperial Cancer Research Fund

Publish with **BMC** and your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours - you keep the copyright



BioMedcentral.com

Submit your manuscript here:

<http://www.biomedcentral.com/manuscript/>

editorial@biomedcentral.com