



Draft Genome Sequences of *Bacillus megaterium* KU143, *Microbacterium testaceum* KU313, and *Pseudomonas protegens* AS15, Isolated from Stored Rice Grains

Jin-Ju Jeong,^a Hyeon-Ji Moon,^a Duleepa Pathiraja,^b Byeonghyeok Park,^b In-Geol Choi,^b  Ki Deok Kim^a

^aLaboratory of Plant Disease and Biocontrol, Department of Biosystems and Biotechnology, Korea University, Seoul, South Korea

^bDepartment of Biotechnology, Korea University, Seoul, South Korea

ABSTRACT *Bacillus megaterium* KU143, *Microbacterium testaceum* KU313, and *Pseudomonas protegens* AS15 from stored rice grains exhibited antifungal activity against *Aspergillus* and *Penicillium* spp. predominant in stored rice. Here, we report their bacterial draft genomes, which contain genes related to biotic and abiotic stress management, as well as antimicrobial and insecticidal traits.

Previously, 460 bacterial strains were isolated from stored rice grains obtained from rice-processing complexes from 11 different regions in South Korea (1, 2). Among them, *Bacillus megaterium* KU143, *Microbacterium testaceum* KU313, and *Pseudomonas protegens* AS15 were selected as potential biocontrol agents (3–5). These bacterial strains showed significant biocontrol activity against *Aspergillus flavus* growth and aflatoxin production as well as the predominant fungi (*A. candidus*, *A. fumigatus*, *Penicillium fellutanum*, and *Penicillium islandicum*) in stored rice grains (3–5). Furthermore, *B. megaterium* KU143 and *P. protegens* AS15 produced volatile compounds that could reduce conidial germination and germ tube elongation, as well as mycelial growth of the predominant fungi (4, 5). Here, we present the draft genome sequences of these strains, KU143, KU313, and AS15, belonging to different genera and exhibiting significant biocontrol activity against rice fungal contamination.

Genome sequencing of bacterial strains KU143, KU313, and AS15 was performed using the Illumina MiSeq platform at the Computational and Synthetic Biology Laboratory, Korea University (Seoul, South Korea). Totals of 336,941, 955,850, and 531,468 paired-end reads (40.3-, 158.3-, and 47.2-fold coverage) for KU143, KU313, and AS15, respectively, were generated from paired-end sequencing of the genomic library with an average insert size of 500 bp. Low-quality reads were trimmed with a quality threshold of Q20, and the trimmed reads were subjected to *de novo* assembly using the SPAdes assembler (6). The reads were assembled to 98, 79, and 52 scaffolds for KU143, KU313, and AS15, respectively, with total lengths and G+C contents as shown in Table 1. The maximum lengths and N_{50} values of the contigs were 542,591 and 192,911 bp for KU143, 768,312 and 756,039 bp for KU313, and 1,020,915 and 290,736 bp for AS15, respectively. The genomes were annotated with the NCBI Prokaryotic Genome Annotation Pipeline (PGAP). The 4,995, 3,298, and 6,040 coding sequences of KU143, KU313, and AS15 showed 48.94, 29.46, and 60.18% sequence similarities to known genes in the NCBI database, respectively. In addition, retrieved numbers of tRNA, 5S rRNA, 16S rRNA, and 23S rRNA sequences of the strains are shown in Table 1.

In general, the genomes of *B. megaterium* KU143, *M. testaceum* KU313, and *P. protegens* AS15 contain genes related to biocontrol traits (i.e., siderophore, polyketide, phosphate solubilization, motility, and biofilm formation) that may be significant factors in the host colonization and protection against pathogenic infections (7–10).

Received 21 April 2018 Accepted 24 April 2018 Published 31 May 2018

Citation Jeong J-J, Moon H-J, Pathiraja D, Park B, Choi I-G, Kim KD. 2018. Draft genome sequences of *Bacillus megaterium* KU143, *Microbacterium testaceum* KU313, and *Pseudomonas protegens* AS15, isolated from stored rice grains. Genome Announcements 6:e00468-18. <https://doi.org/10.1128/genomeA.00468-18>.

Copyright © 2018 Jeong et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Ki Deok Kim, kidkim@korea.ac.kr.

TABLE 1 Summary of genome sequencing and GenBank accession and version numbers of *B. megaterium* KU143, *M. testaceum* KU313, and *P. protegens* AS15

Bacterial strain	Genome size (bp)	G+C content (%)	No. of coding sequences	No. of tRNAs	No. of rRNAs			GenBank accession no.	Version no.
					5S	16S	23S		
<i>B. megaterium</i> KU143	5,022,643	38.04	4,995	102	15	22	20	POTF00000000	POTF01000000
<i>M. testaceum</i> KU313	3,623,185	69.51	3,298	46	4	1	2	PPEE00000000	PPEE01000000
<i>P. protegens</i> AS15	6,756,833	63.52	6,040	60	7	1	1	PPEF00000000	PPEF01000000

Additionally, biotic and abiotic stress management genes (i.e., superoxide dismutase and catalase) were found in all the examined bacterial strains (11, 12). In particular, the genome of strain AS15 has several genes related to antimicrobial or insecticidal compounds (i.e., hydrogen cyanide synthase, ATP-dependent zinc metalloprotease, and chitinase) (13–15). Therefore, the genomes of these bacterial strains belonging to different genera may help to increase our understanding of the characteristics of their biocontrol of fungal contamination in stored rice grains.

Accession number(s). This whole-genome shotgun project has been deposited in DDBJ/EMBL/GenBank under the accession and version numbers listed in Table 1.

ACKNOWLEDGMENT

Jin-Ju Jeong was supported by the Global PhD Program through the National Research Foundation of Korea funded by the Ministry of Education (2015-034526), South Korea.

REFERENCES

- Oh JY, Jee SN, Nam Y, Lee H, Ryou MI, Kim KD. 2007. Populations of fungi and bacteria associated with samples of stored rice in Korea. *Mycobiology* 35:36–38. <https://doi.org/10.4489/MYCO.2007.35.1.036>.
- Lee SY, Oh JY, Ryou MI, Kim KD. 2007. Biological control of the rice storage fungi *Aspergillus* and *Penicillium* species by antagonistic bacteria originated from rice. *Plant Pathol J* 23:328.
- Mannaa M, Oh JY, Kim KD. 2017. Microbe-mediated control of *Aspergillus flavus* in stored rice grains with a focus on aflatoxin inhibition and biodegradation. *Ann Appl Biol* 171:376–392. <https://doi.org/10.1111/aab.12381>.
- Mannaa M, Oh JY, Kim KD. 2017. Biocontrol activity of volatile-producing *Bacillus megaterium* and *Pseudomonas protegens* against *Aspergillus flavus* and aflatoxin production on stored rice grains. *Mycobiology* 45: 213–219. <https://doi.org/10.5941/MYCO.2017.45.3.213>.
- Mannaa M, Kim KD. 2018. Biocontrol activity of volatile-producing *Bacillus megaterium* and *Pseudomonas protegens* against *Aspergillus* and *Penicillium* spp. predominant in stored rice grains: study II. *Mycobiology* 46:52–63. <https://doi.org/10.1080/12298093.2018.1454015>.
- Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko SI, Pham S, Prjibelski AD, Pyshkin AV, Sirotkin AV, Vyahhi N, Tesler G, Alekseyev MA, Pevzner PA. 2012. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *J Comput Biol* 19:455–477. <https://doi.org/10.1089/cmb.2012.0021>.
- Arasu MV, Duraipandiyar V, Ignacimuthu S. 2013. Antibacterial and antifungal activities of polyketide metabolite from marine *Streptomyces* sp. AP-123 and its cytotoxic effect. *Chemosphere* 90:479–487. <https://doi.org/10.1016/j.chemosphere.2012.08.006>.
- Khamna S, Yokota A, Lumyong S. 2009. Actinomycetes isolated from medicinal plant rhizosphere soils: diversity and screening of antifungal compounds, indole-3-acetic acid and siderophore production. *World J Microbiol Biotechnol* 25:649. <https://doi.org/10.1007/s11274-008-9933-x>.
- Sang MK, Kim KD. 2012. The volatile-producing *Flavobacterium johnsoniae* strain GSE09 shows biocontrol activity against *Phytophthora capsici* in pepper. *J Appl Microbiol* 113:383–398. <https://doi.org/10.1111/j.1365-2672.2012.05330.x>.
- Sang MK, Jeong JJ, Kim J, Kim KD. 2018. Growth promotion and root colonisation in pepper plants by phosphate-solubilising *Chryseobacterium* sp. strain ISE14 that suppresses *Phytophthora* blight. *Ann Appl Biol* 172:208–223. <https://doi.org/10.1111/aab.12413>.
- Gill SS, Anjum NA, Gill R, Yadav S, Hasanuzzaman M, Fujita M, Mishra R, Sabat SC, Tuteja N. 2015. Superoxide dismutase—mentor of abiotic stress tolerance in crop plants. *Environ Sci Pollut Res* 22:10375–10394. <https://doi.org/10.1007/s11356-015-4532-5>.
- Singh S, Prasad SM. 2014. Growth, photosynthesis and oxidative responses of *Solanum melongena* L. seedlings to cadmium stress: mechanism of toxicity amelioration by kinetin. *Sci Hortic* 176:1–10. <https://doi.org/10.1016/j.scienta.2014.06.022>.
- Jang JY, Yang SY, Kim YC, Lee CW, Park MS, Kim JC, Kim IS. 2013. Identification of orfamide A as an insecticidal metabolite produced by *Pseudomonas protegens* F6. *J Agric Food Chem* 61:6786–6791. <https://doi.org/10.1021/jf401218w>.
- Flury P, Aellen N, Ruffner B, Péchy-Tarr M, Fataar S, Metla Z, Dominguez-Ferreras A, Bloemberg G, Frey J, Goesmann A, Raaijmakers JM, Duffy B, Höfte M, Blom J, Smits THM, Keel C, Maurhofer M. 2016. Insect pathogenicity in plant-beneficial pseudomonads: phylogenetic distribution and comparative genomics. *ISME J* 10:2527–2542. <https://doi.org/10.1038/ismej.2016.5>.
- Flury P, Vesga P, Péchy-Tarr M, Aellen N, Dennert F, Hofer N, Kupferschmid KP, Kupferschmid P, Metla Z, Ma Z, Siegfried S, de Weert S, Bloemberg G, Höfte M, Keel CJ, Maurhofer M. 2017. Antimicrobial and insecticidal: cyclic lipopeptides and hydrogen cyanide produced by plant-beneficial *Pseudomonas* strains CHA0, CMR12a, and PCL1391 contribute to insect killing. *Front Microbiol* 8:100. <https://doi.org/10.3389/fmicb.2017.00100>.