



## Data Article

# Data for distribution of various species of fecal coliforms in urban, rural and private drinking water sources in ten years period – A case study: Kermanshah, Iran

Reza Davoodi<sup>a</sup>, Meghdad Pirsahab<sup>a</sup>, Kamaladdin Karimyan<sup>b,c</sup>, Vinod Kumar Gupta<sup>d</sup>, Ali Reza Takhtshahi<sup>e</sup>, Hooshmand Sharafi<sup>f</sup>, Masoud Moradi<sup>a,\*</sup>

<sup>a</sup> Research Center for Environmental Determinants of Health, Kermanshah University of Medical Sciences, Kermanshah, Iran

<sup>b</sup> Environmental Health Research Center, Kurdistan University of Medical Sciences, Sanandaj, Iran

<sup>c</sup> Department of Environmental Health Engineering, School of Public Health, Tehran University of Medical Sciences, Tehran, Iran

<sup>d</sup> Department of Applied Chemistry, University of Johannesburg, Johannesburg, South Africa

<sup>e</sup> Water and Wastewater Company of Kermanshah Province, Kermanshah, Iran

<sup>f</sup> Student Research Committee, Kermanshah University of Medical Sciences, Kermanshah, Iran

## ARTICLE INFO

### Article history:

Received 20 January 2018

Received in revised form

12 April 2018

Accepted 16 April 2018

Available online 23 April 2018

### Keywords:

Fecal coliforms

Distribution

Drinking water sources

Kermanshah

## ABSTRACT

This study was aimed to investigate the distribution of various species of fecal coliform in urban, rural and private drinking water sources of Kermanshah, in the west of Iran. For this study, data of ten years period (2006–2016) assessments of microbial quality regarding various species of *Fecal coliforms* was taken from health centers associated with urban, rural and private resources of Kermanshah city. A total number of 8643 samples were taken, 1851 samples from rural, 365 from urban and 4834 from private resources. The results showed that *Fecal coliforms*, *Escherichia coli* (*E. coli*) had the widest distribution in all urban, rural and private water resources (22.3%, 45.9% and 34%, respectively). Moreover, *E. coli* (47.5%) and *Klebsiella* (0.4%) had, respectively, the highest and lowest distribution in all months considered. Based on the results,

\* Corresponding author.

E-mail address: [Mahfooz60@gmail.com](mailto:Mahfooz60@gmail.com) (M. Moradi).

*E.coli* exists mostly in water resources; it is therefore of particular importance in the monitoring of water resources.

© 2018 Published by Elsevier Inc. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

## Specifications Table

Subject area	Environmental Sciences
More specific subject area	Environmental Health
Type of data	Tables and figures
How data was acquired	For this study, data of the 10 years period (2006–2016) assessments of the microbial quality in terms of various species of fecal coliforms was taken from health centers associated with urban, rural and private sources of Kermanshah city.
Data format	Analyzed
Experimental factors	The presence of some coliforms indicate fecal contamination. The term "IMVIC" is an acronym for each of these tests. "I" is for indole test; "M" is for methyl red test; "V" is for Voges–Proskauer test, and "C" is for citrate test. The IMVIC test was not performed in about 3% of the all samples. Therefore, the results of these samples were not included in the study.
Experimental features	The above parameters that mentioned in abstract part, were analyzed according to the standards for water and wastewater treatment handbook.
Data source location	Kermanshah city, Iran
Data accessibility	Data are included in this article

## Value of the data

- Monitoring the quality of drinking water resources especially in term of microbial quality is necessary [1–7], due to the variety and wide range of drinking water sources in the Kermanshah city.
- The obtained data can assist in identifying contaminated resources and planning to adopt a long and short-term strategy for safe water supply.
- The obtained data revealed that rural water resources had more distribution of the various fecal coliform species particularly *E. coli* due to lack of water resources protection, poor sanitation, and improper disinfection and as well as more exposed to environmental pollutants.
- The data of present study show that *Enterobacter aerogenes*, *Enterobacter agglomerans*, and *Klebsiella* were more abundant species in the cold months, while *Citrobacter freundii*, *E. coli* and *Enterobacter cloacea* were more abundant species in the warm months.

## 1. Data

Table 1 shows the distribution of fecal coliforms in urban, rural and private water resources of Kermanshah city based on IMVIC test. *E. coli* (22.3%) and *Klebsiella* (2%) were the most and least bacteria existent in urban water resources, respectively. In rural water sources, *Escherichia coli* (45.9%) and *Enterobacter cloacea* (2.6%) and in private sources *E. coli* (34%) and *Klebsiella* (1.3%) had the most and least existent, respectively.

**Table 1**

Distribution of fecal coliforms based on different sources of drinking water.

Bacteria	Distribution (%)		
	Urban sources	Rural sources	Private sources
<i>Citrobacter freundii</i>	14	13.7	14.8
<i>Escherichia coli</i>	22.3	45.9	34
<i>Enterobacter aerogenes</i>	3.3	7.6	6.3
<i>Enterobacter agglomerans</i>	2.5	8.7	4.1
<i>Enterobacter cloacea</i>	4.3	2.6	2.5
<i>Klebsiella</i>	2	3.8	1.3
No Fecal coliform	51.6	17.7	37
Total	100	100	100
Number of samples	1851	4834	2001

Tables 2 and 3 present fecal coliforms distribution concerning months of the year and the amount of residual chlorine in water sources of the city. The results showed that *E. coli* with an average of 38.1% and *Klebsiella* with 2.8% had the highest and lowest distributions, respectively. The results also showed a significant decrease of distribution of fecal coliforms with increasing residual chlorine, while a decreasing trend is observed from the dose of 0.8 mg/L.

## 2. Materials and methods

### 2.1. Study area

Kermanshah city with 24,500 m<sup>2</sup> area and an altitude of 1200 m above sea level, locates in 47° and 4" east and 19° and 34" north in the west of Iran (Fig. 1). The city has a population over 1,945,227 people. Kermanshah has a population of more than a million people, that the main source of drinking water provided from underground water supplies [8–13]. The above-mentioned water sources are threatened by various sources of pollution (especially urban, industrial and hospital wastewater) [14–19].

### 2.2. Measurement and data collection

For this study, data of the ten years period (2006–2016) assessments of the microbial quality in terms of various species of *Fecal coliforms* was taken from health centers associated with urban, rural and private sources of Kermanshah city. Given that the study was conducted based on census, all results of measuring samples during the 10 years (8643 samples) were analyzed. The number of samples in rural, urban and private resources was 1851, 365 and 4834, respectively. The method to identify various species of fecal coliforms was according to standard methods [20–27]. It is necessary to clarify that the IMVIC test (for determination of various species of fecal coliform), was not performed in about 3% of the all samples. Therefore, the results of these samples were not included in the study.

**Table 2**

Distribution of fecal coliform based on the months of year in drinking water of Kermanshah city.

Seasons	Months	Number of samples	Distribution (%)						
			<i>Citrobacter freundii</i>	<i>Escherichia coli</i>	<i>Enterobacter aerogenes</i>	<i>Enterobacter agglomerans</i>	<i>Enterobacter cloacea</i>	<i>Klebsiella</i>	No Fecal coliform
Spring	April	437	14.2	36.2	5.5	0	5.5	0.7	38
	May	805	10.8	41.7	5.7	2.5	0.6	4.3	24.3
	June	960	12.9	40.6	5.1	10.1	6.6	0.8	23.9
Summer	July	967	14.9	45	5.9	5.6	3.2	2.8	22.6
	August	1066	19.7	36.1	6	7.7	2.8	6.2	21.5
	September	834	17	47.5	7.6	0.9	2.8	0.4	23.9
Fall	October	761	13.6	33.4	8.8	7.7	2.8	0.4	28.8
	November	708	8.9	38.6	12.6	3.4	0.1	1.7	34.7
	December	666	13.7	31.8	2.1	15.5	0.2	0.5	36.3
Winter	January	553	4.7	31.5	4.9	11.9	6.3	4	36.7
	February	497	11.7	27.4	7.2	6	0.6	6.4	40.6
	March	433	7	37.9	3.9	4.4	1.6	0.9	44.3
Average	-	12.42	38.1	6.4	6.06	3	2.8	29.3	

**Table 3**

Distribution of *Fecal coliforms* regarding the amount of residual chlorine in water resources of Kermanshah city.

Residual chlorine range (mg/L)	Number of samples	Distribution (%)							Total
		<i>Citrobacter freundii</i>	<i>Escherichia coli</i>	<i>Enterobacter aerogenes</i>	<i>Enterobacter agglomerans</i>	<i>Enterobacter cloacea</i>	<i>Klebsiella</i>	<i>No fecal coliform</i>	
0	5837	15.2	42.6	7.1	7.5	3.3	3.5	21.1	100
0.5–0	1376	13.5	34.6	5.9	5	3.1	2.2	37.6	100
0.5–0.8	821	11.4	26.8	5	4	2.5	1.8	46.8	100
More than 0.8	609	9	19	3	2	2.4	0.8	63.4	100
Average		11.27	38.1	6.4	4.62	3	2.8	29.1	100



**Fig. 1.** Map of the study area (Kermanshah city, Iran).

## Acknowledgment

The authors gratefully acknowledge the Research Council of Kermanshah University of Medical Sciences (Grant no.: 96402) for the financial support. In addition, the authors wish to acknowledge the invaluable cooperation of rural and urban health centers and Water and Wastewater Company of Kermanshah province, for providing necessary help.

## Transparency document. Supplementary material

Transparency data associated with this article can be found in the online version at <https://doi.org/10.1016/j.dib.2018.04.053>.

## References

- [1] M. Ahamadabadi, M. Saeidi, S. Rahdar, M.R. Narooie, A. Salimi, V. Alipour, R. Khaksefidi, M.M. Baneshi, H. Biglari, Assessment of the chemical quality of groundwater resources in Chabahar City using GIS software in 2016, *Res. J. Appl. Sci. 11* (2016) 1399–1403.
- [2] S. Ghasemi, A. Mohseni-Bandpei, M. Ghaderpoori, Y. Fakhri, H. Keramati, M. Taghavi, B. Moradi, K. Karimyan, Application of modified maize hull for removal of Cu (II) ions from aqueous solutions, *Environ. Prot. Eng. 43* (2017) 93–103.
- [3] D. Jalili Naghan, M. Darvish Motavalli, N. Mirzaei, A. Javid, H.R. Ghaffari, M. Ahmadpour, M. Moradi, K. Sharafi, Efficiency comparison of alum and ferric chloride coagulants in removal of dye and organic material from industrial wastewater – a case study, *Bulg. Chem. Commun. 47* (2015) 206–210.
- [4] K. Sharafi, M. Fazlzaadeh, M. Pirsahab, M. Moradi, A. Azari, H. Sharafi, K. Dindarloo, H.R. Ghafari, Wastewater disinfection using sodium dichloroisocyanate (NaDCC) and sodium hypochlorine (NaOCl): modeling, optimization and comparative analysis, *Desalin. Water Treat. 66* (2017) 221–228.
- [5] M. Pirsahab, K. Sharafie, S. Naderi, H.R. Ghafari, T. Khosravi, Role of reverse osmosis membranes on the concentration fluctuations of heavy metals in used water by dialysis instrument of hemodialysis patients, *Ann. Trop. Med. Public Health 10* (2017) 1019–1023.
- [6] M.D. Motavalli, D.J. Naghan, N. Mirzaei, S.A. Haghghi, Z. Hosseini, H. Sharafi, K. Sharafi, The reusing feasibility of wastewater treatment plant (conventional activated sludge) effluent of tomato paste factory for agricultural irrigation-a case study, *Int. J. Pharm. Technol. 7* (2015) 9672–9679.
- [7] H. Biglari, M. Saeidi, V. Alipour, S. Rahdar, Y. Sohrabi, R. Khaksefidi, M. Narooie, A. Zarei, M. Ahamadabadi, Prospect of disinfection byproducts in water resources of Zabol, *Int. J. Pharm. Technol. 8* (2016) 17856–17865.
- [8] M. Pirsahab, T. Khodadadi, Z. Bonyadi, K. Sharafi, T. Khosravi, Evaluation of pesticide residues 2,4-D, Atrazine and Alachlor concentration in drinking water well of Mahidasht district-Kermanshah, Iran, 2010–2011, *World Appl. Sci. J. 23* (2013) 1530–1537.

- [9] M. Pirsahab, H. Ghaffari, K. Sharafi, Application of response surface methodology for efficiency analysis of strong non-selective ion exchange resin column (A 400 E) in nitrate removal from groundwater, *Int. J. Pharm. Technol.* 8 (2016) 11023–11034.
- [10] M. Pirsahab, T. Khosravi, K. Sharafi, L. Babajani, M. Rezaei, Measurement of heavy metals concentration in drinking water from source to consumption site in Kermanshah—Iran, *World Appl. Sci. J.* 21 (2013) 416–423.
- [11] M. Pirsahab, T. Khosravi, M. Fazlzadeh, K. Sharafie, Effects of loading rate, resin height, and bed volume on nitrate removal from drinking water by non-selective strong anion exchange resin (A400E), *Desalin. Water Treat.* 89 (2017) 127–135.
- [12] N. Mirzaei, H.R. Ghaffari, K. Karimyan, F.M. Moghadam, A. Javid, K. Sharafi, Survey of effective parameters (water sources, seasonal variation and residual chlorine) on presence of thermotolerant coliforms bacteria in different drinking water resources, *Int. J. Pharm. Technol.* 7 (2015) 9680–9689.
- [13] M. Pirsahab, K. Sharafi, E. Ahmadi, M. Moradi, Prevalence of the waterborne diseases (diarrhea, dysentery, typhoid, and hepatitis A) in West of Iran during 5 years (2006–2010), *Ann. Trop. Med. Public Health* 10 (2017) 1524–1528.
- [14] K. Sharafi, M. Pirsahab, T. Khosravi, Performance comparison of activated sludge systems with conventional and extended aeration in removal of protozoan cysts and parasite egg from wastewater: a case study, *Iran. Int. J. Appl. Environ. Sci.* 7 (2012) 281–287.
- [15] H. Biglari, M. Afsharnia, V. Alipour, R. Khosravi, K. Sharafi, A.H. Mahvi, A review and investigation of the effect of nanophotocatalytic ozonation process for phenolic compound removal from real effluent of pulp and paper industry, *Environ. Sci. Pollut. Res.* 24 (2017) 4105–4116.
- [16] K. Sharafi, M. Pirsahab, T. Khosravi, A. Dargahi, M. Moradi, M.T. Savadpour, Fluctuation of organic substances, solids, protozoan cysts, and parasite egg at different units of a wastewater integrated stabilization pond (full scale treatment plant): a case study, *Iran. Desalin. Water Treat.* 57 (2016) 4913–4919.
- [17] N. Azizi, J. Amini, K. Karimyan, H. Ghaffari, H. Sharafi, Management of collection, treatment and disposal of hospital wastewater-case study: Hamadan and Kermanshah Province' hospitals (2014), *Int. J. Pharm. Technol.* 8 (2016) 12924–12933.
- [18] A. Almasi, K. Sharafi, S. Hazrati, M. Fazlzadehdavil, A survey on the ratio of effluent algal BOD concentration in primary and secondary facultative ponds to influent raw BOD concentration, *Desalin. Water Treat.* 53 (2015) 3475–3481.
- [19] K. Sharafi, M. Pirsahab, R. Davoodi, H.R. Ghaffari, M. Fazlzadeh, M. Karimaei, M. Miri, K. Dindarloo, A. Azari, H. Arfaeinia, Quantitative microbial risk assessment of Giardia cyst and Ascaris egg in effluent of wastewater treatment plants used for agriculture irrigation – a case study, *Desalin. Water Treat.* 80 (2017) 142–148.
- [20] M. Afsharnia, H. Biglari, S.S. Rasouli, A. Karimi, M. Kianmehr, Sono-electrocoagulation of fresh leachate from municipal solid waste; simultaneous applying of iron and copper electrodes, *Int. J. Electrochem. Sci.* 13 (2018) 472–484.
- [21] K. Sharafi, M. Moradi, A. Karami, T. Khosravi, Comparison of the efficiency of extended aeration activated sludge system and stabilization ponds in real scale in the removal of protozoan cysts and parasite ova from domestic wastewater using Bailenger method: a case study, *Kermanshah, Iran. Desalin. Water Treat.* 55 (2015) 1135–1141.
- [22] M. Pirsahab, T. Khosravi, K. Sharafi, M. Mouradi, Comparing operational cost and performance evaluation of electrodialysis and reverse osmosis systems in nitrate removal from drinking water in Golshahr, Mashhad, *Desalin. Water Treat.* 57 (12) (2016) 5391–5397.
- [23] K. Sharafi, A.M. Mansouri, A.A. Zinatizadeh, M. Pirsahab, Adsorptive removal of methylene blue from aqueous solutions by pumice powder: process modelling and kinetic evaluation, *Environ. Eng. Manag. J.* 14 (2015) 1067–1078.
- [24] Y. Sohrabi, A.H. Nafez, S.S. Chargineh, D. Abdollahzadeh, H. Biglari, Analyzing chemical parameters of drinking water of urban water system—Kermanshah—Iran, *Bulg. Chem. Commun.* 49 (2017) 256–262.
- [25] M. Pirsahab, Z. Rezai, A.M. Mansouri, A. Rastegar, A. Alahabadi, A.R. Sani, K. Sharafi, Preparation of the activated carbon from India shrub wood and their application for methylene blue removal: modeling and optimization, *Desalin. Water Treat.* 57 (2016) 5888–5902.
- [26] E. Azizi, M. Fazlzadeh, M. Ghayebzadeh, L. Hemati, M. Beikmohammadi, H.R. Ghaffari, H.R. Zakeri, K. Sharafi, Application of advanced oxidation process ( $H_2O_2/UV$ ) for removal of organic materials from pharmaceutical industry effluent, *Environ. Prot. Eng.* 43 (2017) 183–191.
- [27] W.E. Federation, American Public Health Association, Standard Methods for the Examination of Water and Wastewater, American Public Health Association (APHA), Washington, DC, USA, 2005.