



Distributed and dynamic intracellular organization of extracellular information

Alejandro A. Granados^{a,b,1}, Julian M. J. Pietsch^{b,c,1}, Sarah A. Cepeda-Humerez^d, Iseabail L. Farquhar^{b,c}, Gašper Tkačik^d, and Peter S. Swain^{b,c,2}

^aDepartment of Bioengineering, Imperial College London, London SW7 2AZ, United Kingdom; ^bSynthSys–Synthetic and Systems Biology, University of Edinburgh, Edinburgh EH9 3BF, United Kingdom; ^cSchool of Biological Sciences, University of Edinburgh, Edinburgh EH9 3BF, United Kingdom; and ^dInstitute of Science and Technology Austria, 3400 Klosterneuburg, Austria

Edited by Eric D. Siggia, The Rockefeller University, New York, NY, and approved April 27, 2018 (received for review September 21, 2017)

Although cells respond specifically to environments, how environmental identity is encoded intracellularly is not understood. Here, we study this organization of information in budding yeast by estimating the mutual information between environmental transitions and the dynamics of nuclear translocation for 10 transcription factors. Our method of estimation is general, scalable, and based on decoding from single cells. The dynamics of the transcription factors are necessary to encode the highest amounts of extracellular information, and we show that information is transduced through two channels: Generalists (Msn2/4, Tod6 and Dot6, Maf1, and Sfp1) can encode the nature of multiple stresses, but only if stress is high; specialists (Hog1, Yap1, and Mig1/2) encode one particular stress, but do so more quickly and for a wider range of magnitudes. In particular, Dot6 encodes almost as much information as Msn2, the master regulator of the environmental stress response. Each transcription factor reports differently, and it is only their collective behavior that distinguishes between multiple environmental states. Changes in the dynamics of the localization of transcription factors thus constitute a precise, distributed internal representation of extracellular change. We predict that such multidimensional representations are common in cellular decision-making.

cell signaling | mutual information | time series | transcription factors | stress

All organisms sense their environment and internally represent the information gained to elicit a change in behavior (1). Much is understood about such representations in neural systems (2), but single cells must perform an analogous task (1, 3), encoding intracellularly the information about extracellular environments, and yet little is known about the nature of their encoding.

The activation of transcription factors is thought to provide an internal representation of a cell's environment (4–10), but how information is encoded dynamically, whether information is spread across multiple factors, and how information is read downstream all remain unclear (Fig. 1A). We do know that the biochemical implementation of such representations is likely to be stochastic (11) and that the same biochemistry may be used to sense disparate environments. Furthermore, cells typically have just “one shot” at mounting the appropriate response from these internal representations, with competition being unforgiving for those that delay, at least among microbes (12–14). Here, we use information theory to investigate how eukaryotic cells answer these challenges.

To do so, we turn to budding yeast and to environmental changes for which we expect information encoding to be key: stresses that compromise growth and evoke adaptive gene expression (15). In yeast, extracellular changes are sensed by signaling networks that regulate the activity of transcription factors, often by their translocation either into or out of the nucleus (16), analogous to p53 and NF- κ B in mammalian cells (17, 18). We therefore consider the movement of these transcription factors

as a cell's internal representation of an environmental transition. The translocations are dynamic and stochastic, and the information available from the full time series of the response could be substantially higher than that available from any temporal snapshot (9) (Fig. 1A).

Tens of transcription factors translocate in yeast (16), and we focus on a representative subset: Msn2 and its paralog Msn4, which drive the environmental stress response; Mig1 and its paralog Mig2, which respond to low glucose; Hog1 (a kinase), which responds to hyperosmotic stress; Yap1, which responds to oxidative stress; Sfp1, which promotes, and Dot6 and its paralog Tod6, which repress the biogenesis of ribosomes; and Maf1, which represses the synthesis of tRNAs. We include Dot6 and Tod6, which are little studied, to determine if our approach can help determine their physiological importance. Some of these factors (Msn2/4, Mig1/2, and Dot6/Tod6) have pulsatile dynamics, with stochastic bursts of nuclear localization even without stress (19).

We consider environmental shifts from rich medium (2% glucose) into carbon stress (0.1% glucose), hyperosmotic, or oxidative stress. Using fluorescent tagging and microfluidics (20), we measure the degree of nuclear localization of the transcription factors in hundreds of single cells both before and after the stress is applied (Fig. 1B).

Significance

To thrive in diverse environments, cells must represent extracellular change intracellularly despite stochastic biochemistry. Here, we introduce a quantitative framework for investigating the organization of information within a cell. Combining single-cell measurements of intracellular dynamics with a scalable methodology for estimating mutual information between time series and a discrete input, we demonstrate that extracellular information is encoded in the dynamics of the nuclear localization of transcription factors and that information is lost with alternative static statistics. Any one transcription factor is usually insufficient, but the collective dynamics of multiple transcription factors can represent complex extracellular change. We therefore show that a cell's internal representation of its environment can be both distributed across diverse proteins and dynamically encoded.

Author contributions: A.A.G., J.M.J.P., and P.S.S. designed the study; S.A.C.-H. and G.T. developed the estimation algorithm; A.A.G., J.M.J.P., and I.L.F. performed the experiments; A.A.G., J.M.J.P., S.A.C.-H., and P.S.S. analyzed the data; and A.A.G., J.M.J.P., S.A.C.-H., I.L.F., G.T., and P.S.S. wrote the paper.

The authors declare no conflict of interest.

This article is a PNAS Direct Submission.

Published under the PNAS license.

¹A.A.G. and J.M.J.P. contributed equally to this work.

²To whom correspondence should be addressed. Email: peter.swain@ed.ac.uk.

This article contains supporting information online at www.pnas.org/lookup/suppl/doi:10.1073/pnas.1716659115/-DCSupplemental.

Published online May 21, 2018.




Fig. 1. Intracellular responses carry information on extracellular change. (A) Environmental transitions trigger dynamic intracellular responses (purple and blue time series), which may be the only indications to the cell that the environment has altered. Stochasticity can make intracellular responses a poor readout of the new environment, and a probabilistic representation of the possible environmental state based on the instantaneous intracellular response may vary over time. (B) Our approach compares single-cell time series of nuclear translocation (Sfp1 is shown) in different environments. The transcription factors are tagged with green fluorescent protein to reveal their dynamics (insets and *SI Appendix*, Fig. S2). (C) We estimate the mutual information (MI) between the time series and the environment by using 70% of the data to train a classifier to classify the time series into the two environments (each colored line in the heat map is a single cell, with greater nuclear localization indicated by yellow and less by blue). With the remaining data, we determine the confusion matrix (the probability of correctly and incorrectly identifying the environment from a time series) and, from this matrix, a lower bound on the mutual information. (D) By increasing the length of the time series used by the classifier (each time series has the same initial point), we estimate both the probability of correct and incorrect predictions and the mutual information as a function of the duration of the response. Here, the actual environment is 0.1% glucose (green area indicates the fraction of correctly classified time series). At $t = 0$, the two environments cannot be distinguished, and the best prediction is a random guess, which has a prediction probability of 0.5, corresponding to 0 bits of information. Error bars are SD across biological replicates ($n = 6$).

Results

To quantify the information available to the cell, we develop a general and scalable methodology to estimate the mutual information between the time series of cellular responses and the state of the extracellular environment (Fig. 1C and *SI Appendix*, Figs. S4–S7). Mutual information, a measure of statistical dependency (21), allows us both to capture the effects of biochemical stochasticity, which can drive individual responses far from the average, and to avoid a priori assumptions about which

features of the response are relevant, such as its magnitude or duration.

Our method involves training a classifier to predict the state of the environment from the time series of as few as 100 cells. The classifier's output on the test data can then be used to estimate the mutual information (Fig. 1C). Formally, this approach provides a lower bound on the true information (*SI Appendix*), but, biologically, we are quantifying the information that a cell could plausibly recover and act upon after observing a single time series of its response. By varying the duration of the time series used by the classifier, we can determine how quickly cells accumulate information (Fig. 1D). In addition, errors made by the classifier indicate environments that are likely to be confused, giving insight into tasks potentially challenging for the cell. Although the final estimate of mutual information is determined by the signal-to-noise ratio of the data, this noise may not be all biological, again making our estimates a lower bound.

Mutual information is also determined by the choice of input distribution. Our focus is to use mutual information to quantify the signal-to-noise ratios in the single-cell time series, and therefore we choose a uniform distribution, which does not favor any one input. When we repeat the estimations using the input distribution that maximizes the mutual information, we see few changes (*SI Appendix*, Fig. S18). Indeed, in the true (natural) input distribution, stresses may not even occur one at a time as we have assumed.

Detecting Environmental Change. Considering transitions into one of three environments, all of which reduce growth (Fig. 2A), the 10 transcription factors have diverse dynamics (Fig. 2B). The mutual information we calculate addresses whether an environmental transition can be detected from a typical time series of nuclear localization and is a number between 0 bits (indicating no detectable statistical differences between the dynamics of localization before and after the transition) and 1 bit (the dynamics of localization before and after the transition are distinct).

The glucose specialists Mig1/2 perform almost optimally in carbon stress with almost the maximum possible information of 1 bit (Fig. 2C). We note that different transcription factors encode information in different ways. For example, Dot6 and Sfp1 have dissimilar dynamics (Fig. 2B), yet encode the same amount of bits (Fig. 2C). Paralogs, however, do not necessarily carry equal information (compare Tod6 and Dot6).

Information can be encoded within minutes of the environmental transition with no trade-off: The speed of encoding typically increases the more information is encoded. Defining the encoding delay as the time for the mutual information to reach 50% of its maximum, information plateaus earliest in the time series of Mig1/2 (Fig. 2D). Fast responders are therefore typically more accurate, at least for such large stresses.

These general observations hold for transitions into osmotic and oxidative stress (Fig. 2D), establishing a hierarchy: In terms of the information and encoding delays, specialists (in blue) are followed by the environmental stress response (in pink), which in turn is followed by the others. The details of this hierarchy, however, are stress-specific, indicating that the dynamics of the transcription factors may encode not only the presence but also the nature of the environmental transition.

Detecting the Nature of Environmental Change. We therefore extend our method to calculate the mutual information between a single-cell time series and the four environmental states: rich medium (before the environmental transition) and the three stresses (after the transition). Not only do we estimate the mutual information (Fig. 3A), but we can also predict how a typical time series is likely to be classified as a function of




Fig. 2. A hierarchy of information encoding, both in bits and encoding delays, holds across environmental transitions. (A) From population measurements (SI Appendix, Fig. S3), each new environment—carbon stress (0.1% glucose), osmotic stress (0.4 M NaCl), or oxidative stress (0.5 mM H₂O₂)—reduces growth compared with growth in rich medium (2% glucose). (B) For 10 transcription factors, we quantify nuclear localization across four environments using a step change from rich medium to stress at $t = 0$. The median and interquartile range are shown. (C) In response to carbon stress, the mutual information (MI) shows a hierarchy. The maximum possible information is 1 bit (dotted line). Mean and SD of two experiments per transcription factor are shown. (D) The hierarchy's order is maintained for transitions into other stresses: Specialists (blue) encode the most information and are fastest, followed by the environmental stress response (pink).

the duration of the environment (Fig. 3B and SI Appendix, Figs. S11 and S12).

Although no single transcription factor reaches the maximum of 2 bits, the time series of Msn2, Msn4, and, unexpectedly, Dot6, carry sufficient information to identify three environmental categories (for example, two environmental states and the remaining two states lumped together) (Fig. 3A). We observe, however, that the information is instead “spread” so that all environmental states are eventually classified with a >80% accuracy (Fig. 3B, Dot6). In contrast, an ideal specialist should perfectly discriminate one environmental state and lump together the remaining states to encode 0.8 bits (SI Appendix). Indeed, Hog1 and Yap1 do encode this much information (Fig. 3A), and their signaling networks operate nearly optimally in these high stresses. After only a few minutes, both specialists unequivocally identify their associated stress and never report false positives (Fig. 3B, Yap1).

Conditioning the mutual information on the identity of the environmental states delineates specialists from the other transcription factors (Fig. 3A, Inset, and SI Appendix, Fig. S15), which we term generalists because they encode information on multiple types of stress. Nevertheless, these groups are not mutually exclusive: Mig1 is not only a specialist for carbon stress, but also carries information on the other environmental states at late times, particularly osmotic stress, for which the probability of correctly identifying the environment is more than twice the 25% probability of a random choice (Fig. 3B).

Detecting the Magnitude of Environmental Change. In such high stresses, specialists appear unnecessary because the generalists

identify stress so well, but this situation changes if we consider transitions into stresses of lower magnitude (Fig. 3C and SI Appendix, Fig. S10). From rich medium, we apply four different levels of the same type of stress and estimate the mutual information between the time series of translocation and the five environmental states.

Specialists now outperform generalists. Considering the mutual information between the time series and all pairs of the different levels of stress (Fig. 3D and SI Appendix, Fig. S13), we see that distinguishing between adjacent levels is most challenging, and generalists, but not specialists, can often only identify high stress.

Generalists and specialists also encode information differently: Generalists often use their entire time series, whereas specialists only do so to distinguish stresses of lower magnitude. By calculating the mutual information between summary features of the single-cell time series and the environmental state (SI Appendix, Figs. S16 and S17), we find that the amplitude of a specialist's initial translocation can identify its associated stress if that stress is sufficiently severe (yellow dots in Fig. 3A), explaining specialists' short encoding delays. For transitions into stresses of lower magnitude, however, information is encoded in the dynamics of the specialists' response (yellow dots in Fig. 3C). Generalists can encode twice the amount of information in their time series compared with the highest information encoded by any single time point, and both the timing of their initial translocation, particularly for Msn2 and Dot6, and the shape of the times series can be important (SI Appendix).




Fig. 3. Generalists can identify the nature of large environmental transitions, but only specialists distinguish smaller changes. (A) The mutual information (MI) between a time series of translocation and the four environmental states shows that multiple transcription factors can encode the nature of the environment (errors are SD over bootstrap replicates). The highest mutual information that can be found by using a single time point is always lower (yellow dots). (A, Inset) Conditioning the mutual information on the environmental states differentiates the specialists (in color). (B) Our methodology also estimates the typical probability of inferring each environmental state from a time series. The true state is marked with hash lines. Inference generally improves with the duration of the new environment, and rich medium and oxidative stress are confused most often. (C) The mutual information between a time series of translocation and four levels of one type of stress and rich medium. Specialists perform best and encode with their entire time series (compare yellow dots with those in A). (D) Comparing the mutual information between the time series and all pairs of environments with the same type of stress at different levels, generalists can identify high stress, but poorly distinguish adjacent levels compared with specialists. (E) The more a stress reduces growth, the more information is typically encoded. Comparing the mutual information from C with the mean fitness penalty (averaged over the different levels of stress) for each type of stress, there is a strong correlation (>0.9) for all generalists except Sfp1. Triangles, squares, and circles denote oxidative, osmotic, and carbon stress, respectively. Errors are SD.

For three of the four generalists considered, there is a substantial correlation between the amount of mutual information encoded and the severity of the stress (estimated by its reduction of growth compared with growth in rich medium) (Fig. 3E). This correlation may reflect that generalists are typically involved in regulating growth, through, for example, affecting translation (16). Specialists, in contrast, do not encode more information if their cognate stress is more severe (Fig. 3E).

Generalists vs. Specialists. To better understand the differences between generalists and specialists, we ask how the transcription factors are organized within the cell's network of signal transduction. Using data on the substrates of kinases (22), we confirm (16) that the generalists are either directly or indirectly targets of protein kinase A (which has isoforms Tpk1-3) and TORC1 or its downstream kinase Sch9 (S6 kinase) (Fig. 4A). The generalists therefore do respond to the cell's potential for growth: Protein kinase A orchestrates the cell's response to the availability of glucose (16), and TORC1 controls the response to the availability of nitrogen (16). Similarly, Mig1 is sensitive to the levels of cellular ATP through its regulation by AMP kinase (16). In contrast, the

specialists Hog1 and Yap1 are mostly embedded in their own signaling networks.

We quantify the redundancy between pairs of transcription factors to determine if the cell's organization of information reflects the signaling network. If regulated by the same upstream signaling, two transcription factors may be completely redundant, so that when paired together, the amount of information does not increase above that from any one factor alone. By concatenating two time series (SI Appendix, Fig. S19), we can estimate the mutual information from simultaneously observing two transcription factors and so their redundancy (Fig. 4B).

Plotting the redundancy (Fig. 4B), we see a network similar to the network of signal transduction: The generalists are together in a core, from which the specialists are distinct. Msn2/4 and Dot6 appear to coordinate the behavior of specialists with the general environmental stress response, having a substantial degree of redundancy with the highest number of transcription factors, including each other (size of the nodes in Fig. 4B). Specialists are not redundant with other specialists, but each is redundant with a distinct subset of the core generalists: Yap1 is




Fig. 4. Collectively, transcription factors provide an internal representation of complex environments. (A) In the intracellular signaling network, generalists are colocated, and specialists are distinct. Edges between kinase and substrate are weighted by the evidence for that interaction (22). Clusters of highly interconnected components have different colors, and we include only the transcription factors in our study. (B) The redundancy in information between pairs of transcription factors reflects the structure of the signaling network in A. Edges are proportional to a pair's redundancy, $1 - MI_{12}/(MI_1 + MI_2)$ (23), and the size of each node is determined by the number of its edges. Redundancy was calculated from an average of two datasets (six experiments per transcription factor). (C) A generalist and a specialist typically encode the most information out of all possible pairs of transcription factors. For each pair, time series from one transcription factor were concatenated randomly with the time series of another to calculate the mutual information (MI). G, generalist; S, specialist. (D) Only through combinations of transcription factors can information on complex environments be encoded. Different colors represent different numbers of environmental states (*SI Appendix*): orange, seven states; red, five states; blue, four states; and purple, two states. Plotted lines are maxima calculated from all combinations of concatenated time series for a given number of transcription factors. (E) Cells transduce information through two types of channels—specialists and generalists—and four transcription factors are sufficient to encode all of the available information. The direct channels (specialists) respond to extracellular change; the indirect channels (generalists) respond to intracellular change and so to broader categories of extracellular change.

partly redundant with Msn2/4 but not Dot6; Hog1 and Mig2 with Dot6 but not Msn2/4; and Mig1 is partly redundant with all three.

The redundancies imply that pairing a generalist with a specialist is best (Fig. 4C), and indeed such pairs typically encode the highest information (*SI Appendix*, Fig. S20). With its distinct signal transduction (Fig. 4A), a specialist can identify the environmental state that is most poorly distinguished by a generalist. For example, Msn2 is best paired with Mig2 (*SI Appendix*, Fig. S22).

As environments become more complex, multiple transcription factors are needed to generate an internal representation. Pooling the data to consider environments with different states (Fig. 4D), the maximum mutual information plateaus as the numbers of transcription factors increase, with four sufficing to generate ~95% of the information. This increase comes both from the distinct dynamics of the transcription factors (24), such as differences in timing (*SI Appendix*, Fig. S21), and from decreasing the effects of stochasticity by averaging the multiple readouts.

Discussion

In summary, we have shown that transcription factors can encode enough information in the dynamics of their nuclear translocations to unambiguously report an environmental change if that change is sufficiently large, that the nature of the change can also be encoded although with some degree of error, that how the information is encoded alters for changes of different magnitudes, and that no single transcription factor can accurately encode both the nature and magnitude of environmental change.

Information is transduced through two channels of specialists and generalists. Specialists are faster and can better identify a transition into their associated stress than generalists, but the variety of environments experienced by cells makes having a specialist for every environment implausible. We postulate that generalists avoid this constraint by providing an indirect channel that responds not to the extracellular signals sensed by specialists (25, 26), but to intracellular signals (27, 28), such as changes in cAMP, uncharged tRNAs, and the availability of amino acids (16) (Fig. 4E). By detecting physiological

perturbations, generalists respond to broader ranges of stress (*SI Appendix, Fig. S23*) and are agnostic to the environment's precise nature. Generalists are therefore necessarily slower than specialists because they must wait for the environment to modify intracellular biochemistry. Indeed, we conjecture that the stochastic pulsing of the generalists in constant environments (19) is a response to spontaneous fluctuations in intracellular physiology.

Consistent with more recent interpretations (29), our data do not support a distinct environmental stress response controlled by *Msn2/4*, but show that aspects of the behavior of *Msn2/4* are present in the dynamics of multiple transcription factors, such as *Dot6*, *Sfp1*, and *Maf1*. These latter factors act to determine rates of translation, consistent with the push–pull relationship between stress and growth (12). In particular, we have demonstrated that *Dot6*, although not *Tod6*, encodes the nature of environmental change in its dynamics to an accuracy almost comparable with *Msn2/4*, implying that *Dot6* may play a similar role in cellular physiology. We find, too, that *Mig1/2*, although considered a glucose specialist, encodes information on osmotic stress and might better be classed as a generalist. Indeed, *Mig1/2* is activated by AMP kinase [*Snf1*, which responds to levels of ADP (16)], consistent with our proposal that generalists respond to an environmental change's intracellular effects.

Finally, our results show that it is only through the collective dynamics of multiple transcription factors that cells can

encode sufficient information to generate specific downstream responses (15, 30). Paralleling discoveries in neuroscience, we expect that such multidimensional internal representations are widespread within cellular biology and that their failures in encoding information, by causing dysfunctional decision-making, instigate deleterious behaviors and disease (31).

Materials and Methods

Time-Lapse Microscopy. BY strains with fluorescently tagged transcription factors (32) are grown in ALCATRAS microfluidic devices (20) in synthetic complete medium with 2% glucose for at least 3 h and then exposed to stress for 5 h (*SI Appendix*). Switching between media is via an external mixer and syringe pumps. Bright-field and fluorescence images are acquired every 2.5 min, and cells are segmented by using the DISCO algorithm (33).

Estimating the Mutual Information. Our algorithm involves (*SI Appendix*): (i) using principal component analysis to delineate a basis for the time series; (ii) training a linear support vector machine to classify time series in this basis; (iii) calculating a confusion matrix using the test data; and (iv) estimating (a lower bound on) the mutual information by interpreting the entries of the confusion matrix as conditional probabilities.

ACKNOWLEDGMENTS. We thank S. Jaramillo-Riveri, P. Thomas, M. Voliotis, E. Wallace, and the P.S.S. laboratory for critical comments and Reiko Tanaka for her support and advice (A.A.G.). This work was supported by the Biotechnology and Biological Sciences Research Council (J.M.J.P., I.F., and P.S.S.), the Engineering and Physical Sciences Research Council (EPSRC) (A.A.G.), and Austrian Science Fund Grant FWF P28844 (to G.T.).

1. Tkačik G, Bialek W (2016) Information processing in living systems. *Annu Rev Condens Matter Phys* 7:89–117.
2. Gold JI, Shadlen MN (2007) The neural basis of decision making. *Annu Rev Neurosci* 30:535–574.
3. Perkins TJ, Swain PS (2009) Strategies for cellular decision-making. *Mol Syst Biol* 5:326.
4. Cai L, Dalal CK, Elowitz MB (2008) Frequency-modulated nuclear localization bursts coordinate gene regulation. *Nature* 455:485–490.
5. Cheong R, Rhee A, Wang CJ, Nemenman I, Levchenko A (2011) Information transduction capacity of noisy biochemical signaling networks. *Science* 334:354–358.
6. Hao N, O'Shea EK (2011) Signal-dependent dynamics of transcription factor translocation controls gene expression. *Nat Struct Mol Biol* 19:31–39.
7. Hao N, Budnik BA, Gunawardena J, O'Shea EK (2013) Tunable signal processing through modular control of transcription factor translocation. *Science* 339:460–464.
8. Purvis JE, Lahav G (2013) Encoding and decoding cellular information through signaling dynamics. *Cell* 152:945–956.
9. Selimkhanov J, et al. (2014) Systems biology. Accurate information transmission through dynamic biochemical signaling networks. *Science* 346:1370–1373.
10. Goulev Y, et al. (2017) Nonlinear feedback drives homeostatic plasticity in H₂O₂ stress response. *eLife* 6:e23971.
11. Elowitz MB, Levine AJ, Siggia ED, Swain PS (2002) Stochastic gene expression in a single cell. *Science* 297:1183–1186.
12. López-Maury L, Marguerat S, Bähler J (2008) Tuning gene expression to changing environments: From rapid responses to evolutionary adaptation. *Nat Rev Genet* 9:583–593.
13. Mitchell A, Wei P, Lim WA (2015) Oscillatory stress stimulation uncovers an Achilles' heel of the yeast MAPK signaling network. *Science* 350:1379–1383.
14. Granados AA, et al. (2017) Distributing tasks via multiple input pathways increases cellular survival in stress. *Elife* 6:e21415.
15. Gasch AP, et al. (2000) Genomic expression programs in the response of yeast cells to environmental changes. *Mol Biol Cell* 11:4241–4257.
16. Broach JR (2012) Nutritional control of growth and development in yeast. *Genetics* 192:73–105.
17. Purvis JE, et al. (2012) p53 dynamics control cell fate. *Science* 336:1440–1444.
18. Ashall L, et al. (2009) Pulsatile stimulation determines timing and specificity of NF- κ B-dependent transcription. *Science* 324:242–246.
19. Dalal CK, Cai L, Lin Y, Rahbar K, Elowitz MB (2014) Pulsatile dynamics in the yeast proteome. *Curr Biol* 24:2189–2194.
20. Crane MM, Clark IBN, Bakker E, Smith S, Swain PS (2014) A microfluidic system for studying ageing and dynamic single-cell responses in budding yeast. *PLoS One* 9:e100042.
21. Shannon CE, Weaver W (1999) *The Mathematical Theory of Communication* (Univ of Illinois Press, Urbana, IL).
22. Sharifpoor S, et al. (2011) A quantitative literature-curated gold standard for kinase-substrate pairs. *Genome Biol* 12:R39.
23. MacKay DJC (2003) *Information Theory, Inference, and Learning Algorithms* (Cambridge Univ Press, Cambridge, UK).
24. Dubuis JO, Tkacik G, Wieschaus EF, Gregor T, Bialek W (2013) Positional information, in bits. *Proc Natl Acad Sci USA* 110:16301–16308.
25. Delaunay A, Isnard AD, Toledano MB (2000) H₂O₂ sensing through oxidation of the Yap1 transcription factor. *EMBO J* 19:5157–5166.
26. Reiser V, Raitt DC, Saito H (2003) Yeast osmosensor *Sln1* and plant cytokinin receptor *Cre1* respond to changes in turgor pressure. *J Cell Biol* 161:1035–1040.
27. Dechant R, Saad S, Ibáñez AJ, Peter M (2014) Cytosolic pH regulates cell growth through distinct GTPases, *Arf1* and *Gtr1*, to promote *Ras/PKA* and *TORC1* activity. *Mol Cell* 55:409–421.
28. Filteau M, et al. (2015) Systematic identification of signal integration by protein kinase A. *Proc Natl Acad Sci USA* 112:4501–4506.
29. Gasch AP, et al. (2017) Single-cell RNA sequencing reveals intrinsic and extrinsic regulatory heterogeneity in yeast responding to stress. *PLoS Biol* 15:e2004050.
30. Hansen AS, O'Shea EK (2013) Promoter decoding of transcription factor dynamics involves a trade-off between noise and control of gene expression. *Mol Syst Biol* 9:704.
31. Luo Q, Beaver JM, Liu Y, Zhang Z (2017) Dynamics of p53: A master decider of cell fate. *Genes* 8:66.
32. Huh WK, et al. (2003) Global analysis of protein localization in budding yeast. *Nature* 425:686–691.
33. Bakker E, Swain PS, Crane MM (2018) Morphologically constrained and data informed cell segmentation of budding yeast. *Bioinformatics* 34:88–96.