



# Draft Genome Sequences of Two *Sporothrix schenckii* Clinical Isolates Associated with Human Sporotrichosis in Colombia

Oscar M. Gomez,<sup>a,b</sup> Laura C. Alvarez,<sup>b,c</sup> Jose F. Muñoz,<sup>a,d</sup> Elizabeth Misas,<sup>a,b</sup> Juan E. Gallo,<sup>a,e</sup> Maria del P. Jimenez,<sup>c</sup> Myrtha Arango,<sup>c</sup> Juan G. McEwen,<sup>a,c</sup> Orville Hernandez,<sup>a,f</sup> Oliver K. Clay<sup>a,g</sup>

<sup>a</sup>Cellular and Molecular Biology Unit, Corporación para Investigaciones Biológicas, Medellín, Colombia

<sup>b</sup>Institute of Biology, Universidad de Antioquia, Medellín, Colombia

<sup>c</sup>School of Medicine, Universidad de Antioquia, Medellín, Colombia

<sup>d</sup>Broad Institute of MIT and Harvard, Cambridge, Massachusetts, USA

<sup>e</sup>GenomaCES, Universidad CES, Medellín, Colombia

<sup>f</sup>MICROBA Research Group, School of Microbiology, Universidad de Antioquia, Medellín, Colombia

<sup>g</sup>School of Medicine and Health Sciences, Universidad del Rosario, Bogotá, Colombia

**ABSTRACT** *Sporothrix schenckii* is a thermodimorphic fungal pathogen with a high genetic diversity. In this work, we present the assembly and similarity analysis of the whole-genome sequences of two clinical isolates from Colombia of *S. schenckii sensu stricto*.

*Sporothrix schenckii* is a thermodimorphic fungal pathogen and is the etiologic agent of sporotrichosis, a subcutaneous mycosis that affects mammals. The genus *Sporothrix* includes at least four human-pathogenic species: *S. schenckii sensu stricto*, *S. brasiliensis*, *S. globosa*, and *S. luriei* (1–3). *S. schenckii sensu stricto* presents a worldwide distribution and high genetic diversity, and five lineages (A to E) have been described for this species to date (4). In South America, the areas where the pathogen is most endemic include Peru, Colombia, Venezuela, and Brazil (4, 5). The primary mode of infection is traumatic inoculation with contaminated decaying plant material (4). Sporotrichosis has a wide spectrum of clinical manifestations, ranging from cutaneous forms to systemic presentations (6). The diagnosis is based on fungal growth and microscopic observation; however, these methods are time consuming, have low sensitivity, and do not allow species differentiation. Nevertheless, continued efforts have improved sensitivity and specificity for diagnostic molecular tests (7). The genus *Sporothrix* is exceptional in the fungal kingdom due to its high occurrence of outbreaks in areas where the pathogen is endemic and the characteristic differences in those outbreaks (4). Given the wide genetic diversity of the *S. schenckii* complex and the rapid emergence of sporotrichosis in different countries, the study of the genotypes involved in the different outbreaks is crucial for developing public health strategies to control the disease (5). Despite the efforts of the community studying this mycosis, genomic data reflecting the pathogen's diversity remain limited. Thus, we obtained isolates from two clinical cases of cutaneous sporotrichosis in human Colombian patients and sequenced the complete genomes.

Genomic DNA for sequencing was prepared from yeast culture, using phenol-chloroform extraction. Library preparation and 150-bp paired-end sequencing were performed using the Illumina HiSeq 2000 platform, generating ~14 million paired-end reads per strain. The reads were assembled using SOAPdenovo2 version 2.04 and GapCloser for SsEM7 and SPAdes version 3.10 for SsMS1. The assembled scaffolds generated by the two strains were aligned and oriented with MAUVE software. The genomes of the assemblies were processed using the QUAST version 4.5 program, and assembly statistics are shown in Table 1.

Received 3 May 2018 Accepted 7 May 2018 Published 14 June 2018

**Citation** Gomez OM, Alvarez LC, Muñoz JF, Misas E, Gallo JE, Jimenez MDP, Arango M, McEwen JG, Hernandez O, Clay OK. 2018. Draft genome sequences of two *Sporothrix schenckii* clinical isolates associated with human sporotrichosis in Colombia. Genome Announcements 6:e00495-18. <https://doi.org/10.1128/genomeA.00495-18>.

**Copyright** © 2018 Gomez et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Juan G. McEwen, [mcewen@une.net.co](mailto:mcewen@une.net.co).

**TABLE 1** Summary of assembly statistics

Sample ID	Genome size (Mb)	Coverage (×)	No. of scaffolds	Scaffold $N_{50}$ (bp)	Largest scaffold size (Mb)	Scaffold $L_{50}$ (no.)	GC content (%)	GenBank accession no.
SsMS1	32.7	127	124	625,938	1.78	18	54.89	PGUU00000000
SsEM7	32.9	128	181	1,079,862	3.72	9	54.81	NTMI00000000

To identify the genotype of the *S. schenckii* isolates, we first identified ITS, CAL, TEF1, and TEF3 sequences in the assembly using BLAST searches. This was followed by maximum likelihood based phylogenetic reconstruction using IQtree version 1.4.4 software, which included sequences of these markers from DDBJ/EMBL/GenBank as described by Zhang et al. in 2015. FigTree was used for tree visualization. The two isolates were classified as *S. schenckii sensu stricto* lineage A, a subgroup consisting of isolates of hyperendemic zones of South America, while the two genomes available for this species reported by Cuomo et al. in 2014 (8) and Teixeira et al. in 2015 (9) are from lineage E, a subgroup that represents isolates from USA. In this work, we report the first two genomes of lineage A of *S. schenckii*.

**Accession number(s).** The whole-genome sequences of both strains were deposited at DDBJ/ENA/GenBank under the accession numbers cited in Table 1.

## ACKNOWLEDGMENTS

This research was supported by Colciencias via the grant “A Gene Atlas for Human Pathogenic Fungi” (122256934875) and by the Universidad de Antioquia via the grant “Sostenibilidad 2016/2017” (SE84160128) and the CODI grant “Análisis Fenotípico y Filogenético de Aislamientos Clínicos Colombianos, Recuperados de Pacientes con Esporotricosis Cutánea Fija y Linfocutánea” (2015-7042).

## REFERENCES

- Marimon R, Gené J, Cano J, Trilles L, Dos Santos Lazéra M, Guarro J. 2006. Molecular phylogeny of *Sporothrix schenckii*. J Clin Microbiol 44: 3251–3256. <https://doi.org/10.1128/JCM.00081-06>.
- Marimon R, Cano J, Gené J, Sutton DA, Kawasaki M, Guarro J. 2007. *Sporothrix brasiliensis*, *S. globosa*, and *S. mexicana*, three new *Sporothrix* species of clinical interest. J Clin Microbiol 45:3198–3206. <https://doi.org/10.1128/JCM.00808-07>.
- Rodrigues AM, de Hoog GS, de Camargo ZP. 2016. *Sporothrix* species causing outbreaks in animals and humans driven by animal-animal transmission. PLoS Pathog 12:e1005638. <https://doi.org/10.1371/journal.ppat.1005638>.
- Zhang Y, Hagen F, Stielow B, Rodrigues AM, Samerpitak K, Zhou X, Feng P, Yang L, Chen M, Deng S, Li S, Liao W, Li R, Li F, Meis JF, Guarro J, Teixeira M, Al-Zahrani HS, de Camargo ZP, Zhang L, de Hoog G. 2015. Phylogeography and evolutionary patterns in *Sporothrix* spanning more than 14 000 human and animal case reports. Persoonia 35:1–20. <https://doi.org/10.3767/003158515X687416>.
- Rodrigues AM, de Hoog S, de Camargo ZP. 2013. Emergence of pathogenicity in the *Sporothrix schenckii* complex. Med Mycol 51:405–412. <https://doi.org/10.3109/13693786.2012.719648>.
- Ramos-e-Silva M, Vasconcelos C, Carneiro S, Cestari T. 2007. Sporotrichosis. Clin Dermatol 25:181–187. <https://doi.org/10.1016/j.clindermatol.2006.05.006>.
- Rodrigues AM, de Hoog GS, de Camargo ZP. 2015. Molecular diagnosis of pathogenic *Sporothrix* species. PLoS Negl Trop Dis 9:e0004190. <https://doi.org/10.1371/journal.pntd.0004190>
- Cuomo CA, Rodriguez-Del Valle N, Perez-Sanchez L, Abouelleil A, Goldberg J, Young S, Zeng Q, Birren BW. 2014. Genome sequence of the pathogenic fungus *Sporothrix schenckii* (ATCC 58251). Genome Announc 2(3):e00446-14. <https://doi.org/10.1128/genomeA.00446-14>.
- Teixeira MM, de Almeida LGP, Kubitschek-Barreira P, Alves FL, Kioshima ES, Abadio AKR, Fernandes L, Derengowski LS, Ferreira KS, Souza RC, Ruiz JC, de Andrade NC, Paes HC, Nicola AM, Albuquerque P, Gerber AL, Martins VP, Peconick LDF, Neto AV, Chaucanez CB, Silva PA, Cunha OL, de Oliveira FF, dos Santos TC, Barros AL, Soares MA, de Oliveira LM, Marini MM, Villalobos-Duno H, Cunha MML, de Hoog S, da Silveira JF, Henrissat B, Niño-Vega GA, Cisalpino PS, Mora-Montes HM, Almeida SR, Stajich JE, Lopes-Bezerra LM, Vasconcelos ATR, Felipe MSS. 2014. Comparative genomics of the major fungal agents of human and animal sporotrichosis: *Sporothrix schenckii* and *Sporothrix brasiliensis*. BMC Genomics 15:943. <https://doi.org/10.1186/1471-2164-15-943>.