



HHS Public Access

Author manuscript

Clin Cancer Res. Author manuscript; available in PMC 2018 November 01.

Published in final edited form as:

Clin Cancer Res. 2017 November 01; 23(21): 6733–6743. doi:10.1158/1078-0432.CCR-17-1397.

Biochemical, Molecular, and Clinical Characterization of Succinate Dehydrogenase Subunit A Variants of Unknown Significance

Amber E Bannon,

Cell and Developmental Biology, Heinrich Lab, Oregon Health and Science University, Portland, Oregon

Jason Kent,

Cell and Developmental Biology, Heinrich Lab, Oregon Health and Science University, Portland, Oregon

Isaac Forquer,

Portland VA Medical Center and Oregon Health & Science University

Ajia Town,

Heinrich Lab, Oregon Health and Science University, Portland, Oregon

Lillian R. Klug,

Cancer Biology, Heinrich Lab, Oregon Health and Science University, Portland, Oregon

Kelly McCann,

Department of Medicine, Division of Hematology-Oncology, University of California Los Angeles David Geffen School of Medicine

Carol Beadling,

Oregon Health and Science University, Portland, Oregon

Oliver Harismendy,

Division of Biomedical Informatics, Department of Medicine, Moores UCSD Cancer Center, University of California San Diego, La Jolla, CA, USA

Jason K. Sicklick,

Division of Surgical Oncology, Department of Surgery, Moores UCSD Cancer Center, University of California San Diego, La Jolla, CA, USA

Christopher Corless,

Knight Cancer Institute and Department of Pathology, Oregon Health and Science University, Portland, Oregon

Ujwal Shinde, and

Department of Biochemistry and Molecular Biology, MRB 631, Oregon Health and Science University, Portland, Oregon

Corresponding author: Amber E. Bannon, R&D 19, Building 103 Room E223, 3710 SW US Veteran's Hospital Road, Portland, Oregon 97239, Phone number-(503) 330-2332, bannon@ohsu.edu.

Michael C. Heinrich

Department of Medicine; Department of Cell, Developmental, and Cancer Biology, Portland VA Health Care System and OHSU Knight Cancer Institute, Portland, OR

Abstract

Purpose—Patients who inherit a pathogenic loss-of-function genetic variant involving one of the four succinate dehydrogenase (SDH) subunit genes have up to an 86% chance of developing one or more cancers by the age of 50. If tumors are identified and removed early in these high-risk patients, they have a higher potential for cure. Unfortunately, many alterations identified in these genes are variants of unknown significance (VUS), confounding the identification of high-risk patients. If we could identify misclassified SDH VUS as benign or pathogenic SDH mutations, we could better select patients for cancer screening procedures and remove tumors at earlier stages.

Experimental Design—In this study, we combine data from clinical observations, a functional yeast model, and a computational model to determine the pathogenicity of 22 *SDHA* VUS. We gathered *SDHA* VUS from two primary sources: The OHSU Knight Diagnostics Laboratory and the literature. We used a yeast model to identify the functional effect of a VUS on mitochondrial function with a variety of biochemical assays. The computational model was used to visualize variants' effect on protein structure.

Results—We were able to draw conclusions on functional effects of variants using our three-prong approach to understanding VUS. We determined that 16 (73%) of the alterations are actually pathogenic, causing loss of SDH function, and six (27%) have no effect upon SDH function.

Conclusions—We thus report the reclassification of the majority of the VUS tested as pathogenic, and highlight the need for more thorough functional assessment of inherited SDH variants.

Keywords

GI stromal tumor; succinate dehydrogenase; variants of unknown significance; mutation; paraganglioma

Introduction

SDH, succinate dehydrogenase, also known as complex II of the electron transport chain (ETC), is a four-subunit complex encoded by nuclear genes (*SDHA*, *SDHB*, *SDHC*, and *SDHD*, collectively referred to as *SDHx*). The assembled SDH complex localizes to the inner membrane of the mitochondria and links the tricarboxylic acid (TCA) cycle to the ETC, making SDH function critical for aerobic respiration (1).

Loss-of-function mutations affecting the SDH complex predispose patients to develop multiple cancers, including gastrointestinal stromal tumor (GIST), paraganglioma, pheochromocytoma, renal cell carcinoma, Hodgkin lymphoma, chronic lymphocytic leukemia, thyroid cancer, pituitary adenomas, and neuroendocrine tumors of the pancreas (2-6). Tumor formation due to SDH-deficiency requires the complete loss of function of at

least one SDHx subunit (e.g. A, B, C, or D), causing destabilization and loss of enzymatic function of the entire SDH complex (7). There are several genetic mechanisms that can lead to SDH-deficiency. Typically, loss of function of an SDHx subunit is the result of a combination of an inactivating germline mutation (first hit) with a somatic loss of heterozygosity or other inactivating mutation affecting the other allele (second hit). Less commonly, loss of SDH complex occurs due to somatic inactivation of both alleles of a given complex subunit or SDH assembly factor. Finally, SDH-deficiency can be caused by an *SDHC* epimutation, defined as hypermethylation of the *SDHC* promoter, which leads to repression of *SDHC* transcription and depletion of *SDHC* protein levels, without a known underlying heritable cause (8).

Importantly, germline loss-of-function genetic *SDHx* variants are associated with a high lifetime risk of developing the aforementioned malignancies. For example, the chance of a patient with germline loss-of-function *SDHD* variant of developing one or more primary tumors by the age of 50 was reported to be 86% (9,10). Therefore, if we could identify high-risk patients through genetic testing and follow them serially with specialized screening tests, early tumor detection may lead to curative surgical resection before the tumors are metastatic/incurable. Early detection is crucial since there are no effective medical treatments for patients with advanced SDH-deficient cancers.

Currently, an SDH-deficient tumor is identified by measuring SDHB protein abundance using immunohistochemistry (IHC); absence of SDHB protein is indicative of loss of SDH function. However, it can be challenging to determine the underlying cause of SDH-deficiency in a tumor lacking SDHB expression. Clinical sequencing panels may turn up missense mutations in SDHx genes, but many of these are VUS. In addition, such panels can miss large intragenic deletions, and are not designed to identify epigenetic silencing of the *SDHC* promoter. Some SDHB, C, and D VUS have been functionally characterized to determine their effect on function, and thus their pathogenicity in tumors like paraganglioma and pheochromocytoma. However, the study of the functional consequences of *SDHA* VUS has lagged behind that of other SDHx subunits (7).

GIST is a heterogeneous group of tumors that arise from the interstitial cells of Cajal (ICC). However there are several different driver genes that when mutated give rise to GIST (11). The molecular classification of GIST is especially important because of the treatment implications of the different genetic drivers. The majority of GISTs have an activating receptor tyrosine kinase (RTK) mutation but about 13% of GIST lack RTK mutations (RTK-WT). Most RTK-WT GIST are SDH-deficient as assessed by immunohistochemistry (IHC) for SDHB. *SDHA* pathogenic variants are found in 47% of SDH-deficient GIST, and the majority of these *SDHA* mutations are germline, and thus heritable, variants (12). However, some the *SDHA* variants we find in GIST are VUS. A universal problem in the field is that these variants are rarely seen with a complete clinical and pathogenic annotation making it difficult to draw conclusions on functional effect from clinical data alone. Our study gathered these VUS from two primary sources; OHSU and the literature (12,13). We then combine data from clinical observations, a functional yeast model, and a computational model to understand the effects of *SDHA* VUS identified in GIST specimens on SDH complex function. Historically, yeast have been a robust system for identifying the assembly

and enzymatic activity of SDH (14-16). In addition, yeast have proven to be an ideal model to study the functional effect of *SDHB/C/D* variants on SDH complex activity (17-20). Yeast are able to survive without functional mitochondria (e.g. lacking SDH complex activity) if they are provided a fermentable carbon source; thus, they provide a unique model system to study the biochemical effects of *SDHA* VUS (21).

Based on our findings, we discriminated between *SDHA* VUS that affected or did not affect SDH complex activity, and thus, their potential for pathogenicity. These data will aid clinicians' ability to provide genetic counseling and tumor surveillance to patients with germline inheritance of these specific *SDHA* variants.

Materials and methods

Yeast strains and vectors

All *Saccharomyces cerevisiae* strains used in this study were derivatives of BY4741 (MATa *his3 1 leu2 0 met15 0 ura3 0*). The *SDHI* (yeast homolog to *SDHA*) deletion strain (*sdh1*⁻) was purchased from ATCC (catalog #4004998). The *sdh1*⁻ was constructed as part of the *Saccharomyces* Genome Deletion Project by homologous recombination using the *KanMX4* cassette (22). We verified the deletion of *SDHI* using PCR mapping of the *SDHI* locus with primer pairs recommended by the *Saccharomyces* Genome Deletion Project.

An amplicon containing the WT *SDHI* (including the native *SDHI* promoter and 3' UTR) was generated from WT BY4741 and cloned into the pRS416 plasmid (23) (ATCC) and expressed in the *sdh1*⁻ strain. The *SDHI* point mutations were introduced by QuikChange mutagenesis PCR system (Agilent Technology, cat #200521). All mutations were confirmed by Sanger sequencing. Yeast strains were transformed using Frozen EZ Yeast Transformation II (Zymo Research, catalog # T2001). Strains were grown in synthetic complete medium lacking uracil to maintain plasmid selection with either 2% glucose or 3% glycerol as the carbon source.

Alignment of multiple species' succinate dehydrogenase flavoprotein subunit

Cluster Omega was used to align the protein sequences of SDH flavoproteins including *E. Coli* (P0AC41), yeast (Q00711), human (P31040), and pig (Q0QF01).

Immunoblotting

Intact mitochondria were isolated using a previously described method (24). Steady-state levels of mitochondrial proteins were resolved on SDS-PAGE, transferred to nitrocellulose membrane, probed using the indicated primary antibodies and visualized using Amersham enhanced chemiluminescence (ECL) Western Blotting Detection Reagent (GE life sciences, catalog #RPN2106) with horseradish peroxidase-conjugated secondary antibodies (BioRad, catalog #1662408EDU). We used previously described polyclonal rabbit antibodies for immunodetection of Sdh1 and Sdh2 (25). Anti-porin was purchased from ThermoFisher Scientific (catalog # 459500).

Analysis of Sdh1-bound flavin adenine dinucleotide (FAD) and total mitochondrial FAD

Levels of FAD covalently bound to Sdh1 were analyzed as previously described (14,26). Briefly, mitochondrial proteins were resolved on SDS-PAGE and the gel was placed in a 10% acetic acid solution for 20 minutes to oxidize flavin. FAD was visualized upon exposure to UV light using a Bio-Rad ChemiDoc MP Imaging System.

Oxygen consumption assay

Yeast strains were grown to confluency in glucose-based media and then switched to glycerol-based media for 12 hours. Two million cells were plated onto commercially available microplates with oxygen sensors (Oxoplate; PreSens catalog #OP96U) (27). Kinetic reading of oxygen consumption was measured using a spectrofluorometer. A Student's t-test was used to determine statistical significance between a variant of interest and yeast complemented with WT Sdh1.

Computational modeling of Sdh1 variants

A model of yeast Sdh1 (Q00711) from the Swiss Model repository was refined using Yasara Homology Modelling to include the liganded flavin adenine dinucleotide (FAD) interactions with the peptide chain, followed by optimization of the loop and side-chains interactions. Side-chain rotamers were fine-tuned considering electrostatic and knowledge-based packing interactions as well as solvation effects. An unrestrained high-resolution refinement with explicit solvent molecules was run, using YAMBER, a second-generation self-parameterizing force field derived from the AMBER force field. Clinically identified variants were introduced in the homology model, and the structures were minimized as described earlier. The variants were compared with the wild-type (WT) model of *E. coli* or yeast Sdh1 using PYMOL.

FoldX analysis

FoldX was used to measure the effect of point mutations on the stability of the Sdh1 yeast model (28). Except as noted, we used the default software settings. The move neighbors setting was turned off, and the average ΔG was calculated after three runs.

Mutation analysis

We gathered *SDHA* VUS from two primary sources: The OHSU Knight Diagnostics Laboratory (Portland, OR) and the literature(12,13). The majority of the variants pulled from the literature were found in a review highlighting the need for a functional model to characterize *SDHx* variants of unknown significance (12). The remaining literature variants are from a paper identifying novel causes of GIST that were WT for any known oncogenic driver of GIST (13). A collaboration with the OHSU Knight Diagnostics Laboratory, which offers a targeted exome panel to identify genetic drivers in GIST which includes all four *SDHx* subunits (<https://www.ohsu.edu/custom/knight-diagnostic-labs/home/test-details?id=GeneTrails+GIST+Genotyping+Panel>), lead to identification of several novel variants. All of the variants came from tumor samples that lacked any known oncogenic driver of GIST (e.g. no KIT mutations).

Results

Clinical analysis

All the known clinical information on the *SDHA* VUS identified in this study including SDHA/B IHC, tumor mutant allele fraction, other disease references, population data from the ExAC database (29), and ClinVar clinical significance (30) is listed in Table 1. Unfortunately, there were few variants with complete clinical and pathological annotation, emphasizing the challenge of trying to understand the functional effects of these variants from available clinical data alone. Currently, SDHx subunit testing remains uncommon for GIST and only a limited number of reference laboratories offer this testing. However, these laboratories usually do not have access to full clinical annotation and/or prior SDHB IHC testing results. This limitation applies to the published cases as well as the cases from the Knight Diagnostic laboratory. We listed all available additional clinical information for tumors with VUS in supplemental table 2.

Functional studies in yeast

To understand the functional consequences of VUS in human *SDHA*, we used complementation studies in a yeast strain lacking Sdh1 (*sdh1*). The flavoprotein subunit of the SDH complex is highly conserved across all species, including yeast Sdh1 (ySdh1) and human SDHA (hSDHA) (Supplemental Figure 1). For simplicity, we use yeast nomenclature throughout this study and reference the corresponding human variant in all of the tables.

As a positive control, we generated a WT *SDHI* amplicon from BY4741 that included the promoter and the 3' UTR which was cloned into a pRS416 plasmid and expressed in *sdh1* yeast. We used *sdh1* as our negative control. *Sdh1* do not have a functional SDH complex and were unable to utilize their mitochondria for canonical TCA or ETC pathways, allowing growth on a fermentable carbon source (glucose), but preventing growth with a non-fermentable carbon source such as glycerol (Figure 1). Complementation with WT *SDHI* (+Sdh1) restored growth on glycerol (Figure 1). Using this strategy, we tested the ability of plasmids encoding individual VUS to complement the *sdh1* strain and restore a normal growth phenotype. The two loss-of-function controls (yR19X and yH90S), as well as 16 of the variants shown in Figure 1 (yG97R, yT134M, yR179W, yG251R, yH287Y, yR303C, yY399C, yG410R, yC431F, yG432E, yR444C, yR444H, yA447E, yR458W, yR582G, yH601Y) were unable to grow on glycerol, indicating that these variants result in loss of oxidative phosphorylation (Figure 1A). In contrast, six of the variants (yN109S, yR162H, yR186W, yT501I, yW605FF, yV633I) were able to grow on glycerol, indicating these variants had no effect on the Sdh1 function as their phenotype was the same as WT Sdh1 (Figure 1B).

As a secondary measure of SDH complex activity, we measured oxygen consumption to assess oxidative phosphorylation, a surrogate for measuring the ability of the SDH complex to transfer electrons to the ETC. In concordance with our previous results, the same 18 variants that were unable to grow on glycerol consumed significantly less ($p < 0.0001$) oxygen than WT Sdh1 (Figure 1C). Variants that grew on glycerol had similar oxygen consumption to WT Sdh1, confirming that these variants do not disrupt the ETC (Figure

1D). To ensure oxygen consumption was due to the ETC, azide was added to each cell strain, and this promptly abolished oxygen consumption for all controls and variants.

To better understand how each variant affects Sdh1 protein function, we measured the covalent attachment of FAD to Sdh1, a process known as flavination. We also measured the protein abundance of Sdh1 and Sdh2 (Figure 2). Sdh1 flavination is critical for the catalytic activity of Sdh1; without covalent attachment of flavin, the Sdh1 protein is unable to oxidize succinate (14). As a positive control for a variant that inhibited insertion of FAD into Sdh1, we used the previously described Sdh1 H90S variant (14). Variants that complemented *sdh1* yeast in our functional assays had similar levels of flavinated Sdh1, total Sdh1, and total Sdh2 as yeast complemented with WT Sdh1 (Figure 2D). However, variants that were unable to perform oxidative phosphorylation had differential effects on the levels of flavinated Sdh1 and total Sdh1 protein (Figure 2 A–C). Some loss-of-function variants inhibited flavination without affecting the abundance of Sdh1 (H90S, G97R, T134M, R444C, H601Y) while others caused a marked decrease in Sdh1 protein abundance (R179W, G251R, H287Y, R303C, Y399C, G410R, C431F, G432E, R444H, A447E, R458W, R582G). All of the loss-of-function variants resulted in a decrease in Sdh2 protein abundance, consistent with loss of a functional SDH complex as described previously (25). In a subset of no effect and loss-of-function variants, we confirmed that Sdh1 mRNA expression was not significantly different from WT.

Based on all of the above results, we characterized 16 variants (73%) as causing loss of function and six variants (27%) as having no effect on SDH function (Table 2). All the loss-of-function variants were unable to grow on glycerol, had decreased oxygen consumption, and showed decreased Sdh2 protein abundance. In contrast, the no effect group had similar results to cells complemented with WT Sdh1 protein in these assays.

Computational modeling

We next performed computational modeling to predict the potential structural implications of each variant. Using the two groups (loss-of-function and no effect) identified in our yeast model, we further characterized each variant by visualizing the location in the Sdh1 protein. Both groups, loss-of-function and no effect, were located throughout the four domains of Sdh1, suggesting that there are not “hotspot” areas or domains for pathogenic variants, unlike the situation for some proteins. Within the loss-of-function group, 12 variants (yH90S (loss-of-function control), yG97R, yG251R, yH287Y, yR303C, yY399C, yG432E, yR444C, yR444H, yA447E, yR582G, yH601Y) were identified as being involved in cofactor (FAD) or substrate (succinate) binding in the Sdh1 active site (Figure 3, Table 2, and Supplemental Video 1). Structural studies from the homologous protein quinol:fumarate reductase (QFR) in *E. coli* suggests an important catalytic function for most of these amino acids (31). In addition to the amino acids directly interacting with the flavin binding pocket, it has been previously shown that the C-terminal domain of Sdh1 is crucial for the flavination of Sdh1 (25). Two variants located in the C-terminal domain of Sdh1, yR582G and yH601Y, both inhibited flavination of Sdh1. To visualize the substrate (succinate) and cofactor (FAD), which were not visualized using the yeast structure, these Sdh1 variants were modeled using the *E. Coli* crystal structure of Sdh1.

The other six loss-of-function variants (yR19X (control-not picture in computational models because it is an early truncation of the protein), yT134M, yR179W, yG410R, yG431F, yR458W) are not located in the active site of Sdh1. Changes in protein structure based on each of these variants are visualized in Figure 4A and Supplemental Video 1. Further, FoldX analyses of the variants compared to WT provided insight on how the variants affect stability (Supplemental Table 2) (28). yT134M, yR179W, yG410R, and yG431F all have positive

G (change in free energy) indicating these changes are highly destabilizing. Some of these also were associated with decreased Sdh1 protein abundance (Figure 2).

Interestingly, all the variants with no functional effects were localized to the surface of the protein (Figure 4B, Supplemental Video 1) where it would be predicted that they would be less likely to affect protein structure and function. Notably, these mutations did not map into any of the known critical domains of Sdh1 or predicted Sdh2 interaction sites.

Discussion

Molecular testing plays an important role in the clinical management of GIST, including decision making about appropriate medical and surgical therapy. SDH deficiency is a potential cause of GIST in tumors lacking known oncogenic drivers such as gain-of-function KIT mutations. To attempt to identify the cause of SDH deficiency in such GIST, it is necessary to use multi-gene sequencing panels to search for SDHx pathogenic mutations, however, as the routine use of multi-gene sequencing panels has expanded, there has also been an increase in reported VUS. Due to the multiple potential causes of SDH-deficiency, it is difficult to determine if a newly discovered VUS is responsible for the defect, or if a different genetic mechanism is responsible for the loss of SDH complex activity. This problem is exacerbated in GIST as loss of SDHA protein is the most common cause of SDH deficiency in GIST, but SDHA variants have not been extensively characterized at the functional level. Finally, many SDHA VUS are also found in the germline, meaning that they can be inherited. Knowing a patient has a loss-of-function *SDHA* variant in a tumor dramatically changes clinical decision making concerning screening recommendations for tumor syndromes, which affects both the patient and other family members. Since there are no effective medical treatments for advanced SDH-deficient tumors, early detection of tumors allowing curative surgical resection is crucial.

A major problem in the field has been the lack of a validated model system to assess the biochemical function of *SDHA* VUS. Currently, there are no human *SDHA*-deficient cell lines and assessing SDH complex activity in tumor samples is difficult. Previously it has been shown that a yeast model can be used to evaluate the pathogenic significance of *SDHB* mutations (17,18). Extending these studies, we report the successful use of a yeast model to characterize *SDHA* variants found in GIST tumors as either causing loss of function or having no biochemical consequence. Together with computational modeling, structural homology, and patient data, we have drawn conclusions on how and why the *SDHA* variants we studied affect SDH function, providing clinicians with information to guide genetic counseling of patients and family members who harbor one of these germline VUS.

Using our yeast model, we characterized variants as either causing loss of protein function or having no effect. All variants in the loss-of-function group were unable to grow on a non-fermentable carbon source (glycerol). In addition, these variants also failed to consume oxygen, indicating defective oxidative phosphorylation. Interestingly, these variants had differential effects on Sdh1 flavination and/or protein abundance. As a group, all of the loss-of-function variants were associated with a dramatic decrease in Sdh2 protein abundance, which is in concordance with current clinical testing to determine SDH-deficiency in tumors—assessment of SDHB (human equivalent to ySdh2) expression using IHC (2,32-36). Our data provides further evidence supporting the role of SDHB IHC as the most reliable clinical test to identify SDH deficiency in a tumor. Notably, we identified four novel loss-of-function variants that did not affect Sdh1 protein abundance (yG97R, yT134M, yR444C, yH601Y). These variants prevent flavination of Sdh1 making Sdh1 dysfunctional but still expressed at normal levels. Our findings are consistent with previous reports where SDHA expression was retained in some *SDHA*-mutant tumors that lack SDHB expression (37), confirming that not all dysfunctional SDHA (or Sdh1) variants lead to decreased protein expression of this subunit. There are three clinical samples (R188W, G260R, A454E) that we characterized as loss-of-function variants but are associated with normal *sdh1* in our yeast model. Based on these results and coupled with the fact the SDHA IHC is not universally available, we advocate for the use of SDHB IHC to identify tumors that are SDH-deficient.

The conclusions drawn from our yeast model are largely supported by our computational modeling results. For many variants, we could find confirmatory structural modeling evidence that the amino acids affected by these mutations played a role in the catalytic site of Sdh1. The computational model and energy minimization also allowed us to explain why some variants could reduce Sdh1 protein abundance. Furthermore, all the variants that did not affect protein function were on the surface of the protein as visualized by our computational model, suggesting that they would not interact with the catalytic mechanism of Sdh1.

Based on our biochemical data, there are still several tumors that have no known mechanism for loss of SDH activity (Table 1). One of these tumors had the hR195W (yR186W) variant with an allele frequency of 66.7% and negative SDHB IHC. This tumor is likely driven by a different mechanism of SDH-deficiency. The rest of the unexplained tumors had no IHC data available, so it is possible that these GISTs are not driven by SDH-deficiency and instead belong to a different subtype of GIST (11). Alternatively, these GISTs may be SDH-deficient by a different genetic mechanism than the *SDHA* variant identified using targeted exome sequencing. There are several weaknesses of various targeted exome sequencing panels including potential lack of coverage for regions of certain *SDHx* genes, failure to detect large genomic deletions involving SDHx subunits, as well as the inability to measure hypermethylation of the *SDHC* promoter. In addition, mutations that inactivate newly identified SDH assembly factors (SDHAF3, 4) are typically not included in clinical cancer gene sequencing panels (15,16). Given the limitations of using de-identified results from clinical testing, we did not have access to residual tumor samples to perform additional IHC or genomic testing.

We used a yeast model to characterize 22 *SDHA* VUS. This data revealed 16 (73%) of *SDHA* VUS as loss-of-function (and therefore pathogenic), highlighting the importance of understanding such variants to provide better clinical recommendations for genetic counselors concerning family screening and early detection protocols. However, our approach using a functional yeast model paired with computational modeling can distinguish between *SDHA* VUS that cause loss of function and those that have no biochemical effect, allowing us to discriminate between pathogenic and non-pathogenic variants.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

Thank you to Arin McKinley, Diana Griffith, Janice Patterson, and Ashley Young for their constant support during the completion of this project. The advice and feedback from Amanda McCullough, Maureen Hoatlin, and David Koeller were extremely beneficial for experimental design and interpretation of results. Also, thank you to Dennis Winge for his generous gift of primary antibodies for Sdh1 and Sdh2.

Financial support

Amber E Bannon-The author acknowledge of the support the OHSU OCTRI TL1TR000129

Oliver Harismendy-The authors acknowledge the support of NIH R21CA177519, U01CA196406

Jason K. Sicklick-The authors acknowledge the support of NIH K08CA168999, R21CA192072, and P30CA023100

Michael C. Heinrich-GIST Cancer Research Fund (MCH), Life Raft Group (MCH), and a Merit Review Grant from the Department of Veterans Affairs (MCH, 2I01BX000338-05)

References

1. Bezawork-Geleta A, Rohlena J, Dong L, Pacak K, Neuzil J. Mitochondrial Complex II: At the Crossroads. *Trends in biochemical sciences*. 2017; doi: 10.1016/j.tibs.2017.01.003
2. Janeway KA, Kim SY, Lodish M, Nose V, Rustin P, Gaal J, et al. Defects in succinate dehydrogenase in gastrointestinal stromal tumors lacking KIT and PDGFRA mutations. *Proceedings of the National Academy of Sciences of the United States of America*. 2011; 108(1): 314–8. DOI: 10.1073/pnas.1009199108 [PubMed: 21173220]
3. Burnichon N, Abermil N, Buffet A, Favier J, Gimenez-Roqueplo AP. The genetics of paragangliomas. *European annals of otorhinolaryngology, head and neck diseases*. 2012; 129(6): 315–8. DOI: 10.1016/j.anorl.2012.04.007
4. Baysal BE. A recurrent stop-codon mutation in succinate dehydrogenase subunit B gene in normal peripheral blood and childhood T-cell acute leukemia. *PLoS One*. 2007; 2(5):e436.doi: 10.1371/journal.pone.0000436 [PubMed: 17487275]
5. Beamer LC. Cowden syndrome: what oncology nurses need to know about increased risk of developing certain cancers. *Oncol Nurs Forum*. 2014; 41(5):555–7. DOI: 10.1188/14.ONF.555-557 [PubMed: 25158662]
6. Niemeijer ND, Papatomas TG, Korpershoek E, de Krijger RR, Oudijk L, Morreau H, et al. Succinate Dehydrogenase (SDH)-Deficient Pancreatic Neuroendocrine Tumor Expands the SDH-Related Tumor Spectrum. *The Journal of clinical endocrinology and metabolism*. 2015; 100(10):E1386–93. DOI: 10.1210/jc.2015-2689 [PubMed: 26259135]
7. van Nederveen FH, Gaal J, Favier J, Korpershoek E, Oldenburg RA, de Bruyn EM, et al. An immunohistochemical procedure to detect patients with paraganglioma and pheochromocytoma with germline SDHB, SDHC, or SDHD gene mutations: a retrospective and prospective analysis.

- The Lancet Oncology. 2009; 10(8):764–71. DOI: 10.1016/s1470-2045(09)70164-0 [PubMed: 19576851]
8. Killian JK, Miettinen M, Walker RL, Wang Y, Zhu YJ, Waterfall JJ, et al. Recurrent epimutation of SDHC in gastrointestinal stromal tumors. *Science translational medicine*. 2014; 6(268): 268ra177.doi: 10.1126/scitranslmed.3009961
 9. Ricketts CJ, Forman JR, Rattenberry E, Bradshaw N, Laloo F, Izatt L, et al. Tumor risks and genotype-phenotype-proteotype analysis in 358 patients with germline mutations in SDHB and SDHD. *Human mutation*. 2010; 31(1):41–51. DOI: 10.1002/humu.21136 [PubMed: 19802898]
 10. Neumann HP, Pawlu C, Peczkowska M, Bausch B, McWhinney SR, Muresan M, et al. Distinct clinical features of paraganglioma syndromes associated with SDHB and SDHD gene mutations. *Jama*. 2004; 292(8):943–51. DOI: 10.1001/jama.292.8.943 [PubMed: 15328326]
 11. Bannon AE, Klug LR, Corless CL, Heinrich MC. Using molecular diagnostic testing to personalize the treatment of patients with gastrointestinal stromal tumors. *Expert review of molecular diagnostics*. 2017; doi: 10.1080/14737159.2017.1308826
 12. Evenepoel L, Papatomas TG, Krol N, Korpershoek E, de Krijger RR, Persu A, et al. Toward an improved definition of the genetic and tumor spectrum associated with SDH germ-line mutations. *Genetics in medicine : official journal of the American College of Medical Genetics*. 2015; 17(8): 610–20. DOI: 10.1038/gim.2014.162 [PubMed: 25394176]
 13. Shi E, Chmielecki J, Tang CM, Wang K, Heinrich MC, Kang G, et al. FGFR1 and NTRK3 actionable alterations in “Wild-Type” gastrointestinal stromal tumors. *Journal of translational medicine*. 2016; 14(1):339.doi: 10.1186/s12967-016-1075-6 [PubMed: 27974047]
 14. Hao HX, Khalimonchuk O, Schraders M, Dephoure N, Bayley JP, Kunst H, et al. SDH5, a gene required for flavination of succinate dehydrogenase, is mutated in paraganglioma. *Science*. 2009; 325(5944):1139–42. DOI: 10.1126/science.1175689 [PubMed: 19628817]
 15. Van Vranken JG, Bricker DK, Dephoure N, Gygi SP, Cox JE, Thummel CS, et al. SDHAF4 promotes mitochondrial succinate dehydrogenase activity and prevents neurodegeneration. *Cell metabolism*. 2014; 20(2):241–52. DOI: 10.1016/j.cmet.2014.05.012 [PubMed: 24954416]
 16. Na U, Yu W, Cox J, Bricker DK, Brockmann K, Rutter J, et al. The LYR factors SDHAF1 and SDHAF3 mediate maturation of the iron-sulfur subunit of succinate dehydrogenase. *Cell metabolism*. 2014; 20(2):253–66. DOI: 10.1016/j.cmet.2014.05.014 [PubMed: 24954417]
 17. Panizza E, Ercolino T, Mori L, Rapizzi E, Castellano M, Opocher G, et al. Yeast model for evaluating the pathogenic significance of SDHB, SDHC and SDHD mutations in PHEO-PGL syndrome. *Human molecular genetics*. 2013; 22(4):804–15. DOI: 10.1093/hmg/ddt487 [PubMed: 23175444]
 18. Goffrini P, Ercolino T, Panizza E, Giache V, Cavone L, Chiarugi A, et al. Functional study in a yeast model of a novel succinate dehydrogenase subunit B gene germline missense mutation (C191Y) diagnosed in a patient affected by a glomus tumor. *Human molecular genetics*. 2009; 18(10):1860–8. DOI: 10.1093/hmg/ddp102 [PubMed: 19261679]
 19. Szeto SS, Reinke SN, Sykes BD, Lemire BD. Ubiquinone-binding site mutations in the *Saccharomyces cerevisiae* succinate dehydrogenase generate superoxide and lead to the accumulation of succinate. *The Journal of biological chemistry*. 2007; 282(37):27518–26. DOI: 10.1074/jbc.M700601200 [PubMed: 17636259]
 20. Alston CL, Ceccatelli Berti C, Blakely EL, Olahova M, He L, McMahon CJ, et al. A recessive homozygous p.Asp92Gly SDHD mutation causes prenatal cardiomyopathy and a severe mitochondrial complex II deficiency. *Human genetics*. 2015; 134(8):869–79. DOI: 10.1007/s00439-015-1568-z [PubMed: 26008905]
 21. Fontanesi F, Diaz F, Barrientos A. Evaluation of the mitochondrial respiratory chain and oxidative phosphorylation system using yeast models of OXPHOS deficiencies. *Current protocols in human genetics*. 2009; Chapter 19:Unit19.5. doi: 10.1002/0471142905.hg1905s63
 22. Winzeler EA, Shoemaker DD, Astromoff A, Liang H, Anderson K, Andre B, et al. Functional characterization of the *Saccharomyces cerevisiae* genome by gene deletion and parallel analysis. *Science*. 1999; 285(5429):901–6. [PubMed: 10436161]

23. Sikorski RS, Hieter P. A system of shuttle vectors and yeast host strains designed for efficient manipulation of DNA in *Saccharomyces cerevisiae*. *Genetics*. 1989; 122(1):19–27. [PubMed: 2659436]
24. Muller F, Crofts AR, Kramer DM. Multiple Q-cycle bypass reactions at the Qo site of the cytochrome bc1 complex. *Biochemistry*. 2002; 41(25):7866–74. [PubMed: 12069575]
25. Kim HJ, Jeong MY, Na U, Winge DR. Flavinylation and assembly of succinate dehydrogenase are dependent on the C-terminal tail of the flavoprotein subunit. *The Journal of biological chemistry*. 2012; 287(48):40670–9. DOI: 10.1074/jbc.M112.405704 [PubMed: 23043141]
26. Bafunno V, Giancaspero TA, Brizio C, Bufano D, Passarella S, Boles E, et al. Riboflavin uptake and FAD synthesis in *Saccharomyces cerevisiae* mitochondria: involvement of the Flx1p carrier in FAD export. *The Journal of biological chemistry*. 2004; 279(1):95–102. DOI: 10.1074/jbc.M308230200 [PubMed: 14555654]
27. Kitanovic A, Bonowski F, Heigwer F, Ruoff P, Kitanovic I, Ungewiss C, et al. Acetic acid treatment in *S. cerevisiae* creates significant energy deficiency and nutrient starvation that is dependent on the activity of the mitochondrial transcriptional complex Hap2-3-4-5. *Frontiers in oncology*. 2012; 2:118.doi: 10.3389/fonc.2012.00118 [PubMed: 23050242]
28. Schymkowitz J, Borg J, Stricher F, Nys R, Rousseau F, Serrano L. The FoldX web server: an online force field. *Nucleic acids research*. 2005; 33:W382–8. (Web Server issue). DOI: 10.1093/nar/gki387 [PubMed: 15980494]
29. Lek M, Karczewski KJ, Minikel EV, Samocha KE, Banks E, Fennell T, et al. Analysis of protein-coding genetic variation in 60,706 humans. *Nature*. 2016; 536(7616):285–91. DOI: 10.1038/nature19057 [PubMed: 27535533]
30. Landrum MJ, Lee JM, Benson M, Brown G, Chao C, Chitipiralla S, et al. ClinVar: public archive of interpretations of clinically relevant variants. *Nucleic acids research*. 2016; 44(D1):D862–8. DOI: 10.1093/nar/gkv1222 [PubMed: 26582918]
31. Iverson TM. Catalytic mechanisms of complex II enzymes: a structural perspective. *Biochimica et biophysica acta*. 2013; 1827(5):648–57. DOI: 10.1016/j.bbabi.2012.09.008 [PubMed: 22995215]
32. Boikos SA, Pappo AS, Killian JK, LaQuaglia MP, Weldon CB, George S, et al. Molecular Subtypes of KIT/PDGFRA Wild-Type Gastrointestinal Stromal Tumors: A Report From the National Institutes of Health Gastrointestinal Stromal Tumor Clinic. *JAMA oncology*. 2016; 2(7): 922–8. DOI: 10.1001/jamaoncol.2016.0256 [PubMed: 27011036]
33. Gill AJ, Chou A, Vilain RE, Clifton-Bligh RJ. “Pediatric-type” gastrointestinal stromal tumors are SDHB negative (“type 2”) GISTs. *The American journal of surgical pathology*. 2011; 35(8):1245–7. author reply 7–8. DOI: 10.1097/PAS.0b013e3182217b93 [PubMed: 21753704]
34. Gill AJ, Benn DE, Chou A, Clarkson A, Muljono A, Meyer-Rochow GY, et al. Immunohistochemistry for SDHB triages genetic testing of SDHB, SDHC, and SDHD in paraganglioma-pheochromocytoma syndromes. *Human pathology*. 2010; 41(6):805–14. DOI: 10.1016/j.humpath.2009.12.005 [PubMed: 20236688]
35. Gaal J, Stratakis CA, Carney JA, Ball ER, Korpershoek E, Lodish MB, et al. SDHB immunohistochemistry: a useful tool in the diagnosis of Carney-Stratakis and Carney triad gastrointestinal stromal tumors. *Modern pathology : an official journal of the United States and Canadian Academy of Pathology, Inc.* 2011; 24(1):147–51. DOI: 10.1038/modpathol.2010.185
36. Doyle LA, Nelson D, Heinrich MC, Corless CL, Hornick JL. Loss of succinate dehydrogenase subunit B (SDHB) expression is limited to a distinctive subset of gastric wild-type gastrointestinal stromal tumours: a comprehensive genotype-phenotype correlation study. *Histopathology*. 2012; 61(5):801–9. DOI: 10.1111/j.1365-2559.2012.04300.x [PubMed: 22804613]
37. Miettinen M, Killian JK, Wang ZF, Lasota J, Lau C, Jones L, et al. Immunohistochemical loss of succinate dehydrogenase subunit A (SDHA) in gastrointestinal stromal tumors (GISTs) signals SDHA germline mutation. *The American journal of surgical pathology*. 2013; 37(2):234–40. DOI: 10.1097/PAS.0b013e3182671178 [PubMed: 23282968]
38. Italiano A, Chen CL, Sung YS, Singer S, DeMatteo RP, LaQuaglia MP, et al. SDHA loss of function mutations in a subset of young adult wild-type gastrointestinal stromal tumors. *BMC cancer*. 2012; 12:408.doi: 10.1186/1471-2407-12-408 [PubMed: 22974104]

39. Pantaleo MA, Nannini M, Astolfi A, Biasco G. A distinct pediatric-type gastrointestinal stromal tumor in adults: potential role of succinate dehydrogenase subunit A mutations. *The American journal of surgical pathology*. 2011; 35(11):1750–2. DOI: 10.1097/PAS.0b013e318230a523 [PubMed: 21997697]
40. Korpershoek E, Favier J, Gaal J, Burnichon N, van Gessel B, Oudijk L, et al. SDHA immunohistochemistry detects germline SDHA gene mutations in apparently sporadic paragangliomas and pheochromocytomas. *The Journal of clinical endocrinology and metabolism*. 2011; 96(9):E1472–6. DOI: 10.1210/jc.2011-1043 [PubMed: 21752896]
41. Heinrich M, Rankin C, Blanke CD, Demetri GD, Borden EC, Ryan CW, et al. Correlation of Long-term Results of Imatinib in Advanced Gastrointestinal Stromal Tumors With Next-Generation Sequencing Results: Analysis of Phase 3 SWOG Intergroup Trial S0033. *JAMA oncology*. 2017; doi: 10.1001/jamaoncol.2016.6728
42. Birch-Machin MA, Taylor RW, Cochran B, Ackrell BA, Turnbull DM. Late-onset optic atrophy, ataxia, and myopathy associated with a mutation of a complex II gene. *Annals of neurology*. 2000; 48(3):330–5. [PubMed: 10976639]
43. Pantaleo MA, Astolfi A, Urbini M, Nannini M, Paterini P, Indio V, et al. Analysis of all subunits, SDHA, SDHB, SDHC, SDHD, of the succinate dehydrogenase complex in KIT/PDGFRA wild-type GIST. *European journal of human genetics : EJHG*. 2014; 22(1):32–9. DOI: 10.1038/ejhg.2013.80 [PubMed: 23612575]
44. Alston CL, Davison JE, Meloni F, van der Westhuizen FH, He L, Hornig-Do HT, et al. Recessive germline SDHA and SDHB mutations causing leukodystrophy and isolated mitochondrial complex II deficiency. *Journal of medical genetics*. 2012; 49(9):569–77. DOI: 10.1136/jmedgenet-2012-101146 [PubMed: 22972948]
45. Burnichon N, Briere JJ, Libe R, Vescovo L, Riviere J, Tissier F, et al. SDHA is a tumor suppressor gene causing paraganglioma. *Human molecular genetics*. 2010; 19(15):3011–20. DOI: 10.1093/hmg/ddq206 [PubMed: 20484225]
46. Baysal BE, Lawrence EC, Ferrell RE. Sequence variation in human succinate dehydrogenase genes: evidence for long-term balancing selection on SDHA. *BMC biology*. 2007; 5:12. doi: 10.1186/1741-7007-5-12 [PubMed: 17376234]
47. Chapman KB, Solomon SD, Boeke JD. SDH1, the gene encoding the succinate dehydrogenase flavoprotein subunit from *Saccharomyces cerevisiae*. *Gene*. 1992; 118(1):131–6. [PubMed: 1511876]
48. Cecchini G, Schroder I, Gunsalus RP, Maklashina E. Succinate dehydrogenase and fumarate reductase from *Escherichia coli*. *Biochimica et biophysica acta*. 2002; 1553(1–2):140–57. [PubMed: 11803023]

Translational relevance

Molecular testing plays an important role in the clinical management of gastrointestinal stromal tumors (GIST), including decision making about the most appropriate medical or surgical therapy. As the routine use of multi-gene sequencing panels has expanded, there has also been an increase in reported variants of unknown significance (VUS) of the *SDHA* gene. In many cases, these *SDHA* VUS are present in the germline and are therefore potentially heritable by other family members. In order to understand the functional consequences of these variants, we combined clinical observations, data from a functional yeast model, and computational modeling to classify these *SDHA* VUS as having no effect or causing loss of function. These results will be helpful for appropriate genetic counseling of individuals with these germline variants. In addition, our data highlight the limitations of SDHA immunohistochemistry in clinical testing of tumors with SDHA VUS.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript




Figure 1. ySdh1 variants causing loss of function are unable to use glycerol as a sole carbon source or consume oxygen

Yeast were grown to a confluency of 2×10^7 /mL and then serially diluted and plated onto either glucose or glycerol media plates (Left to right: highest to lowest). All variants, regardless of SDH complex function, can grow on glucose but only those variants that complement *sdh1* and restore SDH complex activity are able to grow on glycerol and consume oxygen. A. Loss-of-function variants that are unable to grow with glycerol as their sole carbon source. B. Variants with no effect are able to grow with glycerol as their sole carbon source. C. Sdh1 variants that consume significantly less oxygen than cells

complemented with WT Sdh1 (+Sdh1 control). D. Variants that consume oxygen at a similar rate to the +Sdh1 control. **** p-value <0.0001

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript




Figure 2. Abundance of flavinated Sdh1, total Sdh1, and total Sdh2 in mitochondrial lysates from yeast expressing each variant

Western blotting of mitochondrial lysates from each variant are shown. Positive and negative control lanes are included on each western blot (WT, SDH1delta, +SDH1). WT and +Sdh1 show bands at expected size for each protein. Sdh1 does not show bands for Sdh1-FAD, Sdh1, or Sdh2, but does have normal expression of the mitochondrial marker, porin. A-C. Loss-of-function variants are shown in numerical order by altered amino acid residue. D. No effect variants are shown in numerical order by altered amino acid residue and have with no reduction in flavin, Sdh1, or Sdh2 expression compared to WT controls. E. Sdh2 protein abundance was calculated relative to Sdh1 and then normalized to porin (control for mitochondrial protein loading). The no effect variants are labeled in black, while the loss-of-function variants are in gray. No effect variants have Sdh2 protein abundance more similar to complemented Sdh1 and loss-of-function variants have decreased Sdh2 similar to *sdh1* . F. Ratio of Sdh1/Sdh2 protein abundance in no effect variants.




Figure 3. Variants involving the active site cause loss of Sdh1 function

A ribbon representation of Sdh1 (*E. coli* model PBD file 2WP9) is shown with WT protein carbons colored gray and the variant carbons colored red. The FAD is shown as teal-colored spheres while the dicarboxylic acid substrate (succinate analog) is depicted with yellow-colored spheres. In each case, the view has been rotated so that residues of interest are clearly observed. References detailing structural implications of each variant can be found in Table 2.




Figure 4. Structural consequences of variants visualized in a yeast model (yeast model PDB file 1ORZ)

A. Variants not in the active site causing loss-of-function. WT Sdh1 protein is represented as a gray ribbon with the amino acid of interest labeled as a gray stick and the re-folded variant Sdh1 protein as a red ribbon with the amino acid of interest indicated by a red stick. In each case, the view has been rotated such that residues of interest are clearly observed. Potential structural implications of each variant can be found in Table 2. B. Variants that do not affect protein function are located on the surface of Sdh1. The surface area of WT Sdh1 is shown

in gray, the variant residue is labeled red, and the surface area of Sdh2 is labeled in green. In each case, the view has been rotated such that residues of interest are clearly observed.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Table 1

Clinical variants of human *SDHA* with unknown significance

All of the genetic variants were found in GIST, except for hG260R which was found in a paraganglioma. A summary of available clinical data for each variant is listed, including source of the variant, clinical IHC results for SDHB/SDHA, frequency of the variant in tumor, other *SDHA* variants found in the same tumor, information on functional effect of other variants in the tumor, other disease references to the variant of interest in the literature, population allele frequency, and results from our yeast model. The no effect variants are highlighted in grey for clarity. Loss of function (LOF); Not available (N/A); Not Done (ND); neg (negative-SDHx absent); pos (positive-SDHx expression).

hSDHA mutation	ySdh1 mutation	Source	SDHB/SDHA IHC	Tumor mutant allele fraction	Other disease references	Allele frequency from EXAC database	ClinVar clinical significance	Conclusions drawn
R31X	R19X	Control	Neg/neg (38)	N/A	GIST (38,39) Paraganglioma (40)	1.647×10^{-4}	Pathogenic/Likely pathogenic	LOF (control)
H99S	H90S	Control (14)	N/A	N/A	None	N/A	N/A (H99Y Likely pathogenic)	LOF (control)
G106R	G97R	OHSU	N/A	92%	Novel	N/A	N/A	LOF
N118S	N109S	OHSU	N/A	50.7%	Novel	N/A	N/A	No effect
T143M	T134M	Literature (13)	N/A	22%	Novel	N/A (T143R reported 8.31×10^{-6})	Uncertain significance	LOF
R171H	R162H	OHSU	N/A	34%	Novel	8.24×10^{-6}	Uncertain significance	No effect
R171H	R162H	OHSU	N/A	40.3%	Novel	8.24×10^{-6}	Uncertain significance	No effect
R188W	R179W	Literature (12)	Neg/pos [30]	N/A	None	N/A (R188Q reported 1.65×10^{-5})	N/A	LOF
R195W	R186W	OHSU	Neg/ND	66.7%	GIST (37)	N/A (R195Q reported 8.23×10^{-6})	N/A	No effect
G260R	G251R	Literature (12)	Pos/pos [32]	N/A	None	N/A	Uncertain significance	LOF
H296Y	H287Y	Literature (13)	N/A	87%	Novel	N/A	N/A	LOF
R312C	R303C	OHSU	N/A	34.5%	Novel	N/A	N/A (R312P Uncertain significance)	LOF
R408C	Y399C	OHSU	N/A	52.8%	GIST(41); late onset neurodegenerative disease (42)	N/A	N/A	LOF
G419R	G410R	OHSU	N/A	77%	GIST (43)	N/A	N/A	LOF
C438F	C431F	OHSU	N/A	48.5%	Novel	N/A	N/A	LOF

G439E	G432E	OHSU	N/A	87.6%	Novel	N/A	N/A	N/A	LOF
R451C	R444C	Literature (13)	N/A	18%	Complex II deficiency (42)	N/A	N/A (R451H reported 8.23×10^{-6})	N/A	LOF
R451H	R444H	OHSU	N/A	47.5%	Novel	N/A	8.23×10^{-6}	N/A	LOF
R451H	R444H	OHSU	N/A	39.5%	Novel	N/A	8.23×10^{-6}	N/A	LOF
A454E	A447E	Literature (12)	Neg/pos [31]	N/A	None	N/A	N/A	N/A (A454T Uncertain significance)	LOF
R465W	R458W	OHSU	N/A	50%	Novel	N/A	8.23×10^{-6}	Uncertain significance	LOF
T508I	T501I	OHSU	N/A	50.6%	Cardiomyopathy and Leukodystrophy (44)	N/A	7.60×10^{-4}	Conflicting interpretations of pathogenicity	No effect
R589G	R582G	OHSU	N/A	40.2%	Paraganglioma (45)	N/A	8.29×10^{-6}	Uncertain significance	LOF
H625W	H601Y	Literature (12)	Neg/neg [39]	N/A	Pituitary adenoma and Pheochromocytoma/paraganglioma	N/A	N/A	N/A	LOF
Y629F	W605F	OHSU	N/A	99.6%		N/A	1.52×10^{-1}	Benign/likely benign	No effect
V657I	V633I	OHSU	N/A	95.1%	Not pathogenic (46)	N/A	1.30×10^{-1}	Likely benign	No effect

Table 2
Consistent findings across multiple functional assays for each variant in our yeast model

A summary of results including the growth on glycerol, oxygen consumption, Sdh2 protein abundance, consequences of changing the WT amino acid using computational modeling and literature search for studies in other species' flavoprotein are tabulated along with our classification of each variant based on our yeast model. The no effect variants are highlighted in grey for clarity. Numerical values for oxygen consumption were taken from Figure 1C and D. Numerical values for Sdh2 protein were taken from Figure 2E.

hSDHA mutation	ySdh1 mutation	Growth phenotype on glycerol	Oxygen consumption (% of complemented)	Sdh2 protein (% of complemented)	Structural implications of variants	Group name
R31X (control)	R19X (control)	Unable to grow	Decreased (20%)	Decreased (0%)	Truncates mitochondrial targeting sequence (47)	Loss-of-function (control)
H99S (control)	H90S (control)	Unable to grow	Decreased (-2%)	Decreased (15%)	Inhibits covalent bond to FAD (14)	Loss-of-function (control)
G106R	G97R	Unable to grow	Decreased (7%)	Decreased (14%)	Distorts the active site (31)	Loss-of-function
N118S	N109S	Growth	Similar (108%)	Similar (83%)	Surface of protein	No effect
T143M	T134M	Unable to grow	Decreased (3%)	Decreased (35%)	No obvious disturbances but causes loss of function	Loss-of-function
R171H	R162H	Growth	Similar (84%)	Similar (78%)	Surface of protein	No effect
R188W	R179W	Unable to grow	Decreased (21%)	Decreased (0%)	Disrupts salt bridge with D108	Loss-of-function
R195W	R186W	Growth	Similar (91%)	Similar (82%)	Surface of protein	No effect
G260R	G251R	Unable to grow	Decreased (27%)	Decreased (0%)	Obstructs flavin binding pocket	Loss-of-function
H296Y	H287Y	Unable to grow	Decreased (4%)	Decreased (11%)	Inhibits succinate binding (31)	Loss-of-function
R312C	R303C	Unable to grow	Decreased (4%)	Decreased (3%)	Contributes to proper orientation and activation of the flavin isoalloxazine ring to facilitate formation of the covalent FAD bond and disrupts salt bridge (48)	Loss-of-function
R408C	Y399C	Unable to grow	Decreased (9%)	Decreased (4%)	In flavin binding site	Loss-of-function
G419R	G410R	Unable to grow	Decreased (19%)	Decreased (2%)	Distorts helix	Loss-of-function
C438F	C431F	Unable to grow	Decreased (19%)	Decreased (2%)	Bulky change disrupts helix	Loss-of-function
G439E	G432E	Unable to grow	Decreased (-2%)	Decreased (3%)	Obstructs flavin binding pocket	Loss-of-function
R451C	R444C	Unable to grow	Decreased (12%)	Decreased (6%)	Disrupts flavin binding, succinate binding and proton shuttle necessary for catalytic activity (31)	Loss-of-function
R451H	R444H	Unable to grow	Decreased (3%)	Decreased (0%)	Disrupts flavin binding, succinate binding and proton shuttle necessary for catalytic activity (31)	Loss-of-function
A454E	A447E	Unable to grow	Decreased (11%)	Decreased (0%)	Lines succinate binding site (31)	Loss-of-function
R465W	R458W	Unable to grow	Decreased (5%)	Decreased (21%)	Disrupts salt bridge with E136	Loss-of-function

hSDHA mutation	γSdhI mutation	Growth phenotype on glycerol	Oxygen consumption (% of complemented)	Sdh2 protein (% of complemented)	Structural implications of variants	Group name
T508I	T501I	Growth	Similar (113%)	Similar (75%)	Surface of protein	No effect
R589G	R582G	Unable to grow	Decreased (12%)	Decreased (0%)	Inhibits C-terminal flavination (25)	Loss-of-function
H625W	H601Y	Unable to grow	Decreased (18%)	Decreased (2%)	Inhibits C-terminal flavination	Loss-of-function
Y629F	W605F	Growth	Similar (113%)	Similar (101%)	Surface of protein	No effect
V657I	V633I	Growth	Similar (108%)	Similar (82%)	Surface of protein	No effect