



BP180 dysfunction triggers spontaneous skin inflammation in mice

Yang Zhang^{a,b,1}, Bin-Jin Hwang^{c,1}, Zhen Liu^{b,d}, Ning Li^b, Kendall Lough^e, Scott E. Williams^e, Jinbo Chen^{b,f}, Susan W. Burette^b, Luis A. Diaz^b, Maureen A. Su^{c,g}, Shengxiang Xiao^{a,2}, and Zhi Liu^{b,c,h,2}

^aDepartment of Dermatology, The Second Affiliated Hospital, School of Medicine, Xi'an Jiaotong University, Xi'an, 710004 Shaanxi, China; ^bDepartment of Dermatology, School of Medicine, University of North Carolina at Chapel Hill, Chapel Hill, NC 27599; ^cDepartment of Microbiology and Immunology, School of Medicine, University of North Carolina at Chapel Hill, Chapel Hill, NC 27599; ^dGuangdong Center for Adverse Drug Reactions of Monitoring, 510000 Guangzhou, China; ^eDepartment of Pathology and Laboratory Medicine, School of Medicine, University of North Carolina at Chapel Hill, Chapel Hill, NC 27599; ^fWuhan No. 1 Hospital, The Fourth Affiliated Hospital, Tongji Medical College, Huazhong University of Science and Technology, 430022 Wuhan, China; ^gDepartment of Pediatrics, School of Medicine, University of North Carolina at Chapel Hill, Chapel Hill, NC 27599; and ^hLineberger Comprehensive Cancer Center, University of North Carolina at Chapel Hill, Chapel Hill, NC 27599

Edited by Valerie Horsley, Yale University, New Haven, CT, and accepted by Editorial Board Member Ruslan Medzhitov May 9, 2018 (received for review December 18, 2017)

BP180, also known as collagen XVII, is a hemidesmosomal component and plays a key role in maintaining skin dermal/epidermal adhesion. Dysfunction of BP180, either through genetic mutations in junctional epidermolysis bullosa (JEB) or autoantibody insult in bullous pemphigoid (BP), leads to subepidermal blistering accompanied by skin inflammation. However, whether BP180 is involved in skin inflammation remains unknown. To address this question, we generated a BP180-dysfunctional mouse strain and found that mice lacking functional BP180 (termed Δ NC16A) developed spontaneous skin inflammatory disease, characterized by severe itch, defective skin barrier, infiltrating immune cells, elevated serum IgE levels, and increased expression of thymic stromal lymphopoietin (TSLP). Severe itch is independent of adaptive immunity and histamine, but dependent on increased expression of TSLP by keratinocytes. In addition, a high TSLP expression is detected in BP patients. Our data provide direct evidence showing that BP180 regulates skin inflammation independently of adaptive immunity, and BP180 dysfunction leads to a TSLP-mediated itch. The newly developed mouse strain could be a model for elucidation of disease mechanisms and development of novel therapeutic strategies for skin inflammation and BP180-related skin conditions.

collagen XVII | hemidesmosome | atopic dermatitis | TSLP | keratinocyte

BP180, also known as collagen XVII, is a 180-kDa transmembrane glycoprotein of the hemidesmosome anchoring basal keratinocytes into the underneath basal membrane of the skin (1). The intracellular region of BP180 is linked to the intermediate filament network, and its extracellular portion is anchored into the basement membrane zone (BMZ) through interacting with extracellular matrix proteins (2, 3). Dysfunction of BP180 by gene mutations can lead to junctional epidermolysis bullosa (JEB) (4, 5), a subepidermal blistering disease in humans. In the skin autoimmune subepidermal blistering disease bullous pemphigoid (BP), anti-BP180 autoantibodies attack and impair function of BP180 autoantigen in basal keratinocytes causing dermal-epidermal separation (6). Pathogenic anti-BP180 autoantibodies mainly target the juxtamembranous noncollagenous NC16A domain (7). These findings in JEB and BP establish BP180 as a key cell-cell matrix adhesion molecule in the skin. However, whether BP180 is involved in other biological processes and pathological conditions is largely unknown.

Both BP and JEB share some features of skin inflammation. BP is characterized by skin infiltration of immune cells, increased IgE, and pruritus (1, 8). JEB is a very rare genetic disease, and some JEB patients developed atopic dermatitis (AD)-like skin inflammation including inflammatory cell infiltration and elevated serum IgE (9). Snauwaert et al. (10) found that itch occurred in 100% of the JEB patients in their study. Previously, another group generated a BP180 dysfunctional mouse strain (termed Δ NC14A

mice), which shows an increased itch, eosinophil influx, and serum IgE (11). However, potential molecular and/or cellular mechanisms underlying the BP180 dysfunction-associated phenotypes were not determined.

To uncover functions of BP180 beside its cell-matrix adhesion property, we generated a BP180 dysfunctional mouse model by removal of the NC16A domain (termed Δ NC16A mice). Mice expressing the NC16A-truncated BP180 developed subepidermal blistering with severe skin inflammation and itch. The spontaneous skin inflammation and itch in Δ NC16A mice were neither dependent on histamine nor adaptive immunity, but dependent on TSLP. Moreover, high TSLP expression is also found in BP patients. This report demonstrates that BP180 dysfunction leads to TSLP-mediated skin inflammation.

Results

Generation of Δ NC16A Mice. Human and mouse BP180 lack immune cross-reactivity (12). To study the immunopathogenesis of

Significance

BP180 is a cell-cell matrix adhesion molecule and is critical for attachment of epidermis to underlying dermis. We generated a dysfunctional BP180 mouse strain (Δ NC16A mice). The Δ NC16A mice at 8 weeks start developing spontaneous itch and skin inflammation characterized by increased skin thickness, infiltrating immune cells, increased levels of IgE in circulation, proinflammatory mediators in the skin, and impaired skin barrier. Bullous pemphigoid (BP) patients also present increased TSLP in lesional skin and circulation. Our findings suggest that BP180 regulates pruritic skin inflammation. Elucidation of molecular and cellular mechanisms underlying the role of BP180 in skin inflammation should help better understanding of pathophysiology of skin inflammation and identify therapeutic targets for skin inflammatory diseases such as BP and atopic dermatitis.

Author contributions: Y.Z., B.-J.H., N.L., S.E.W., J.C., L.A.D., M.A.S., S.X., and Zhi Liu designed research; Y.Z., B.-J.H., Zhen Liu, and S.W.B. performed research; Y.Z., K.L., S.E.W., and S.X. contributed new reagents/analytic tools; Y.Z., B.-J.H., Zhen Liu, N.L., S.W.B., and Zhi Liu analyzed data; and Y.Z., B.-J.H., N.L., S.E.W., J.C., M.A.S., S.X., and Zhi Liu wrote the paper.

The authors declare no conflict of interest.

This article is a PNAS Direct Submission. V.H. is a guest editor invited by the Editorial Board.

Published under the [PNAS license](#).

¹Y.Z. and B.-J.H. contributed equally to this work.

²To whom correspondence may be addressed. Email: xiao_sx@163.com or zhi_liu@med.unc.edu.

This article contains supporting information online at www.pnas.org/lookup/suppl/doi:10.1073/pnas.1721805115/-DCSupplemental.

Published online June 4, 2018.




Fig. 2. $\Delta NC16A$ mice exhibit aberrant itch, increased TSLP level, and defective skin barrier. Compared with WT mice, $\Delta NC16A$ mice starting 8 wk old exhibited a significantly increased epidermis thickness (A), itch (scratch motion) (B), serum IgE levels (C), and TSLP level in ear skin (D) ($n = 8$ per group). (E) Toluidine blue penetration assay showed no barrier defect in E18.5 embryos of $\Delta NC16A$ and WT mice ($n = 8$ per group). Adult $\Delta NC16A$ mice started to show a significant reduction in barrier function at 8~12 wk after birth, by Even Blue (F) ($n = 6$ per group) and FITC-conjugated BSA permeability assays (G). (Scale bars: 50 μm .) * $P < 0.05$, Student's t test, graphs A, B, and F show mean \pm SE.

infiltration (including neutrophils, eosinophils, T cells, and mast cells) (SI Appendix, Fig. S3 B–F), increased serum IgE, high TSLP expression, and impaired skin barrier ($n = 8$) (SI Appendix, Fig. S3 G–I). Therefore, BP180 dysfunction in basal keratinocytes is sufficient to promote skin inflammation.

Skin Inflammation Occurs in $\Delta NC16A$ Mice in the Absence of Adaptive Immunity. Two hypotheses concerning the mechanism of skin inflammation have been proposed: One hypothesis proposes that the primary defect resides in an immunologic disturbance that causes skin inflammation; the other hypothesis proposes that an intrinsic defect in the epithelial cells (for example, keratinocytes) leads to skin inflammation (16, 17). A previous report suggested that BP180 dysfunction may trigger a spontaneous autoimmune response, which leads to BP-like skin inflammation (11). To assess whether adaptive immunity plays a major role in the spontaneous skin inflammation in our model, we crossed the $\Delta NC16A$ mice with T cell- and B cell-deficient $Rag1^{-/-}$ mice to generate adaptive immunity-deficient $\Delta NC16A$ (termed $Rag1^{-/-}\Delta NC16A$) (SI Appendix, Fig. S4 A and B). $Rag1^{-/-}\Delta NC16A$ and $\Delta NC16A$ mice were then compared for development of skin inflammation at 4, 8, and 12 wk old using $Rag1^{-/-}\Delta NC16A$ mice as a negative control (SI Appendix, Fig. S4C). $Rag1^{-/-}\Delta NC16A$ and $\Delta NC16A$ mice developed the same degree of skin inflammation including skin lesion, increased epidermal hyperplasia, increased infiltration of neutrophils, eosinophils, and mast cells into the dermis (SI Appendix, Fig. S4C) and pruritus ($n = 6$) (Fig. 4A). Similar to $\Delta NC16A$ mice, $Rag1^{-/-}\Delta NC16A$ mice had significantly increased TSLP levels in the lesional skin (Fig. 4B) and a defective skin barrier ($n = 6$) (Fig. 4 C and D). These results suggest that adaptive immunity plays a minimal role in skin inflammation in $\Delta NC16A$ mice.

Pruritus in $\Delta NC16A$ Mice Is Independent on IgE and Histamine but Dependent on TSLP. Pruritus in skin inflammation is orchestrated by the complex interplay of numerous mediators (18). A previous report showed that there is increased IgG and IgE in the circulation of BP180 dysfunctional mice, and the authors suggested that the increased IgE may be the cause of spontaneous itch and skin inflammation triggered by BP180 dysfunction (11). $\Delta NC16A$ mice exhibited significantly elevated IgE levels, but $Rag1^{-/-}\Delta NC16A$, which lack antibodies, and $\Delta NC16A$ mice showed similar degree of pruritus, strongly suggesting that IgE is not involved in the itch phenotype. To further support this conclusion, B cell-deficient $\Delta NC16A$ mice ($Ighm^{tm1Lcgt}\Delta NC16A$) were generated and examined for itch (SI Appendix, Fig. S5). $\Delta NC16A$ mice lacking B cells, hence lacking IgE, still showed the spontaneous pruritic skin inflammation and increased level of TSLP similar to $\Delta NC16A$ and $Rag1^{-/-}\Delta NC16A$ mice (Fig. 4 E and F).

Induction of pruritus can be generally divided into two categories: histaminergic and nonhistaminergic (15). We measured




Fig. 3. Skin $\Delta NC16A$ mice develop skin inflammation with increased immune cell infiltration, epidermal thickness, itch, serum IgE, TSLP, and defective barrier. (A) The ears of skin $\Delta NC16A$ mice started showing skin lesions clinically and histologically at 4 wk after tamoxifen treatment compared with control mice (Cre^{-}). Immune staining of ear skin also showed that skin $\Delta NC16A$ mice (Cre^{+}) have significantly increased infiltrating immune cells at 4 wk after tamoxifen treatment. Toluidine blue (TB) staining indicated that there was a significant increase in mast cells in skin $\Delta NC16A$ mice starting at 1 wk after tamoxifen treatment (A and E). (Original magnifications: 200 \times) Compared with Cre^{-} mice, Cre^{+} mice show increased epidermal thickness (B), spontaneous itch (C), serum IgE level (D), and TSLP (F) in skin starting at 2 wk after tamoxifen treatment ($n = 8$ per group). (G) Evans Blue dye penetration assay showed barrier defect in skin $\Delta NC16A$ but not control mice starting 1 wk after tamoxifen treatment. (H) FITC-conjugated BSA permeability assays also revealed barrier defect in skin $\Delta NC16A$ but not control mice starting 2 wk after tamoxifen treatment ($n = 6$ per group). (Scale bars: 50 μm .) * $P < 0.05$, Student's t test, graphs B, C, E, and G show mean \pm SE.




Fig. 4. Itch in $\Delta NC16A$ mice is independent on IgE and histamine but dependent on TSLP. (A) Similar to $\Delta NC16A$ mice, $Rag1^{-/-}\Delta NC16A$ mice exhibited similar degree of itch (A), increased TSLP in skin (B), and barrier defect determined by Evans Blue penetration assay (C) and FITC-conjugated BSA permeability assays (D) ($n = 6$ per group). (Scale bars: 50 μm .) (E and F) B cell-deficient $\Delta NC16A$ mice exhibited a similar degree of itch and TSLP level as $\Delta NC16A$ mice. (G) Serum histamine levels in $\Delta NC16A$, $K14Cre/\Delta NC16A$, and WT mice were similar. (H) $\Delta NC16A$ mice at 8 wk old were administered orally with H1R and H4R antagonist, respectively. The blockade of H1R and/or H4R had no effect on itch. (I) The ear of skin $\Delta NC16A$ mice were injected with TSLP neutralizing antibody or IgG2a control antibody (20 μg per ear). Meanwhile, 8-wk-old WT mice were injected with TSLP after injection with the same amount of TSLP-neutralizing antibody or IgG2a control antibody. The mice were videotaped in the absence of antibody and 1 h after antibody injection for 15 min. TSLP blockade significantly reduced itch in skin $\Delta NC16A$ mice compared with the control antibody-treated skin $\Delta NC16A$ mice. * $P < 0.05$, Student's t test, $n = 6$ per group. Graphs A, C, E, and G–I show mean \pm SE.

serum histamine level of $\Delta NC16A$, $K14Cre/\Delta NC16A$, and WT mice and found no difference among these three groups of mice (Fig. 4G). We then used antagonists of the histamine receptors H1R and H4R, two HRs expressed and identified as potential mediators of pruriception in the skin (19). To demonstrate that the histamine response can be blocked by these HR antagonists, the 8-wk-old WT mice were injected with histamine after HR antagonist was administered directly. This administration reduced the itch, while neither separate nor concomitant blockade of H1R and H4R resulted in significant reduction of scratching in $\Delta NC16A$ mice ($n = 6$) (Fig. 4H). Therefore, these results suggest that itch in $\Delta NC16A$ mice is independent on IgE and histamine.

Numerous studies have reported that TSLP acts as a master switch of skin inflammation (20–22). We found that both concentration of TSLP and the number of scratches were significantly increased in skin $\Delta NC16A$ (Fig. 3F) and $\Delta NC16A$ lacking both T and B cells ($Rag1^{-/-}$) and B cells ($Ighm^{tm1cgn}$) (Fig. 4B and F). To determine whether TSLP is required for itch in $\Delta NC16A$ mice, first, WT mice were injected with anti-TSLP neutralizing antibody or control antibody before TSLP injection to confirm the efficiency of neutralizing antibody. Then, TSLP neutralizing antibody and control antibody were injected into the ear of skin $\Delta NC16A$ mice 2 wk after tamoxifen treatment. Anti-TSLP antibody treatment reduced scratching in $\Delta NC16A$ mice compared with control antibody ($n = 6$) (Fig. 4I). Taken together, these results suggest that TSLP, but not IgE or

histamine pathway, is the mechanism underlying severe itch in $\Delta NC16A$ mice.

$\Delta NC16A$ Promotes the Release of TSLP from Keratinocytes In Vitro and in Vivo. TSLP is expressed predominantly by epithelial cells in the thymus, lung, skin, intestine, and tonsils as well as stromal cells and mast cells, but is not found in most hematopoietic cell types and endothelial cells. Our data showed that dysfunction of BP180 can induce the expression of TSLP even in the absence of T cells and B cells (Fig. 4B). Increased TSLP was seen in both skin $\Delta NC16A$ and $K14Cre/\Delta NC16A$ mice, suggesting that keratinocytes could be the cellular source of TSLP production. Immunohistochemical staining confirmed this is the case: TSLP was mainly expressed in epidermal keratinocytes in skin $\Delta NC16A$ compared with WT (Fig. 5A). To further confirm this immune localization finding, we generated primary keratinocyte culture from WT and $\Delta NC16A$ mice and compared their capacity of producing and releasing TSLP. $\Delta NC16A$ keratinocytes stimulated with mouse TNF α showed a significantly increased TSLP level in cell culture medium compared with WT keratinocytes ($n = 6$) (Fig. 5B). These data supported that BP180 dysfunction promotes the release of TSLP by keratinocytes.

BP Patients Exhibit High TSLP Expression. To sustain our conclusions and make our animal model findings more clinically relevant, we investigated whether increased TSLP expression is associated with BP, a skin disease associated with dysfunctional BP180 caused by autoantibody insult. TSLP expression at protein level was determined in serum, blister fluid, and lesional skin of BP patients. A significantly increased expression of TSLP was seen in serum of BP patients compared with healthy control ($n = 12$) (Fig. 5C). BP blister fluids also had significantly higher level of




Fig. 5. Increased TSLP is produced by keratinocytes in $\Delta NC16A$ mice and BP patients. (A) Compared with WT, ear skin from skin $\Delta NC16A$ mice showed a higher expression of TSLP mainly in epidermis. (B) WT and $\Delta NC16A$ keratinocytes were stimulated with mouse TNF α for 24 h. $\Delta NC16A$ keratinocytes released significantly higher amount of TSLP in culture medium than WT keratinocytes ($n = 6$ per group). (C–E) The serum, blister fluids, and lesional skin from BP patients exhibited an increased level of TSLP compared with normal control, blister fluids from herpes zoster, and normal control, respectively ($n = 12$ per group). (F) The increased expression of TSLP in BP patients is mainly seen in epidermis ($n = 9$ per group). (Original magnifications: A, 200 \times ; B, 400 \times .) (Scale bars: A, 100 μm ; F, 50 μm .) * $P < 0.05$. Student's t test, graph B shows mean \pm SE.

TSLP than those of herpes zoster, a skin blistering disease without itch ($n = 12$) (Fig. 5D). In addition, a high level of TSLP was found in BP lesional skin protein extracts compared with control by ELISA ($n = 12$) (Fig. 5E). Like $\Delta NC16A$ mice, TSLP was detected mainly in keratinocytes especially in the lower layers of the epidermis of BP lesional skin by immunohistochemical staining, while TSLP was below detection limit in normal control skin ($n = 9$) (Fig. 5F). These results suggest that BP180 dysfunction in BP also leads to increased expression of TSLP.

Discussion

BP180 is well documented as a key cell–cell matrix adhesion molecule in the skin; however, other biological functions remain largely unknown (15). BP180 mutations lead to partial or complete BP180 functional loss, which cause most commonly non-Herlitz JEB (23, 24). From a study based on 43 European JEB patients with BP180 mutations, the patient group exhibited a wide clinical variability from mild to severe phenotypes (24). In general, the milder forms of JEB are associated with missense or splice-site mutations and the presence of truncated BP180 protein in skin (24). In this study, we generated a mouse model with the deletion of *NC16A* domain (termed $\Delta NC16A$ mice). Because our $\Delta NC16A$ mice still express a truncated form of BP180, we consider $\Delta NC16A$ mice as a BP180 dysfunctional mouse model. $\Delta NC16A$ mice developed severe spontaneous skin inflammation characterized by immune cell infiltration, increased concentration of IgE and TSLP, pruritus, and defective skin barrier. We further showed that the skin inflammation in $\Delta NC16A$ mice is caused by *NC16A* domain deletion in basal keratinocytes since skin- and basal keratinocyte-specific $\Delta NC16A$ mice phenocopy $\Delta NC16A$ mice. These findings suggest that BP180 also plays an important role in regulation of skin inflammation.

Dysfunction of BP180 can be caused by BP180 gene mutations in JEB or anti-BP180 autoantibody in BP (4, 6). JEB is a rare autosomal recessive disorder caused by defects in any one of six genes including *BPAG2*, *LAMA3*, *LAMB3*, *LAMC2*, *ITGB4*, and *ITGA6*, which encode the hemidesmosomal proteins BP180, laminin-332, and integrin $\alpha 6\beta 4$, respectively (5, 25, 26). Patients with JEB develop subepidermal blisters within the lamina lucida of the BMZ, and itch is one of the most bothersome symptoms in JEB (10, 27). Case reports indicate that JEB patients with BP180 gene mutations show increased itch and skin inflammation (28, 29). BP is the most common skin autoimmune blistering disease characterized by inflammatory cell infiltration, anti-BP180 IgG and IgE autoantibodies, and pruritus (8). Some rare childhood BP cases shared some clinical similarity with AD (30, 31), and patients can suffer from AD and BP at the same time (30). Taken together, these clinical observations, previous animal model studies, and our BP180 dysfunctional mouse models suggest that dysfunction of BP180 can lead to skin inflammation.

Persistent pruritus is the most important symptom in skin inflammatory diseases. Some JEB patients suffer from severe itch with skin influx of proinflammatory cells (10). Transmission of pruritus can be divided into two categories: histaminergic and nonhistaminergic (15). Both histamine receptors, H1R and H4R, have been identified as potential mediators of itch in the skin (19). We rule out the histamine-dependent pathway as a cause of severe itch in $\Delta NC16A$ mice by showing that $\Delta NC16A$ mice did not have increased histamine levels (Fig. 4G), and blockade of H1R and H4R did not reduce severity of itch in $\Delta NC16A$ mice (Fig. 4H).

Recent work has highlighted TSLP in various inflammatory diseases, including skin inflammation (22). TSLP is highly expressed in acute and chronic lesions in patients with skin inflammation (32). A mutation that increases TSLP expression in the skin has direct consequences on the development of a severe atopic disease in both humans and mice (33). Interestingly, cytokines, which are commonly found at high levels in lesional skin of patients with skin inflammation, including IL-1 β , TNF α , IL-4, and IL-13, can synergize to induce TSLP expression by

keratinocytes (13), suggesting a positive feedback loop to amplify inflammation and itch. Significantly increased levels of TSLP (Figs. 2D, 3F, and 4B and F), IL-1 β , and IL-13 (SI Appendix, Fig. S1) were present in the skin of $\Delta NC16A$ mice, which correlated with severity of itch. More importantly, blockade of TSLP activity using TSLP neutralizing antibody reduced scratching in $\Delta NC16A$ mice (Fig. 4I). Thus, TSLP in the skin is required for severe itch in $\Delta NC16A$ mice.

TSLP is primarily expressed by epithelial cells in the gut, lungs, ocular tissue, thymus, epidermal keratinocytes in the skin, and also some other types of cells, such as mast cells, cancer cells, basophils, and dendritic cells (DCs) (20). In our study, we rule out T and B cells as a cellular source of TSLP since *Rag1*^{-/-} $\Delta NC16A$ mice show no change in skin TSLP levels and itch (Fig. 4). Both whole body and basal keratinocyte-specific $\Delta NC16A$ mice show significantly increased TSLP and itch. $\Delta NC16A$ primary keratinocytes produce and release significantly more TSLP compared with normal control keratinocytes when stimulated with TNF α (Fig. 5B). These in vivo and in vitro results suggest that basal keratinocytes are the cellular origin of the increased TSLP, which is caused by BP180 dysfunction and initiates itch. However, it is still unclear how BP180 dysfunction promotes the increased expression and secretion of TSLP by keratinocytes. It is possible that some cytokines (such as TNF α and IL-1 β) and/or skin infection may be a trigger to initiate TSLP expression since these proinflammatory mediators are able to induce TSLP production and secretion by keratinocytes (34). Future studies should address this important issue and identify the exact molecular interactions/pathways involved in this process.

A role for TSLP in the development of AD was hypothesized when high levels of TSLP were found in the lesional skin of AD patients (32) and in a variety of AD-like mouse models (35–37). Patients with Netherton syndrome, a severe ichthyosis in which affected individuals have a significant predisposition to AD, have increased levels of TSLP in their skin (33). However, the association of TSLP with itch in BP and JEB has not been investigated; therefore, our findings that BP patients have increased TSLP in the lesional skin and mainly express in epidermal keratinocytes represent a demonstration implicating TSLP in skin inflammation and itch in BP. Future studies should provide direct evidence whether JEB with itch are also associated with an increased TSLP level in the skin. Epithelial cells can directly communicate to cutaneous sensory neurons via TSLP to promote itch (38). However, whether TSLP effects on itch is directly through its binding to TSLPR on neurons or indirectly through other pruritic mediators in $\Delta NC16A$ mice remains to be determined.

Previously, another group also generated a BP180 dysfunction mice termed $\Delta NC14A$ mice (11). The $\Delta NC14A$ mice show phenotypes similar to $\Delta NC16A$ mice, including increased itch, eosinophil influx into skin, and increased concentration of serum IgE (11). Based on the increased IgE found in the $\Delta NC14A$ mice, the authors proposed that BP180 dysfunction may trigger autoimmunity, which leads to skin inflammation (11). Because adaptive immunity has been considered as a key factor in the development of skin inflammation (16, 17), it is possible that the components of adaptive immunity may play a role in the skin inflammation triggered by BP180 dysfunctional, especially elevated IgE. This possibility was tested by the *Rag1*^{-/-} $\Delta NC16A$ and B cell-deficient $\Delta NC16A$ mice. Although both mouse strains lack IgE, they continue to scratch and develop skin inflammation as much as $\Delta NC16A$ mice (Fig. 4 and SI Appendix, Fig. S5). Therefore, we ruled out that the increased IgE or any component of adaptive immunity is the cause of severe itch in $\Delta NC16A$ mice. These findings suggest that T and B cells are not required for spontaneous skin inflammation in $\Delta NC16A$ mice, and local deficiency of BP180 function in basal keratinocytes is necessary and sufficient to drive skin inflammatory pathology.

The onset of spontaneous skin inflammation and itch is different between the whole body $\Delta NC16A$ mice (starting at 8 wk old) and *skin* $\Delta NC16A$ mice (starting 2 wk after tamoxifen treatment). We speculate that a buildup of skin microbiota/skin

infection is necessary for the BP180 dysfunction-caused skin inflammation and itch, and in the SPF environment in an animal facility, it may take 8–10 wk to build up sufficient level of skin microbiome/infection for its effects. The previously described *ΔNC14A* mice also start the disease at 10–12 wk old (11). This speculation is further supported by clinical practice—antibiotic treatment improves BP, AD, and JEB (39–41). Another possibility is that skin barrier impairment caused by BP180 dysfunction is age-dependent, which works alone or in concert with skin microbiota/infection for the skin inflammation and itch. Future studies should address these scenarios.

In summary, we developed a BP180 dysfunction (*ΔNC16A*) mouse model to investigate the role of BP180 in vivo. Our results demonstrate that BP180, a cell–cell matrix adhesion molecule of the hemidesmosome, plays an important role in regulating skin inflammation. Dysfunction of BP180 in basal keratinocytes leads to increased TSLP expression, itch, immune cell infiltration, and defective skin barrier. More importantly, BP patients also show high expression of TSLP in keratinocytes, indicating that BP and JEB patients may have a higher risk to develop skin inflammation with itch (28, 29). Our animal model could also provide a research

angle for better understanding the mechanisms of BP180 dysfunction-related skin inflammatory diseases including BP and JEB. Future investigations to elucidate the regulatory pathways underlying the molecular link between BP180 and skin inflammation should help identify new therapeutic targets for skin inflammatory diseases associated with altered BP180 expression.

Materials and Methods

Details of mouse generation and human tissue collection, itch analysis, cytokine analysis and blockade, barrier function measurement, qPCR, immunohistology, cell culture, and statistical analysis are described in the *SI Appendix*. All of the animal experiments were approved by the local ethics committees of the Institutional Animal Care and Use Committee at the University of North Carolina at Chapel Hill. After obtaining informed consent from patients and normal healthy individuals, human samples were taken and the study was approved by the local ethics committees of the Second Affiliated Hospital, School of Medicine, Xi'an Jiaotong University.

ACKNOWLEDGMENTS. This work was supported in part by NIH Grants R01 AI40768 and R01 AR070276 (to Zhi Liu), NS079683 (to M.A.S.), DOD CA140238 (to M.A.S.), 2016LCZX-03 (to S.X.), and China Scholarship Council (to Y.Z.).

- Hammers CM, Stanley JR (2016) Mechanisms of disease: Pemphigus and bullous pemphigoid. *Annu Rev Pathol* 11:175–197.
- Koster J, Geerts D, Favre B, Borradori L, Sonnenberg A (2003) Analysis of the interactions between BP180, BP230, plectin and the integrin alpha6beta4 important for hemidesmosome assembly. *J Cell Sci* 116:387–399.
- Has C, Kern JS (2010) Collagen XVII. *Dermatol Clin* 28:61–66.
- Bauer JW, Lanschuetzer C (2003) Type XVII collagen gene mutations in junctional epidermolysis bullosa and prospects for gene therapy. *Clin Exp Dermatol* 28:53–60.
- Intong LRA, Murrell DF (2012) Inherited epidermolysis bullosa: New diagnostic criteria and classification. *Clin Dermatol* 30:70–77.
- Liu Z, Diaz LA, Giudice GJ (1994) Autoimmune response against the bullous pemphigoid 180 autoantigen. *Dermatology* 189(Suppl 1):34–37.
- Liu Z, et al. (2008) Subepidermal blistering induced by human autoantibodies to BP180 requires innate immune players in a humanized bullous pemphigoid mouse model. *J Autoimmun* 31:331–338.
- Amber KT, Murrell DF, Schmidt E, Joly P, Borradori L (2018) Autoimmune subepidermal bullous diseases of the skin and mucosae: Clinical features, diagnosis, and management. *Clin Rev Allergy Immunol* 54:26–51.
- Sibaud V, Roul S, Leaute-Labreze C, Memeguzi G, Taïeb A (2002) Atopic dermatitis: Therapeutic challenge in an infant with dystrophic epidermolysis bullosa. *Br J Dermatol* 147:350–352.
- Snauwaert JLL, et al. (2014) Burden of itch in epidermolysis bullosa. *Br J Dermatol* 171:73–78.
- Hurskainen T, et al. (2015) Deletion of the major bullous pemphigoid epitope region of collagen XVII induces blistering, autoimmunization, and itching in mice. *J Invest Dermatol* 135:1303–1310.
- Liu Z, et al. (1993) A passive transfer model of the organ-specific autoimmune disease, bullous pemphigoid, using antibodies generated against the hemidesmosomal antigen, BP180. *J Clin Invest* 92:2480–2488.
- Boguniewicz M, Leung DY (2011) Atopic dermatitis: A disease of altered skin barrier and immune dysregulation. *Immunol Rev* 242:233–246.
- Dharmage SC, et al. (2014) Atopic dermatitis and the atopic march revisited. *Allergy* 69:17–27.
- Papoiu ADP, Coghill RC, Kraft RA, Wang H, Yosipovitch G (2012) A tale of two itches. Common features and notable differences in brain activation evoked by cowhage and histamine induced itch. *Neuroimage* 59:3611–3623.
- Bieber T (2008) Atopic dermatitis. *N Engl J Med* 358:1483–1494.
- Pasparakis M, Haase I, Nestle FO (2014) Mechanisms regulating skin immunity and inflammation. *Nat Rev Immunol* 14:289–301.
- Bautista DM, Wilson SR, Hoon MA (2014) Why we scratch an itch: The molecules, cells and circuits of itch. *Nat Neurosci* 17:175–182.
- Rossbach K, et al. (2009) Histamine H4 receptor antagonism reduces hapten-induced scratching behaviour but not inflammation. *Exp Dermatol* 18:57–63.
- Ziegler SF (2012) Thymic stromal lymphopoietin and allergic disease. *J Allergy Clin Immunol* 130:845–852.
- He R, et al. (2008) TSLP acts on infiltrating effector T cells to drive allergic skin inflammation. *Proc Natl Acad Sci USA* 105:11875–11880.
- Zhang Y, Zhou B (2012) Functions of thymic stromal lymphopoietin in immunity and disease. *Immunol Res* 52:211–223.
- Fine J-D (2010) Inherited epidermolysis bullosa. *Orphanet J Rare Dis* 5:12.
- Kiriti D, et al. (2011) Molecular mechanisms of phenotypic variability in junctional epidermolysis bullosa. *J Med Genet* 48:450–457.
- Ng Y-Z, et al. (2012) Fibroblast-derived dermal matrix drives development of aggressive cutaneous squamous cell carcinoma in patients with recessive dystrophic epidermolysis bullosa. *Cancer Res* 72:3522–3534.
- Sawamura D, Nakano H, Matsuzaki Y (2010) Overview of epidermolysis bullosa. *J Dermatol* 37:214–219.
- Cohn HI, Teng JMC (2016) Advancement in management of epidermolysis bullosa. *Curr Opin Pediatr* 28:507–516.
- Mabuchi E, et al. (2007) Oral steroid improves bullous pemphigoid-like clinical manifestations in non-Herlitz junctional epidermolysis bullosa with COL17A1 mutation. *Br J Dermatol* 157:596–598.
- Cifuentes L, et al. (2013) A case of junctional epidermolysis bullosa with prurigo-like lesions and reduction of collagen XVII and filaggrin. *Br J Dermatol* 169:195–198.
- Kamiya K, Aoyama Y, Nishio E, Horio A, Tokura Y (2016) Management of erythematous skin lesions in bullous pemphigoid associated with atopic dermatitis. *J Dermatol* 43:1102–1103.
- Kuenzli S, et al. (2004) Childhood bullous pemphigoid: Report of a case with life-threatening course during homeopathy treatment. *Pediatr Dermatol* 21:160–163.
- Soumelis V, et al. (2002) Human epithelial cells trigger dendritic cell mediated allergic inflammation by producing TSLP. *Nat Immunol* 3:673–680.
- Briot A, et al. (2009) Kallikrein 5 induces atopic dermatitis-like lesions through PAR2-mediated thymic stromal lymphopoietin expression in Netherton syndrome. *J Exp Med* 206:1135–1147.
- Kumari V, Babina M, Hazzan T, Worm M (2015) Thymic stromal lymphopoietin induction by skin irritation is independent of tumour necrosis factor- α , but supported by interleukin-1. *Br J Dermatol* 172:951–960.
- Yoo J, et al. (2005) Spontaneous atopic dermatitis in mice expressing an inducible thymic stromal lymphopoietin transgene specifically in the skin. *J Exp Med* 202:541–549.
- Li M, et al. (2006) Topical vitamin D3 and low-calcemic analogs induce thymic stromal lymphopoietin in mouse keratinocytes and trigger an atopic dermatitis. *Proc Natl Acad Sci USA* 103:11736–11741.
- Li M, et al. (2005) Retinoid X receptor ablation in adult mouse keratinocytes generates an atopic dermatitis triggered by thymic stromal lymphopoietin. *Proc Natl Acad Sci USA* 102:14795–14800.
- Wilson SR, et al. (2013) The epithelial cell-derived atopic dermatitis cytokine TSLP activates neurons to induce itch. *Cell* 155:285–295.
- Chalmers JR, et al.; BLISTER study group (2015) A randomized controlled trial to compare the safety and effectiveness of doxycycline (200 mg daily) with oral prednisolone (0.5 mg kg⁻¹ daily) for initial treatment of bullous pemphigoid: A protocol for the bullous pemphigoid steroids and tetracyclines (BLISTER) trial. *Br J Dermatol* 173:227–234.
- Lakhani F, Lee K, Lio PA (2017) Case series study of the efficacy of compounded antibacterial, steroid, and moisturizer in atopic dermatitis. *Pediatr Dermatol* 34:322–325.
- Goldstein AM, Davenport T, Sheridan RL (1998) Junctional epidermolysis bullosa: Diagnosis and management of a patient with the Herlitz variant. *J Pediatr Surg* 33:756–758.