



Published in final edited form as:

J Org Chem. 2018 March 16; 83(6): 3220–3225. doi:10.1021/acs.joc.8b00104.

Prediction of ^{19}F NMR Chemical Shifts for Fluorinated Aromatic Compounds

Carla Saunders[‡], Mohammad B. Khaled[§], Jimmie D. Weaver III[§], and Dean J. Tantillo^{‡,*}


[‡]Department of Chemistry, University of California–Davis, Davis, CA 95616

[§]Department of Chemistry, Oklahoma State University, Stillwater, OK 74078

Abstract

Scaling factors are reported for use in predicting ^{19}F NMR chemical shifts for fluorinated (hetero)aromatic compounds with relatively low levels of theory. Our recommended scaling factors were developed using a curated data set of >50 compounds, with 100 individual ^{19}F shifts spanning a range of >150 ppm. With a maximum deviation of 6.5 ppm between experimental and computed shifts, or 4% of the range tested, these scaling factors allow for the assignment of chemical shifts to specific fluorines in multifluorinated aromatics. The utility of this approach is highlighted by several structural reassignments.

Graphical Abstract



Introduction

The incorporation of fluorine atoms in organic molecules (e.g., materials,¹ agrochemicals, 2–3 pharmaceuticals⁴) is becoming increasingly common, as a result of the unique electronic and steric properties of fluorine atoms.^{5–12} Consequently, ^{19}F NMR is applied ever more frequently in structural assignments. Fortunately for chemists working with fluorine, there

*Corresponding Author: djtantillo@ucdavis.edu.

ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge on the ACS Publications website.

Energies for all computed structures, table with shifts and experimental solvents for all compounds used (PDF)

Coordinates for all computed structures (mol2)

are many appeals of using ^{19}F NMR, e.g., the NMR active isotope is 100% naturally abundant, has a gyromagnetic ratio of about 0.94 times ^1H , and is spin $\frac{1}{2}$.¹³ Nonetheless, ^{19}F spectra can be difficult to interpret, especially when multiple fluorine atoms are present in a molecule. Some of us have been involved in developing methods for regioselective defluorination of polyfluorinated aromatics, methods that produce exactly the sort of compounds subject to assignment difficulties.^{5–9, 14} Useful resources are available to assist organic chemists in the interpretation of ^{19}F NMR, but it remains challenging to reliably assign all ^{19}F shifts within a multifluorinated molecule.¹³ Here we endeavor to provide a readily accessible method for reliably predicting ^{19}F shifts of fluorinated aromatic compounds.

Predicting ^1H and ^{13}C NMR chemical shifts with quantum chemical methods (especially density functional theory [DFT]) is now commonplace.^{15–18} Often, linear scaling methods are used with such approaches to remove systematic errors in computations.^{19–20} In doing so, absolute isotropic shieldings are computed for a “training set” of molecules and compared to experimental chemical shifts. When tight linear correlations between computed and experimental data are found, simple scaling factors (slope and y-intercept for the correlation line, particular to the specific theoretical method used) fall out, and can then be used for predicting chemical shifts for molecules not included in the training set. This approach has been explored previously for ^{19}F chemical shifts, including fluorinated aromatic compounds,^{21–27,28} but we provide scaling factors here that work with relatively low level theoretical methods accessible to non-experts. We also demonstrate the applicability of these scaling factors for heterocycles (common in bioactive compounds) and highlight their use in assigning structures of multifluorinated compounds.

Our training set consisted of approximately 50 compounds, with 100 individual ^{19}F shifts spanning a range of >150 ppm (Chart 1). This set contains fluorinated aromatics, many with multiple fluorine atoms in the same molecule, and mono- and bicyclic heteroaromatics. A variety of issues made compiling this data set difficult, including variations in reported shifts across the literature and the use of different reference compounds.²⁷ The compounds in Chart 1 were chosen not just for the variety of environments in which their fluorine atoms reside, but because their ^{19}F shifts were supported by multiple published reports or a single report with sufficient experimental details to make us confident in their validity.^{27, 29–42}

Methods

Geometry optimizations were run using B3LYP/6-31+G(d,p) in the gas phase using *Gaussian09*, and resulting structures were confirmed as minima using frequency calculations.^{43–47,26} These structures were subjected to NMR (GIAO)^{48–54} calculations at B3LYP/6-31+G(d,p), B3LYP/6-311+G(2d,p) and B3LYP/6-31G levels in the gas phase. B3LYP/6-31+G(d,p)/B3LYP/6-31+G(d,p) calculations were repeated using the SMD implicit solvent model for experimental solvents in NMR calculations.⁵⁵ Experimental shifts were plotted against calculated isotropic shielding values, and scaling factors were derived according to equation (1):

$$\sigma_{Experimental} = slope * \theta_{Isotropic} + intercept \quad (1)$$

These scaling factors were applied to all each test compound using equation (2):

$$\frac{\theta_{Isotropic} - intercept}{slope} = \sigma_{Scaled} \quad (2)$$

While the compounds in Chart 1 were chosen, in part, for their lack of conformational flexibility, some compounds to which the developed scaling factors were applied were conformationally flexible. For such compounds, conformational searches were run with *Spartan10*, and then optimized in *Gaussian09* with B3LYP/6-31+G(d,p) in the gas phase.^{26,56-57} Conformers within 3 kcal/mol of the lowest energy conformer were included in the NMR calculations, with their contributions weighted with a Boltzmann distribution based on their relative free energies.

Results and Discussion

Using B3LYP/6-31+G(d,p)//B3LYP/6-31+G(d,p), the maximum error on any one ¹⁹F shift was 6.5 ppm, with a mean absolute deviation (MAD) of 2.1 ppm over a range of 153 ppm (Figure 1, middle). Performance was similar when NMR calculations were repeated with a larger basis set (B3LYP//6-311+G(2d,p)//B3LYP/6-31+G(d,p)): maximum error of 6.6 ppm and MAD of 1.7 ppm (Figure 1, right). Use of a smaller basis set (B3LYP/6-31G//B3LYP/6-31+G(d,p)) resulted in a much higher maximum deviation (28 ppm) and a much higher MAD of 4.0 ppm (Figure 1, left). On the basis of these results, we recommend the B3LYP/6-31+G(d,p)//B3LYP/6-31+G(d,p) for rapid predictions of ¹⁹F chemical shifts.

Although the ¹⁹F spectra for compounds used in the training set were taken in a variety of solvents, the majority were taken in either CDCl₃ or CCl₄ (a full list of the solvents is available in the Supporting Information). To test the effect of solvent on our predictions, the gas phase optimized structures were subjected to NMR calculations in continuum solvent (SMD) and new scaling factors were determined (Figure 2). For the 45 ¹⁹F shifts from our test set that were measured in chloroform, gas phase NMR calculations had a maximum error of 8.2 ppm, while continuum chloroform calculations had a maximum error of 7.2 ppm. The MAD also decreased slightly, from 2.9 to 2.6 ppm. For the 27 shifts determined in carbon tetrachloride, the maximum error decreased from 5.1 to 4.6 ppm, while the MAD remained the same at 1.5 ppm. Although including solvent in the calculations did provide small improvements in prediction accuracy, gas phase calculations appear to be sufficient.

To put our scaling factors to the test, we applied them to several multifluorinated compounds synthesized in one of our labs (Chart 2). The original assignment of one structure was found to be in error on the basis of our predicted ¹⁹F shifts. The shifts for the corrected structure (Chart 2) are completely consistent with the experimental spectrum.

After discovering this misassignment, a group of 13 substituted benzene and pyridine compounds with various substitution patterns (Chart 3) were examined to confirm, or correct, the published structures.^{7, 9, 12} Although many of the spectra match experiment, two of the structures had large errors (Chart 3, red). Based on the correction described above, revised structures were tested and these match the experimental values within the error expected for our scaling factors. These straightforward applications of computational ¹⁹F shift prediction highlight its utility in avoiding mistakes in assignments for highly fluorinated compounds.

Conclusion

We have developed scaling factors aimed at fast and accurate prediction of ¹⁹F shifts for fluoroaromatics. In particular, the methods described here allow for the assignment of challenging structures bearing multiple fluorine atoms. The application of these scaling factors does not require advanced computations and, as a result, we hope that it will be applied broadly.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We gratefully acknowledge support from the National Science Foundation (CHE-1565933, CHE-0957416 and CHE-030089 [via XSEDE]).

References

1. Berger R, Resnati G, Metrangolo P, Weber E, Hulliger J. Organic fluorine compounds: a great opportunity for enhanced materials properties. *Chemical Society Reviews*. 2011; 40(7):3496–3508. [PubMed: 21448484]
2. Fujiwara T, O'Hagan D. Successful fluorine-containing herbicide agrochemicals. *Journal of Fluorine Chemistry*. 2014; 167(Supplement C):16–29.
3. Theodoridis, G. Chapter 4 Fluorine-Containing Agrochemicals: An Overview of Recent Developments. In: Tressaud, A., editor. *Advances in Fluorine Science*. Vol. 2. Elsevier; 2006. p. 121-175.
4. Purser S, Moore PR, Swallow S, Gouverneur V. Fluorine in medicinal chemistry. *Chemical Society Reviews*. 2008; 37(2):320–330. [PubMed: 18197348]
5. Senaweera SM, Singh A, Weaver JD. Photocatalytic Hydrodefluorination: Facile Access to Partially Fluorinated Aromatics. *Journal of the American Chemical Society*. 2014; 136(8):3002–3005. [PubMed: 24547940]
6. Senaweera SM, Weaver JD. Selective Perfluoro- and Polyfluoroarylation of Meldrum's Acid. *The Journal of Organic Chemistry*. 2014; 79(21):10466–10476. [PubMed: 25271542]
7. Singh A, Kubik JJ, Weaver JD. Photocatalytic C-F alkylation; facile access to multifluorinated arenes. *Chemical Science*. 2015; 6(12):7206–7212. [PubMed: 29861956]
8. Arora A, Weaver JD. Visible Light Photocatalysis for the Generation and Use of Reactive Azolyl and Polyfluoroaryl Intermediates. *Accounts of Chemical Research*. 2016; 49(10):2273–2283. [PubMed: 27682342]
9. Senaweera S, Weaver JD. Dual C–F, C–H Functionalization via Photocatalysis: Access to Multifluorinated Biaryls. *Journal of the American Chemical Society*. 2016; 138(8):2520–2523. [PubMed: 26890498]

10. Cybulski MK, Nicholls JE, Lowe JP, Mahon MF, Whittlesey MK. Catalytic Hydrodefluorination of Fluoroarenes Using Ru(IME4)2L2H2 (IME4 = 1,3,4,5-Tetramethylimidazol-2-ylidene; L2 = (PPh3)2, dppe, dppp, dppm) Complexes. *Organometallics*. 2017; 36(12):2308–2316.
11. Liu Y, Zhou Y, Zhao Y, Qu J. Synthesis of gem-Difluoroallylboronates via FeCl2-Catalyzed Boration/ β -Fluorine Elimination of Trifluoromethyl Alkenes. *Organic Letters*. 2017; 19(4):946–949. [PubMed: 28139930]
12. Singh A, Fennell CJ, Weaver JD. Photocatalyst size controls electron and energy transfer: selectable E/Z isomer synthesis via C-F alkenylation. *Chemical Science*. 2016; 7(11):6796–6802. [PubMed: 28042465]
13. Dolbier, WR. *Guide to Fluorine NMR for Organic Chemists*. Wiley; 2009.
14. Muir M, Baker J. A simple calculational model for predicting the site for nucleophilic substitution in aromatic perfluorocarbons. *Journal of Fluorine Chemistry*. 2005; 126(5):727–738.
15. Lodewyk MW, Siebert MR, Tantillo DJ. Computational Prediction of ¹H and ¹³C Chemical Shifts: A Useful Tool for Natural Product, Mechanistic, and Synthetic Organic Chemistry. *Chemical Reviews*. 2012; 112(3):1839–1862. [PubMed: 22091891]
16. Grimblat N, Sarotti AM. Computational Chemistry to the Rescue: Modern Toolboxes for the Assignment of Complex Molecules by GIAO NMR Calculations. *Chemistry – A European Journal*. 2016; 22(35):12246–12261.
17. Di Micco S, Chini MG, Riccio R, Bifulco G. Quantum Mechanical Calculation of NMR Parameters in the Stereostructural Determination of Natural Products. *European Journal of Organic Chemistry*. 2010; 2010(8):1411–1434.
18. Willoughby PH, Jansma MJ, Hoye TR. A guide to small-molecule structure assignment through computation of (¹H and ¹³C) NMR chemical shifts. *Nature Protocols*. 2014; 9:643. [PubMed: 24556787]
19. Jain R, Bally T, Rablen PR. Calculating Accurate Proton Chemical Shifts of Organic Molecules with Density Functional Methods and Modest Basis Sets. *The Journal of Organic Chemistry*. 2009; 74(11):4017–4023. [PubMed: 19435298]
20. Bally T, Rablen PR. Quantum-Chemical Simulation of ¹H NMR Spectra. 2. Comparison of DFT-Based Procedures for Computing Proton–Proton Coupling Constants in Organic Molecules. *The Journal of Organic Chemistry*. 2011; 76(12):4818–4830. [PubMed: 21574622]
21. Fadeev DS, Chuikov IP, Mamatyuk VI. Non-empirical and DFT calculations of ¹⁹F and ¹³C chemical shifts in the NMR spectra of substituted pentafluorobenzenes. *Journal of Structural Chemistry*. 2013; 54(1):180–185.
22. Ghiviriga I, Zhang L, Martinez H, Contreras RH, Tormena CF, Nodin L, Dolbier WR. ¹⁹F chemical shifts, coupling constants and conformational preferences in monosubstituted perfluoroparacyclophanes. *Magnetic Resonance in Chemistry*. 2011; 49(3):93–105. [PubMed: 21246624]
23. Kasireddy C, Bann JG, Mitchell-Koch KR. Demystifying fluorine chemical shifts: electronic structure calculations address origins of seemingly anomalous ¹⁹F-NMR spectra of fluorohistidine isomers and analogues. *Physical Chemistry Chemical Physics*. 2015; 17(45):30606–30612. [PubMed: 26524669]
24. Sanders LK, Oldfield E. Theoretical Investigation of ¹⁹F NMR Chemical Shielding Tensors in Fluorobenzenes. *The Journal of Physical Chemistry A*. 2001; 105(34):8098–8104.
25. Shaghghi H, Ebrahimi H, Tafazzoli M, Jalali-Heravi M. A survey of wave function effects on theoretical calculation of gas phase ¹⁹F NMR chemical shifts using factorial design. *Journal of Fluorine Chemistry*. 2010; 131(1):47–52.
26. Sternberg U, Klipfel M, Grage SL, Witter R, Ulrich AS. Calculation of fluorine chemical shift tensors for the interpretation of oriented ¹⁹F-NMR spectra of gramicidin A in membranes. *Physical Chemistry Chemical Physics*. 2009; 11(32):7048–7060. [PubMed: 19652840]
27. Hogben MG, Graham WAG. Chemical shifts and coupling constants in pentafluorophenyl derivatives. I. Correlations of chemical shifts, coupling constants, and π -electronic interactions. *Journal of the American Chemical Society*. 1969; 91(2):283–291.
28. Saielli G, Bini R, Bagno A. Computational ¹⁹F NMR. 2. Organic compounds. *RSC Advances*. 2014; 4(78):41605–41611.

29. He K-H, Tan F-F, Zhou C-Z, Zhou G-J, Yang X-L, Li Y. Acceptorless Dehydrogenation of N-Heterocycles by Merging Visible-Light Photoredox Catalysis and Cobalt Catalysis. *Angewandte Chemie International Edition*. 2017; 56(11):3080–3084. [PubMed: 28156039]
30. Yuan X, Yao J-F, Tang Z-Y. Decarboxylative Fluorination of Electron-Rich Heteroaromatic Carboxylic Acids with Selectfluor. *Organic Letters*. 2017; 19(6):1410–1413. [PubMed: 28256137]
31. Boden N, Emsley JW, Feeney J, Sutcliffe LH. The effects of π -electron distribution and intramolecular electric fields on the ^{19}F N.M.R. shielding in substituted perfluorobenzenes. *Molecular Physics*. 1964; 8(2):133–149.
32. Plevy RG, Rendell RW, Tatlow JC. Fluorinations with complex metal fluorides. Part 6 [1] fluorination of yridine and related compounds with caesium tetrafluorocobaltate(III) [2]. *Journal of Fluorine Chemistry*. 1982; 21(2):159–169.
33. Coe PL, Holton AG, Tatlow JC. Fluorinations with potassium tetrafluorocobaltate(III) Part VII. Further investigations on the fluorination of pyridine. *Journal of Fluorine Chemistry*. 1982; 21(2): 171–189.
34. Gribble GW, Keavy DJ, Olson ER, Rae ID, Staffa A, Herr TE, Ferraro MB, Contreras RH. Fluorine deshielding in the proximity of a methyl group. An experimental and theoretical study. *Magnetic Resonance in Chemistry*. 1991; 29(5):422–432.
35. Burdon J, Chivers GE, Tatlow JC. Highly fluorinated heterocycles. Part III. The preparation and reactions of some polyfluorofurans. *Journal of the Chemical Society C: Organic*. 1970; (16):2146–2151.
36. Burdon J, Campbell JG, Parsons IW, Tatlow JC. Highly fluorinated heterocycles. Part V. The preparation and some reactions of tetrafluorothiophen and some polyfluorothiophens. *Journal of the Chemical Society C: Organic*. 1971; (0):352–355.
37. Baasner B, Klauke E. A new route to the synthesis of 5-fluorouracil. *Journal of Fluorine Chemistry*. 1989; 45(3):417–430.
38. Chambers DR, Hall WC, Hutchinson J, Millar WR. Polyhalogenated heterocyclic compounds. Part 42.1 Fluorinated nitrogen heterocycles with unusual substitution patterns. *Journal of the Chemical Society, Perkin Transactions 1*. 1998; (10):1705–1714.
39. Schlosser M, Bobbio C, Rausis T. Regiochemically Flexible Substitutions of Di-, Tri-, and Tetrahalopyridines: The Trialkylsilyl Trick. *The Journal of Organic Chemistry*. 2005; 70(7):2494–2502. [PubMed: 15787535]
40. Bobbio C, Rausis T, Schlosser M. Removal of Fluorine from and Introduction of Fluorine into Polyhalopyridines: An Exercise in Nucleophilic Heteroatomic Substitution. *Chemistry – A European Journal*. 2005; 11(6):1903–1910.
41. Božilović J, Bats JW, Engels JW. Synthesis and structure of fluorindole nucleosides. *Canadian Journal of Chemistry*. 2007; 85(4):283–292.
42. Lemal DM, Akashi M, Lou Y, Kumar V. Tetrafluorothiophene S,S-Dioxide: A Perfluorinated Building Block. *The Journal of Organic Chemistry*. 2013; 78(24):12330–12337. [PubMed: 24313897]
43. Becke AD. A new mixing of Hartree–Fock and local density-functional theories. *The Journal of Chemical Physics*. 1993; 98(2):1372–1377.
44. Becke AD. Density-functional thermochemistry. III. The role of exact exchange. *The Journal of Chemical Physics*. 1993; 98(7):5648–5652.
45. Lee C, Yang W, Parr RG. Development of the Colle-Salvetti correlation-energy formula into a functional of the electron density. *Physical Review B*. 1988; 37(2):785–789.
46. Stephens PJ, Devlin FJ, Chabalowski CF, Frisch MJ. Ab Initio Calculation of Vibrational Absorption and Circular Dichroism Spectra Using Density Functional Force Fields. *The Journal of Physical Chemistry*. 1994; 98(45):11623–11627.
47. Tirado-Rives J, Jorgensen WL. Performance of B3LYP Density Functional Methods for a Large Set of Organic Molecules. *J Chem Theory Comput*. 2008; 4(2):297–306. [PubMed: 26620661]
48. London F. Théorie quantique des courants interatomiques dans les combinaisons aromatiques. *J Phys Radium*. 1937; 8(10):397–409.
49. McWeeny R. Perturbation Theory for the Fock-Dirac Density Matrix. *Physical Review*. 1962; 126(3):1028–1034.

50. Ditchfield R. Self-consistent perturbation theory of diamagnetism. *Molecular Physics*. 1974; 27(4): 789–807.
51. Wolinski K, Hinton JF, Pulay P. Efficient implementation of the gauge-independent atomic orbital method for NMR chemical shift calculations. *Journal of the American Chemical Society*. 1990; 112(23):8251–8260.
52. Cheeseman JR, Trucks GW, Keith TA, Frisch MJ. A comparison of models for calculating nuclear magnetic resonance shielding tensors. *The Journal of Chemical Physics*. 1996; 104(14):5497–5509.
53. Bagno A, Saielli G. Addressing the stereochemistry of complex organic molecules by density functional theory-NMR. *Wiley Interdisciplinary Reviews: Computational Molecular Science*. 2015; 5(2):228–240.
54. de Albuquerque AC, Ribeiro DJ, de Amorim MB. Structural determination of complex natural products by quantum mechanical calculations of (13)C NMR chemical shifts: development of a parameterized protocol for terpenes. *J Mol Model*. 2016; 22(8):183. [PubMed: 27424297]
55. Marenich AV, Cramer CJ, Truhlar DG. Universal Solvation Model Based on Solute Electron Density and on a Continuum Model of the Solvent Defined by the Bulk Dielectric Constant and Atomic Surface Tensions. *The Journal of Physical Chemistry B*. 2009; 113(18):6378–6396. [PubMed: 19366259]
56. Spartan'10. Wavefunction, Inc; Irvine, CA, USA: 2010.
57. Gaussian 09, Revision D. 01. Gaussian, Inc; Wallingford, CT, USA: 2013.




Figure 1. Correlation data for two NMR computational methods (all using B3LYP/6-31+G(d,p) geometries): Left: B3LYP/6-31G Middle: B3LYP/6-31+G(d,p); Right: B3LYP/6-311+G(2d,p). Mono-cyclic aromatic compounds (group **A**, Chart 1) are shown in green, non-heterocyclic compounds (group **B**) are shown in grey, and bicyclic-heterocyclic compounds (group **C**) are shown in yellow.







Figure 2. Experimental and computed shifts for the training set used for optimization and NMR calculation at B3LYP/6-31+G(d,p) – structures taken in chloroform (top) and calculated in gas (left) and chloroform (right), structures taken in carbon tetrachloride (bottom). Monocyclic aromatic compounds (group A) are shown in green, non-heterocyclic compounds (group B) are shown in grey, and bicyclic-heterocyclic compounds (group C) are shown in yellow.

**Chart 1.**

Experimental shifts of compounds used as training set to develop scaling factors: **A**: substituted mono-cyclic heterocycles, **B**: substituted non-heterocyclic aromatics, **C**: substituted bicyclic heterocycles.

**Chart 2.**

Initial compounds explored, including a structure that was corrected by computations (highlighted in red). Computed shifts and error from experiment are shown for each aromatic fluorine.

**Chart 3.**

Fluorinated aromatic compounds for which assignments were not certain. Numbers in parentheses are numbering from the original paper. The calculated shifts and error from experimental shift are shown for each aromatic fluorine.