

# Calcineurin Regulatory Subunit Calcium-Binding Domains Differentially Contribute to Calcineurin Signaling in *Saccharomyces cerevisiae*

Sean Connolly,\* Devona Quasi-Woode,\* Laura Waldron,<sup>†</sup> Christian Eberly,\* Kerri Waters,\* Eric M. Muller,<sup>†</sup> and Tami J. Kingsbury\*<sup>1</sup>

\*Department of Physiology, Center for Regenerative Medicine and Stem Cell Biology and the Marlene and Stewart Greenbaum Cancer Center, University of Maryland School of Medicine, Baltimore, Maryland 21201 and <sup>†</sup>Department of Biology, Iona College, New Rochelle, New York 10801

ORCID ID: 0000-0003-3095-3788 (T.J.K.)

**ABSTRACT** The protein phosphatase calcineurin is central to Ca<sup>2+</sup> signaling pathways from yeast to humans. Full activation of calcineurin requires Ca<sup>2+</sup> binding to the regulatory subunit CNB, comprised of four Ca<sup>2+</sup>-binding EF hand domains, and recruitment of Ca<sup>2+</sup>-calmodulin. Here we report the consequences of disrupting Ca<sup>2+</sup> binding to individual Cnb1 EF hand domains on calcineurin function in *Saccharomyces cerevisiae*. Calcineurin activity was monitored via quantitation of the calcineurin-dependent reporter gene, *CDRE-lacZ*, and calcineurin-dependent growth under conditions of environmental stress. Mutation of EF2 dramatically reduced *CDRE-lacZ* expression and failed to support calcineurin-dependent growth. In contrast, Ca<sup>2+</sup> binding to EF4 was largely dispensable for calcineurin function. Mutation of EF1 and EF3 exerted intermediate phenotypes. Reduced activity of EF1, EF2, or EF3 mutant calcineurin was also observed in yeast lacking functional calmodulin and could not be rescued by expression of a truncated catalytic subunit lacking the C-terminal autoinhibitory domain either alone or in conjunction with the calmodulin binding and autoinhibitory segment domains. Ca<sup>2+</sup> binding to EF1, EF2, and EF3 in response to intracellular Ca<sup>2+</sup> signals therefore has functions in phosphatase activation beyond calmodulin recruitment and displacement of known autoinhibitory domains. Disruption of Ca<sup>2+</sup> binding to EF1, EF2, or EF3 reduced Ca<sup>2+</sup> responsiveness of calcineurin, but increased the sensitivity of calcineurin to immunophilin-immunosuppressant inhibition. Mutation of EF2 also increased the susceptibility of calcineurin to hydrogen peroxide inactivation. Our observations indicate that distinct Cnb1 EF hand domains differentially affect calcineurin function *in vivo*, and that EF4 is not essential despite conservation across taxa.

**KEYWORDS** calcineurin; calcium signaling; phosphatase; EF hand domain; Crz1

**C**ALCINEURIN (formerly protein phosphatase 2B, now PPP3C) is a ubiquitously expressed Ca<sup>2+</sup>-regulated serine/threonine protein phosphatase first isolated from the bovine brain, where it accounts for 1% of the total protein content (Klee *et al.* 1988). Calcineurin activation by intracellular Ca<sup>2+</sup> signals orchestrates cell responses to developmental cues, environmental stimuli, and intracellular stress to affect

cell proliferation, differentiation, and death (Shibasaki and McKeon 1995; Wang *et al.* 1999; Kahl and Means 2003; Zayzafoon 2006). Studies in diverse model organisms have shown that calcineurin regulates fundamental cell processes to control development, behavior, life span, and adaptive responses (Saneyoshi *et al.* 2002; Nishiyama *et al.* 2007; Dwivedi *et al.* 2009; Mair *et al.* 2011; Nakai *et al.* 2011; Lee *et al.* 2013; Kujawski *et al.* 2014; Deng *et al.* 2015). Calcineurin is also essential for the life cycle, host cell invasion, and virulence of human pathogens, including *Plasmodium falciparum*, *Toxoplasma gondii*, *Cryptococcus neoformans*, *Aspergillus fumigatus*, the causative agents of malaria, toxoplasmosis, and cryptococcal meningitis, and invasive aspergillosis, respectively (Odum *et al.* 1997; Cruz *et al.* 2001; Juvvadi *et al.* 2013; Paul *et al.* 2015; Philip and Waters 2015; Chow *et al.* 2017).

Copyright © 2018 by the Genetics Society of America

doi: <https://doi.org/10.1534/genetics.118.300911>

Manuscript received March 13, 2018; accepted for publication May 2, 2018; published Early Online May 7, 2018.

Supplemental material available at Figshare: <https://doi.org/10.25386/genetics.6229040>.

<sup>1</sup>Corresponding author: Department of Physiology, Center for Regenerative Medicine and Stem Cell Biology and the Marlene and Stewart Greenbaum Cancer Center, University of Maryland School of Medicine, Rm. S103C HSF II, 20 Penn St., Baltimore, MD 21201. E-mail: [tkingsbury@som.umaryland.edu](mailto:tkingsbury@som.umaryland.edu)

In mammals, calcineurin activation of transcription factor EB (TFEB) in response to lysosomal  $\text{Ca}^{2+}$  signals regulates autophagy initiated by oxidative stress or nutrient deprivation (Medina *et al.* 2015; Tong and Song 2015; Zhang *et al.* 2016; Tseng *et al.* 2017). Calcineurin activation of nuclear factor of activated T cells transcription (NFAT) factors promotes development and function of the immune, cardiovascular, nervous, and musculoskeletal systems (Crabtree and Olson 2002; Horsley and Pavlath 2002; Schulz and Yutzey 2004; Wu *et al.* 2007; Nguyen and Di Giovanni 2008; Musson *et al.* 2012; Patel *et al.* 2015; Peiris and Keating 2018). Alterations in calcineurin activity contribute to human diseases such as cardiac hypertrophy (Bueno *et al.* 2002; Wilkins and Molkenkin 2004), cancer (Buchholz and Ellenrieder 2007; Peuker *et al.* 2016; Wang *et al.* 2017), neurodegeneration (Mukherjee *et al.* 2010; Mukherjee and Soto 2011; Qu *et al.* 2012; Luo *et al.* 2014; Aufschnaiter *et al.* 2017; Shah *et al.* 2017), and mental illness (Manji *et al.* 2003; Mathieu *et al.* 2008; Forero *et al.* 2016). Calcineurin inhibitors FK506 (tacrolimus) and cyclosporin A are widely used clinically to prevent allograft rejection in transplant patients and treat inflammatory skin diseases (Musson *et al.* 2012; Azzi *et al.* 2013; Nygaard *et al.* 2017).

Intracellular  $\text{Ca}^{2+}$  transients exhibit distinct spatial and temporal patterns, the amplitude, frequency, and duration of which are integrated by calcineurin to shape cellular responses (Dolmetsch *et al.* 1997; Berridge *et al.* 2003). Multiple mechanisms coordinately modulate calcineurin signaling in cells (Musson and Smit 2011). The requirement for two distinct EF hand  $\text{Ca}^{2+}$ -binding proteins, CNB and calmodulin, to activate calcineurin enables the activation and inactivation of calcineurin within the narrow window of intracellular  $\text{Ca}^{2+}$  concentrations. Interactions with additional cellular factors such as the immunophilins FKBP12 and cyclophilin A (Cardenas *et al.* 1994), Bcl-2 (Shibasaki *et al.* 1997; Erin *et al.* 2003), superoxide dismutase 1 (Wang *et al.* 1996; Agbas *et al.* 2007), heat shock proteins (Someren *et al.* 1999; Imai and Yahara 2000), RCANs (Kingsbury and Cunningham 2000; Rothermel *et al.* 2003; Hilioti *et al.* 2004), CIB1 (Heineke *et al.* 2010), KIF1B $\beta$  (Li *et al.* 2016a), and AKAP79 (Li *et al.* 2011) impose additional layer of regulation via modulating phosphatase activity, conformation, localization, and ability to interact with either calmodulin or substrates. Post-translational modifications of calcineurin, including CNA phosphorylation, which can regulate calcineurin activity *in vitro*, presumably tune calcineurin function *in vivo* (Hashimoto and Soderling 1989; Martensen *et al.* 1989; Calalb *et al.* 1990; Juvvadi *et al.* 2013). CNB myristylation, conserved from yeast to humans, reduces  $\text{Ca}^{2+}$  responsiveness of calcineurin in yeast (Connolly and Kingsbury 2012). Reversible substrate docking and substrate competition further shape calcineurin signaling in cells (Roy *et al.* 2007; Roy and Cyert 2009; Li *et al.* 2011).

Calcineurin is highly conserved across taxa. Calcineurin is a heterodimer consisting of a ~60 kDa catalytic subunit (CNA) and a smaller ~19 kDa regulatory subunit (CNB), comprised of four EF hand  $\text{Ca}^{2+}$ -binding domains (Klee *et al.* 1998)

(hereafter EF1, EF2, EF3, or EF4). *In vitro* analysis of recombinant mammalian calcineurin revealed that EF1 and EF2 have a lower affinity for  $\text{Ca}^{2+}$  than EF3 and EF4, suggesting  $\text{Ca}^{2+}$  binding to EF1 and EF2 mediates responsiveness to changes in intracellular  $\text{Ca}^{2+}$  (Feng and Stemmer 2001). EF3 and EF4 are predicted to constitutively bind  $\text{Ca}^{2+}$  within cells, consistent with a structural role for these sites. Maximal stimulation of calcineurin phosphatase activity requires  $\text{Ca}^{2+}$ , CNB, and  $\text{Ca}^{2+}$ /calmodulin (Klee *et al.* 1998).  $\text{Ca}^{2+}$  binding to EF1 and EF2 induces conformational changes in CNB that trigger conformational changes in CNA, leading to partial phosphatase activity. Binding of  $\text{Ca}^{2+}$ /calmodulin to CNA results in full enzyme activation via displacing the autoinhibitory segment (AIS) from the LxVP docking pocket and the C-terminal autoinhibitory domain (AID) from the catalytic site (Wang *et al.* 2008; Li *et al.* 2016b).

To begin to understand the requirements for calcineurin activation *in vivo*, we took advantage of the budding yeast *Saccharomyces cerevisiae*, in which CNB is encoded by a single gene (*CNB1*). Two CNA encoding genes are present: *CNA1* and *CNA2* (Cyert *et al.* 1991). Calcineurin is not essential in *S. cerevisiae*, except under conditions of environmental or cellular stress (Iida *et al.* 1990, 1994; Cyert *et al.* 1991; Cyert and Thorner 1992; Moser *et al.* 1996; Fischer *et al.* 1997; Paidhungat and Garrett 1997; Withee *et al.* 1997; Cyert 2003). Here, we investigated the consequences of introducing mutations into the  $\text{Ca}^{2+}$ -binding domains of *Cnb1* to elucidate the requirement for specific EF hand domains in promoting calcineurin function in yeast. Our results demonstrate that EF4 is largely dispensable for calcineurin activity in response to intracellular  $\text{Ca}^{2+}$  signals, as previously reported for mammalian calcineurin *in vitro* (Feng and Stemmer 1999, 2001). In contrast, EF1, EF2, and EF3 each contribute to calcineurin activation, even in the absence of the bipartite CNA autoinhibitory elements and calmodulin binding. Our findings reveal an additional function of  $\text{Ca}^{2+}$  binding to CNB in the activation of calcineurin *in vivo*, beyond calmodulin recruitment: AIS and AID displacement.

## Materials and Methods

### Yeast culture

All yeast strains used in this study were derived from W303-1A (Table 1). Yeast strains were grown at 30° in standard rich media (YPD) or synthetic media (SD) (Clontech or Sigma, St. Louis, MO). Agar, amino acids and salts were purchased from Difco and Sigma. YPD medium was buffered to pH 5.5 by addition of 0.5 M succinic acid for experiments requiring the addition of extracellular  $\text{CaCl}_2$ . Yeast culture and transformations were conducted using standard techniques (Sherman 1991; Gietz *et al.* 1995).

### Cloning and mutagenesis

QuikChange site-directed mutagenesis (Stratagene, La Jolla, CA) was conducted according to manufacturer's instructions in pYDZ3 (Zhu *et al.* 1995). For EF1, EF2, and EF4

**Table 1 Yeast strains used in this study**

Yeast strain	Relevant genotype	Source
K603	<i>MATa cnb1::LEU2</i>	Cunningham and Fink (1994)
TKY102	<i>MATa cnb1::LEU2 cna1Δ cna2::HIS3</i>	This study
K687	<i>MATa cnb1::LEU2 cmd1-3</i>	Cunningham laboratory
K650	<i>MATa cnb1 vcx1</i>	Cunningham laboratory
DMY22	<i>MATa cnb1 crz1 pmc1</i>	Cunningham laboratory

mutagenesis, the GAT encoding aspartic acid was mutated to GCT, to encode alanine. For EF3, GAC was mutated to GCC to convert the aspartic acid to alanine. *CNB1* alleles were cloned into pRS313 (Sikorski and Hieter 1989) (centromere (CEN)) or pYO323 (Qadota *et al.* 1992) (2  $\mu$ m), using *Bam*HI-*Xho*I. Genomic DNA corresponding to *CNA1* was PCR amplified (forward: AGGATCCGCTCTTCGTCAACAATTGGTTC; reverse: ACTCGAGCATTGGTTACAAGTCCCTAAC) and cloned into pRS314 *Bam*HI-*Xho*I. Truncated alleles were generated by PCR with *CNA1* forward primer combined with either reverse GGACTGAAGGTTTGAATGAAACGCAACTGTGATAAATCTCA (*Cna1ΔAID*) or GCGATATTAGAAGATGAAACCCAACCTGTGATAAATCTCA (*CNA1ΔCBD*). All constructs were verified by DNA sequencing (Supplemental Material, Table S1).

#### Ion tolerance assays

Ion tolerance assays were performed as previously described (Cunningham and Fink 1996), with minor modifications. *cnb1Δ* yeast strains transformed with CEN-based *CNB1* alleles were precultured in SD-HIS medium. Ion tolerance assays were conducted in YPD supplemented with a range of NaCl, LiCl, or MnCl<sub>2</sub> concentrations. YPD (pH 5.5) medium was used for CaCl<sub>2</sub> assays to maintain salt solubility. Yeast were grown as 0.18 ml cultures in 96-well, flat-bottomed dishes in a 30° incubator without shaking for 36–48 hr. The OD<sub>600</sub> was measured using a BioTek  $\mu$ Quant platereader.

#### $\beta$ -Galactosidase assays

Yeast cultures were grown overnight in SD dropout media (–HIS –URA for *cnb1Δ* yeast or –HIS –URA –LEU –TRP for *cna1Δ cna2Δ cnb1Δ* yeast). Yeast strains were pelleted and resuspended in YPD (pH 5.5) before inoculation into media supplemented with CaCl<sub>2</sub>, FK506, cyclosporine A (CsA),  $\alpha$ -factor (Genscript), or H<sub>2</sub>O<sub>2</sub> as indicated. For  $\alpha$ -factor experiments yeast were diluted to 0.2 OD. Cells were incubated at 30° with shaking for 4 hr (3 hr for experiments with  $\alpha$ -factor). After incubation, 0.9 ml of yeast was harvested by centrifugation and resuspended into 720  $\mu$ l of Z buffer (60 mM Na<sub>2</sub>HPO<sub>4</sub>, 40 mM NaH<sub>2</sub>PO<sub>4</sub>, 10 mM KCl, 1 mM MgSO<sub>4</sub>, 50 mM  $\beta$ -mercaptoethanol, pH 7.0) + 0.0075% SDS. Cells were lysed by addition of 50  $\mu$ l of chloroform and vortexing for 10 sec. Reactions were initiated by addition of 180  $\mu$ l of ONPG (4 mg/ml in Z buffer) and terminated by the addition of 450  $\mu$ l 1 M Na<sub>2</sub>CO<sub>3</sub>.  $\beta$ -Galactosidase activity was quantitated by measuring the OD<sub>600</sub> of the yeast culture used for the assay and the OD<sub>420</sub> of the reaction product.

$\beta$ -Galactosidase activity was calculated as follows: (1000  $\times$  OD<sub>420</sub>)/OD<sub>600</sub>  $\times$  time (min)  $\times$  volume (ml) = units of  $\beta$ -galactosidase activity.

#### Protein analysis

Yeast were grown overnight in SD-HIS media, diluted into YPD, and grown an additional 4 hr before harvesting for protein. Cells were pelleted and washed once in PBS. Cell pellets were lysed with lithium acetate and NaOH as previously described (Zhang *et al.* 2011), except that 1 M LiOAc was used. Pellets were resuspended in 2 $\times$  Laemmli buffer, boiled for 5 min, and cleared by centrifugation before separation by PAGE using Novex Wedgewell 16% Tris-Glycine gels run in 1 $\times$  SDS Tris-Glycine buffer (Invitrogen, Carlsbad, CA). When indicated, protein lysates were for incubated for 10 min with either 1 mM CaCl<sub>2</sub> or 2 mM EGTA before gel loading. Western blotting was conducted using anti-*Cnb1* (Zhu *et al.* 1995) at 1:5000 in TBS-Tween as primary antibody and HRP-conjugated anti-rabbit at 1:10,000 as secondary antibody (Jackson ImmunoResearch Laboratories). Anti-PGK at 1:2000 (Invitrogen) was used as a loading control.


#### Data availability

All yeast strains used in this study are available upon request. Table S1 contains list of plasmids utilized in this study. All plasmids generated in this study will be deposited at Addgene for distribution. Data necessary to confirm findings of this article are present within article, figures, and tables. Supplemental material available at Figshare: <https://doi.org/10.25386/genetics.6229040>.

## Results

### Generation of *CNB1* mutant alleles

The Ca<sup>2+</sup>-binding regulatory subunit calcineurin B is encoded by a single gene *CNB1* in *S. cerevisiae* (Cyert and Thorner 1992). To investigate the contribution of specific *Cnb1* EF hand domains to calcineurin function (Figure 1A), we generated a series of *CNB1* alleles expressed from low-copy CEN-based vectors in which an individual EF hand domain harbored a single amino acid mutation designed to disrupt Ca<sup>2+</sup> binding. EF hand domains are comprised of two  $\alpha$ -helices separated by a short loop that binds Ca<sup>2+</sup>. Within the Ca<sup>2+</sup>-binding loop, amino acids 1, 3 5, 7, 9, and 12 (X, Y, Z, –Y, –X, and –Z, respectively) coordinate Ca<sup>2+</sup> (Nakayama and Kretsinger 1994). Using site-directed mutagenesis, the aspartic acid located at position X was mutated to alanine (Figure 1B) in EF1, EF2, EF3, or EF4. EGTA depletion of Ca<sup>2+</sup> from EF3 and EF4, but not EF1 or EF2, reduces the electrophoretic mobility of mammalian CNB in high-percentage SDS-PAGE (Feng and Stemmer 1999). We observed that treatment of yeast cell lysates treated with EGTA similarly reduced *Cnb1* electrophoretic mobility in high-percentage SDS-PAGE (Figure 1C). In the presence of Ca<sup>2+</sup>, multiple species of *Cnb1* were observed in high-percentage gels. Treatment with EGTA resulted in the loss of the faster



**Figure 1** Site-directed mutagenesis of yeast calcineurin B EF hand domains. (A) Schematic of Cnb1. (B) Sequence of the four EF hand domains of Cnb1. The Ca<sup>2+</sup>-binding loop is denoted in bold lettering. Amino acids that bind Ca<sup>2+</sup> are enlarged. The aspartate residue (D) mutated to alanine in each EF hand is indicated by an arrow. (C) Electrophoretic mobility of Cnb1 mutant proteins. Protein lysates from *cnb1Δ* yeast expressing the wild type (WT), Cnb1 EF hand mutants vs. empty vector (EV) were treated with 1 mM CaCl<sub>2</sub> or 2 mM EGTA before SDS-PAGE and Western blot analysis using anti-Cnb1 antibody (Zhu *et al.* 1995).

migrating species. In the presence of Ca<sup>2+</sup>, Cnb1 harboring mutations in either EF3 (Cnb1mutEF3) or EF4 (Cnb1mutEF4) lacked the faster migrating species, consistent with the D/A mutations disrupting Ca<sup>2+</sup> binding. In contrast, the mobility of Cnb1mutEF1 and Cnb1mutEF2 was comparable to the fast migrating species observed for wild-type Cnb1, suggesting that mutation of EF1 or EF2 neither altered Cnb1 mobility nor disrupted Ca<sup>2+</sup> binding to EF3 or EF4.


**Disruption of EF1, EF2, and EF3 reduces calcineurin function:** Calcineurin mediates yeast responses to Ca<sup>2+</sup> signals triggered by extracellular stimuli and intracellular stress. Intracellular Ca<sup>2+</sup> signals stimulate calcineurin, which dephosphorylates the transcription factor Crz1. Crz1 then translocates to the nucleus to mediate calcineurin-dependent changes in gene expression (Cyert 2003; Thewes 2014). Calcineurin activity can be readily quantitated by measuring the expression of the calcineurin-dependent reporter gene *CDRE-lacZ* (calcineurin-dependent response element), composed of four copies of the Crz1-binding site (Stathopoulos and Cyert 1997). In the absence of calcineurin, *CDRE-lacZ* expression is not stimulated by Ca<sup>2+</sup> signaling. Wild-type vs. mutant *CNB1* alleles were transformed into *cnb1Δ* yeast (Y203) harboring the *CDRE-lacZ* reporter gene. Yeast transformed with empty vector were used as negative controls. To stimulate Ca<sup>2+</sup> signaling, yeast were treated with 100 mM extracellular CaCl<sub>2</sub> for 4 hr before harvesting for quantitative ortho-nitrophenyl-beta-galactoside (ONPG) assays to measure β-galactosidase activity. Yeast expressing wild-type Cnb1 exhibited a robust activation of *CDRE-lacZ* activity in response to 100 mM CaCl<sub>2</sub>

(Figure 2A). Yeast expressing Cnb1mutEF4 were most similar, with ~80% of the stimulated *CDRE-lacZ* activity observed in yeast expressing wild-type Cnb1. In contrast, yeast expressing Cnb1mutEF2 had *CDRE-lacZ* activity levels similar to yeast transformed with empty vector. Yeast expressing Cnb1mutEF1 and Cnb1mutEF3 exhibited intermediate levels of *CDRE-lacZ* activity. Overexpression of Cnb1 mutants from high-copy 2 μm expression constructs (Figure S1) was unable to rescue the reduced *CDRE-lacZ* activity levels observed upon mutation of EF1, EF2, or EF3 (Figure 2A). In addition, increased expression of wild-type Cnb1 did not significantly increase *CDRE-lacZ* levels following stimulation with 100 mM CaCl<sub>2</sub>, indicating Cnb1 is not limiting for Crz1 activation. A similar pattern of *CDRE-lacZ* activity was observed in yeast lacking the calcineurin-dependent vacuolar Ca<sup>2+</sup> transporters *Vcx1* and *Pmc1*, indicating that reduced ability of calcineurin to stimulate Crz1 was not dependent on altered function of these transporters (Figure 1, B and C), although transporter deletion enhanced our ability to detect Cnb1mutEF2-dependent *CDRE-lacZ* activity above background levels, presumably due to increased cytosolic Ca<sup>2+</sup> levels (Cunningham and Fink 1994; Cui *et al.* 2009).

**Mutation of EF1, EF2, or EF3 reduces Ca<sup>2+</sup> responsiveness of calcineurin:** Our observation that the absence of *Vcx1* enhanced our ability to detect Cnb1mutEF2 activity above background suggested that mutant Cnb1 had reduced Ca<sup>2+</sup> responsiveness. We therefore assayed *CDRE-lacZ* activity across a range of extracellular CaCl<sub>2</sub> from 0 to 100 mM to determine the relative Ca<sup>2+</sup> sensitivity of the EF hand mutants. Assays were conducted in the *cnb1Δ vcx1Δ* (Y250) background, where we could consistently detect Ca<sup>2+</sup> stimulated Cnb1mutEF2-dependent *CDRE-lacZ* activity, albeit at very low levels (~5% wild type). Differences in the ability of Ca<sup>2+</sup> to activate the mutant calcineurin would be observed as a shift in the Ca<sup>2+</sup> dose-response curve. For each mutant, the *CDRE-lacZ* activity measured at 100 mM CaCl<sub>2</sub> was set to 100% activity, and *CDRE-lacZ* activity at the remaining CaCl<sub>2</sub> concentrations was plotted as a percentage of the activity at 100 mM CaCl<sub>2</sub>. As shown in Figure 2D, calcineurin comprised of Cnb1mutEF1, Cnb1mutEF2, or Cnb1mutEF3 required higher levels of Ca<sup>2+</sup> for stimulation of *CDRE-lacZ* expression than wild-type Cnb1 or Cnb1mutEF4. Thus, the Cnb1mutEF1, Cnb1mutEF2, or Cnb1mutEF3 mutants have reduced sensitivity to elevated intracellular Ca<sup>2+</sup>.

#### **EF1, EF2, and EF3 mutants disrupt calcineurin responses to physiologic stimuli**

Calcineurin mediates yeast responses to environmental stimuli, including extracellular stress and mating pheromone. Calcineurin activation of Crz1 is required for yeast to grow in the presence of increasing levels of extracellular Mn<sup>2+</sup>, Li<sup>+</sup>, and Na<sup>+</sup>. We therefore tested the ability of *cnb1Δ* yeast (Y203) transformed with wild-type vs. mutant *CNB1* alleles to grow in the presence of high levels of MnCl<sub>2</sub>, LiCl, or NaCl (Figure 3, A–C). *cnb1Δ* yeast transformed with wild-type



**Figure 2** Cnb1 EF hand mutants reduce calcineurin-stimulated reporter gene expression. (A–C) The *CDRE-lacZ* reporter gene was introduced into *cnb1Δ* (A), *cnb1Δ vcx1Δ* (B), or *cnb1Δ pmc1Δ vcx1Δ* (C) yeast expressing either wild-type or mutant Cnb1, expressed from low-copy centromere-based plasmids (CEN) or high-copy 2 μm plasmids (2 μm). β-Galactosidase activity was quantitated after 4 hr stimulation in YPD (pH 5.5) supplemented with 100 mM CaCl<sub>2</sub>. Data are normalized to the activity measured in yeast transformed with empty vector in the absence of stimulation. (D) *CDRE-lacZ* activity was measured following 4 hr stimulation at 0, 10, 25, 50, and 100 mM added extracellular CaCl<sub>2</sub> in *cnb1Δ vcx1Δ* yeast. For each Cnb1 mutant construct, the *CDRE-lacZ* activity at each CaCl<sub>2</sub> concentration is plotted as a percentage of the activity obtained at 100 mM CaCl<sub>2</sub> (100%). For all assays, data plotted are the average of four independent yeast transformants ± SD.

*CNB1* or empty vector were used as positive and negative controls, respectively. Yeast expressing Cnb1mutEF4 exhibited a similar level of ion resistant growth as yeast expressing wild-type Cnb1 in the presence of Mn<sup>2+</sup>, Li<sup>+</sup>, and Na<sup>+</sup>. In contrast, yeast expressing Cnb1mutEF1, Cnb1mutEF2, or Cnb1mutEF3 exhibited reduced ion tolerance, becoming growth inhibited at lower concentrations of extracellular MnCl<sub>2</sub>, LiCl, or NaCl than yeast expressing wild-type Cnb1. In each case, Cnb1mutEF2 expressing yeast were the most sensitive to growth inhibition. In the presence of NaCl, Cnb1mutEF2 expressing yeast were indistinguishable from *cnb1Δ* yeast transformed with empty vector. These results demonstrate that EF1, EF2, and EF3 are required for calcineurin activity in response to environmental stress due to high levels of Mn<sup>2+</sup>, Li<sup>+</sup>, and Na<sup>+</sup>.

We next tested whether Cnb1 mutants exhibited reduced calcineurin activity in response to mating pheromone. Haploid yeast cells of opposite mating type sense each other through plasma membrane receptors, grow toward each other, and ultimately fuse at the tips of extended cytoplasmic projections. Exposure of a mating type haploid yeast to α-factor mating pheromone triggers Ca<sup>2+</sup> signals and calcineurin-mediated stimulation of Crz1 (Matheos *et al.* 1997; Stathopoulos and Cyert 1997). Wild-type vs. mutant *CNB1* alleles were transformed into a *cnb1Δ* (Y203) yeast harboring the *CDRE-lacZ* reporter gene. Yeast transformed with empty vector were used as negative controls. Ca<sup>2+</sup> signals were induced by treating yeast with α-factor mating pheromone for 3 hr before harvesting for quantitative ONPG assays to measure β-galactosidase activity. Yeast expressing either wild-type Cnb1 or Cnb1mutEF4 stimulated *CDRE-lacZ* expression ~19-fold in response to α-factor (Figure 3D). In contrast, yeast expressing Cnb1mutEF2 were similar to yeast transformed with empty vector. Yeast expressing Cnb1mutEF1 and Cnb1mutEF3 had an intermediate response, exhibiting ~fivefold stimulation of *CDRE-lacZ* activity following α-factor treatment. As observed following stimulation with high


CaCl<sub>2</sub>, overexpression of wild-type Cnb1 or Cnb1mutEF4 from high-copy 2 μm vectors did not increase *CDRE-lacZ* activity, and overexpression of Cnb1EF1, Cnb1mutEF2, or Cnb1mutEF3 did not rescue *CDRE-lacZ* activity.

#### EF1, EF2, and EF3 function beyond autoinhibitory domain displacement

The carboxyl terminal domain of CNA contains an AID that binds to the active site of calcineurin. Ca<sup>2+</sup> binding to CNB increases the interaction of calcineurin with Ca<sup>2+</sup>/calmodulin, which leads to displacement of the AID during phosphatase activation (Yang and Klee 2000) (Figure 4A). Deletion of the Cna1 AID (Cna1-ΔAID) generates a constitutively active calcineurin that can be robustly stimulated by Ca<sup>2+</sup> signaling (Figure 4C). We tested whether deletion of the AID could restore calcineurin activity in Cnb1 mutants. *cnb1Δ cna1Δ cna2Δ* (TKY102) yeast harboring the *CDRE-lacZ* reporter gene and expressing Cna1-ΔAID were transformed with wild-type vs. mutant *CNB1* alleles. Calcineurin stimulation of *CDRE-lacZ* activity was tested following 4 hr in the presence of 100 mM extracellular CaCl<sub>2</sub> before harvesting for ONPG assays. In the absence of stimulation, only yeast expressing wild-type Cnb1 or Cnb1mutEF4 exhibited the expected increase in basal *CDRE-lacZ* activity (Figure 4C). As observed in the presence of wild-type endogenous Cna1 and Cna2, Cnb1mutEF1, Cnb1mutEF2, and Cnb1mutEF3 exhibited reduced *CDRE-lacZ* activity compared to Cnb1mutEF4 and wild-type Cnb1. Thus, the reduced calcineurin activity observed upon mutation of either EF1, EF2, or EF3 is not solely due to a defect in displacement of the AID in response to Cnb1 Ca<sup>2+</sup> binding.

#### Mutation of EF1, EF2, or EF3 impairs calmodulin-independent calcineurin function

Full activation of calcineurin requires Ca<sup>2+</sup> binding to Cnb1 and recruitment of Ca<sup>2+</sup>/calmodulin. In the absence of the



**Figure 3** Mutation of EF1, EF2, and EF3 reduces calcineurin responses to physiologic stimuli. (A–C) Wild-type Cnb1, Cnb1mutEF1, Cnb1mutEF2, Cnb1mutEF3, or Cnb1mutEF4 were expressed in *cnb1Δ* yeast to assess the ability of the mutants to promote ion-resistant growth. Yeast strains were grown in YPD supplemented with increasing concentrations of MnCl<sub>2</sub> (A), LiCl (B), or NaCl (C). Following incubation for 2 days at 30°, the optical density of cultures was measured at 600 nm and plotted as a function of ion concentration. Data plotted are the average of four independent yeast transformants ± SE. (D) *CDRE-lacZ* activity was measured following 3 hr stimulation with 20 μM α-factor in a *cnb1Δ* yeast. Cnb1 was expressed from either low-copy centromere-based plasmids (CEN) or high-copy 2 μm plasmids (2 μm). For all assays, data plotted are the average of four independent yeast transformants ± SD.

Ca<sup>2+</sup>/calmodulin interaction, Crz1 activation in response to Ca<sup>2+</sup> signals is dramatically reduced. To determine whether Cnb1 EF hand mutants alter calmodulin-independent calcineurin activity in cells, *cnb1Δ cmd1-3* (K687) yeast harboring the *CDRE-lacZ* reporter were transformed with wild-type vs. mutant *CNB1* alleles. *cmd1-3* yeast express calmodulin that cannot bind to Ca<sup>2+</sup> and fails to stimulate calcineurin activity (Geiser *et al.* 1991). Stimulation of *cmd1-3* yeast expressing wild-type Cnb1 with 100 mM extracellular CaCl<sub>2</sub> for 4 hr resulted in less than fourfold stimulation of *CDRE-lacZ* expression (Figure 4D). Cnb1EFmut4 resulted in twofold stimulation in *CDRE-lacZ* activity following Ca<sup>2+</sup> stimulation in *cmd1-3* background, whereas Cnb1mutEF1, Cnb1mutEF2, or Cnb1mutEF3 were unable to stimulate *CDRE-lacZ* expression in the absence of Ca<sup>2+</sup>/calmodulin.

Calmodulin-independent calcineurin function was further tested in *cnb1Δ cna1Δ cna2Δ* (TKY102) yeast expressing Cna1 protein truncated before the calmodulin-binding domain (Cna1-ΔCBD). Truncation of the catalytic subunit upstream of the calmodulin-binding domain also eliminates the AIS and the AID of Cna1, resulting in a constitutively active enzyme that can be further stimulated by Ca<sup>2+</sup> binding to Cnb1. Stimulation of yeast expressing Cna1-ΔCBD with either wild-type Cnb1 or Cnb1mutEF4 by addition of 100 mM CaCl<sub>2</sub> to the media resulted in similar levels of *CDRE-lacZ* activity (Figure 4E). Yeast expressing Cnb1mutEF1 or Cnb1mutEF3 had lower levels of both basal and stimulated *CDRE-lacZ* activity, consistent with impaired calmodulin-independent calcineurin function. Yeast expressing Cnb1mutEF2 resembled the empty vector negative control in the presence and absence of stimulation.


#### Cnb1 mutants increase sensitivity to immunosuppressant inhibition

Calcineurin is the target of the immunosuppressants FK506 and CsA. Interaction of calcineurin with immunophilin drug

complexes has previously been shown to require Ca<sup>2+</sup> and CNB (Li and Handschumacher 1993; Milan *et al.* 1994). We therefore tested the ability of FK506 and CsA to inhibit calcineurin activity in yeast expressing the Cnb1 EF mutants. Yeast were stimulated with 100 mM CaCl<sub>2</sub> in the presence or absence of increasing doses of FK506 or CsA for 4 hr before harvesting to assay *CDRE-lacZ* activity. As shown in Figure 5, yeast expressing Cnb1mutEF1, Cnb1mutEF2, or Cnb1mutEF3 were more sensitive to inhibition by FK506 (Figure 5A) and CsA (Figure 5B) than yeast expressing wild-type Cnb1. Cnb1mutEF4 expressing yeast resembled wild type in the presence of FK506, but exhibited a small increase in sensitivity to inhibition by CsA. Estimation of the concentration of drug at which *CDRE-lacZ* stimulation was reduced by 50%, suggested that Cnb1mutEF1, Cnb1mutEF2 and Cnb1mutEF3 expressing yeast were 2.7-, 6.4-, and 4.0-fold more sensitive to FK506 inhibition than wild-type Cnb1 expressing yeast, respectively. In the presence of CsA, EF1, EF2, EF3, and EF4 mutant calcineurin was 3.0, 13, 4.5, and 1.8 times as sensitive as wild-type calcineurin, respectively.

#### Cnb1 mutants exhibit increased sensitivity to inactivation by oxidative stress

Calcineurin activity is sensitive to oxidative inactivation (Wang *et al.* 1996; Namgaladze *et al.* 2002; Musson and Smit 2011). Since the reduced activity we observe in the EF mutants could reflect increased inactivation, we investigated the sensitivity of calcineurin to hydrogen peroxide in yeast expressing wild-type vs. mutant Cnb1. *Cnb1Δ vcx1Δ* yeast harboring the indicated *CNB1* alleles and the *CDRE-lacZ* reporter gene were stimulated with 100 mM CaCl<sub>2</sub> for 4 hr in either the absence or presence of 2 mM H<sub>2</sub>O<sub>2</sub> before harvesting for ONPG assays. H<sub>2</sub>O<sub>2</sub> had no effect on the levels of *CDRE-lacZ* β-galactosidase activity in the absence of added exogenous CaCl<sub>2</sub> (data not shown). In the presence of



**Figure 4** Removal of Cna1 autoinhibitory domains or calcineurin interaction does not bypass the requirement for Cnb1 EF1, EF2, or EF3. (A) Cartoon model of calcineurin activation by Ca<sup>2+</sup> and calcineurin. (B) Schematic of wild type (WT) and Cna1 truncations. (C–E) *cnb1Δ cna1Δ cna2Δ* yeast harboring the *CDRE-lacZ* reporter expressing either Cna1-ΔAID (C), *cnb1Δ cmd1-3* (D), or Cna1-ΔCBD (E) were transformed with WT vs. mutant Cnb1 constructs. Each strain was grown to log phase and exposed for 4 hr to 100 mM CaCl<sub>2</sub> in YPD medium (pH 5.5) before β-galactosidase activity measurement. Error bars represent the SD of four independent transformants. BBH, CNB binding helix.

100 mM CaCl<sub>2</sub>, calcineurin composed of wild-type Cnb1 or Cnb1mutEF4 was inhibited only 10–15% by inclusion of H<sub>2</sub>O<sub>2</sub> in the media (Figure 6). In contrast, yeast expressing Cnb1mutEF1, Cnb1mutEF2, or Cnb1mutEF3 were inhibited ~45, 72, and 35%, respectively, by the presence of H<sub>2</sub>O<sub>2</sub>. Thus, mutation of EF1, EF2, or EF3 increased the sensitivity of calcineurin to inhibition by H<sub>2</sub>O<sub>2</sub>, with mutation of EF2 being the most dramatically affected. Although our data reveal that EF1, EF2, and EF3 mutant calcineurin are more sensitive to oxidative stress, we were unable to reverse reduced activity of mutants by inclusion of reduced glutathione in the media (S Connolly and TJ Kingsbury unpublished data), suggesting that increased oxidation of calcineurin in the absence of induced oxidative stress does not account for the reduced activity observed in these mutants.


#### **Cnb1 EF1, EF2, and EF3 mutants impair *Crz1*-independent calcineurin function**

To determine whether the reduced function of Cnb1mutEF1, Cnb1mutEF2, or Cnb1mutEF3 calcineurin extends to additional phosphatase substrates, we tested the ability of the Cnb1 mutants to inhibit the Ca<sup>2+</sup>/H<sup>+</sup> exchanger, *Vcx1*. In wild-type yeast, the vacuolar ATPase *Pmc1* maintains Ca<sup>2+</sup> homeostasis and enables growth in the presence of high extracellular CaCl<sub>2</sub>. In the absence of *Pmc1*, disrupting calcineurin function increases Ca<sup>2+</sup> tolerance due to loss of calcineurin-mediated inhibition of *Vcx1* (Cunningham and Fink 1996). Calcineurin inhibition of *Vcx1* is independent

of *Crz1*. To test the ability of Cnb1 EF hand mutants to inhibit *Vcx1* activity, *cnb1 pmc1Δ crz1Δ* (Figure 7) were transformed with wild-type vs. mutant Cnb1 expression constructs and growth quantitated after 2 days culture in the presence of increasing concentrations of CaCl<sub>2</sub>. Yeast expressing either Cnb1mutEF1 or Cnb1mutEF2 exhibited increased Ca<sup>2+</sup>-resistant growth similar to that observed in yeast lacking Cnb1, suggesting an impaired ability to inhibit *Vcx1*. Cnb1mutEF3 displayed an intermediate level of Ca<sup>2+</sup>-resistant growth, while Cnb1mutEF4 resembled yeast expressing wild-type Cnb1. These results show that the reduced activity of EF1, EF2, or EF3 mutant Cnb1 calcineurin is not limited to a single physiologic substrate.

#### **Discussion**

CNB interaction with CNA is essential for calcineurin activation *in vitro* and function *in vivo* (Cyert and Thorner 1992; Nakamura *et al.* 1993; Kawamura and Su 1995; Tokoyoda *et al.* 2000). Yeast in which the *CNB1* has been disrupted exhibit calcineurin-deficient phenotypes such as failure to recover from mating pheromone and ion sensitivity (Cyert 2003). To begin to understand the requirements for Ca<sup>2+</sup> binding to CNB in promoting calcineurin function *in vivo*, we investigated the functional consequences of individually mutating each of the Ca<sup>2+</sup>-binding domains of Cnb1. Here we show that mutation of Cnb1 Ca<sup>2+</sup>-binding domains EF1, EF2, and EF3 reduces calcineurin function in yeast. Using a



**Figure 5** EF hand mutants increase calcineurin immunophilin-immunosuppressant sensitivity. The ability of immunosuppressants to inhibit calcineurin-dependent stimulation of *CDRE-lacZ* activity was assayed across a range of (A) FK506 (nM) or (B) CsA ( $\mu\text{M}$ ) concentrations as indicated. *cnb1Δ vcx1Δ* yeast expressing wild-type (WT) or mutant *cnb1* were stimulated for 4 hr in YPD supplemented with 100 mM  $\text{CaCl}_2$  in the presence or absence of FK506 or CsA.  $\text{Ca}^{2+}$ -stimulated *CDRE-lacZ* activity measured in the absence of immunosuppressant drugs for each strain containing a *cnb1* construct was set to 100%, and the  $\beta$ -galactosidase activity observed in the presence of drugs was plotted as a corresponding percentage. Data plotted represent the average of four independent yeast transformants  $\pm$  SD.

calcineurin-dependent reporter gene, we show that mutation of EF1, EF2, or EF3 decreased total calcineurin activity, with mutation of EF2 exhibiting the most dramatic reduction in response to  $\text{Ca}^{2+}$  signaling stimulated by either high extracellular  $\text{Ca}^{2+}$  levels or  $\alpha$ -factor stimulation. Consistent with reduced calcineurin stimulation of *Crz1* activity, *Cnb1* EF1, EF2, and EF3 mutants exhibited reduced calcineurin-mediated ion-tolerant growth, which is primarily but not exclusively mediated by *Crz1*. Thus, multiple distinct sources of  $\text{Ca}^{2+}$  signaling each revealed reduced calcineurin function upon mutation of EF1, EF2, or EF3. Mutation of EF1, EF2, or EF3 also impaired the ability of calcineurin to inhibit the vacuolar  $\text{Ca}^{2+}/\text{H}^+$  exchanger *Vcx1*, a *Crz1*-independent calcineurin function, demonstrating that the observed defect in calcineurin was not specific to *Crz1* substrate. In contrast, EF4 was found to be largely dispensable for calcineurin function


in wild-type yeast. Our findings are consistent with previous *in vitro* analysis of recombinant mammalian calcineurin (Feng and Stemmer 1999, 2001) despite the dynamic complexity of intracellular  $\text{Ca}^{2+}$  signals, which are in part shaped by calcineurin-dependent processes, and additional calcineurin regulatory interactions presumably occurring within the context of a cell. To our knowledge, our data are the first demonstration of the functional role of EF1, EF2, and EF3 in conferring calcineurin responsiveness to changes in intracellular  $\text{Ca}^{2+}$  levels.

Consistent with impaired calcineurin function observed with *Cnb1* EF1, EF2, or EF3 mutants, we found reduced sensitivity to activation across a range of  $\text{Ca}^{2+}$  signals, indicating that these EF hand domains determine calcineurin responsiveness to increases in intracellular  $\text{Ca}^{2+}$ . The reduced  $\text{Ca}^{2+}$  sensitivity of *Cnb1* EF1 and EF2 mutant calcineurin is consistent with their proposed role in binding  $\text{Ca}^{2+}$  in response to intracellular  $\text{Ca}^{2+}$  transients based on their lower affinity for  $\text{Ca}^{2+}$  *in vitro* (Feng and Stemmer 1999, 2001; Yang and Klee 2000; Gallagher *et al.* 2001). In contrast, EF3 and EF4 are higher affinity  $\text{Ca}^{2+}$ -binding sites predicted to be constitutively bound to  $\text{Ca}^{2+}$  in cells. The reduced  $\text{Ca}^{2+}$  sensitivity of *Cnb1* EF3 mutant calcineurin reveals an essential role for EF3 in mediating calcineurin responses to changes in intracellular  $\text{Ca}^{2+}$ . It is unclear how reduced  $\text{Ca}^{2+}$  affinity of EF3 causes reduced  $\text{Ca}^{2+}$  sensitivity of calcineurin, although it is possible that mutation of EF3 lowers the affinity of EF1 or EF2 for  $\text{Ca}^{2+}$  due to communication between the sites (Gallagher *et al.* 2001). Mutation of EF4 did not alter phosphatase  $\text{Ca}^{2+}$  sensitivity.

Full activation of calcineurin during  $\text{Ca}^{2+}$  signaling requires both  $\text{Ca}^{2+}$  binding to *Cnb1* and the recruitment of  $\text{Ca}^{2+}$ /calmodulin (Klee *et al.* 1998). In the inactive state, the CBD interacts with the CNB binding helix and the AID acts as a pseudosubstrate blocking the catalytic site. Recent crystal structure of the full-length  $\beta$ -isoform of mammalian CNA has revealed an additional inhibitory domain, called the autoinhibitory segment (AIS), located at the carboxy terminus of the CBD, which interacts with a hydrophobic domain previously shown to mediate substrate and immunophilin interaction (Li *et al.* 2016b). Existence of an additional AID was previously implicated from CNA mutant analysis (Perrino *et al.* 1995; Wang *et al.* 2008). *In vitro* functional analysis further suggested that AID and AIS cooperate to inhibit calcineurin and suggested an updated model for calcineurin activation (Li *et al.* 2016b). In this model,  $\text{Ca}^{2+}$  binding to CNB EF1 and EF2 induces a conformational change in CNB, which leads to movement of a large domain of CNA and subsequent stimulation of basal activity. Recruitment of  $\text{Ca}^{2+}$ /calmodulin causes dissociation of AIS from the substrate docking domain and displacement of the AID from the catalytic site, likely via a reorientation rather than displacement, allowing full enzyme activation.

The reduced *in vitro* activity of recombinant mammalian calcineurin comprised of mutant EF1, EF2, or EF3 CNB also reduced calcineurin activity in the absence of calmodulin. Using the *cmd1-3* mutant yeast background, which expresses






**Figure 6**  $\text{Ca}^{2+}$ -binding domains modulate the hydrogen peroxide sensitivity of calcineurin. The *CDRE-lacZ* plasmid was transformed into *cnb1Δ vcx1Δ* yeast expressing either wild-type (WT) or mutant *cnb1* as indicated.  $\beta$ -Galactosidase assays were conducted after 4 hr stimulation of yeast strains in YPD (pH 5.5) supplemented with 100 mM  $\text{CaCl}_2$  in the presence or absence of 2 mM  $\text{H}_2\text{O}_2$ . Data are plotted as the percentage  $\text{Ca}^{2+}$ -stimulated activity observed in the presence of  $\text{H}_2\text{O}_2$  relative to activity observed in cultures lacking  $\text{H}_2\text{O}_2$ . Data plotted are the average of four independent yeast transformants  $\pm$  SD.

a form of calmodulin that cannot bind  $\text{Ca}^{2+}$  and therefore does not stimulate calcineurin (Moser *et al.* 1996), we found that EF1, EF2, or EF3 *Cnb1* mutant calcineurin failed to stimulate calmodulin-independent calcineurin activity. Since binding of  $\text{Ca}^{2+}$ /calmodulin is necessary to displace AID and AIS (Li *et al.* 2016b), we further tested whether removal of the *Cna1* C-terminal regulatory CBD, AIS or AID could bypass the requirement for EF1, EF2, or EF3. Removal of the *Cna1* AID alone or in combination with the CBD and AIS was not able to restore wild-type levels of calcineurin activity. Further, *Cna1* truncation to additionally remove the CBD and AIS (*Cna1*- $\Delta$ CBD) revealed that mutation of EF1, EF2, or EF3 impairs transmission of an activation signal from *Cnb1* to *Cna1* that is independent of conformational changes involving the CBD, AIS, or AIDs of *Cna1*. Although our findings suggest an additional role, they do not rule out the possibility that reduced calmodulin interaction or ineffective displacement of the bipartite *Cna1* AIDs contribute to the reduced activity of *Cnb1* EF1, EF2, or EF3 mutant calcineurin in the context of full-length *Cna1* and functional calmodulin. Indeed, mutation of EF2 has previously been reported to reduce the interaction of recombinant mammalian calcineurin with  $\text{Ca}^{2+}$ /calmodulin *in vitro* (Feng and Stemmer 2001).

Interaction of calcineurin with immunophilin-immunosuppressant complexes requires  $\text{Ca}^{2+}$  and CNB (Cardenas *et al.* 1994; Milan *et al.* 1994; Rodríguez *et al.* 2009). *Cnb* EF1, EF2, and EF3 mutant calcineurin had increased sensitivity to inhibition by immunophilin-immunosuppressant complexes despite their reduced activity and  $\text{Ca}^{2+}$  responsiveness. These findings suggest that mutation of specific EF hands alters the conformation of calcineurin leading to enhanced



**Figure 7** *Cnb1* mutants impede calcineurin-mediated inhibition of the vacuolar  $\text{H}^+/\text{Ca}^{2+}$  exchanger *Vcx1*. *cnb1Δ pmc1Δ crz1Δ* yeast expressing wild-type (WT) vs. mutant *Cnb1* were grown for 2 days in YPD (pH 5.5) in the presence of the increasing concentrations of extracellular  $\text{CaCl}_2$  at 30°. Optical density ( $\text{OD}_{600}$ ) was measured for each culture and plotted as four individual transformants  $\pm$  SD.

interaction with immunophilin-immunosuppressant complexes. Since these complexes share an overlapping binding site with substrates (Rodríguez *et al.* 2009), our observations suggest that substrate interactions may be altered in EF1, EF2, and EF3 *Cnb1* mutants, contributing to either reduced binding or ineffective dephosphorylation due to inappropriate positioning of phosphorylated residues with respect to the catalytic site. Altered conformation and increased accessibility of the active site is also suggested by the increased sensitivity of EF1, EF2, and EF3 mutant calcineurin to oxidative inactivation. *In vitro* analysis of mammalian CNB has demonstrated that  $\text{Ca}^{2+}$  binding to CNB reduces the Michaelis-Menten constant ( $K_m$ ) for substrate binding (Perrino *et al.* 1995).

In summary, our findings reveal striking conservation in the roles of *Cnb1* EF hand domains in regulating calcineurin function between yeast *in vivo* and mammalian calcineurin *in vitro*, and extend those findings to reveal that EF1, EF2, and EF3 stimulate phosphatase activity independent of known conformational changes induced upon  $\text{Ca}^{2+}$  binding that lead to calmodulin binding and relief of bipartite auto-inhibition mediated by C-terminus of CNA. Further analysis of these mutants may shed light on how CNB stimulates calcineurin activity following  $\text{Ca}^{2+}$  binding within the context of interacting cellular factors that may function to shape calcineurin activity downstream of  $\text{Ca}^{2+}$  signaling, or development of novel strategies to inhibit calcineurin activity in cells.

## Acknowledgments

We thank Kyle Cunningham at Johns Hopkins University for yeast strains, Martha Cyert at Stanford University for the *CDRE-lacZ* expression plasmid, Joseph Heitman at Duke University for pYDZ3 and anti-*Cnb1* antibody, and Stephen

Liggett at University of South Florida for use of equipment. S.C. and T.J.K. were supported by University of Maryland Department of Physiology funding and National Institutes of Health grants NS058464 and Eunice Kennedy Shriver National Institute of Child Health and Human Development/Office of Research on Women's Health/National Institute of Diabetes and Digestive and Kidney Diseases (K12 HD 43489). Project funding was also provided by University of Maryland School of Medicine Marlene and Stewart Greenebaum Cancer Center Developing Shared Service Pilot grant (National Cancer Institute P30CA134274-02S2). L.W. and E.M.M. were supported by an American Express endowed fund to the Iona College Honors Program.

## Literature Cited

- Agbas, A., D. Hui, X. Wang, V. Tek, A. Zaidi *et al.*, 2007 Activation of brain calcineurin (Cn) by Cu-Zn superoxide dismutase (SOD1) depends on direct SOD1-Cn protein interactions occurring in vitro and in vivo. *Biochem. J.* 405: 51–59. <https://doi.org/10.1042/BJ20061202>
- Aufschnaiter, A., V. Kohler, and S. Buttner, 2017 Taking out the garbage: cathepsin D and calcineurin in neurodegeneration. *Neural Regen. Res.* 12: 1776–1779. <https://doi.org/10.4103/1673-5374.219031>
- Azzi, J. R., M. H. Sayegh, and S. G. Mallat, 2013 Calcineurin inhibitors: 40 years later, can't live without. *J. Immunol.* 191: 5785–5791. <https://doi.org/10.4049/jimmunol.1390055>
- Berridge, M. J., M. D. Bootman, and H. L. Roderick, 2003 Calcium signalling: dynamics, homeostasis and remodelling. *Nat. Rev. Mol. Cell Biol.* 4: 517–529. <https://doi.org/10.1038/nrm1155>
- Buchholz, M., and V. Ellenrieder, 2007 An emerging role for Ca<sup>2+</sup>/calcineurin/NFAT signaling in cancerogenesis. *Cell Cycle* 6: 16–19. <https://doi.org/10.4161/cc.6.1.3650>
- Bueno, O. F., E. van Rooij, J. D. Molkentin, P. A. Doevendans, and L. J. De Windt, 2002 Calcineurin and hypertrophic heart disease: novel insights and remaining questions. *Cardiovasc. Res.* 53: 806–821. [https://doi.org/10.1016/S0008-6363\(01\)00493-X](https://doi.org/10.1016/S0008-6363(01)00493-X)
- Calalb, M. B., R. L. Kincaid, and T. R. Soderling, 1990 Phosphorylation of calcineurin: effect on calmodulin binding. *Biochem. Biophys. Res. Commun.* 172: 551–556. [https://doi.org/10.1016/0006-291X\(90\)90708-U](https://doi.org/10.1016/0006-291X(90)90708-U)
- Cardenas, M. E., C. Hemenway, R. S. Muir, R. Ye, D. Fiorentino *et al.*, 1994 Immunophilins interact with calcineurin in the absence of exogenous immunosuppressive ligands. *EMBO J.* 13: 5944–5957.
- Chow, E. W., S. A. Clancey, R. B. Billmyre, A. F. Averette, J. A. Granek *et al.*, 2017 Elucidation of the calcineurin-Crz1 stress response transcriptional network in the human fungal pathogen *Cryptococcus neoformans*. *PLoS Genet.* 13: e1006667. <https://doi.org/10.1371/journal.pgen.1006667>
- Connolly, S., and T. Kingsbury, 2012 Regulatory subunit myristoylation antagonizes calcineurin phosphatase activation in yeast. *J. Biol. Chem.* 287: 39361–39368. <https://doi.org/10.1074/jbc.M112.366617>
- Crabtree, G. R., and E. N. Olson, 2002 NFAT signaling: choreographing the social lives of cells. *Cell* 109: S67–S79. [https://doi.org/10.1016/S0092-8674\(02\)00699-2](https://doi.org/10.1016/S0092-8674(02)00699-2)
- Cruz, M. C., D. S. Fox, and J. Heitman, 2001 Calcineurin is required for hyphal elongation during mating and haploid fruiting in *Cryptococcus neoformans*. *EMBO J.* 20: 1020–1032. <https://doi.org/10.1093/emboj/20.5.1020>
- Cui, J., J. A. Kaandorp, O. O. Ositelu, V. Beaudry, A. Knight *et al.*, 2009 Simulating calcium influx and free calcium concentrations in yeast. *Cell Calcium* 45: 123–132. <https://doi.org/10.1016/j.ceca.2008.07.005>
- Cunningham, K. W., and G. R. Fink, 1994 Calcineurin-dependent growth control in *Saccharomyces cerevisiae* mutants lacking PMC1, a homolog of plasma membrane Ca<sup>2+</sup> ATPases. *J. Cell Biol.* 124: 351–363. <https://doi.org/10.1083/jcb.124.3.351>
- Cunningham, K. W., and G. R. Fink, 1996 Calcineurin inhibits VCX1-dependent H<sup>+</sup>/Ca<sup>2+</sup> exchange and induces Ca<sup>2+</sup> ATPases in *Saccharomyces cerevisiae*. *Mol. Cell. Biol.* 16: 2226–2237. <https://doi.org/10.1128/MCB.16.5.2226>
- Cyert, M. S., 2003 Calcineurin signaling in *Saccharomyces cerevisiae*: how yeast go crazy in response to stress. *Biochem. Biophys. Res. Commun.* 311: 1143–1150. [https://doi.org/10.1016/S0006-291X\(03\)01552-3](https://doi.org/10.1016/S0006-291X(03)01552-3)
- Cyert, M. S., and J. Thorner, 1992 Regulatory subunit (CNB1 gene product) of yeast Ca<sup>2+</sup>/calmodulin-dependent phosphoprotein phosphatases is required for adaptation to pheromone. *Mol. Cell. Biol.* 12: 3460–3469. <https://doi.org/10.1128/MCB.12.8.3460>
- Cyert, M. S., R. Kunisawa, D. Kaim, and J. Thorner, 1991 Yeast has homologs (CNA1 and CNA2 gene products) of mammalian calcineurin, a calmodulin-regulated phosphoprotein phosphatase. *Proc. Natl. Acad. Sci. USA* 88: 7376–7380 [corrigenda: *Proc. Natl. Acad. Sci. USA* 89: 4220 (1992)]. <https://doi.org/10.1073/pnas.88.16.7376>
- Deng, H., A. A. Gerencser, and H. Jasper, 2015 Signal integration by Ca<sup>2+</sup> regulates intestinal stem-cell activity. *Nature* 528: 212–217. <https://doi.org/10.1038/nature16170>
- Dolmetsch, R. E., R. S. Lewis, C. C. Goodnow, and J. I. Healy, 1997 Differential activation of transcription factors induced by Ca<sup>2+</sup> response amplitude and duration. *Nature* 386: 855–858. <https://doi.org/10.1038/386855a0>
- Dwivedi, M., H. O. Song, and J. Ahnn, 2009 Autophagy genes mediate the effect of calcineurin on life span in *C. elegans*. *Autophagy* 5: 604–607. <https://doi.org/10.4161/auto.5.5.8157>
- Erin, N., S. K. Bronson, and M. L. Billingsley, 2003 Calcium-dependent interaction of calcineurin with Bcl-2 in neuronal tissue. *Neuroscience* 117: 541–555. [https://doi.org/10.1016/S0306-4522\(02\)00933-8](https://doi.org/10.1016/S0306-4522(02)00933-8)
- Feng, B., and P. M. Stemmer, 1999 Interactions of calcineurin A, calcineurin B, and Ca<sup>2+</sup>. *Biochemistry* 38: 12481–12489. <https://doi.org/10.1021/bi990492w>
- Feng, B., and P. M. Stemmer, 2001 Ca<sup>2+</sup> binding site 2 in calcineurin-B modulates calmodulin-dependent calcineurin phosphatase activity. *Biochemistry* 40: 8808–8814. <https://doi.org/10.1021/bi0025161>
- Fischer, M., N. Schnell, J. Chattaway, P. Davies, G. Dixon *et al.*, 1997 The *Saccharomyces cerevisiae* CCH1 gene is involved in calcium influx and mating. *FEBS Lett.* 419: 259–262. [https://doi.org/10.1016/S0014-5793\(97\)01466-X](https://doi.org/10.1016/S0014-5793(97)01466-X)
- Forero, D. A., L. Herteleer, S. De Zutter, K. F. Norrback, L. G. Nilsson *et al.*, 2016 A network of synaptic genes associated with schizophrenia and bipolar disorder. *Schizophr. Res.* 172: 68–74. <https://doi.org/10.1016/j.schres.2016.02.012>
- Gallagher, S. C., Z. H. Gao, S. Li, R. B. Dyer, J. Trewthella *et al.*, 2001 There is communication between all four Ca<sup>2+</sup>-bindings sites of calcineurin B. *Biochemistry* 40: 12094–12102. <https://doi.org/10.1021/bi0025060>
- Geiser, J. R., D. van Tuinen, S. E. Brockerhoff, M. M. Neff, and T. N. Davis, 1991 Can calmodulin function without binding calcium? *Cell* 65: 949–959. [https://doi.org/10.1016/0092-8674\(91\)90547-C](https://doi.org/10.1016/0092-8674(91)90547-C)
- Gietz, R. D., R. H. Schiestl, A. R. Willems, and R. A. Woods, 1995 Studies on the transformation of intact yeast cells by the LiAc/SS-DNA/PEG procedure. *Yeast* 11: 355–360. <https://doi.org/10.1002/yea.320110408>
- Hashimoto, Y., and T. R. Soderling, 1989 Regulation of calcineurin by phosphorylation. Identification of the regulatory site

- phosphorylated by Ca<sup>2+</sup>/calmodulin-dependent protein kinase II and protein kinase C. *J. Biol. Chem.* 264: 16524–16529.
- Heineke, J., M. Auger-Messier, R. N. Correll, J. Xu, M. J. Benard *et al.*, 2010 CIB1 is a regulator of pathological cardiac hypertrophy. *Nat. Med.* 16: 872–879. <https://doi.org/10.1038/nm.2181>
- Hilioti, Z., D. A. Gallagher, S. T. Low-Nam, P. Ramaswamy, P. Gajer *et al.*, 2004 GSK-3 kinases enhance calcineurin signaling by phosphorylation of RCNs. *Genes Dev.* 18: 35–47. <https://doi.org/10.1101/gad.1159204>
- Horsley, V., and G. K. Pavlath, 2002 NFAT: ubiquitous regulator of cell differentiation and adaptation. *J. Cell Biol.* 156: 771–774. <https://doi.org/10.1083/jcb.200111073>
- Iida, H., Y. Yagawa, and Y. Anraku, 1990 Essential role for induced Ca<sup>2+</sup> influx followed by [Ca<sup>2+</sup>]<sub>i</sub> rise in maintaining viability of yeast cells late in the mating pheromone response pathway. A study of [Ca<sup>2+</sup>]<sub>i</sub> in single *Saccharomyces cerevisiae* cells with imaging of fura-2. *J. Biol. Chem.* 265: 13391–13399.
- Iida, H., H. Nakamura, T. Ono, M. S. Okumura, and Y. Anraku, 1994 MID1, a novel *Saccharomyces cerevisiae* gene encoding a plasma membrane protein, is required for Ca<sup>2+</sup> influx and mating. *Mol. Cell. Biol.* 14: 8259–8271. <https://doi.org/10.1128/MCB.14.12.8259>
- Imai, J., and I. Yahara, 2000 Role of HSP90 in salt stress tolerance via stabilization and regulation of calcineurin. *Mol. Cell. Biol.* 20: 9262–9270. <https://doi.org/10.1128/MCB.20.24.9262-9270.2000>
- Juvvadi, P. R., C. Gehrke, J. R. Fortwendel, F. Lamoth, E. J. Soderblom *et al.*, 2013 Phosphorylation of Calcineurin at a novel serine-proline rich region orchestrates hyphal growth and virulence in *Aspergillus fumigatus*. *PLoS Pathog.* 9: e1003564. <https://doi.org/10.1371/journal.ppat.1003564>
- Kahl, C. R., and A. R. Means, 2003 Regulation of cell cycle progression by calcium/calmodulin-dependent pathways. *Endocr. Rev.* 24: 719–736. <https://doi.org/10.1210/er.2003-0008>
- Kawamura, A., and M. S. Su, 1995 Interaction of FKBP12–FK506 with calcineurin A at the B subunit-binding domain. *J. Biol. Chem.* 270: 15463–15466. <https://doi.org/10.1074/jbc.270.26.15463>
- Kingsbury, T. J., and K. W. Cunningham, 2000 A conserved family of calcineurin regulators. *Genes Dev.* 14: 1595–1604.
- Klee, C. B., G. F. Draetta, and M. J. Hubbard, 1988 Calcineurin. *Adv. Enzymol. Relat. Areas Mol. Biol.* 61: 149–200.
- Klee, C. B., H. Ren, and X. Wang, 1998 Regulation of the calmodulin-stimulated protein phosphatase, calcineurin. *J. Biol. Chem.* 273: 13367–13370. <https://doi.org/10.1074/jbc.273.22.13367>
- Kujawski, S., W. Lin, F. Kitte, M. Bormel, S. Fuchs *et al.*, 2014 Calcineurin regulates coordinated outgrowth of zebrafish regenerating fins. *Dev. Cell* 28: 573–587. <https://doi.org/10.1016/j.devcel.2014.01.019>
- Lee, J. I., S. Mukherjee, K. H. Yoon, M. Dwivedi, and J. Bandyopadhyay, 2013 The multiple faces of calcineurin signaling in *Caenorhabditis elegans*: development, behaviour and aging. *J. Biosci.* 38: 417–431. <https://doi.org/10.1007/s12038-013-9319-6>
- Li, H., A. Rao, and P. G. Hogan, 2011 Interaction of calcineurin with substrates and targeting proteins. *Trends Cell Biol.* 21: 91–103. <https://doi.org/10.1016/j.tcb.2010.09.011>
- Li, S., S. M. Fell, O. Surova, E. Smedler, K. Wallis *et al.*, 2016a The 1p36 tumor suppressor KIF1Bbeta is required for calcineurin activation, controlling mitochondrial fission and apoptosis. *Dev. Cell* 36: 164–178. <https://doi.org/10.1016/j.devcel.2015.12.029>
- Li, S. J., J. Wang, L. Ma, C. Lu, J. Wang *et al.*, 2016b Cooperative autoinhibition and multi-level activation mechanisms of calcineurin. *Cell Res.* 26: 336–349. <https://doi.org/10.1038/cr.2016.14>
- Li, W., and R. E. Handschumacher, 1993 Specific interaction of the cyclophilin-cyclosporin complex with the B subunit of calcineurin. *J. Biol. Chem.* 268: 14040–14044.
- Luo, J., L. Sun, X. Lin, G. Liu, J. Yu *et al.*, 2014 A calcineurin- and NFAT-dependent pathway is involved in alpha-synuclein-induced degeneration of midbrain dopaminergic neurons. *Hum. Mol. Genet.* 23: 6567–6574. <https://doi.org/10.1093/hmg/ddu377>
- Mair, W., I. Morantte, A. P. Rodrigues, G. Manning, M. Montminy *et al.*, 2011 Lifespan extension induced by AMPK and calcineurin is mediated by CRTC-1 and CREB. *Nature* 470: 404–408. <https://doi.org/10.1038/nature09706>
- Manji, H. K., I. I. Gottesman, and T. D. Gould, 2003 Signal transduction and genes-to-behaviors pathways in psychiatric diseases. *Sci. STKE* 2003: pe49. <https://doi.org/10.1126/stke.2003.207.pe49>
- Martensen, T. M., B. M. Martin, and R. L. Kincaid, 1989 Identification of the site on calcineurin phosphorylated by Ca<sup>2+</sup>/CaM-dependent kinase II: modification of the CaM-binding domain. *Biochemistry* 28: 9243–9247. <https://doi.org/10.1021/bi00450a002>
- Matheos, D. P., T. J. Kingsbury, U. S. Ahsan, and K. W. Cunningham, 1997 Tcn1p/Crz1p, a calcineurin-dependent transcription factor that differentially regulates gene expression in *Saccharomyces cerevisiae*. *Genes Dev.* 11: 3445–3458. <https://doi.org/10.1101/gad.11.24.3445>
- Mathieu, F., S. Miot, B. Etain, M. A. El Khoury, F. Chevalier *et al.*, 2008 Association between the PPP3CC gene, coding for the calcineurin gamma catalytic subunit, and bipolar disorder. *Behav. Brain Funct.* 4: 2. <https://doi.org/10.1186/1744-9081-4-2>
- Medina, D. L., S. Di Paola, I. Peluso, A. Armani, D. De Stefani *et al.*, 2015 Lysosomal calcium signalling regulates autophagy through calcineurin and TFEB. *Nat. Cell Biol.* 17: 288–299. <https://doi.org/10.1038/ncb3114>
- Milan, D., J. Griffith, M. Su, E. R. Price, and F. McKeon, 1994 The latch region of calcineurin B is involved in both immunosuppressant-immunophilin complex docking and phosphatase activation. *Cell* 79: 437–447. [https://doi.org/10.1016/0092-8674\(94\)90253-4](https://doi.org/10.1016/0092-8674(94)90253-4)
- Moser, M. J., J. R. Geiser, and T. N. Davis, 1996 Ca<sup>2+</sup>-calmodulin promotes survival of pheromone-induced growth arrest by activation of calcineurin and Ca<sup>2+</sup>-calmodulin-dependent protein kinase. *Mol. Cell. Biol.* 16: 4824–4831. <https://doi.org/10.1128/MCB.16.9.4824>
- Mukherjee, A., and C. Soto, 2011 Role of calcineurin in neurodegeneration produced by misfolded proteins and endoplasmic reticulum stress. *Curr. Opin. Cell Biol.* 23: 223–230. <https://doi.org/10.1016/j.ceb.2010.12.006>
- Mukherjee, A., D. Morales-Scheihing, D. Gonzalez-Romero, K. Green, G. Tagliatalata *et al.*, 2010 Calcineurin inhibition at the clinical phase of prion disease reduces neurodegeneration, improves behavioral alterations and increases animal survival. *PLoS Pathog.* 6: e1001138. <https://doi.org/10.1371/journal.ppat.1001138>
- Musson, R. E., and N. P. Smit, 2011 Regulatory mechanisms of calcineurin phosphatase activity. *Curr. Med. Chem.* 18: 301–315. <https://doi.org/10.2174/092986711794088407>
- Musson, R. E., C. M. Cobbaert, and N. P. Smit, 2012 Molecular diagnostics of calcineurin-related pathologies. *Clin. Chem.* 58: 511–522. <https://doi.org/10.1373/clinchem.2011.167296>
- Nakai, Y., J. Horiuchi, M. Tsuda, S. Takeo, S. Akahori *et al.*, 2011 Calcineurin and its regulator sra/DSCR1 are essential for sleep in *Drosophila*. *J. Neurosci.* 31: 12759–12766. <https://doi.org/10.1523/JNEUROSCI.1337-11.2011>
- Nakamura, T., Y. Liu, D. Hirata, H. Namba, S. Harada *et al.*, 1993 Protein phosphatase type 2B (calcineurin)-mediated, FK506-sensitive regulation of intracellular ions in yeast is an important determinant for adaptation to high salt stress conditions. *EMBO J.* 12: 4063–4071.
- Nakayama, S., and R. H. Kretsinger, 1994 Evolution of the EF-hand family of proteins. *Annu. Rev. Biophys. Biomol. Struct.* 23: 473–507. <https://doi.org/10.1146/annurev.bb.23.060194.002353>
- Namgaladze, D., H. W. Hofer, and V. Ullrich, 2002 Redox control of calcineurin by targeting the binuclear Fe(2+)-Zn(2+) center at the enzyme active site. *J. Biol. Chem.* 277: 5962–5969. <https://doi.org/10.1074/jbc.M111268200>

- Nguyen, T., and S. Di Giovanni, 2008 NFAT signaling in neural development and axon growth. *Int. J. Dev. Neurosci.* 26: 141–145. <https://doi.org/10.1016/j.ijdevneu.2007.10.004>
- Nishiyama, T., N. Yoshizaki, T. Kishimoto, and K. Ohsumi, 2007 Transient activation of calcineurin is essential to initiate embryonic development in *Xenopus laevis*. *Nature* 449: 341–345. <https://doi.org/10.1038/nature06136>
- Nygaard, U., M. Deleuran, and C. Vestergaard, 2017 Emerging treatment options in atopic dermatitis: topical therapies. *Dermatology* 233: 333–343. <https://doi.org/10.1159/000484407>
- Odom, A., S. Muir, E. Lim, D. L. Toffaletti, J. Perfect *et al.*, 1997 Calcineurin is required for virulence of *Cryptococcus neoformans*. *EMBO J.* 16: 2576–2589. <https://doi.org/10.1093/emboj/16.10.2576>
- Paidhungat, M., and S. Garrett, 1997 A homolog of mammalian, voltage-gated calcium channels mediates yeast pheromone-stimulated Ca<sup>2+</sup> uptake and exacerbates the *cdc1(Ts)* growth defect. *Mol. Cell. Biol.* 17: 6339–6347. <https://doi.org/10.1128/MCB.17.11.6339>
- Patel, A., N. Yamashita, M. Ascano, D. Bodmer, E. Boehm *et al.*, 2015 RCAN1 links impaired neurotrophin trafficking to aberrant development of the sympathetic nervous system in Down syndrome. *Nat. Commun.* 6: 10119. <https://doi.org/10.1038/ncomms10119>
- Paul, A. S., S. Saha, K. Engelberg, R. H. Jiang, B. I. Coleman *et al.*, 2015 Parasite calcineurin regulates host cell recognition and attachment by apicomplexans. *Cell Host Microbe* 18: 49–60. <https://doi.org/10.1016/j.chom.2015.06.003>
- Peiris, H., and D. J. Keating, 2018 The neuronal and endocrine roles of RCAN1 in health and disease. *Clin. Exp. Pharmacol. Physiol.* 45: 377–383. <https://doi.org/10.1111/1440-1681.12884>
- Perrino, B. A., L. Y. Ng, and T. R. Soderling, 1995 Calcium regulation of calcineurin phosphatase activity by its B subunit and calmodulin. Role of the autoinhibitory domain. *J. Biol. Chem.* 270: 7012. <https://doi.org/10.1074/jbc.270.1.340>
- Peuker, K., S. Muff, J. Wang, S. Kunzel, E. Bosse *et al.*, 2016 Epithelial calcineurin controls microbiota-dependent intestinal tumor development. *Nat. Med.* 22: 506–515. <https://doi.org/10.1038/nm.4072>
- Philip, N., and A. P. Waters, 2015 Conditional degradation of plasmodium calcineurin reveals functions in parasite colonization of both host and vector. *Cell Host Microbe* 18: 122–131. <https://doi.org/10.1016/j.chom.2015.05.018>
- Qadota, H., I. Ishii, A. Fujiyama, Y. Ohya, and Y. Anraku, 1992 *RHO* gene products, putative small GTP-binding proteins, are important for activation of the *CAL1/CDC43* gene product, a protein geranyltransferase in *Saccharomyces cerevisiae*. *Yeast* 8: 735–741. <https://doi.org/10.1002/yea.320080906>
- Qu, J., R. Matsouaka, R. A. Betensky, B. T. Hyman, and C. L. Grosskreutz, 2012 Calcineurin activation causes retinal ganglion cell degeneration. *Mol. Vis.* 18: 2828–2838.
- Rodríguez, A., J. Roy, S. Martínez-Martínez, M. D. López-Maderuelo, P. Niño-Moreno *et al.*, 2009 A conserved docking surface on calcineurin mediates interaction with substrates and immunosuppressants. *Mol. Cell* 33: 616–626. <https://doi.org/10.1016/j.molcel.2009.01.030>
- Rothermel, B. A., R. B. Vega, and R. S. Williams, 2003 The role of modulatory calcineurin-interacting proteins in calcineurin signaling. *Trends Cardiovasc. Med.* 13: 15–21. [https://doi.org/10.1016/S1050-1738\(02\)00188-3](https://doi.org/10.1016/S1050-1738(02)00188-3)
- Roy, J., and M. S. Cyert, 2009 Cracking the phosphatase code: docking interactions determine substrate specificity. *Sci. Signal.* 2: re9. <https://doi.org/10.1126/scisignal.2100re9>
- Roy, J., H. Li, P. G. Hogan, and M. S. Cyert, 2007 A conserved docking site modulates substrate affinity for calcineurin, signaling output, and in vivo function. *Mol. Cell* 25: 889–901. <https://doi.org/10.1016/j.molcel.2007.02.014>
- Saneyoshi, T., S. Kume, Y. Amasaki, and K. Mikoshiba, 2002 The Wnt/calcium pathway activates NF-AT and promotes ventral cell fate in *Xenopus* embryos. *Nature* 417: 295–299. <https://doi.org/10.1038/417295a>
- Schulz, R. A., and K. E. Yutzey, 2004 Calcineurin signaling and NFAT activation in cardiovascular and skeletal muscle development. *Dev. Biol.* 266: 1–16. <https://doi.org/10.1016/j.ydbio.2003.10.008>
- Shah, S. Z., T. Hussain, D. Zhao, and L. Yang, 2017 A central role for calcineurin in protein misfolding neurodegenerative diseases. *Cell. Mol. Life Sci.* 74: 1061–1074. <https://doi.org/10.1007/s00018-016-2379-7>
- Sherman, F., 1991 Getting started with yeast. *Methods Enzymol.* 194: 3–21. [https://doi.org/10.1016/0076-6879\(91\)94004-V](https://doi.org/10.1016/0076-6879(91)94004-V)
- Shibasaki, F., and F. McKeon, 1995 Calcineurin functions in Ca(2+)-activated cell death in mammalian cells. *J. Cell Biol.* 131: 735–743. <https://doi.org/10.1083/jcb.131.3.735>
- Shibasaki, F., E. Kondo, T. Akagi, and F. McKeon, 1997 Suppression of signalling through transcription factor NF-AT by interactions between calcineurin and Bcl-2. *Nature* 386: 728–731. <https://doi.org/10.1038/386728a0>
- Sikorski, R. S., and P. Hieter, 1989 A system of shuttle vectors and yeast host strains designed for efficient manipulation of DNA in *Saccharomyces cerevisiae*. *Genetics* 122: 19–27.
- Someren, J. S., L. E. Faber, J. D. Klein, and J. A. Tumlin, 1999 Heat shock proteins 70 and 90 increase calcineurin activity in vitro through calmodulin-dependent and independent mechanisms. *Biochem. Biophys. Res. Commun.* 260: 619–625. <https://doi.org/10.1006/bbrc.1999.0800>
- Stathopoulos, A. M., and M. S. Cyert, 1997 Calcineurin acts through the CRZ1/TCN1-encoded transcription factor to regulate gene expression in yeast. *Genes Dev.* 11: 3432–3444. <https://doi.org/10.1101/gad.11.24.3432>
- Thewes, S., 2014 Calcineurin-Crz1 signaling in lower eukaryotes. *Eukaryot. Cell* 13: 694–705. <https://doi.org/10.1128/EC.00038-14>
- Tokoyoda, K., Y. Takemoto, T. Nakayama, T. Arai, and M. Kubo, 2000 Synergism between the calmodulin-binding and autoinhibitory domains on calcineurin is essential for the induction of their phosphatase activity. *J. Biol. Chem.* 275: 11728–11734. <https://doi.org/10.1074/jbc.275.16.11728>
- Tong, Y., and F. Song, 2015 Intracellular calcium signaling regulates autophagy via calcineurin-mediated TFEB dephosphorylation. *Autophagy* 11: 1192–1195. <https://doi.org/10.1080/15548627.2015.1054594>
- Tseng, H. H. L., C. T. Vong, Y. W. Kwan, S. M. Lee, and M. P. M. Hoi, 2017 Lysosomal Ca(2+) signaling regulates high glucose-mediated interleukin-1beta secretion via transcription factor EB in human monocytic cells. *Front. Immunol.* 8: 1161. <https://doi.org/10.3389/fimmu.2017.01161>
- Wang, H., Y. Du, B. Xiang, W. Lin, X. Li *et al.*, 2008 A renewed model of CNA regulation involving its C-terminal regulatory domain and CaM. *Biochemistry* 47: 4461–4468. <https://doi.org/10.1021/bi702539e>
- Wang, H. G., N. Pathan, I. M. Ethell, S. Krajewski, Y. Yamaguchi *et al.*, 1999 Ca<sup>2+</sup>-induced apoptosis through calcineurin dephosphorylation of BAD. *Science* 284: 339–343. <https://doi.org/10.1126/science.284.5412.339>
- Wang, X., V. C. Culotta, and C. B. Klee, 1996 Superoxide dismutase protects calcineurin from inactivation. *Nature* 383: 434–437. <https://doi.org/10.1038/383434a0>
- Wang, X., X. Peng, X. Zhang, H. Xu, C. Lu *et al.*, 2017 The emerging roles of CIB1 in cancer. *Cell. Physiol. Biochem.* 43: 1413–1424. <https://doi.org/10.1159/000481873>
- Wilkins, B. J., and J. D. Molkentin, 2004 Calcium-calcineurin signaling in the regulation of cardiac hypertrophy. *Biochem. Biophys. Res. Commun.* 322: 1178–1191. <https://doi.org/10.1016/j.bbrc.2004.07.121>

- Withee, J. L., J. Mulholland, R. Jeng, and M. S. Cyert, 1997 An essential role of the yeast pheromone-induced Ca<sup>2+</sup> signal is to activate calcineurin. *Mol. Biol. Cell* 8: 263–277. <https://doi.org/10.1091/mbc.8.2.263>
- Wu, H., A. Peisley, I. A. Graef, and G. R. Crabtree, 2007 NFAT signaling and the invention of vertebrates. *Trends Cell Biol.* 17: 251–260. <https://doi.org/10.1016/j.tcb.2007.04.006>
- Yang, S. A., and C. B. Klee, 2000 Low affinity Ca<sup>2+</sup>-binding sites of calcineurin B mediate conformational changes in calcineurin A. *Biochemistry* 39: 16147–16154. <https://doi.org/10.1021/bi001321q>
- Zayzafoon, M., 2006 Calcium/calmodulin signaling controls osteoblast growth and differentiation. *J. Cell. Biochem.* 97: 56–70. <https://doi.org/10.1002/jcb.20675>
- Zhang, T., J. Lei, H. Yang, K. Xu, R. Wang *et al.*, 2011 An improved method for whole protein extraction from yeast *Saccharomyces cerevisiae*. *Yeast* 28: 795–798. <https://doi.org/10.1002/yea.1905>
- Zhang, X., L. Yu, and H. Xu, 2016 Lysosome calcium in ROS regulation of autophagy. *Autophagy* 12: 1954–1955. <https://doi.org/10.1080/15548627.2016.1212787>
- Zhu, D., M. E. Cardenas, and J. Heitman, 1995 Myristoylation of calcineurin B is not required for function or interaction with immunophilin-immunosuppressant complexes in the yeast *Saccharomyces cerevisiae*. *J. Biol. Chem.* 270: 24831–24838. <https://doi.org/10.1074/jbc.270.42.24831>

Communicating editor: A. Mitchell