



Published in final edited form as:

J Org Chem. 2018 July 06; 83(13): 7109–7120. doi:10.1021/acs.joc.8b00094.

Comparative Ascaroside Profiling of *Caenorhabditis* Exometabolomes Reveals Species-Specific (ω) and ($\omega - 2$)-Hydroxylation Downstream of Peroxisomal β -Oxidation

Chuanfu Dong^{†,‡,iD}, Douglas K. Reilly[§], Célia Bergame^{||}, Franziska Dolke[†], Jagan Srinivasan^{§,iD}, and Stephan H. von Reuss^{*,†,||,iD}

[†]Department of Bioorganic Chemistry, Max Planck Institute for Chemical Ecology, Hans-Knoell Strasse 8, D-07745 Jena, Germany [‡]Department for Integrative Evolutionary Biology, Max Planck Institute for Developmental Biology, Max-Planck-Ring 9, D-72076 Tübingen, Germany [§]Department of Biology and Biotechnology, Worcester Polytechnic Institute, 60 Prescott Street, Worcester, Massachusetts 01605, United States ^{||}Laboratory of Bioanalytical Chemistry, University of Neuchâtel, Avenue de Bellevaux 51, CH-2000 Neuchâtel, Switzerland

Abstract

Chemical communication in nematodes such as the model organism *Caenorhabditis elegans* is modulated by a variety of glycosides based on the dideoxysugar L-ascarylose. Comparative ascaroside profiling of nematode exometabolome extracts using a GC-EIMS screen reveals that several basic components including ascr#1 (asc-C7), ascr#2 (asc-C6-MK), ascr#3 (asc-C9), ascr#5 (asc- ω C3), and ascr#10 (asc-C9) are highly conserved among the *Caenorhabditis*. Three novel side chain hydroxylated ascaroside derivatives were exclusively detected in the distantly related *C. nigoni* and *C. afra*. Molecular structures of these species-specific putative signaling molecules were elucidated by NMR spectroscopy and confirmed by total synthesis and chemical correlations. Biological activities were evaluated using attraction assays. The identification of (ω)- and ($\omega - 2$)-hydroxyacyl ascarosides demonstrates how GC-EIMS-based ascaroside profiling facilitates the detection of novel ascaroside components and exemplifies how species-specific hydroxylation of ascaroside aglycones downstream of peroxisomal β -oxidation increases the structural diversity of this highly conserved class of nematode signaling molecules.

Graphical abstract

*Corresponding Author. Phone: 0041 (0)32-718-2510; Fax: 0041 (0)32-718-2511; stephan.vonreuss@unine.ch.

ORCID

Chuanfu Dong: 0000-0003-3043-7257

Jagan Srinivasan: 0000-0001-5449-7938

Stephan H. von Reuss: 0000-0003-4325-5495


ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge on the ACS Publications website at DOI: 10.1021/acs.joc.8b00094.

Supporting figures as indicated in the text; NMR spectra of isolated ascarosides (10–12) and synthetic compounds (10a, 10b, 12, and 15–29) (PDF)

The authors declare no competing financial interest.



INTRODUCTION

Research with the model organism *Caenorhabditis elegans* and the development of novel analytical techniques promoted the characterization of ascarosides, a modular glycolipid library based on the 3,6-dideoxysugar L-ascarylose linked to fatty acid derived aglycones (Scheme 1).¹ In *C. elegans*, ascarosides modulate a large diversity of biological responses, including dauer formation,^{2–6} reproduction,^{6,7} stress resistance,^{7,8} lifespan,^{8,9} and behavior.^{10–14} The diversity of biological responses modulated by ascarosides is paralleled by their large structural diversity. Even small changes in molecular structures,^{7,14,15} synergistic effects,^{10,11,16} and variations in ascaroside compositions^{7,17,18} can dramatically alter their biological activity. Homologous series originate from chain shortening of very long chain precursors upon peroxisomal β -oxidation (Scheme 1A) to furnish ascarosides carrying acyl (**1**, asc-C#), enoyl (**2**, asc- C#), and (3*R*)-hydroxyacyl (**3**, asc- β OH-C#) aglycones.^{19–23} 2-Ketoalkyl (**6**, asc-C#-MK) and 2-hydroxyalkyl (**7**, asc-C#-OH) aglycones are presumably produced via decarboxylation and subsequent reduction of labile 3-ketoacyl intermediates (**4**, X = OH). Furthermore, (ω)-ascarosides such as asc- ω C3 (**8**, asc τ #5) demonstrate that (ω – 1) and (ω)-linked components enter the peroxisomal β -oxidation cycle. Downstream of the peroxisomal β -oxidation cycle, the resulting basic ascaroside skeletons serve as scaffolds for the attachment of additional structural units derived from primary metabolic pathways to furnish a modular library of species-specific signaling components such as the indole ascaroside IC-asc-C5 (**9**, icas#9) (Scheme 1B).^{5,12,16,22,24}

Ascaroside signaling is highly conserved in intraspecies nematode communication^{16,25–29} and also involved in intergenotypic competition^{30,31} and cross-kingdom interactions,^{32–34} demonstrating that ascarosides represent key regulators of nematode chemical ecology. Considering the small amounts of ascarosides present in nematode exometabolomes, the large diversity of homologous structures, and the complexity of the background matrix, mass spectrometric techniques are indispensable for comprehensive ascaroside analysis. Mass spectrometric screens that employ specific fragment ions as markers are capable to highlight putative nematode-derived ascarosides and their biosynthetic precursors and mask the background matrix. We previously developed an HPLC-ESI(-)-MS/MS precursor ion screen that employs an ascarylose-derived fragment ion at m/z 73.1 [$C_3H_5O_2$]⁻ for the selective detection of known as well as yet unidentified components.²² ESI(-)-MS/MS precursor ion screening has been employed in various studies^{22,25–27,29,33,34} but ultimately requires a triple quadrupole instrument. Considering the importance of ascaroside signaling

in nematode chemical ecology, we recently developed a complementary GC-EIMS technique³⁵ that employs an ascarylose-derived K1-fragment ion at m/z 130.1 [$C_6H_{14}OSi$]⁺ along with A1 and A2 fragment ions at m/z 275.1 [$C_{12}H_{27}O_3Si_2$]⁺ and m/z 185.1 [$C_9H_{17}O_2Si$]⁺ as characteristic markers to facilitate selective ascarioside profiling in trimethylsilyl (TMS) derivatized crude nematode exometabolome extracts (Scheme 2). Furthermore, aglycone-specific fragment ions for a rearranged oxonium ion (J1) at $[M - 173]^+$ and a carbocation (J2) at $[M - 291]^+$ facilitate the identification of compound specific side chains. Here, we employ the GC-EIMS screen for comparative ascarioside profiling in a variety of *Caenorhabditis* species to demonstrate its potential for the discovery of novel ascarioside components in crude unfractionated nematode exometabolome extracts.

RESULTS AND DISCUSSION

Following a phylometabolomic approach, we performed a GC-EIMS-based comparative analysis of exometabolome extracts from 13 *Caenorhabditis* species with a strong focus on the *Elegans* group that harbors the model organism *C. elegans*.^{36,37} Liquid cultures were established in S-medium, and propagating nematodes were fed with concentrated *E. coli* OP50 for 7 days, after which cultures were starved for another 7 days.³⁸ The media supernatant representing the exometabolome was collected, lyophilized, and extracted with methanol. Crude exometabolome extracts were converted into their TMS derivatives using *N*-methyl-*N*-(trimethylsilyl) trifluoroacetamide (MSTFA) and subsequently analyzed by GC-EIMS to show a large diversity of primary and secondary metabolites, some of which could be tentatively identified using the NIST 14 mass spectral library (Figure 1A). Putative ascariosides were detected by inspection of the extracted ion chromatograms for the highly characteristic ascarylose-derived K1 fragment ion at m/z 130.1 [$C_6H_{14}OSi$]⁺ (Figures 1B and S1). Individual ascarioside structures were identified based on their aglycone-specific J1 $[M - 173]^+$ and J2 $[M - 291]^+$ fragment ion signals and Kovats retention indices by comparison with a collection of more than 200 components that carry ($\omega - 1$)- or (ω)-linked acyl (**1**), enoyl (**2**), (3*R*)-hydroxyacyl (**3**), 2-ketoalkyl (**6**), or 2-hydroxyalkyl (**7**) side chains previously characterized in *C. elegans* wild-type and peroxisomal β -oxidation mutants.³⁵

These analyses demonstrate that basic ascariosides with side chains ranging from 3 to 11 carbons are highly conserved in all *Caenorhabditis* wild-type isolates tested, although their relative compositions vary significantly between the different species (Figure 2). A set of five compounds, asc-C7 (**1**, $n = 2$, X = OH; ascr#1, daumone#1),² asc-C9 (**1**, $n = 4$, X = OH; ascr#10),¹² asc-C9 (**2**, $n = 4$, X = OH; ascr#3, daumone#3),³ asc-C6-MK (**6**, $n = 2$; ascr#2, daumone#2),³ and asc- ω C3 (**8**, asc#5)⁴ consistently represent the dominating components and, taken together, account for more than 60% of the total ascariosides identified. An exception is the *C. portoensis* metabolome that predominantly contains asc-C11 (**1**, $n = 6$, X = OH; ascr#18). Furthermore, asc-C6-MK (**6**, $n = 2$; ascr#2) is absent in *C. nigoni*, *C. japonica*, *C. afra*, and *C. portoensis*, whereas asc- ω C3 (**8**, asc#5) is absent in *C. nigoni*, *C. japonica*, and *C. portoensis*. Asc-C6-OH (**7**, $n = 2$; ascr#6) was exclusively detected alongside asc-C6-MK (**6**, $n = 2$; ascr#2), supporting the assumption of a common biosynthetic origin (Scheme 1). While these basic ascariosides are common in most *Caenorhabditis* species tested, GC-EIMS-based ascarioside profiling also revealed some highly species-specific components, including a yet unidentified component with a J2

fragment at m/z 186.1 from *C. elegans*, asc- β OH-C13 (**3**, $n = 8$, X = OH; bhas#22) from *C. japonica*, and asc-C5-EA (**1**, $n = 0$, X = NHCH₂CH₂OH; easc#9) from *C. portoensis*.³⁵

In addition, we observed 3 putative ascarosides that accounted for around 10% of the total ascarosides detected in the exometabolome of *C. nigoni* (Figures 2 and 1B) but did not match any of the more than 200 basic ascaroside structures previously identified in *C. elegans* wild-type and mutant metabolomes.³⁵ Inspection of their 70 eV EIMS spectra (Figure 3) revealed dominating K1-fragment ion signals at m/z 130.1 [C₆H₁₄OSi]⁺, along with A1-fragments at m/z 275.2 [C₁₂H₂₇O₃Si₂]⁺ and A2-fragments at m/z 185.1 [C₉H₁₇O₂Si]⁺ that are characteristic for the ascarylose unit (Scheme 2). Several aglycone-specific signals for J1 fragment ions at m/z 433.2 [C₁₉H₄₁O₅Si₃]⁺, J2 fragment ions at m/z 315.2 [C₁₅H₃₁O₃Si₂]⁺, and (J2 – TMSOH) fragment ions at m/z 225.1 [C₁₂H₂₁O₂Si]⁺ indicated nine carbon side chains with one additional unit of unsaturation and one trimethylsilyloxy moiety for compounds **10** and **12**. The third compound **11** displayed diagnostic signals for a J1 fragment at m/z 435.3 [C₁₉H₄₃O₅Si₃]⁺, a J2 fragment at m/z 317.2 [C₁₅H₃₃O₃Si₂]⁺, and a (J2 – TMSOH) fragment at m/z 227.1 [C₁₂H₂₃O₂Si]⁺ indicative for a trimethylsilyloxy-substituted nine carbon side chain. Comparison with the known (3*R*)-hydroxylated asc- β OH-C9 (**3**, $n = 4$, X = OH; bhas#10), previously characterized in the exometabolome of *Panagrellus redivivus*,^{26,35} demonstrated that both compounds are different (Figure S2) and excluded a β -oxidation-derived 3-hydroxyacyl aglycone due to the lack of the characteristic fragment ion at m/z 233.1 [C₉H₂₁O₃Si₂]⁺ derived from α -cleavage. However, the identification of a homologous fragment ion signal at m/z 289.2 [C₁₃H₂₉O₃Si₂]⁺ suggested a 7-hydroxyacyl structure for asc-7OH-C9 (**11**) from *C. nigoni*, demonstrating how EIMS fragmentation can aid in structure assignment.

In conclusion, comparative GC-EIMS-based ascaroside screening revealed three species-specific, side chain hydroxylated compounds in the *C. nigoni* exometabolome. The same components with molecular ion signals at m/z 317.2 [M – H][–] for **10** and **12** and m/z 319.2 [M – H][–] for **11** were also detected using the HPLC-ESI(–)-MS/MS precursor ion screen for m/z 73.1 [C₃H₅O₂][–] (Figure 1C), demonstrating that the GC-EIMS and HPLC-MS/MS methods complement each other. However, in contrast to GC-EIMS, the MS/MS precursor ion screen also revealed additional derivatives, including large amounts of indole ascarosides such as IC-asc-C5 (**9**, icas#9), a male attractant in *C. nigoni*,¹⁶ demonstrating that GC-EIMS is restricted to the most basic ascaroside compounds.

To identify the molecular structures of the species-specific ascarosides (**10–12**), the exometabolome extract of 1.6 L *C. nigoni* liquid culture supernatant was fractionated by solid phase extraction (RP-C₁₈-SPE) using a 10%-stepwise gradient of aqueous methanol as eluent. Fractions were screened for ascarosides by GC-EIMS (Figure S3) and ¹H NMR spectroscopy (Figure S4) using the K1 fragment ion at m/z 130.1 [C₆H₁₄OSi]⁺ and the anomeric proton at approximately δ_{H} 4.65 ppm (s, 1H) as characteristic markers, respectively. These analyses confirmed the assignment of several known ascarosides and traced the target compounds to a fraction eluted with 40% methanol (Figure S3) that contained predominantly asc-C7 (**1**, $n = 2$, X = OH; asc#1) along with a diversity of additional metabolites such as indole acetic acid (IAA, auxine) and anthranilic acid (Figure

S5). Subsequent separation by semipreparative HPLC using a C18 column furnished fractions of sufficient purity to facilitate structure assignment.

The molecular formula of $C_{15}H_{26}O_7$ for compound **10** (~275 μg) was established by HR-EIMS. Inspection of one- and two-dimensional NMR spectra (^1H NMR, *dqf*-COSY, HSQC) confirmed an α -configured ascarylose moiety along with an $(\omega - 1)$ -linked α,β -unsaturated C9 side chain (Table 1). Furthermore, the $(\omega - 2)$ -position of the additional hydroxy group was deduced based on *dqf*-COSY correlations from the terminal (ω) -methyl group at δ_{H} 1.14 ppm (d, $J = 6.3$ Hz, 3H), δ_{C} 14.5 ppm to the $(\omega - 1)$ -oxymethine group at δ_{H} 3.74 ppm (dq, $J = 3.9$ Hz, $J = 6.3$ Hz, 1H), δ_{C} 75.5 ppm and further on to the $(\omega - 2)$ -position at δ_{H} 3.53 ppm (m, 1H), δ_{C} 74.8 ppm. While all $(\omega - 1)$ -linked homologous ascarysides that have been identified so far share the same (*R*)-configuration at the penultimate carbon,¹ the stereochemistry of the $(\omega - 2)$ -hydroxymethine group of asc-7OH-C9 (**10**) could not be unambiguously assigned based solely on the vicinal *H,H*-coupling constant of 3.9 Hz.³⁹

Compound **11** (~110 μg) with a molecular formula of $C_{15}H_{28}O_7$ according to ESI-HRMS exhibits almost identical NMR data for the ascarylose unit and the (ω) -part of the side chain but displays a triplet signal at 2.21 (t, $J = 7.6$ Hz, 2H) instead of the signals for an α,β -unsaturated enoyl moiety (Table 1), thus suggesting the corresponding dihydro-derivative structure asc-7OH-C9 (**11**).

Compound **12** (~130 μg) with a molecular formula of $C_{15}H_{26}O_7$ according to ESI-HRMS was obtained as a 1:1 mixture with asc-7OH-C9 (**11**). Comparative analysis of their *dqf*-COSY spectra indicated an ascarylose unit with considerably different chemical shifts (Table 1). Furthermore, an α,β -unsaturated side chain was identified due to δ_{H} 5.81 (d, $J_{\text{E}} = 15.6$ Hz, 1H) and 6.95 (dt, $J_{\text{E}} = 15.6$ Hz, $J = 6.9$ Hz, 1H), along with a hydroxymethylene group at δ_{H} 3.50 (dd, $^2J = 11.7$ Hz, $^3J = 5.6$ Hz, 1H), 3.60 (dd, $^2J = 11.7$ Hz, $^3J = 4.2$ Hz, 1H), and δ_{C} 64.6 ppm that displayed vicinal *H,H*-coupling correlation to the $(\omega - 1)$ -oxymethine proton at δ_{H} 3.69 (m, 1H) and δ_{C} 78.6 ppm, thus indicating the terminal (ω) -position for the hydroxylation of the aglycone in asc-9OH-C9 (**12**). In conclusion, analysis of one and two-dimensional NMR spectra revealed three novel side chain modified ascarysides that carry additional hydroxy functions at the 7-position $(\omega - 2)$ and the 9-position (ω) , the structures of which were finally established by total synthesis and chemical correlations.

Both diastereomeric $(\omega - 2)$ -hydroxy ascarysides (*7R,8R*)-*threo*-**10a** and (*7S,8R*)-*erythro*-**10b** were synthesized as shown in Scheme 3. (*R*)-Methyl lactate (**14**) was converted to the *para*-methoxybenzyl (PMB) ether **15** and reduced to the aldehyde **16** using DIBAL-H. Addition of 4-pentenylmagnesium bromide afforded (*6R,7R*)-*threo*-6-hydroxy-7-PMB-*O*-1-octene (**17**) with a diastereomeric excess (de) of 92% due to asymmetric induction via a Cram chelate complex.⁴⁰ Esterification of **17** with benzoyl chloride and pyridine or benzoic acid under Mitsunobu conditions (PPh₃, DIAD) afforded the diastereomeric benzoates (*6R,7R*)-*threo*-**18a** or (*6S,7R*)-*erythro*-**18b**, respectively, in excellent diastereomeric purities of de > 99% after column chromatography (Figure S6). Cross metathesis with ethyl acrylate using Grubbs second generation catalyst⁴¹ furnished the corresponding ethyl 7-Bz-*O*-8-PMB-*O*-(2*E*)-nonenoates (*7R,8R*)-*threo*-**19a** or (*7S,8R*)-*erythro*-**19b** that were subsequently deprotected using 2,3-dichloro-5,6-dicyano-1,4-benzoquinone (DDQ)⁴² to afford (*7R,8R*)-

threo-20a or *(7S,8R)-erythro-20b*. Coupling with 2,4-di-*O*-benzoyl protected ascarose via the trichloroacetimidate route⁴³ furnished the diastereomeric ascarosides (**22a** or **22b**) that were finally deprotected using alkaline hydrolysis to give the desired ($\omega - 2$)-hydroxy ascarosides *(7R,8R)-threo-10a* or *(7S,8R)-erythro-10b*. In addition, an undesired intramolecular cyclization product (**23a** or **23b**) was obtained. Comparison of the NMR (Figure S7) and GC-EIMS (Figure S8) data of diastereomeric *(7R,8R)-threo-10a* and *(7S,8R)-erythro-10b* with those of the natural product (**10**) isolated from the *C. nigoni* exometabolome confirmed its *(7R,8R)-threo* configuration. Furthermore, the *(7R,8R)-threo*-configuration of the dihydro-derivative asc-7OH-C9 (**11**) from *C. nigoni* was unambiguously established by comparison of the mass spectra and GC retention times with those of the hydrogenation products of *(7R,8R)-threo-10a* and *(7S,8R)-erythro-10b* (Figure S9).

The (ω)-hydroxylated ascaroside asc-9OH- C9 (**12**) was synthesized as shown in Scheme 4. Copper(I)-catalyzed addition of pentenyl magnesium bromide to *tert*-butyldimethylsilyl (*S*)-glycidyl ether (**24**) afforded (*S*)-8-*tert*-butyldimethylsilyloxy-7-hydroxy-1-octene (**25**). Cross metathesis with ethyl acrylate using Grubbs second generation catalyst⁴¹ gave the corresponding (8*S*,2*E*)-ethyl 8-hydroxy-9-*tert*-butyldimethylsilyloxy-2-nonenoate (**26**). Coupling of **26** to 2,4-di-*O*-benzoyl-ascarsyl trichloroacetimidate (**21**) furnished **27**, which was subsequently deprotected using tetrabutylammonium fluoride (TBAF) to give the alcohol **28**, followed by alkaline hydrolysis to give asc-9OH- C9 (**12**) identical to the natural product from *C. nigoni* as shown by comparison of the GC-EIMS and NMR data (Figures S8 and S10). In addition, an undesired intramolecular cyclization product **29** with an oxacyclooctan ring was obtained.

Using the synthetic compounds as reference standards, *threo*-asc-7OH- C9 (**10a**), *threo*-asc-7OH-C9 (**11a**), and asc-9OH- C9 (**12**) could also be identified in the exometabolome extract of *Caenorhabditis afra* (sp. 7) strain JU1286 from Ghana (Figures S11 and S12), a member of the distantly related Japonica group (Figure 2),^{36,37} suggesting that (ω) and ($\omega - 2$)-hydroxylation in *C. nigoni* and *C. afra* has a polyphyletic origin. Targeted screening for homologous compounds using GC-EIMS and HPLC-MS demonstrated that (ω) and ($\omega - 2$)-hydroxylation in *C. nigoni* and *C. afra* is tightly controlled and exclusively affects C9 and C9 aglycones, indicating that this species-specific modification of ascaroside aglycones occurs downstream of peroxisomal β -oxidation.

Aiming to characterize the biological functions of the novel ascarosides, behavioral response of *C. nigoni* males or females to 1 μ M asc-(*7R*)-OH- C9 (**10a**), asc-(*7S*)-OH- C9 (**10b**), asc-9OH- C9 (**12**), as well as asc- C9 (**2**, $n = 4$, X = OH; ascr#3) was evaluated using a spot attraction assay. While these analyses demonstrate that the common asc- C9 (**2**, $n = 4$, X = OH; ascr#3) acts as a potent male attractant in *C. nigoni*, reminiscent of the attraction of *C. elegans* males,¹⁰ the species-specific ($\omega - 2$) and (ω)-hydroxylated derivatives (**10a**, **10b**, and **12**) are not active (Figure S13). Additional experiments are required to clarify the biological functions of these compounds and unravel the ecological significance of species-specific (ω) and ($\omega - 2$)-hydroxylation in *C. nigoni* and *C. afra*.

CONCLUSION

Our results demonstrate that comparative GC-EIMS-based ascaroside profiling represents a powerful technique to characterize ascaroside diversity and detect novel species-specific components in TMS-derivatized crude nematode exometabolomes. We isolated three new (ω) and ($\omega - 2$)-hydroxylated ascarosides from *C. nigoni* and determined their structures using NMR spectroscopy. Structure assignments were unambiguously established by total synthesis and chemical correlations. Comparative analysis of 13 *Caenorhabditis* species demonstrates that within the *Elegans* group, both (ω)- and ($\omega - 2$)-hydroxylated ascarosides (**10–12**) are highly specific for *C. nigoni* (sp. 9), but the same compounds were also detected in the rather distantly related *C. afra* (sp. 7), a member of the Japonica group, thus suggesting a polyphyletic origin for the hydroxylation steps. In addition, we found that (ω)- and ($\omega - 2$)-hydroxylation in *C. nigoni* and *C. afra* is highly specific for ascarosides carrying C9 and C9 side chains, strongly suggesting that the hydroxylation step occurs downstream of peroxisomal β -oxidation. While traces of (3*R*)-hydroxyacyl ascarosides, intermediates of the peroxisomal β -oxidation cycle, are widespread in nematode exometabolomes, side chain hydroxylation downstream of β -oxidation such as (ω)- and ($\omega - 2$)-hydroxylation in *C. nigoni* and *C. afra* has so far been described only as male-specific ($\omega - 3$)-hydroxylation in *Panagrellus redivivus* (Scheme 1C).²⁶ However, while the dihydroxylated dhas#18 (**13**) represents a male-produced female-attractant in *Panagrellus redivivus*,²⁶ the hydroxy ascarosides (**10–12**) did not attract *C. nigoni* males or females. Additional research is required to elucidate the biological functions of these species-specific components and decipher the ecological significance of (ω)- and ($\omega - 2$)-hydroxylation of ascaroside aglycones as a means to further increase structural diversity of this highly conserved class of nematode signaling molecules.

EXPERIMENTAL SECTION

Preparation of Exometabolome Extracts

Wild-type isolates of 13 *Caenorhabditis* species were cultivated at 23 °C on NGM agar seeded with *E. coli* OP50: *C. elegans* N2 (Bristol), *C. nigoni* (sp. 9) JU1422, *C. briggsae* AF16, *C. sinica* (sp. 5) JU727, *C. remanei* PB4641, *C. tropicalis* (sp. 11) JU1373, *C. wallacei* (sp. 16) JU1904, *C. doughertyi* (sp. 10) JU1771, *C. brenneri* (sp. 4) PB2801, *C. japonica* DF5081, *C. afra* (sp. 7) JU1286, *C. portoensis* (sp. 6) EG4788, and *C. n.* sp. 8 (sp. 8) QX1182. Mixed stage nematodes from five 10 cm plates collected in M9 buffer served as inoculums for liquid cultures grown in 100 mL S-medium at 23 °C and 150 rpm. Concentrated *E. coli* OP50 bacteria pellet from an overnight culture in LB medium at 37 °C and 170 rpm was provided as food from days 1–7, after which the cultures were starved for 7 days. After 14 days, nematodes were separated by centrifugation (5 min at 5000*g*). The filtered supernatant representing the exometabolome was frozen at –80 °C, lyophilized, and extracted with 3 × 100 mL methanol for 12 h each. The combined extract was filtered, concentrated to dryness at 40 °C under reduced pressure, and reconstituted in 1 mL methanol, and aliquots were analyzed by HPLC-HRMS, HPLC-MS/MS precursor ion screening for *m/z* 73.1, and GC-EIMS. All experiments were performed in triplicate.

Preparation of Trimethylsilyl (TMS) Derivatives for GC-EIMS Analysis

Aliquots of crude nematode exometabolome extracts, *C. nigoni* exometabolome fractions, and synthetic ascaroside standards were concentrated to dryness. The residues were treated with 10 μ L *N*-methyl-*N*-(trimethylsilyl) trifluoroacetamide (MSTFA) at 60 °C for 40 min and diluted with 10 μ L DCM, and 1 μ L of the solution was analyzed by GC-EIMS.

Gas Chromatography-Electron Impact-Mass Spectrometry (GC-EIMS)

Separation of volatile TMS derivatives and acquisition of their 70 eV electron impact mass spectra was performed using a Trace GC 2000 series (Thermo Scientific) equipped with a Zebtron ZB-5 Guardian column (15 m, 0.25 mm ID, 0.25 μ m film thickness; with 10 m guardian end) coupled to a single quadrupole ThermoQuest Trace MS (Finnigan). Helium was used as the carrier gas at a flow rate of 1 mL/min. A temperature program starting at 130 °C for 5 min, followed by a linear gradient of +10 °C/min to 350 °C was applied. A total volume of 1 μ L was injected using a 10:1 split ratio and an injector temperature of 250 °C. Electron ionization (EI, 70 eV) mass spectra were acquired from m/z 35–650 amu. Data were analyzed with the Xcalibur 3.1 software (Thermo Fisher Scientific).

Liquid Chromatography-Electrospray Ionization-High Resolution-Mass Spectrometry (HPLC-ESI-HRMS)

HPLC-ESI-HRMS analysis of crude nematode exometabolome extracts and *C. nigoni* exometabolome fractions was performed using a Dionex UltiMate 3000 HPLC instrument coupled to a Bruker Maxis ultrahigh resolution (UHR) qTOF mass spectrometer equipped with an electrospray ionization (ESI) unit operated in positive or negative mode. Chromatographic separations were achieved using an Agilent ZORBAX Eclipse XDB-C18 column (250 \times 3 mm, 5 μ m particle diameter) with a flow rate of 400 μ L/min and gradient elution starting at 3% acetonitrile in 0.5% aqueous acetic acid for 5 min followed by a linear increase to 100% acetonitrile with 0.5% acetic acid within 35 min. Data were analyzed with the Compass DataAnalysis 4.3 software (Bruker).

Liquid Chromatography Electrospray Ionization Precursor Ion Screening

HPLC-MS/MS precursor ion screening for m/z 73.1 was performed using an Agilent 1260 HPLC instrument (Agilent Technologies) coupled to an API5000 Triple Quadrupole LC/MS/MS mass spectrometer (AB Sciex, Darmstadt) equipped with an electrospray ionization (ESI) unit operated in negative mode. A CID energy of –34 was applied. Chromatographic separations were achieved using an Agilent ZORBAX Eclipse XDB-C18 column (50 \times 4.6 mm, 1.8 μ m particle diameter) (Agilent Technologies) with a flow rate of 1.1 mL/min and gradient elution starting at 5% acetonitrile in 0.05% aqueous formic acid followed by a linear increase to 95% acetonitrile with 0.05% formic acid within 10 min. Data were analyzed with the Analyst 1.6 software (AB Sciex).

NMR Spectroscopy

NMR spectra were recorded in CD₃OD or CDCl₃ at 400 MHz for ¹H and 100 MHz for ¹³C using a Bruker AMX400 instrument. Residual solvent signals were used as internal standard with ¹H at 3.31 ppm and ¹³C at 49.05 ppm for CD₃OD or ¹H at 7.26 ppm and ¹³C at 77.16

ppm for CDCl₃. Two-dimensional homonuclear double quantum filtered (*dqf*)-COSY spectra were recorded using phase cycling for coherence selection. For the isolated compounds a total of 32 scans were acquired using a time domain of 8k in F2 (acquisition time of 1.2 s) and 512 increments in F1. For two-dimensional heteronuclear HSQC spectra 96 scans were acquired using a time domain of 1k in F2 and 256 increments in F1. Spectra were zero-filled to 8k × 4k (COSY) or 4k × 2k (HSQC) prior to Fourier transformation, phased manually, and baseline corrected using the Topspin 3.2 (Bruker) and MNova 9.0 (Mestrelab Research) software.

Spot Retention Assay

Assays were performed as described previously.^{10,44} 50–60 larval-stage 4 (L4) worms were segregated by sex and stored at 20 °C for 5 h to overnight to be assayed as young adults. 0.6 μL of vehicle control or ascaroside solution was placed in each scoring region. As the working stock of ascaroside was made in Milli-Q-purified ultrapure H₂O, this was used as the vehicle control. Five animals were placed on each “X” of the assay plate, which was then transferred to a microscope containing a camera and recorded for 20 min. Each sex and compound was assayed over three plates per day on at least three different days.

Isolation of Hydroxy Ascarosides from the *C. nigoni* Exometabolome

Hydroxyacyl ascarosides of *C. nigoni* were isolated from 1.6 L of the liquid culture supernatant. The filtered supernatant was frozen at –80 °C, lyophilized, and the residue extracted with 3 × 100 mL methanol for 12 h each. The filtered extract was concentrated to dryness under reduced pressure and the resulting *C. nigoni* exometabolome extract was adsorbed onto 2 g of Celite and fractionated by reverse phase chromatography on 5 g RP-C₁₈-SPE cartridges (Chromabond, Macherey-Nagel) using increasing concentrations of methanol in water as eluent to afford 10 fractions (20 mL each). Aliquots of 10 μL were concentrated to dryness under reduced pressure, treated with 10 μL MSTFA at 60 °C for 30 min, diluted with 10 μL DCM, and analyzed by GC-EIMS (Figure S3). Fractions were concentrated to dryness under reduced pressure and analyzed by ¹H NMR spectroscopy (Figure S4). The 40% methanol fraction containing the target components according to GC-EIMS was subsequently submitted to semipreparative HPLC using an Agilent HP-1100 HPLC instrument equipped with a Grom-Sil 120 ODS-4 HE column (250 × 8 mm, 5 μm) coupled to a Gilson 206 Abimed fraction collector. A flow rate of 2 mL/min with gradient elution was used starting at 3% acetonitrile in 0.5% aqueous acetic acid for 3 min, followed by a linear increase to 100% acetonitrile with 0.5% acetic acid within 30 min. Aliquots of 10 μL were analyzed by GC-EIMS and HPLC-ESI(–)-HR-MS as described before. Fractions containing the target compounds were concentrated to dryness, dissolved in 650 μL CD₃OD, and analyzed by one- and two-dimensional NMR spectroscopy.

(7R,8R,2E)-threo-8-[(3',6'-Dideoxy-α-L-arabino-hexopyranosyl)-oxy]-7-hydroxy-2-nonenic Acid (threo-Asc-7OH- C9, 10)—Isolated from the *C. nigoni* exometabolome (275 μg, *c* = ~540 nmol/L), for ¹H and ¹³C NMR data see Table 1; HRMS (ESI-TOF) *m/z* (M – H)[–] calcd for C₁₅H₂₅O₇ 317.1606, found 317.1619.

(7R,8R)-threo-8-[(3',6'-Dideoxy- α -L-arabino-hexopyranosyl)oxy]-7-hydroxynonanoic Acid (threo-Asc-7OH-C9, 11)—Isolated from the *C. nigoni* exometabolome (110 μ g, ~215 nmol/L), for ^1H and ^{13}C NMR data see Table 1; HRMS (ESI-TOF) m/z ($\text{M} - \text{H}$) $^-$ calcd for $\text{C}_{15}\text{H}_{27}\text{O}_7$ 319.1762, found 319.1771.

(2E,8S)-8-[(3,6-Dideoxy- α -L-arabino-hexopyranosyl)oxy]-9-hydroxy-2-nonenoic Acid (Asc-9OH-C9, 12)—Isolated from the *C. nigoni* exometabolome (130 μ g, ~255 nmol/L), for ^1H and ^{13}C NMR data see Table 1; HRMS (ESI-TOF) m/z ($\text{M} - \text{H}$) $^-$ calcd for $\text{C}_{15}\text{H}_{25}\text{O}_7$ 317.1606, found 317.1617.

(R)-Methyl 2-(4-methoxybenzyloxy)propanoate (15).⁴⁵—Under argon atmosphere a solution of (*R*)-(+)-methyl 2-hydroxypropanoate (**14**) (1.04 g, 10 mmol) and 4-methoxybenzyl 2,2,2-trichloroacetimidate (2.8 g, 10 mmol) in dry DCM (15 mL) at 0 $^\circ\text{C}$ was treated with trimethylsilyl triflate (10 μL). After being stirred at 0 $^\circ\text{C}$ for 3 h, the reaction was quenched by addition of saturated NaHCO_3 solution (1 mL), and the mixture was diluted with DCM (15 mL), washed with saturated NaHCO_3 solution (2×10 mL), dried over Na_2SO_4 , and concentrated under reduced pressure. The product was isolated by column chromatography (silica gel, 9/1 v/v hexane/ethyl acetate elution, $R_f = 0.22$) to afford **15** (1.38 g, 6.2 mmol, 62%) as a yellowish oil. ^1H NMR (400 MHz, CDCl_3) δ 7.28 (d, $J = 8.7$ Hz, 2H), 6.87 (d, $J = 8.7$ Hz, 2H), 4.61 (d, $J = 11.3$ Hz, 1H), 4.38 (d, $J = 11.3$ Hz, 1H), 4.05 (q, $J = 6.8$ Hz, 1H), 3.79 (s, 3H), 3.75 (s, 3H), δ 1.41 (d, $J = 6.8$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ 173.9, 159.5, 129.72, 129.70, 113.9, 73.7, 71.8, 55.4, 52.0, 18.8.⁴⁵

(R)-2-(4-Methoxybenzyloxy)propanal (16).⁴⁵—Under argon atmosphere a solution of **15** (672 mg, 3 mmol) in dry DCM (10 mL) at -78 $^\circ\text{C}$ was treated dropwise with a 1 M DIBAL-H solution (3.3 mL, 3.3 mmol) in toluene. After stirring at -78 $^\circ\text{C}$ for 30 min the reaction was quenched with methanol (0.5 mL) and saturated sodium potassium tartrate solution (10 mL) and stirred for 1 h and extracted with DCM (2×20 mL). The combined organic phase was dried over Na_2SO_4 and concentrated under reduced pressure. Column chromatography (silica gel, DCM elution) afforded **16** (580 mg, 3 mmol, 100% yield) as a colorless oil that was directly used for the next step. ^1H NMR (400 MHz, CDCl_3) δ 9.63 (d, $J = 1.8$ Hz, 1H), 7.29 (d, $J = 8.7$ Hz, 2H), 6.89 (d, $J = 8.7$ Hz, 2H), 4.57 (d, $J = 11.4$ Hz, 1H), 4.54 (d, $J = 11.4$ Hz, 1H), 3.87 (dq, $J = 1.8$ Hz, $J = 7.0$ Hz, 1H), 3.80 (s, 3H), 1.31 (d, $J = 7.0$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ 203.7, 159.7, 129.8, 129.5, 114.1, 79.3, 71.9, 55.4, 15.4.⁴⁵

(2R,3R)-threo-3-Hydroxy-2-(4-methoxybenzyloxy)-7-octene (17)—Under argon atmosphere, a solution of 4-pentenylmagnesium bromide (6 mmol) in diethyl ether (10 mL) prepared from 5-bromo-1-pentene (900 mg, 6 mmol) and magnesium (150 mg, 6.2 mmol) was cooled to 0 $^\circ\text{C}$ and treated dropwise with **16** (580 mg, 3 mmol) in Et_2O (2 mL) over the course of 5 min. The resulting mixture was stirred at 0 $^\circ\text{C}$ for 1 h and quenched with saturated aqueous NH_4Cl solution (10 mL), and the aqueous layer extracted with ethyl acetate (2×10 mL). The combined organic phases were dried over Na_2SO_4 and concentrated under reduced pressure. The product was isolated by column chromatography (silica gel, 4/1 v/v hexane/ethyl acetate elution, $R_f = 0.42$) to afford **17** (460 mg, 1.74 mmol,

58% yield) with a diastereoisomeric excess of $de = 92\%$ as determined by ^1H NMR spectroscopy. ^1H NMR (400 MHz, CDCl_3) δ 7.25 (d, $J = 8.6$ Hz, 2H), 6.88 (d, $J = 8.6$ Hz, 2H), 5.80 (ddt, $J = 17.2$ Hz, $J = 10.2$ Hz, $J = 6.7$ Hz, 1H), 5.00 (dbr, $J = 17.2$ Hz, 1H), 4.94 (dbr, $J = 10.3$ Hz, 1H), 4.60 (d, $^2J = 11.1$ Hz, 1H), 4.36 (d, $^2J = 11.1$ Hz, 1H), 3.80 (s, 3H), 3.40 (m, 1H), 3.34 (dq, $J = 6.0$, $J = 6.3$ Hz, 1H), 2.09 (m, 2H), 1.62 (m, 1H), 1.47 (m, 1H), 1.44 (m, 2H), 1.17 (d, $J = 6.0$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ 159.4, 138.9, 130.6, 129.5, 114.6, 114.0, 78.2, 74.9, 70.8, 55.3, 33.8, 32.4, 24.9, 15.7.

(2R,3R)-threo-3-Benzoyloxy-2-(4-methoxybenzyloxy)-7-octene (18a)—A solution of **17** (316.5 mg, 1.2 mmol) and dry pyridine (290 μL , 3.6 mmol) in dry DCM (2 mL) at 0°C was treated with benzoyl chloride (280 μL , 2.4 mmol) in dry DCM (1 mL). After stirring at RT for 12 h the mixture was diluted with DCM (10 mL), washed with 1 M HCl (10 mL), saturated aqueous NaHCO_3 solution (10 mL), dried over Na_2SO_4 , and concentrated under reduced pressure. Column chromatography of the residue (silica gel, 9/1 v/v hexane/ethyl acetate elution, $R_f = 0.37$) afforded **(2R,3R)-threo-18a** (386 mg, 1.05 mmol, 87% yield) as a colorless oil. ^1H NMR (400 MHz, CDCl_3) δ 8.09 (d, $J = 7.6$ Hz, 2H), 7.59 (m, 1H), 7.47 (m, 2H), 7.26 (d, $J = 8.5$ Hz, 2H), 6.86 (d, $J = 8.5$ Hz, 2H), 5.79 (ddt, $J = 17.0$ Hz, $J = 10.3$ Hz, $J = 6.7$ Hz, 1H), 5.25 (m, 1H), 5.01 (d, $J = 17.1$ Hz, 1H), 4.96 (d, $J = 10.3$ Hz, 1H), 4.62 (d, $J = 11.5$ Hz, 1H), 4.49 (d, $J = 11.5$ Hz, 1H), 3.81 (s, 3H), 3.74 (dq, $J = 4.9$ Hz, $J = 6.3$ Hz, 1H), 2.10 (m, 2H), 1.77 (m, 2H), 1.46 (m, 2H), 1.23 (d, $J = 6.4$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ 166.4, 159.3, 138.5, 133.0, 130.7, 129.8, 129.4, 129.0, 128.5, 114.9, 113.9, 76.1, 74.6, 70.9, 55.4, 33.7, 29.0, 25.0, 15.5.

(2R,3S)-erythro-3-Benzoyloxy-2-(4-methoxybenzyloxy)-7-octene (18b)—Under argon atmosphere a solution of **17** (294.5 mg, 1.11 mmol), triphenylphosphine (668.8 mg, 2.55 mmol), and benzoic acid (300.4 mg, 2.46 μmol) in dry THF (6 mL) at 0°C was treated with 440 μL diisopropyl azodicarboxylate (DIAD, 440 μL , 2.23 mmol). After stirring at RT for 4 h the solvent was removed under reduced pressure. Column chromatography of the residue (silica gel, 9/1 v/v hexane/ethyl acetate elution, $R_f = 0.37$) afforded **(2R,3S)-erythro-18b** (223.5 mg, 606.6 μmol , 55% yield) as a colorless oil. ^1H NMR (400 MHz, CDCl_3) δ 8.09 (d, $J = 7.6$ Hz, 2H), 7.59 (m, 1H), 7.47 (m, 2H), 7.27 (d, $J = 8.5$ Hz, 2H), 6.85 (d, $J = 8.5$ Hz, 2H), 5.80 (ddt, $J = 17.0$ Hz, $J = 10.2$ Hz, $J = 6.7$ Hz, 1H), 5.28 (m, 1H), 5.03 (d, $J = 17.1$ Hz, 1H), 4.98 (d, $J = 10.3$ Hz, 1H), 4.59 (d, $J = 11.5$ Hz, 1H), 4.54 (d, $J = 11.5$ Hz, 1H), 3.79 (s, 3H), 3.74 (dq, $J = 3.8$ Hz, $J = 6.3$ Hz, 1H), 2.11 (m, 2H), 1.80 (m, 2H), 1.52 (m, 2H), 1.28 (d, $J = 6.4$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ 166.3, 159.2, 138.5, 132.9, 130.68, 130.74, 129.8, 129.4, 128.5, 114.9, 113.9, 76.2, 75.4, 70.8, 55.3, 33.7, 29.3, 25.0, 16.0.

(7R,8R,2E)-threo-Ethyl 7-Benzoyloxy-8-(4-methoxybenzyloxy)-2-nonenoate (19a)—Under argon atmosphere a solution of **18a** (368.5 mg, 1 mmol) and ethyl acrylate (545 μL , 5 mmol) in DCM (30 mL) was treated with Grubbs-II catalyst (50 mg, 59 μmol) and stirred at 40°C for 9 h. The solution was concentrated under reduced pressure and the residue purified by chromatography (silica gel, 4/1 v/v hexane/ethyl acetate elution, $R_f = 0.40$) to afford **19a** (335.7 mg, 762 μmol , 76% yield) as a green oil. ^1H NMR (400 MHz, CDCl_3) δ 8.07 (d, $J = 7.2$ Hz, 2H), 7.57 (m, 1H), 7.47 (m, 2H), 7.25 (d, $J = 8.6$ Hz, 2H), 6.93

(dt, $J = 15.6, J = 7.0$ Hz, 1H), 6.86 (d, $J = 8.6$ Hz, 2H), 5.82 (d, $J = 15.6$ Hz, 1H), 5.24 (m, 1H), 4.62 (d, $J = 11.5$ Hz, 1H), 4.48 (d, $J = 11.5$ Hz, 1H), 4.19 (q, $J = 7.0$ Hz, 2H), 3.81 (s, 3H), 3.73 (dq, $J = 4.6, J = 6.4$ Hz, 1H), 2.23 (m, 2H), 1.78 (m, 2H), 1.52 (m, 2H), 1.30 (t, $J = 6.9$ Hz, 3H), 1.23 (d, $J = 6.4$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ 166.7, 166.4, 159.3, 148.6, 133.1, 130.7, 130.4, 129.8, 129.5, 128.5, 121.8, 113.9, 75.8, 74.4, 70.9, 60.3, 55.4, 32.1, 29.1, 24.2, 15.4, 14.4.

(7S,8R,2E)-erythro-Ethyl 7-Benzoyloxy-8-(4-methoxybenzyloxy)-2-nonenoate

(19b)—Under argon atmosphere a solution of **18b** (223.5 mg, 607 μmol) and ethyl acrylate (330 μL , 3 mmol) in DCM (20 mL) was treated with Grubbs-II catalyst (30 mg, 35 μmol) and stirred at 40 °C for 9 h. The solution was concentrated under reduced pressure and the residue purified by chromatography (silica gel, 4/1 v/v hexane/ethyl acetate elution, $R_f = 0.40$) to afford **19b** (218.2 mg, 495 μmol , 81% yield) as a brownish oil. ^1H NMR (400 MHz, CDCl_3) δ 8.07 (d, $J = 7.2$ Hz, 2H), 7.59 (m, 1H), 7.47 (m, 2H), 7.26 (d, $J = 8.5$ Hz, 2H), 6.94 (dt, $J = 15.6, J = 6.9$ Hz, 1H), 6.84 (d, $J = 8.5$ Hz, 2H), 5.82 (d, $J = 15.6$ Hz, 1H), 5.24 (dt, $J = 9.1$ Hz, $J = 3.9$ Hz, 1H), 4.55 (s, 2H), 4.19 (q, $J = 7.1$ Hz, 2H), 3.79 (s, 3H), 3.72 (dq, $J = 3.9$ Hz, $J = 6.4$ Hz, 1H), 2.25 (m, 2H), 1.79 (m, 2H), 1.56 (m, 2H), 1.29 (t, $J = 7.0$ Hz, 3H), 1.26 (d, $J = 6.4$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ 166.7, 166.2, 159.2, 148.5, 133.0, 130.6, 130.5, 129.7, 129.4, 128.5, 121.8, 113.8, 76.0, 75.3, 70.8, 60.2, 55.3, 32.0, 29.3, 24.2, 16.0, 14.4.

(7R,8R,2E)-threo-Ethyl 7-Benzoyloxy-8-hydroxy-2-nonenoate (20a)—A solution

of **19a** (330 mg, 750 μmol) in DCM (5 mL) was treated with water (260 μL) and 2,3-dichloro-5,6-dicyano-1,4-benzoquinone (187 mg, 825 μmol). After stirring at RT for 1 h the mixture was quenched with water (1 mL) and the aqueous phase extracted with DCM (5×1 mL). The combined organic phase was dried over Na_2SO_4 and concentrated under reduced pressure. Column chromatography (silica gel, 2/1 v/v hexane/ethyl acetate elution, $R_f = 0.42$) afforded **20a** (168.7 mg, 527 μmol , 70% yield) as a yellowish oil.

^1H NMR (400 MHz, CDCl_3) δ 8.05 (m, 2H); 7.57 (m, 1H), 7.45 (m, 2H), 6.91 (dt, $J = 15.7, J = 7.0$ Hz, 1H), 5.80 (d, $J = 15.7$ Hz, 1H), 5.05 (dt, $J = 7.8$ Hz, $J = 5.0$ Hz, 1H), 4.16 (q, $J = 7.2$ Hz, 2H), 3.94 (dq, $J = 4.8, J = 6.4$ Hz, 1H), 2.23 (m, 2H), 1.77 (m, 2H), 1.55 (m, 2H), 1.26 (t, $J = 7.0$ Hz, 3H), 1.23 (d, $J = 6.5$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ 166.7, 166.6, 148.4, 133.3, 130.1, 129.8, 128.6, 121.9, 77.9, 69.0, 60.3, 32.0, 30.2, 24.1, 19.6, 14.4; HRMS (ESI-TOF) m/z ($\text{M} + \text{NH}_4$) $^+$ calcd for $\text{C}_{18}\text{H}_{28}\text{NO}_5$ 338.1962, found 338.1979.

(7S,8R,2E)-erythro-Ethyl 7-Benzoyloxy-8-hydroxy-2-nonenoate (20b)—A

solution of **19b** (218.2 mg, 495 μmol) in DCM (5 mL) was treated with water (260 μL) and 2,3-dichloro-5,6-dicyano-1,4-benzoquinone (124 mg, 545 μmol). After being stirred at RT for 1 h, the mixture was quenched with 1 mL water, and the aqueous phase was extracted with DCM (5×1 mL). The combined organic phase was dried over Na_2SO_4 and concentrated under reduced pressure. Column chromatography (silica gel, 2/1 v/v hexane/ethyl acetate elution, $R_f = 0.42$) afforded **20b** (113.5 mg, 354 μmol , 72% yield) as a yellowish oil. ^1H NMR (400 MHz, CDCl_3) δ 8.05 (m, 2H), 7.57 (m, 1H), 7.45 (m, 2H), 6.91 (dt, $J = 15.7, J = 6.9$ Hz, 1H), 5.80 (d, $J = 15.7$ Hz, 1H), 5.11 (dt, $J = 9.3$ Hz, $J = 3.7$ Hz, 1H), 4.15 (q, $J = 7.2$ Hz, 2H), 4.00 (dq, $J = 3.8, J = 6.4$ Hz, 1H), 2.24 (m, 2H), 1.76 (m, 2H), 1.57

(m, 2H), 1.26 (t, $J = 7.1$ Hz, 3H), 1.23 (d, $J = 6.3$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ 166.7, 166.7, 148.4, 133.2, 130.1, 129.7, 128.5, 121.9, 78.2, 69.3, 60.2, 31.9, 29.2, 24.2, 18.3, 14.3; HRMS (ESI-TOF) m/z ($\text{M} + \text{NH}_4$) $^+$ calcd for $\text{C}_{18}\text{H}_{28}\text{NO}_5$ 338.1962, found 338.1977.

O-(2,4-Di-O-benzoyl-3,6-dideoxy- α -L-arabino-hexopyranosyl)

Trichloroacetimidate (21)—A solution of 2,4-di-*O*-benzoyl-ascarylose (53.5 mg, 150 μmol) in DCM (1 mL) was treated with trichloroacetonitrile (32 μL , 320 μmol) and 1,8-diazabicyclo[5.4.0]-undec-7-ene (5 μL , 33.4 μmol). After being stirred for 30 min, the yellowish solution was concentrated under reduced pressure. Column chromatography of the residue (silica gel, 4/1 v/v hexane/ethyl acetate elution) afforded **20** (61.2 mg, 122.4 μmol , 81% yield) as a colorless oil that was directly used for the next steps.

(7R,8R,2E)-threo-Ethyl 7-Benzoyloxy-8-[(2,4-di-O-benzoyl-3,6-dideoxy- α -L-arabino-hexopyranosyl)oxy]-2-nonenoate (22a)

—A solution of **20a** (14.7 mg, 45.9 μmol) and **21** (15.3 mg, 30.6 μmol) in dry DCM (1 mL) at 0 °C was treated with trimethylsilyl triflate (5 μL) and stirred for 3 h. The reaction was quenched with sat. NaHCO_3 solution (100 μL), dried over Na_2SO_4 , and concentrated under reduced pressure. Column chromatography (silica gel, 4/1 v/v hexane/ethyl acetate elution, $R_f = 0.24$) afforded **22a** (13.8 mg, 20.9 μmol , 68% yield) as a colorless oil. ^1H NMR (400 MHz, CDCl_3) δ 8.12 (m, 2H), 8.10 (m, 2H), 7.80 (d, $J = 7.6$ Hz, 2H), 7.58 (m, 3H), 7.47 (m, 2H), 7.46 (m, 2H), 7.42 (m, 2H), 6.94 (dt, $J = 15.7$, $J = 7.0$ Hz, 1H), 5.83 (d, $J = 15.6$ Hz, 1H), 5.31 (m, 1H), 5.15 (s.br, 1H), 5.10 (ddd, $J = 11.0$, $J = 9.8$ Hz, $J = 4.7$ Hz, 1H), 4.98 (s, 1H), 4.16 (q, $J = 7.1$ Hz, 2H), 4.08 (m, 1H), 4.01 (dq, $J = 9.7$ Hz, $J = 6.3$ Hz, 1H), 2.41 (dt, $J = 13.7$ Hz, $J = 3.8$ Hz, 1H), 2.28 (m, 2H), 2.19 (m, 1H), 1.80 (m, 2H), 1.61 (m, 2H), 1.29 (d, $J = 6.5$ Hz, 3H), 1.26 (t, $J = 7.0$ Hz, 3H), 1.03 (d, $J = 6.2$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ 166.6, 166.1, 165.9, 165.7, 148.3, 133.4, 133.2, 133.2, 130.4, 130.3, 130.0, 129.91, 129.88, 129.7, 128.6, 128.5, 122.0, 93.3, 75.9, 72.5, 71.1, 70.4, 67.3, 60.3, 32.0, 29.9, 29.8, 24.0, 17.8, 15.0, 14.4; HRMS (ESI-TOF) m/z ($\text{M} + \text{NH}_4$) $^+$ calcd for $\text{C}_{38}\text{H}_{46}\text{NO}_{10}$ 676.3116, found 676.3133.

(7S,8R,2E)-erythro-Ethyl 7-Benzoyloxy-8-[(2,4-di-O-benzoyl-3,6-dideoxy- α -L-arabino-hexopyranosyl)oxy]-2-nonenoate (22b)

—A solution of **20b** (14.7 mg, 45.9 μmol) and **21** (15.3 mg, 30.6 μmol) in dry DCM (1 mL) at 0 °C was treated with trimethylsilyl triflate (5 μL) and stirred for 3 h. The reaction was quenched with sat. NaHCO_3 solution (100 μL), dried over Na_2SO_4 and concentrated under reduced pressure. The residue was chromatographed (silica gel, 4/1 v/v hexane/ethyl acetate elution, $R_f = 0.26$) to afford **22b** (12.9 mg, 19.5 μmol , 64% yield) as a colorless oil. ^1H NMR (400 MHz, CDCl_3) δ 8.12 (m, 2H), 8.09 (m, 2H), 7.99 (d, $J = 7.8$ Hz, 2H), 7.58 (m, 3H), 7.46 (m, 6H), 6.96 (dt, $J = 15.8$, $J = 7.2$ Hz, 1H), 5.85 (d, $J = 15.5$ Hz, 1H), 5.22 (dt, $J = 15.2$ Hz, $J = 4.0$ Hz, 1H), 5.15 (s.br, 1H), 5.12 (m, 1H), 4.94 (s, 1H), 4.16 (q, $J = 7.1$ Hz, 2H), 4.16 (dq, $J = 3.8$ Hz, $J = 6.4$ Hz, 1H), 4.13 (m, 1H), 2.41 (dt, $J = 13.2$ Hz, $J = 3.8$ Hz, 1H), 2.30 (m, 2H), 2.19 (ddd, $J = 13.0$ Hz, $J = 11.7$ Hz, $J = 2.8$ Hz, 1H), 1.93 (m, 1H), 1.82 (m, 1H), 1.64 (m, 2H), 1.29 (d, $J = 6.5$ Hz, 3H), 1.26 (t, $J = 7.1$ Hz, 3H), 1.03 (d, $J = 6.2$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ 166.7, 166.2, 165.9, 165.7, 148.4, 133.4, 133.3, 133.2, 130.4, 130.1,

130.0, 129.8, 128.6, 128.5, 122.0, 93.6, 76.6, 72.5, 71.0, 70.6, 67.2, 60.3, 32.1, 29.8, 28.6, 24.4, 17.7, 14.8, 14.4; HRMS (ESI-TOF) m/z ($M + NH_4$)⁺ calcd for C₃₈H₄₆NO₁₀ 676.3116, found 676.3129.

(7R,8R,2E)-threo-8-[(3',6'-Dideoxy- α -L-arabino-hexopyranosyl)-oxy]-7-hydroxy-2-nonenic Acid ((7R,8R)-threo-10a)—A solution of **22a** (13.8 mg, 20.9 μ mol) in methanol (2 mL) was treated with LiOH monohydrate (7.0 mg, 167 μ mol) in H₂O (100 μ L). After stirring for 12 h the reaction mixture was acidified with acetic acid and concentrated under reduced pressure. The product was isolated by a combination of column chromatography on silica gel using a mixture of 15% methanol in dichloromethane with 0.1% acetic acid and solid phase extraction on reverse phase C18 using increasing concentrations of methanol in water as eluent to afford *threo*-**10a** (2.4 mg, 7.5 μ mol, 36% yield) along with the intramolecular cyclization product **23a** (2.5 mg, 7.9 μ mol, 39% yield).

(7R,8R)-threo-10a—¹H NMR (400 MHz, CD₃OD) δ 6.95 (dt, $J = 15.6$ Hz, $J = 7.0$ Hz, 1H), 5.82 (d, $J = 15.6$ Hz, 1H), 4.65 (s, 1H), 3.75 (s.br, 1H), 3.74 (dq, $J = 3.9$ Hz, $J = 6.1$ Hz, 1H), 3.64 (dq, $J = 9.3$ Hz, $J = 6.2$ Hz, 1H), 3.53 (m, 1H), 3.52 (m, 1H), 2.27 (m, 2H), 1.95 (dt, $J = 13.1$ Hz, $J = 3.8$ Hz, 1H), 1.80 (ddd, $J = 13.1$ Hz, $J = 11.0$ Hz, $J = 3.0$ Hz, 1H), 1.70 (m, 1H), 1.54 (m, 3H), 1.22 (d, $J = 6.2$ Hz, 3H), 1.14 (d, $J = 6.2$ Hz, 3H); ¹³C NMR (100 MHz, CD₃OD) δ 170.6, 150.5, 123.2, 97.8, 75.3, 74.9, 71.4, 69.9, 68.3, 35.9, 33.1, 33.0, 25.9, 18.1, 14.7; HRMS (ESI-TOF) m/z ($M - H$)⁻ calcd for C₁₅H₂₅O₇ 317.1606, found 317.1618.

2-((6R)-6-((R)-1-[(3,6-Dideoxy- α -L-arabino-hexopyranosyl)oxy]-ethyl)tetrahydro-2H-pyran-2-yl)acetic Acid (23a)—¹H NMR (400 MHz, CD₃OD) δ 4.66 (s, 1H), 3.76 (s.br, 1H), 3.65 (m, 1H), 3.74 (m, 1H), 3.65 (m, 1H), 3.52 (m, 2H), 2.50 (ddd, $J = 15.0$ Hz, $J = 7.2$ Hz, $J = 2.6$ Hz, 1H), 2.41 (ddd, $J = 15.0$ Hz, $J = 5.6$ Hz, $J = 2.7$ Hz, 1H), 1.95 (dt, $J = 13.1$ Hz, $J = 3.8$ Hz, 1H), 1.81 (ddd, $J = 13.1$ Hz, $J = 11.3$ Hz, $J = 3.0$ Hz, 1H), 1.58 (m, 3H), 1.51 (m, 2H), 1.46 (m, 1H), 1.22 (d, $J = 6.2$ Hz, 3H), 1.14 (d, $J = 6.1$ Hz, 3H); ¹³C NMR (100 MHz, CD₃OD) δ 175.6, 97.8, 79.32/79.37, 75.33/75.38, 75.1, 71.4, 69.9, 68.4, 40.4, 35.9, 35.0, 33.4, 22.83/22.87, 18.1, 14.7; HRMS (ESI-TOF) m/z ($M - H$)⁻ calcd for C₁₅H₂₅O₇ 317.1606, found 317.1614.

(7S,8R,2E)-erythro-8-[(3',6'-Dideoxy- α -L-arabino-hexopyranosyl)-oxy]-7-hydroxy-2-nonenic Acid ((7S,8R)-erythro-10b)—A solution of **22b** (12.9 mg, 19.5 μ mol) in methanol (2 mL) was treated with LiOH monohydrate (6.6 mg, 156 μ mol) in H₂O (100 μ L). After being stirred for 12 h, the reaction mixture was acidified with acetic acid and concentrated under reduced pressure. The product was isolated by a combination of column chromatography on silica gel using a mixture of 15% methanol in dichloromethane with 0.1% acetic acid and solid phase extraction on reverse phase C18 using increasing concentrations of methanol in water as eluent to afford *erythro*-**10b** (2.9 mg, 9.1 μ mol, 46% yield) along with the intramolecular cyclization product **23b** (2.6 mg, 8.2 μ mol, 42% yield).

(7S,8R)-erythro-10b—¹H NMR (400 MHz, CD₃OD) δ 6.96 (dt, *J* = 15.5 Hz, *J* = 7.1 Hz, 1H), 5.82 (d, *J* = 15.5 Hz, 1H), 4.66 (s, 1H), 3.75 (s.br, 1H), 3.66 (m, 1H), 3.63 (dq, *J* = 9.5 Hz, *J* = 6.2 Hz, 1H), 3.53 (m, 1H), 3.52 (m, 1H), 2.27 (m, 2H), 1.95 (dt, *J* = 13.3 Hz, *J* = 3.7 Hz, 1H), 1.80 (ddd, *J* = 13.1 Hz, *J* = 11.1 Hz, *J* = 3.0 Hz, 1H), 1.65 (m, 2H), 1.54 (m, 1H), 1.43 (m, 1H), 1.22 (d, *J* = 6.2 Hz, 3H), 1.15 (d, *J* = 6.2 Hz, 3H); ¹³C NMR (100 MHz, CD₃OD) δ 170.4, 150.6, 123.0, 97.6, 75.6, 75.2, 71.3, 69.8, 68.4, 35.9, 33.6, 33.0, 25.6, 18.1, 14.2; HRMS (ESI-TOF) *m/z* (M - H)⁻ calcd for C₁₅H₂₅O₇ 317.1606, found 317.1615.

2-((6S)-6-((R)-1-[(3,6-Dideoxy-α-L-arabino-hexopyranosyl)oxy]-ethyl)tetrahydro-2H-pyran-2-yl)acetic Acid (23b)—¹H NMR (400 MHz, CD₃OD) δ 4.66 (s, 1H), 3.75 (s.br, 1H), 3.66 (m, 2H), 3.64 (m, 1H), 3.52 (m, 2H), 2.50 (ddd, *J* = 15.0 Hz, *J* = 7.1 Hz, *J* = 2.1 Hz, 1H), 2.41 (ddd, *J* = 15.1 Hz, *J* = 5.4 Hz, *J* = 2.2 Hz, 1H), 1.95 (1H, dt, *J* = 13.1 Hz, *J* = 3.9 Hz, 1H), 1.80 (ddd, *J* = 13.1 Hz, *J* = 11.3 Hz, *J* = 3.1 Hz, 1H), 1.59 (m, 1H), 1.57 (m, 2H), 1.44 (m, 1H), 1.41 (m, 2H), 1.22 (d, *J* = 6.2 Hz, 3H), 1.15 (d, *J* = 6.2 Hz, 3H); ¹³C NMR (100 MHz, CD₃OD) δ 175.7, 97.6, 79.3, 75.57/75.61, 75.4, 71.3, 69.8, 68.4, 40.4, 35.9, 34.9, 34.1, 22.6, 18.1, 14.1; HRMS (ESI-TOF) *m/z* (M - H)⁻ calcd for C₁₅H₂₅O₇ 317.1606, found 317.1622.

(7R,8R)-threo- or (7S,8R)-erythro-8-[(3',6'-Dideoxy-α-L-arabino-hexopyranosyl)oxy]-7-hydroxynonanoic Acid (threo-11a or erythro-11b)—Aliquots (10 μg) of synthetic (*7R,8R*)-*threo*-asc-7OH- C₉ (**10a**) or (*7S,8R*)-*erythro*-asc-7OH- C₉ (**10b**) in methanol (500 μL) were treated with 10% palladium on carbon (10 mg) and hydrogenated under atmospheric pressure for 1 h. The mixture was filtered over a small patch of silica, concentrated to dryness, and the resulting (*7R,8R*)-*threo*-asc-7OH-C₉ (**11a**) or (*7S,8R*)-*erythro*-asc-7OH-C₉ (**11b**) submitted to TMS derivatization for chemical correlation with the natural product isolated from *C. nigoni*.

(7S)-7-tert-Butyldimethylsilyloxy-6-hydroxy-1-octene (25)—Under argon atmosphere, a solution of 4-pentenylmagnesium bromide (1.68 mmol) in THF (2 mL) prepared from 5-bromo-1-pentene (250 mg, 1.68 mmol) and magnesium (45 mg, 1.88 mmol) was added slowly to a mixture of copper(I)iodide (32 mg, 168 μmol) and (*S*)-3-*tert*-butyldimethylsilyloxy-1,2-epoxypropane (**24**) (210 mg, 1.12 mmol) in THF (2 mL) at 0 °C. After being stirred at 0 °C for 3 h, the solution was quenched with saturated ammonium chloride solution and stirred for 1 h, and the aqueous phase extracted with diethyl ether. The organic phase was washed with brine and concentrated under reduced pressure, and the residue was chromatographed (silica gel, DCM elution, *R_f* = 0.43) to afford **25** (273 mg, 1.05 mmol, 94% yield) as a colorless oil. ¹H NMR (400 MHz, CDCl₃) δ 5.76 (ddt, *J* = 17.0 Hz, *J* = 10.3 Hz, *J* = 6.7 Hz, 1H), 4.95 (ddt, *J* = 17.1 Hz, *J* = 2.0 Hz, *J* = 1.5 Hz, 1H), 4.89 (ddt, *J* = 10.2 Hz, *J* = 2.0 Hz, *J* = 1.5 Hz, 1H), 3.58 (m, 2H), 3.36 (m, 1H), 2.02 (dt, *J* = 6.7 Hz, *J* = 7.1 Hz, 2H), 1.26–1.48 (m, 6H), 0.87 (s, 9H), 0.03 (s, 6H); ¹³C NMR (100 MHz, CDCl₃) δ 138.9, 114.4, 71.8, 67.4, 33.8, 32.7, 29.1, 26.0, 25.1, 18.3, -5.3, -5.4.

(2E,8S)-Ethyl 9-tert-Butyldimethylsilyloxy-8-hydroxy-2-nonenoate (26)—Under argon atmosphere, a solution of **25** (258.5 mg, 1 mmol) and ethyl acrylate (545 μL, 5 mmol) in DCM (30 mL) was treated with Grubbs second generation catalyst (50 mg, 58.9 μmol)

and stirred at 40 °C for 9 h. The solution was concentrated under reduced pressure, and the residue was chromatographed (silica gel, 4/1 v/v hexane/ethyl acetate elution, $R_f = 0.51$) to afford **26** (268 mg, 810 μmol , 81% yield) as a yellowish oil. ^1H NMR (400 MHz, CDCl_3) δ 6.95 (dt, $J = 15.7$ Hz, $J = 6.9$ Hz, 1H), 5.80 (d, $J = 15.7$ Hz, 1H), 4.17 (q, $J = 7.1$ Hz, 2H), 3.62 (m, 1H), 3.60 (dd, $J = 10.6$ Hz, $J = 3.3$ Hz, 1H), 3.37 (dd, $J = 10.6$ Hz, $J = 8.3$ Hz, 1H), 2.20 (m, 2H), 1.33–1.54 (m, 6H), 1.27 (t, $J = 7.1$ Hz, 3H), 0.89 (s, 9H), 0.06 (s, 6H); ^{13}C NMR (100 MHz, CDCl_3) δ 166.9, 149.2, 121.6, 71.8, 67.4, 60.3, 32.7, 32.2, 28.3, 26.0, 25.3, 18.4, 14.4, –5.2, –5.3; HRMS (ESI-TOF) m/z ($\text{M} + \text{NH}_4$) $^+$ calcd for $\text{C}_{17}\text{H}_{38}\text{NO}_4\text{Si}$ 348.2565, found 348.2572.

(2E,8S)-Ethyl 8-[(2,4-Di-O-benzoyl-3,6-dideoxy- α -L-arabinohexopyranosyl)oxy]-9-tert-butyltrimethylsilyloxy-2-nonenoate (27)—

Under argon atmosphere, a solution of **26** (30.3 mg, 91.8 μmol) and **21** (30.6 mg, 61.2 μmol) in dry DCM (1 mL) at 0 °C was treated with trimethylsilyl triflate (5 μL) and stirred for 3 h. The reaction was quenched by addition of saturated aqueous NaHCO_3 solution (100 μL), dried over Na_2SO_4 , and concentrated under reduced pressure. The product was isolated by column chromatography (silica gel, 4/1 v/v hexane/ethyl acetate elution, $R_f = 0.50$) to afford **27** (21.7 mg, 32.4 μmol , 53% yield) as a colorless oil. ^1H NMR (400 MHz, CDCl_3) δ 8.10 (d, $J = 8.0$ Hz, 2H), 8.03 (d, $J = 7.9$ Hz, 2H), 7.58 (m, 2H), 7.46 (m, 4H), 6.99 (dt, $J = 15.6$ Hz, $J = 7.0$ Hz, 1H), 5.85 (d, $J = 15.6$ Hz, 1H), 5.22 (s.br, 1H), 5.15 (ddd, $J = 11.4$ Hz, $J = 10.3$ Hz, $J = 4.4$ Hz, 1H), 5.10 (s, 1H), 4.16 (q, $J = 7.1$ Hz, 2H), 4.15 (m, 1H), 3.78 (m, 1H), 3.65 (dd, $J = 10.5$ Hz, $J = 5.9$ Hz, 1H), 3.61 (dd, $J = 10.6$ Hz, $J = 5.0$ Hz, 1H), 2.43 (dt, $J = 13.3$ Hz, $J = 3.9$ Hz, 1H), 2.26 (m, 2H), 2.19 (ddd, $J = 13.5$ Hz, $J = 11.6$ Hz, $J = 3.0$ Hz, 1H), 1.40–1.70 (m, 6H), 1.28 (d, $J = 6.9$ Hz, 3H), 1.26 (t, $J = 7.1$ Hz, 3H), 0.85 (s, 9H), 0.06 (s, 3H), 0.05 (s, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ 166.8, 165.8, 165.7, 149.1, 133.3, 130.2, 130.1, 130.0, 129.8, 128.6, 128.5, 121.7, 96.4, 78.4, 71.0, 70.8, 67.3, 65.5, 60.3, 32.3, 31.8, 29.8, 28.2, 25.9, 25.8, 25.3, 18.3, 18.0, 14.4, –5.3, –5.4; HRMS (ESI-TOF) m/z ($\text{M} + \text{NH}_4$) $^+$ calcd for $\text{C}_{37}\text{H}_{56}\text{NO}_9\text{Si}$ 686.3719, found 686.3735.

(2E,8S)-Ethyl 8-[(2,4-Di-O-benzoyl-3,6-dideoxy- α -L-arabinohexopyranosyl)oxy]-9-hydroxy-2-nonenoate (28)—

Under argon atmosphere a solution of **27** (21.7 mg, 32.4 μmol) in dry THF (1 mL) was treated with 1 M tetrabutylammonium fluoride (50 μL , 50 μmol) in dry THF (1 mL) and stirred at RT for 3 h. The solution was concentrated under reduced pressure, and the residue was separated by column chromatography (silica gel, 2/1 v/v hexane/ethyl acetate elution) to afford **28** (14.0 mg 25.2 μmol , 78% yield). ^1H NMR (400 MHz, CDCl_3) δ 8.10 (d, $J = 7.8$ Hz, 2H), 8.05 (d, $J = 7.7$ Hz, 2H), 7.59 (m, 2H), 7.46 (m, 4H), 6.97 (dt, $J = 15.6$ Hz, $J = 6.9$ Hz, 1H), 5.83 (d, $J = 15.7$ Hz, 1H), 5.20 (s.br, 1H), 5.19 (ddd, $J = 11.3$ Hz, $J = 10.5$ Hz, $J = 4.5$ Hz, 1H), 5.08 (s, 1H), 4.16 (q, $J = 7.1$ Hz, 2H), 4.13 (m, 1H), 3.82 (m, 1H), 3.76 (dd, $J = 12.0$ Hz, $J = 2.9$ Hz, 1H), 3.61 (dd, $J = 12.0$ Hz, $J = 5.6$ Hz, 1H), 2.44 (dt, $J = 13.5$ Hz, $J = 3.9$ Hz, 1H), 2.25 (m, 2H), 2.21 (ddd, $J = 13.5$ Hz, $J = 11.6$ Hz, $J = 3.0$ Hz, 1H), 1.40–1.75 (m, 6H), 1.29 (d, $J = 6.8$ Hz, 3H), 1.26 (t, $J = 7.1$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ 166.8, 166.1, 165.8, 148.9, 133.5, 133.4, 130.1, 130.0, 129.82, 129.79, 128.6, 121.7, 95.9, 79.3, 71.2, 70.6, 67.5, 64.5, 60.3, 32.2, 31.4, 29.8 28.2, 25.3, 18.0, 14.4; HRMS (ESI-TOF) m/z ($\text{M} + \text{NH}_4$) $^+$ calcd for $\text{C}_{31}\text{H}_{42}\text{NO}_9$ 572.2854, found 572.2871.

(2E,8S)-8-[(3,6-Dideoxy- α -L-arabino-hexopyranosyl)oxy]-9-hydroxy-2-nonenoic Acid (Asc-9OH- C9) (12)—A solution of **28** (14.0 mg, 25.3 μ mol) in methanol (2 mL) was treated with LiOH monohydrate (8.5 mg, 202.4 μ mol) in water (100 μ L) and stirred for 12 h. The mixture was acidified with acetic acid and concentrated to dryness under reduced pressure. The residue was separated by column chromatography (silica gel, 15% v/v methanol in dichloromethane with 0.1% acetic acid as eluent) and solid phase extraction (reverse phase C18, 10% stepwise increase of methanol in water as eluent) to afford **12** (2.0 mg, 6.3 μ mol, 25% yield), identical to the natural product from *C. nigoni*, along with its intramolecular cyclization product **28** (5.6 mg, 17.6 μ mol, 69% yield).

Asc-9OH- C9 (12)—Colorless oil, ^1H NMR (400 MHz, CD_3OD) δ 6.95 (dt, J = 15.6 Hz, J = 6.9 Hz, 1H), 5.81 (d, J = 15.6 Hz, 1H), 4.75 (s, 1H), 3.84 (s.br, 1H), 3.69 (m, 1H), 3.67 (m, 1H), 3.60 (dd, J = 11.7 Hz, J = 4.2 Hz, 1H), 3.53 (m, 1H), 3.50 (dd, J = 11.7 Hz, J = 5.6 Hz, 1H), 2.25 (m, 2H), 1.95 (dt, J = 13.0 Hz, J = 3.8 Hz, 1H), 1.78 (ddd, J = 13.0 Hz, J = 11.2 Hz, J = 3.1 Hz, 1H), 1.59 (m, 2H), 1.52 (m, 2H), 1.50 (m, 2H), 1.22 (d, J = 6.1 Hz, 3H); ^{13}C NMR (100 MHz, CD_3OD) δ 170.3, 150.7, 122.9, 99.8, 78.6, 71.4, 69.6, 68.4, 64.6, 35.9, 33.1, 32.8, 29.3, 26.3 18.1; HRMS (ESI-TOF) m/z ($\text{M} - \text{H}$) $^-$ calcd for $\text{C}_{15}\text{H}_{25}\text{O}_7$ 317.1606, found 317.1614.

(7S)-7-[(3,6-Dideoxy- α -L-arabino-hexopyranosyl)oxy]oxocan-2-yl)-acetic Acid (29)—Colorless oil. ^1H NMR (400 MHz, CD_3OD) δ 4.75 (s, 1H), 3.84 (s.br, 1H), 3.69 (m, 1H), 3.68 (m, 1H), 3.65 (m, 1H), 3.53 (m, 1H), 3.52 (m, 1H), 2.49 (dd, J = 15.2 Hz, J = 7.2 Hz, 1H), 2.40 (dd, J = 15.2 Hz, 5.6 Hz, 1H), 1.95 (dt, J = 13.0 Hz, J = 4.0 Hz, 1H), 1.78 (m, 1H), 1.59 (m, 1H), 1.57 (m, 2H), 1.40–1.60 (m, 4H), 1.22 (d, J = 6.2 Hz, 3H), ^{13}C NMR (100 MHz, CD_3OD) δ 175.6, 99.8, 79.29/79.32, 78.73/78.77, 71.37/71.40, 69.6, 68.4, 64.6, 40.4, 36.0, 34.9, 32.8, 26.8, 26.3, 18.2; HRMS (ESI-TOF) m/z ($\text{M} - \text{H}$) $^-$ calcd for $\text{C}_{15}\text{H}_{25}\text{O}_7$ 317.1606, found 317.1619.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

Financial support by the Max Planck Society (MPG), the Jena School of Microbial Communication (JSMC), and the Swiss National Science Foundation (SNSF) is gratefully acknowledged. This work was supported by grants from the NIH (R01DC016058) and startup funds from WPI to JS. We thank Dr. Michael Reichelt (Department for Biochemistry, MPICE, Jena) for acquisition of the MS/MS precursor ion screen and Kerstin Ploss (Department of Bioorganic Chemistry, MPICE, Jena) for technical assistance with the GC-EIMS measurements. All nematode strains were provided by the *Caenorhabditis* Genetics Center, which is funded by NIH Office of Research Infrastructure Programs (Grant P40 OD010440).

References

1. For recent reviews see: Butcher RA. Nat. Prod. Rep. 2017; 34:472. [PubMed: 28386618] Butcher RA. Nat. Chem. Biol. 2017; 13:577. [PubMed: 28514418] von Reuss SH, Schroeder FC. Nat. Prod. Rep. 2015; 32:994. [PubMed: 26059053] Schroeder FC. Chem. Biol. 2015; 22:7. [PubMed: 25484238] Chute CD, Srinivasan J. Semin. Cell Dev. Biol. 2014; 33:18. [PubMed: 24977334] Ludewig AH, Schroeder FC. WormBook. 2013; 18:1.

2. Jeong PY, Jung M, Yim YH, Kim H, Park M, Hong E, Lee W, Kim YH, Kim K, Paik YK. *Nature*. 2005; 433:541. [PubMed: 15690045]
3. Butcher RA, Fujita M, Schroeder FC, Clardy J. *Nat. Chem. Biol.* 2007; 3:420. [PubMed: 17558398]
4. Butcher RA, Ragains JR, Kim E, Clardy J. *Proc. Natl. Acad. Sci. U. S. A.* 2008; 105:14288. [PubMed: 18791072]
5. Butcher RA, Ragains JR, Clardy J. *Org. Lett.* 2009; 11:3100. [PubMed: 19545143]
6. Wharam B, Weldon L, Viney M. *BMC Evol. Biol.* 2017; 17:197. [PubMed: 28830356]
7. Aprison EZ, Ruvinsky I. *PLoS Genet.* 2015; 11:e1005729. [PubMed: 26645097]
8. Ludewig AH, Izrayelit Y, Park D, Malik RU, Zimmermann A, Mahanti P, Fox BW, Bethke A, Doering F, Riddle DL, Schroeder FC. *Proc. Natl. Acad. Sci. U. S. A.* 2013; 110:5522. [PubMed: 23509272]
9. Maures TJ, Booth LN, Benayoun BA, Izrayelit Y, Schroeder FC, Brunet A. *Science*. 2014; 343:541. [PubMed: 24292626]
10. Srinivasan J, Kaplan F, Ajredini R, Zachariah C, Alborn HT, Teal PE, Malik RU, Edison AS, Sternberg PW, Schroeder FC. *Nature*. 2008; 454:1115. [PubMed: 18650807]
11. Pungaliya C, Srinivasan J, Fox BW, Malik RU, Ludewig AH, Sternberg PW, Schroeder FC. *Proc. Natl. Acad. Sci. U. S. A.* 2009; 106:7708. [PubMed: 19346493]
12. Srinivasan J, von Reuss SH, Bose N, Zaslaver A, Mahanti P, Ho MC, O'Doherty OG, Edison AS, Sternberg PW, Schroeder FC. *PLoS Biol.* 2012; 10:e1001237. [PubMed: 22253572]
13. Zhang YK, Sanchez-Ayala MA, Sternberg PW, Srinivasan J, Schroeder FC. *Org. Lett.* 2017; 19:2837. [PubMed: 28513161]
14. Izrayelit Y, Srinivasan J, Campbell SL, Jo Y, von Reuss SH, Genoff MC, Sternberg PW, Schroeder FC. *ACS Chem. Biol.* 2012; 7:1321. [PubMed: 22662967]
15. Hollister KA, Conner ES, Zhang XX, Spell M, Bernard GM, Patel P, de Carvalho ACGV, Butcher RA, Ragains JR. *Bioorg. Med. Chem.* 2013; 21:5754. [PubMed: 23920482]
16. Dong CF, Dolce F, von Reuss SH. *Org. Biomol. Chem.* 2016; 14:7217. [PubMed: 27381649]
17. Aprison EZ, Ruvinsky I. *Curr. Biol.* 2017; 27:2589. [PubMed: 28844646]
18. Aprison EZ, Ruvinsky I. *Curr. Biol.* 2016; 26:2827. [PubMed: 27618262]
19. Butcher RA, Ragains JR, Li W, Ruvkun G, Clardy J, Mak HY. *Proc. Natl. Acad. Sci. U. S. A.* 2009; 106:1875. [PubMed: 19174521]
20. Joo HJ, Yim YH, Jeong PY, Jin YX, Lee JE, Kim H, Jeong SK, Chitwood DJ, Paik YK. *Biochem. J.* 2009; 422:61. [PubMed: 19496754]
21. Joo HJ, Kim KY, Yim YH, Jin YX, Kim H, Kim MY, Paik YK. *J. Biol. Chem.* 2010; 285:29319. [PubMed: 20610393]
22. von Reuss SH, Bose N, Srinivasan J, Yim JJ, Judkins JC, Sternberg PW, Schroeder FC. *J. Am. Chem. Soc.* 2012; 134:1817. [PubMed: 22239548]
23. Zhang X, Feng L, Chinta S, Singh P, Wang Y, Nunnery JK, Butcher RA. *Proc. Natl. Acad. Sci. U. S. A.* 2015; 112:3955. [PubMed: 25775534]
24. Panda O, Akagi AE, Artyukhin AB, Judkins JC, Le HH, Mahanti P, Cohen SM, Sternberg PW, Schroeder FC. *Angew. Chem., Int. Ed.* 2017; 56:4729.
25. Choe A, von Reuss SH, Kogan D, Gasser RB, Platzner EG, Schroeder FC, Sternberg PW. *Curr. Biol.* 2012; 22:772. [PubMed: 22503501]
26. Choe A, Chuman T, von Reuss SH, Dossey AT, Yim J, Ajredini R, Kolawa AA, Kaplan F, Alborn HT, Teal PE, Schroeder FC, Sternberg PW, Edison AS. *Proc. Natl. Acad. Sci. U. S. A.* 2012; 109:20949. [PubMed: 23213209]
27. Bose N, Ogawa A, von Reuss SH, Yim JJ, Sommer RJ, Schroeder FC. *Angew. Chem., Int. Ed.* 2012; 51:12438.
28. Noguez JH, Conner ES, Zhou Y, Ciche TA, Ragains JR, Butcher RA. *ACS Chem. Biol.* 2012; 7:961. [PubMed: 22444073]
29. Chaudhuri J, Bose N, Tandonnet S, Adams S, Zuco G, Kache V, Parihar M, von Reuss SH, Schroeder FC, PiresdaSilva A. *Sci. Rep.* 2016; 5:17676.

30. Bose N, Meyer JM, Yim JJ, Mayer MG, Markov GV, Ogawa A, Schroeder FC, Sommer RJ. *Curr. Biol.* 2014; 24:1536. [PubMed: 24980503]
31. Diaz SA, Brunet V, Lloyd-Jones GC, Spinner W, Wharam B, Viney M. *BMC Evol. Biol.* 2014; 14:46. [PubMed: 24618411]
32. Hsueh YP, Mahanti P, Schroeder FC, Sternberg PW. *Curr. Biol.* 2013; 23:83–86. [PubMed: 23246407]
33. Manosalva P, Manohar M, von Reuss SH, Chen S, Koch A, Kaplan F, Choe A, Micikas RJ, Wang X, Kogel KH, Sternberg PW, Williamson VM, Schroeder FC, Klessig DF. *Nat. Commun.* 2015; 6:7795. [PubMed: 26203561]
34. Zhao L, Zhang X, Wei Y, Zhou J, Zhang W, Qin P, Chinta S, Kong X, Liu Y, Yu H, Hu S, Zou Z, Butcher RA, Sun J. *Nat. Commun.* 2016; 7:12341. [PubMed: 27477780]
35. von Reuss SH, Dolke F, Dong CF. *Anal. Chem.* 2017; 89:10570. [PubMed: 28866881]
36. Kiontke KC, Félix MA, Ailion M, Rockman MV, Braendle C, Pénigault JB, Fitch DHA. *BMC Evol. Biol.* 2011; 11:339. [PubMed: 22103856]
37. Félix MA, Braendle C, Cutter AD. *PLoS One.* 2014; 9:e94723. [PubMed: 24727800]
38. Stiernagle, T. *WormBook. The C. elegans Research Community; 2006. Maintenance of C. elegans.* <http://www.wormbook.org>
39. Matsumori N, Kaneno D, Murata M, Nakamura H, Tachibana K. *J. Org. Chem.* 1999; 64:866. [PubMed: 11674159]
40. (a) Roush WR, Bennett CE, Roberts SE. *J. Org. Chem.* 2001; 66:6389. [PubMed: 11559191] (b) Zhao Z, Hong L, Liu H. *J. Am. Chem. Soc.* 2005; 127:7692. [PubMed: 15913355] (c) Hong L, Zhao Z, Melançon CE, Zhang H, Liu H. *J. Am. Chem. Soc.* 2008; 130:4954. [PubMed: 18345667] (d) Yadav JS, Swamy T, Reddy BVS, Ravinder V. *Tetrahedron Lett.* 2014; 55:4054.
41. Scholl M, Ding S, Lee CW, Grubbs RH. *Org. Lett.* 1999; 1:953. [PubMed: 10823227]
42. (a) Oikawa Y, Yoshioka T, Yonemitsu O. *Tetrahedron Lett.* 1982; 23:885.(b) Horita K, Yoshioka T, Tanaka T, Oikawa Y, Yonemitsu O. *Tetrahedron.* 1986; 42:3021.(c) Chandrasekhar S, Sumithra G, Yadav JS. *Tetrahedron Lett.* 1996; 37:1645.
43. (a) Schmidt RR, Michel J. *Angew. Chem.* 1980; 92:763.(b) Schmidt RR, Michel J. *Angew. Chem., Int. Ed. Engl.* 1980; 19:731.
44. Narayan A, Venkatachalam V, Durak O, Reilly DK, Bose N, Schroeder FC, Samuel AD, Srinivasan J, Sternberg PW. *Proc. Natl. Acad. Sci. U. S. A.* 2016; 113:E1392–401. [PubMed: 26903633]
45. Curti C, Zanardi F, Battistini L, Sartori A, Rassu G, Pinna L, Casiraghi G. *J. Org. Chem.* 2006; 71:8552. [PubMed: 17064033]




Figure 1. Ascaroside profiling of the *C. nigoni* exometabolome. (A) Total ion chromatogram (TIC) of the TMS-derivatized *C. nigoni* crude exometabolome; (B) extracted ion chromatogram (EIC) for the K1 fragment at m/z 130.1 $[C_6H_{14}OSi]^{++}$; and (C) HPLC-ESI-MS/MS precursor ion screen for m/z 73.1 $[C_3H_5O_2]^-$.




Figure 2.
Comparative GC-EIMS ascarioside profiling of 13 *Caenorhabditis* exometabolomes.







Figure 3. GC-EIMS spectra of TMS-derivatized (ω) and ($\omega - 2$) hydroxyacyl ascarosides from *C. nigoni*.



Scheme 1. Ascaroside Diversity Originating from (A) Side Chain Modification and Shortening via Peroxisomal β -Oxidation, (B) Subsequent Attachment of Additional Metabolic Units from Primary Metabolism, or (C) Specific (ω), ($\omega - 2$), or ($\omega - 3$)-Hydroxylation Downstream of Peroxisomal β -Oxidation




Scheme 2. EI-Induced Fragmentation of TMS-Derivatized Ascarosides



Scheme 3. Synthesis of (ω - 2)-Hydroxyacyl Ascarosides *threo-asc-7OH*- C9 (10a**) and *erythro-asc-7OH*- C9 (**10b**)**

Reagents and conditions: (i) 4-methoxybenzyl trichloroacetimidate, trimethylsilyl triflate, CH_2Cl_2 , 0 °C, 3 h, 62%; (ii) DIBAL-H, CH_2Cl_2 , -78 °C, 0.5 h, 100%; (iii): 4-pentenylmagnesium bromide, Et_2O , 0 °C, 1 h, 58%; (iv) benzoyl chloride, pyridine, CH_2Cl_2 , 0 °C, 12 h, 87% (**18a**); (v) benzoic acid, PPh_3 , DIAD, THF, 4 h, 55% (**18b**); (vi): ethyl acrylate, Grubbs second generation catalyst, CH_2Cl_2 , 40 °C, 9 h, 76% (**19a**) and 81% (**19b**); (vii): DDQ, $\text{CH}_2\text{Cl}_2/\text{H}_2\text{O}$, 1 h, 70% (**20a**) and 72% (**20b**); (viii): 2,4-di-*O*-benzoyl-ascarosyl-1-(2,2,2-trichloroacetimidate) (**21**), trimethylsilyl triflate, CH_2Cl_2 , 0 °C, 3 h, 68% (**22a**) and 64% (**22b**); (ix): lithium hydroxide, water, MeOH, 12 h, 36% (**10a**), 46% (**10b**), 39% (**23a**), and 42% (**23b**)



Scheme 4. Synthesis of the (ω)-Hydroxyacyl Ascaroside asc-9OH- C9 (12**)**

Reagents and conditions: (i) 4-pentenylmagnesium bromide, copper(I)iodide, THF, 0 °C, 3 h, 94%; (ii) ethyl acrylate, Grubb's second generation catalyst, CH₂Cl₂, 40 °C, 9 h, 81%; (iii) 2,4-di-*O*-benzoylascarosyl-1-(2,2,2-trichloroacetimidate) (**21**), trimethylsilyl triflate, CH₂Cl₂, 0 °C, 3 h, 53%; (iv) tetrabutylammonium fluoride, THF, 3 h, 78%; (v) lithium hydroxide, water, MeOH, 12 h, 25% (**12**) and 69% (**29**).

Table 1

NMR Data for (ω) and ($\omega - 2$)-Hydroxyacyl Ascarosides (10–12) Isolated from *C. nigoni* (400 MHz, CD₃OD)

position	threo-asc-7OH- C9 (10)			threo-asc-7OH-C9 (11)			asc-9OH- C9 (12)		
	δ_c^a	δ_H^b , mult (J, Hz)		δ_c^a	δ_H^b , mult (J, Hz)		δ_c^a	δ_H^b , mult (J, Hz)	
1	nd	C	–	nd	C	–	nd	C	–
2	124.8	CH	5.83 d (15.5)	37.5	CH ₂	2.21 t (7.6)	122.9	CH	5.81 d (15.6)
3	148.1	CH	6.86 dt (15.5, 7.0)	26.9	CH ₂	1.63 m	150.7	CH	6.95 dt (15.6, 6.9)
4	32.7	CH ₂	2.25 m	30.4	CH ₂	1.40 m	33.1	CH ₂	2.25 m
5	25.6	CH ₂	1.54–1.68 m	27.0	CH ₂	1.40 m	29.3	CH ₂	1.52 m
6	32.8	CH ₂	1.48–1.62 m	32.9	CH ₂	1.48–1.53 m	26.3	CH ₂	1.50 m
7	74.8	CH	3.53 m	75.2	CH	3.52 m	32.8	CH ₂	1.59 m
8	75.5	CH	3.74 dq (3.9, 6.3)	75.6	CH	3.73 dq (3.8, 6.2)	78.6	CH	3.69 m
9	14.5	CH ₃	1.14 d (6.3)	14.6	CH ₃	1.14 d (6.2)	64.6	CH ₂	3.50 dd (11.7, 5.6) 3.60 dd (11.7, 4.2)
1'	97.9	CH	4.65 s	97.8	CH	4.65 s	99.8	CH	4.75 s
2'	69.7	CH	3.76 s.br	69.8	CH	3.76 s.br	69.6	CH	3.84 s.br
3' _{ax}	35.8	CH ₂	1.95 dt (13.0, 3.8)	35.8	CH ₂	1.95 dt (13.0, 3.8)	35.9	CH ₂	1.95 dt (13.0, 3.8)
3' _{eq}			1.80 ddd (13.0, 11.4, 3.0)			1.81 ddd (13.0, 11.4, 3.0)			1.78 ddd (13.0, 11.2, 3.1)
4'	68.2	CH	3.52 ddd (11.3, 9.3, 4.3)	68.4	CH	3.51 ddd (11.4, 9.5, 4.4)	68.4	CH	3.53 m
5'	71.3	CH	3.64 dq (9.3, 6.3)	71.4	CH	3.66 dq (9.5, 6.3)	71.4	CH	3.67 m
6'	17.8	CH ₃	1.22 d (6.2)	17.8	CH ₃	1.22 d (6.2)	18.1	CH ₃	1.22 d (6.1)

^aFrom HSQC spectrum.

^bFrom ¹H NMR and *dgf*-COSY spectra.