

Overexpression of MTH1 and OGG1 proteins in ulcerative colitis-associated carcinogenesis

YOSHITERU KUMAGAE¹, MINAKO HIRAHASHI¹, KATSUMI TAKIZAWA¹,
HIDETAKA YAMAMOTO¹, MASAKI GUSHIMA², MOTOHIRO ESAKI³, TAKAYUKI MATSUMOTO⁴,
MASAFUMI NAKAMURA³, TAKANARI KITAZONO³ and YOSHINAO ODA¹

¹Department of Anatomic Pathology, Graduate School of Medical Sciences, Kyushu University, Fukuoka, Fukuoka 812-8582;

²Department of Medical Gastroenterology, Shimonoseki Hospital, Yamaguchi, Yamaguchi 750-8520; ³Department of Medicine and Clinical Science, Surgery and Oncology, Graduate School of Medical Sciences, Kyushu University, Fukuoka, Fukuoka 812-8582; ⁴Division of Gastroenterology, Department of Internal Medicine, School of Medicine,

Iwate Medical University, Morioka, Iwate 020-8505, Japan

Received December 8, 2016; Accepted September 5, 2017

DOI: 10.3892/ol.2018.8812

Abstract. Oxidative stress, demonstrated by an accumulation of 8-hydroxy-2'-deoxyguanosine (8-OHdG), results in DNA damage, which is normally repaired by base excision repair enzymes including 8-OHdG DNA glycosylase (OGG1) and human MutY homolog (MUTYH), in addition to nucleotide pool sanitizing enzymes including MutT Homolog 1 (MTH1). Abnormalities of this repair system are present in various cancer types. The present study aimed to elucidate the clinicopathological significance of altered expression levels of inducible nitric oxide synthase (iNOS), 8-OHdG, OGG1, MTH1 and MUTYH in ulcerative colitis (UC) and UC-associated neoplasms. Immunohistochemical staining for these markers and p53 in 23 cases of UC-associated neoplasm (Group A, 14 carcinomas and nine dysplasias), 16 cases of UC without neoplasm (Group B) and 17 cases of normal colon specimens (Group C) was performed. Mutation analyses was conducted for KRAS proto-oncogene, GTPase (*K-ras*), tumor protein P53 (*TP53*) and isocitrate dehydrogenase (NADP (+)) 1, cytosolic (*IDH1*) genes. Immunohistochemically, the iNOS, 8-OHdG, OGG1 and MTH1 expression levels were increased in Groups A and B compared with Group C. The OGG1 and MTH1 expression levels in Group A were also increased compared with Group B. Group A and Group B exhibited increased cytoplasmic expression and decreased nuclear expression of MUTYH compared with Group C. Mutations of *K-ras* and

TP53 were detected in 2/21 (9.5%) and 10/22 (45.5%) cases of Group A, respectively. *IDH1* mutation was not detected in any cases. These findings suggest that, as a response to oxidative damage, OGG1 and MTH1 may be upregulated in UC through an inflammatory condition that progresses to cancer formation. Persisting oxidative damage stress may play a role in the pathogenesis of UC-associated tumors.

Introduction

Ulcerative colitis (UC) is an inflammatory bowel disease characterized by periods of inflammatory recurrence and remission events, accompanied by cell death and regeneration of the colonic mucosa. Patients with UC face an increased risk of UC-associated neoplasm (UCAN) including dysplasia and carcinoma (1). The incidence of colorectal dysplasia in UC patients was observed to be 1.9% at 5 years, 5.1% at 15 years, and 9.2% at 25 years after the onset of the UC (2). The elevated risk of UCAN development is associated with several factors such as disease duration, the extent and severity of inflammation, family history of colorectal carcinoma, backwash ileitis, and primary sclerosing cholangitis (3-5).

The pathogenesis of UCAN is thought to be associated with oxidative DNA damage (6). Nitric oxide (NO) is synthesized by nitric oxide synthase (iNOS) and contributes to the pathogenesis of various types of cancer including colonic carcinoma (7-9). It is known that 8-oxoguanine (8-oxoG) is the most stable product of the base damage due to oxidative stress, and 8-oxoG mismatches with adenine residues, leading to a G:C to T:A transversion mutation (10,11). 8-oxoG and 8-hydroxy-2'-deoxyguanosine (8-OHdG) undergo keto-enol tautomerism, which favors the oxidized product 8-oxodG (12). It was reported that the levels of 8-OHdG are elevated in colorectal cancer and UC-associated dysplasia (6,13).

DNA damage is normally repaired by base excision repair (BER) enzymes such as 8-OHdG DNA glycosylase (OGG1) and human MutY homolog (MUTYH) and nucleotide pool sanitizing enzyme such as MutT Homolog 1 (MTH1). MTH1

Correspondence to: Professor Yoshinao Oda, Department of Anatomic Pathology, Graduate School of Medical Sciences, Kyushu University, Maidashi 3-1-1, Higashi-ku, Fukuoka, Fukuoka 812-8582, Japan
E-mail: oda@surgpath.med.kyushu-u.ac.jp

Key words: ulcerative colitis, carcinoma, oxidative stress, OGG1, MTH1, base excision repair enzyme

protein exists in the nucleotide pool and functions to prevent the misincorporation of 8-OHdG by hydrolyzing 8-OH-dGTP to 8-OH-dGMP (14). OGG1 excises 8-OHdG, which has been mispaired with cytosine. MUTYH excises adenines that have been misincorporated opposite 8-OH-G during replication (15). Altered expressions of OGG1, MTH1 and MUTYH have been reported in various types of cancer (14,16-19). In our previous study, we showed that the nuclear expression of MUTYH was lower in UCAN and UC than in the non-inflamed mucosa (18). However, the expression status of OGG1 and MTH1 in UCAN has not been reported to date.

Various types of genetic mutation have been reported in UCAN, including *K-ras* (30%), *TP53* (40-80%), *p16* (100%), *CTNNB1* (45-50%) and *APC* (10-30%) (20,21). Isocitrate dehydrogenase 1 (*IDH1*) catalyzes the oxidative carboxylation of isocitrate to α -ketoglutarate, resulting in the production of nicotinamide adenine dinucleotide phosphate. Mutations of *IDH1* gene lead to an accumulation of 2-hydroxyglutamate that can induce the hypermethylation of DNA CpG islands, resulting in altered gene transcription and genome stability (22,23). A 2014 study showed that adenocarcinomas associated with inflammatory bowel disease had *IDH1* mutations more frequently compared to sporadic colon cancer (23).

Here we attempted to elucidate the mechanisms of DNA damage and repair in UCAN by evaluating the accumulation of oxidative stress markers and expression DNA repair proteins, iNOS, 8-OHdG, OGG1, MTH1 and MUTYH in UC and UC-associated neoplasms. We also examined the frequencies of *K-ras*, *TP53* and *IDH1* gene mutations as candidates of oxidative stress-induced DNA damage.

Materials and methods

Cases and histological evaluation. We examined the surgical specimens of 23 cases of UC-associated neoplasia (Group A) including 14 carcinoma cases (Group A1) and nine dysplasia cases (Group A2), 16 cases of UC patients without neoplasia (Group B) and 17 cases of normal colon (Group C). These lesions were surgically resected at Kyushu University Hospital and its referral hospitals during the 28-year period from 1987 to 2015. The diagnosis of UC had been made based on clinical, endoscopic and histologic findings. All neoplastic lesions were classified according to the criteria proposed by Riddell *et al.* (24). Patients in Group B were in the active phase of ulcerative colitis and underwent surgery because of fulminant or intractable disease. Group B specimens showed diffuse and severe neutrophilic infiltration, goblet cell depletion, cryptitis, and crypt abscess indicating the active phase of UC. As for Group C, we obtained the specimens of non-cancerous tissue at least 10 cm apart from sporadic colorectal cancer lesions of non-UC patients. Group C specimens showed only slight to mild inflammatory infiltrate.

The study was approved by the Institutional Review Boards of Kyushu University Hospital (no. 27-388).

Immunohistochemical analysis. Formalin-fixed and paraffin-embedded tissue specimens were used for immunohistochemical stainings for iNOS, 8-OHdG, OGG1, MTH1, MUTYH and p53. The primary antibodies and immunohistochemical staining procedures are summarized in Table I. In

brief, after the pretreatment, the primary antibody was applied to the specimen and left overnight at 4°C. Sections were then incubated with a biotinylated secondary antibody for 20 min, followed by the use of a streptavidin-biotin-alkaline phosphatase kit (Nichirei, Tokyo) for 8-OHdG. The reaction was visualized with substrate for alkaline phosphatase for 8-OHdG. For other markers, Envision+ (Dako, Glostrup, Denmark) was used as the secondary antibody, and 3,3-diamino-benzidine was used for visualization. Finally, sections were counterstained with Mayer's hematoxylin.

For the evaluation of immunohistochemical staining, we counted a minimum of 500 cells on each slide. 8-OHdG immunoreactivity in the colorectal crypts was evaluated by the labeling index (LI) of nuclear staining (18). We divided the sections into High and Low expression groups using the median as a reference value, and we compared these values with other immunohistochemical scores and the percentage of gene mutation. The expressions of iNOS expression were evaluated as described (25). The expression level was graded based on the combination of the proportion and the intensity of immunoreactive cells. Proportion scores (PS) were rated on the following scale: 0 (no staining) 1 (<1%), 2 (1% to <10%), 3 (10% to <33%), 4 (33-66%) and 5 (>66%). Intensity scores (IS) were rated on the scale 0 (no staining), 1 (weak), 2 (moderate) and 3 (strong). The total score was obtained by adding the PS and the IS.

The MUTYH expression was evaluated as described (18). The proportion of nuclear staining of the colorectal crypts was classified using a three-grade scale: 0 (no staining), 1 (5-50%) and 2 (>50%), and the intensity of cytoplasmic expression was classified with a two-grade scale (1, weak; 2, strong).

The expressions of MTH1 and OGG1 expression were evaluated as described (16). Staining intensity was rated on the following scale: 0 (no staining), 1 (weak), 2 (moderate) and 3 (strong). The proportion of immunoreactive cells was then scored as 0 (no staining), 1 (1-50%) or 2 (51-100%). The final score was calculated by multiplying the IS by the PS, achieving theoretical results ranging from 0 to 6. Immunoreactivity for p53 was regarded as positive when >10% of the cells were stained (26,27).

Mutational analysis of *K-ras*, *TP53* and *IDH1*. For the mutational analysis, we analyzed the cases of 23 patients with UCAN. Genomic DNA was extracted from paraffin-embedded sections using a macrodissection or microdissection technique. For the macrodissection technique, histological areas measuring approx. ≥ 1 cm in dia. were removed by macroscopic dissection with a needle so as to contain >70% tumor cells. For the microdissection technique, the tumor cells were isolated selectively using laser microdissection (AS LMD system; Leica, Nussloch, Germany) and a pressure catapulting system to get at least 1,000 tumor cells per sample for a polymerase chain reaction (PCR) analysis. Subsequently, genomic DNA was extracted from the samples using a QIAamp[®] DNA Micro kit (Qiagen, Tokyo) and DNeasy Blood & Tissue kits (Qiagen), respectively, in accordance with the manufacturer's protocols.

The mutational analysis for *K-ras*, *TP53* and *IDH1* was performed using PCR and direct sequencing. Mutational hot spots of *K-ras* (codon12 and 13), *TP53* (exons 5-9) and *IDH1* (R132) were included in these PCR analyses (22,28,29).

Table I. Antibodies used in immunohistochemical stain.

Antigen	Primary antibodies		Dilution	Source	Antigen retrieval
iNOS	nos typeII	Mouse monoclonal	1:100	BD Biosciences, Franklin Lakes, NJ	Microwaved for 30 min in target retrieval solution high pH (DAKO, Glostrup, Denmark)
8-OHdG	N45.1	Mouse monoclonal	1:20	Japanese aging control institute, Shizuoka, Japan	Proteinase K for 7 min incubate with 4N HCl for 20 min and with 50 mM Tris base for 5 min
OGG1	NB100-106	Rabbit polyclonal	1:250	Novus Biologicals, Littleton, CO,	Microwaved for 30 min in target retrieval solution high pH (DAKO, Glostrup, Denmark)
MTH1	D-2	Rabbit polyclonal	1:400	Abcam, Cambridge, England	Microwaved for 30 min in target retrieval solution high pH (DAKO, Glostrup, Denmark)
MUTYH	ab13698	Rabbit-polyclonal	1:50	Abcam, Cambridge, England	Microwaved for 30 min in target retrieval solution high pH (DAKO, Glostrup, Denmark)
p53 citrate	PAb1801	Mouse monoclonal	1:100	Oncogene Research Products, San Diego, CA.	Microwaved for 10 min in buffer (pH 6.0)

PCR reactions were performed in a thermocycler (Tgradient; Biometra, Göttingen, Germany). The amplified PCR products were then purified using Montage centrifugal filters (Millipore, Bedford, MA, USA). After purification, direct sequencing was performed using an ABI 3500 genetic analyzer (Applied Biosystems, Foster City, CA).

Statistical analysis. We examined the correlations among clinicopathological factors and molecular data using the Mann-Whitney U-test and Fisher's exact test. A P-value <0.05 was considered significant. As the post-hoc test, we re-examined the data using Tukey's test.

Results

Clinicopathological findings. The clinicopathological features of the patients are summarized in Table II. Among the 23 patients of the Group A (UC-associated neoplasm), 13 patients were men and 10 were women, with the median age of 48 years (range 25-77 years). The median disease duration of Group A was 16 years (range 0.5-35 years). The neoplastic lesions in both Group A1 (UC-associated carcinomas) and Group A2 (UC-associated dysplasia) were predominantly located in the left side of the colon and rectum.

Histologically, Group A2 lesions showed atypical columnar cells with hyperchromatic nuclei and mild nuclear stratification as low-grade dysplasia (LGD) (Fig. 1A), and prominent dystrophic goblet cells and/or severely atypical columnar cells with prominent nuclear stratification without vesicular nuclei, as high-grade dysplasia (HGD). Most of the carcinomas of Group A1 were well to moderately differentiated adenocarcinoma (Fig. 1B). Among the 16 patients of Group B (UC without neoplasm), nine patients were men and seven were

women, with the median age of 43 years (range 22-77 years). The median disease duration of Group B was 6 years (range 0.5-31 years). The disease duration of UC was significantly longer in Group A compared to Group B (P=0.0067).

Immunohistochemical analyses

iNOS. The iNOS expression scores in Group A (median: 8) and Group B (median: 7) were significantly higher than that of Group C (median: 5) (P=0.0002, P=0.0034, respectively) (Table III, Figs. 1C and 2A). There was no significant difference in iNOS expression between Group A and Group B, or between Group A1UC+carc and Group A2 (Tables III and IV).

8-OHdG. The labeling indices of 8-OHdG in Group A (median: 49.3) and Group B (median: 53.6) were significantly higher than that in Group C (median: 29.1) (P=0.0002, P=0.0034, respectively) (Table III; Figs. 1D and 2B). There was no significant difference between Group A and Group B, or between Group A1 and Group A2 (Tables III and IV).

We then examined the correlation between the expression of 8-OHdG and other markers. The OGG1 scores of the high 8-OHdG cases (score 2: one case, score 4: five cases, score 6: five cases) and those of the low 8-OHdG cases (score 2: one case, score 4: five cases, score 6: five cases) revealed no significant correlations. The MTH1 scores of the high 8-OHdG cases (score 2: no cases, score 4: six cases, score 6: five cases) and those of the low 8-OHdG cases (score 2: one case, score 4: four cases, score 6: six cases) revealed no significant correlations. No significant correlations with other markers were revealed (Table V).

OGG1. OGG1 expression was observed only in cytoplasmic regions in our cases, and both the intensity and the proportion of




Figure 1. Representative images of (A and B) histology and (C-H) immunohistochemical stainings in the UC-associated neoplasms. (A) The UC-associated low-grade dysplasias (Group A2) were composed of atypical columnar cells with hyperchromatic nuclei and mild nuclear stratification. (B) The UC-associated carcinomas (Group A1) showed well to moderately differentiated adenocarcinoma, invading the stroma. Group A1 lesions showed diffuse and strong cytoplasmic expressions of (C) iNOS, (E) OGG1 and (F) MTH1, and nuclear accumulations of (D) 8-OHdG and (H) p53. (G) Group A1 lesions also exhibited high cytoplasmic staining and low nuclear staining for MUTYH.




Figure 2. The expression levels of (A) iNOS, (B) 8-OHdG, (C) OGG1 and (D) MTH1 in the UC-associated carcinomas (Group A1) and dysplasias (Group A2), the UC cases without neoplasm (Group B) and the normal colon (Group C). (A-D) Group A (i.e., the combined Group A1 and Group A2) and Group B showed a significantly higher expression of each molecule compared to the Group C. (C and D) In addition, Group A showed significantly higher expressions of OGG1 and MTH1 compared to the than Group B. (A-D) There was no difference between Group A1 and A2 in terms of the expression of these molecules.

Table II. Summary of clinicopathological findings in UC-associated neoplasm, UC without neoplasm and normal colon.

Characteristic	Group A1 UC associated carcinoma (n=14)	Group A2 UC associated dysplasia (n=9)	Group A UC associated neoplasm (n=23)	Group B UC without neoplasm (n=16)	Group C normal colon (n=17)	P-value ^a
Age (year)	55.5 (38-77)	43 (25-68)	48 (25-77)	42.5 (22-77)	69 (53-85)	0.2414
Gender						
Male (%)	8 (57.1)	5 (55.6)	13 (52.2)	9 (56.3)	13 (76.4)	0.9866
Female (%)	6 (42.9)	4 (44.4)	10 (47.8)	7 (43.8)	4 (23.6)	
Disease duration (year)	17.5 (0.5-35)	15 (9-23)	16 (0.5-35)	6 (0.5-31)	-	0.0067 ^b
Location						
Left side (%)	9 (64.3)	7 (77.8)	16 (69.6)	-	6 (35.3)	
Right side (%)	4 (28.6)	2 (22.2)	6 (26.1)	-	11 (64.7)	
Unknown (%)	1 (7.1)	-	1 (4.3)	-	-	
Histology						
Well, moderate (%)	11 (78.6)	-	-	-	-	
Poor, mucinous (%)	3 (21.4)	-	-	-	-	
Tumor depth						
pT1 (%)	4 (28.6)	-	-	-	-	
pT2 (%)	5 (35.7)	-	-	-	-	
pT3 or pT4 (%)	5 (35.7)	-	-	-	-	
Lymph vessel permeation						
Negative (%)	7 (50)	-	-	-	-	
Positive (%)	7 (50)	-	-	-	-	
Venous permeation						
Negative (%)	11 (78.6)	-	-	-	-	
Positive (%)	3 (21.4)	-	-	-	-	
Lymph node metastasis						
Negative (%)	9 (64.3)	-	-	-	-	
Positive (%)	5 (35.7)	-	-	-	-	

^aComparison between UC associated neoplasm and UC without neoplasm. ^bStatistically significant. In the parentheses, range is described for age and disease duration, whereas percentage is described for other factors. Outside the parentheses, average years are described for age and disease duration, whereas case numbers are described for other factors.

Table III. Immunohistochemical findings of iNOS, 8-OHdG, OGG1, MTH1, MUTYH and p53 in UC associated neoplasm compared with UC without neoplasm and normal colon.

Variable	Group A UC associated neoplasm (n=23)	Group B UC without neoplasm (n=16)	Group C normal colon (n=17)	P-value Group A vs. Group B	P-value Group A vs. Group C	P-value Group B vs. Group C
iNOS Score	8 (4-8)	7 (4-8)	5 (0-8)	0.1878	0.0002 ^a	0.0034 ^a
8-OHdG Labeling index	49.3 (10.4-78.1)	53.6 (36.7-77)	29.1 (5.4-58)	0.4407	0.0034 ^a	0.0005 ^a
OGG1 Score	4 (2-6)	4 (1-6)	1 (0-4)	0.0227 ^a	<0.0001 ^a	0.0011 ^a
MTH1 Score	6 (2-6)	4 (2-6)	1 (0-4)	0.0012 ^a	<0.0001 ^a	<0.0001 ^a
MUTYH nuclear Low (score 0) (%)	19 (82.7)	9 (56.3)	4 (23.5)	0.1461	0.0003 ^a	0.0799
High (score 1) (%)	4 (17.3)	7 (43.7)	13 (76.5)			
Cytoplasmic Low (score 1) (%)	9 (39.1)	7 (43.8)	15 (88.2)	1.0000	0.0019 ^a	0.0104 ^a
High (score 2) (%)	14 (60.9)	9 (56.3)	2 (11.8)			
p53 protein Positive (%)	11 (47.8)	0 (0)	Not done	0.0009 ^a	Not available	Not available
Negative (%)	12 (52.1)	16 (100)	Not done			

^aStatistically significant. In the parentheses, range is described for iNOS, 8-OHdG, OGG1 and MTH1, whereas percentage is described for other factors. Outside the parentheses, median scores are described for iNOS, 8-OHdG, OGG1 and MTH1, whereas case numbers are described for other factors.

Table IV. Immunohistochemical findings of iNOS, 8-OHdG, OGG1, MTH1, MUTYH and p53 and TP53 mutation in UC associated carcinoma compared with UC associated dysplasia.

Variable	Group A1 UC associated carcinoma (n=14)	Group A2 UC associated dysplasia (n=9)	P-value
iNOS			
Score	8 (4-8)	7 (4-8)	0.7007
8-OHdG			
Labeling index	49.2 (10.4-78.1)	49.3 (23.4-69.1)	0.6821
OGG1			
Score	5 (2-6)	4 (2-6)	0.5623
MTH1			
Score	6 (2-6)	4 (4-6)	0.7209
MUTYH nuclear			
Low (score 0) (%)	11 (78.6)	8 (88.9)	0.5241
High (score 1) (%)	3 (21.4)	1 (11.1)	
Cytoplasmic			
Low (score 1) (%)	2 (14.3)	7 (77.8)	0.0023 ^a
High (score 2) (%)	12 (85.7)	2 (22.2)	
p53 protein			
Positive (%)	4 (28.6)	7 (77.8)	0.0211 ^a
Negative (%)	10 (71.4)	2 (22.2)	
TP53 mutation ^b			
Positive (%)	5 (35.7)	5 (62.5)	0.2248
Negative (%)	9 (64.3)	3 (37.5)	

^aStatistically significant. ^bOne case with unsuccessful PCR for TP53 was excluded (n=22). In the parentheses, range is described for iNOS, 8-OHdG, OGG1 and MTH1, whereas percentage is described for other factors. Outside the parentheses, median scores are described for iNOS, 8-OHdG, OGG1 and MTH1, whereas case numbers are described for other factors.

cytoplasmic expression were different in each case. The OGG1 expression score was significantly higher in Group A (median: 4) compared to Group B (median: 4) and Group C (median: 1) (P=0.0227, P<0.0001, respectively) (Table III; Figs. 1E and 2C). The OGG1 expression score was also significantly higher in Group B than in Group C (P=0.0011). There was no significant difference in OGG1 expression between Group A1 and Group A2 (Table IV).

MTH1. The MTH1 expression score was significantly higher in Group A (median: 6) compared to Group B (median: 4) and Group C (median: 1) (P=0.0012, P<0.0001, respectively) (Table III; Figs. 1F and 2D). The MTH1 expression was also significantly higher in Group B compared to Group C (P<0.0001). There was no significant difference in MTH1 expression between Group A1 and Group A2 (Table IV).

MUTYH. Regarding the nuclear expression of MUTYH, each case was scored as 0 or 1; no cases were judged as score 2. The nuclear expression of MUTYH was positive (score 1) in 4/23 cases (17.3%) in Group A, 7/16 cases (43.7%) in Group B and 13/17 cases (76.5%) in Group C (Table III and Fig. 1G). Group A and Group B showed significantly lower nuclear expression of MUTYH compared to the Group C (P=0.0003 and P=0.0799, respectively). There was no significant difference in MUTYH nuclear expression between Group A1 and Group A2 (Table IV).

As for the cytoplasmic expression of MUTYH, each case was scored 1 or 2; no cases were judged as score 0. High (score 2) cytoplasmic expression of MUTYH was present in 14/23 cases (60.9%) in Group A, 9/16 cases (56.3%) in Group B and 2/17 cases (11.8%) in Group C (Table III and Fig. 1G). Thus, Group A and Group B showed significantly higher cytoplasmic expression of MUTYH compared to the Group C (P=0.019, P=0.0104, respectively). Notably, Group A1 showed significantly higher cytoplasmic expression of MUTYH compared to Group A2 (P=0.0023) (Table IV).

p53. A nuclear accumulation of p53 protein was present in 11/23 cases (47.8%) of Group A, but it was not observed in Group B (0/16 cases) (P=0.0009) (Table III and Fig. 1H). In addition, Group A2 showed a significantly higher frequency of p53 accumulation compared to the Group A1 (P=0.0211) (Table IV).

In Group A2, 7/9 cases (77.8%) showed nuclear accumulation of p53 protein, and 5/8 cases (62.5%) showed TP53 mutation. Four of the five dysplasias (i.e., Group A2) with TP53 mutation revealed nuclear accumulation of p53 protein (4/5 cases, 80%). In Group A1, 4/14 cases (28.6%) showed nuclear accumulation of p53 protein, and 5/14 cases (35.7%) showed TP53 mutation in Group A1. Of the five carcinomas (i.e., Group A1) with TP53 mutation, only one case revealed nuclear accumulation of p53 protein (1/5 cases, 20%). The concordance rate between nuclear accumulation of p53 protein and TP53 mutation was

Table V. The correlations between 8-OHdG accumulation and immunohistochemical features of UC associated neoplasm (n=22)^a.

Immunohistochemistry	8-OHdG accumulation		P-value
	High (n=11)	Low (n=11)	
Age (year)	49 (26-77)	44 (25-68)	0.5759
Gender			
Male (%)	5 (45.5)	8 (72.7)	0.3870
Female (%)	6 (54.5)	3 (27.3)	
Disease duration (year)	17 (9-35)	12 (0.5-24)	0.3832
iNOS			
Score	7 (4-8)	8 (5-8)	0.0819
OGG1			
Score	4 (2-6)	4 (2-6)	0.5482
MTH1			
Score	4 (4-6)	6 (2-6)	0.9714
MUTYH nuclear			
Low (score 0) (%)	8 (72.7)	10 (90.9)	0.5865
High (score 1) (%)	3 (27.3)	1 (9.1)	
Cytoplasmic			
Low (score 1) (%)	3 (27.3)	5 (45.5)	0.6594
High (score 2) (%)	8 (72.7)	6 (54.5)	
p53 protein			
Positive (%)	6 (54.5)	4 (36.4)	0.3918
Negative (%)	5 (45.5)	7 (63.6)	
TP53 mutation			
Positive (%)	4 (36.4)	6 (54.6)	0.6699
Negative (%)	7 (63.6)	5 (45.4)	

^aOne case with unsuccessful PCR for TP53 was excluded. In the parentheses, range is described for iNOS, OGG1 and MTH1, whereas percentage is described for other factors. Outside the parentheses, median scores are described for iNOS, OGG1 and MTH1, whereas case numbers are described for other factors.

not significantly different between the two groups, but in the dysplasias (Group A 2), this rate tended to be higher than that in the carcinomas (Group A1) (P=0.058).

Mutational analysis

TP53. The results of our mutation analysis as well as the clinicopathological data of Group A the UC+ group are summarized in Table VI. The PCR for TP53 was successful in 22 cases of Group A. The TP53 mutation was present in a total of 10 of the 22 (45.5%) cases, including 5/14 cases (35.7%) of Group A1 and 5/8 cases (62.5%) of Group A2. There was no significant difference in the frequency of TP53 mutation between these two groups. Among the 10 cases with TP53 mutation, transversion and transition mutations were present in two and eight cases, respectively.

We then checked the correlation between TP53 mutation and inflammation-associated marker expressions; no significant correlation was observed (Table VII).

K-ras. The mutation analysis of K-ras was successful in 21 cases in Group A. We found that 2 of 21 (9.5%) cases had

K-ras gene mutation at codon 12; one mutation was the transversion type and the other was the transition type (Table VI).

IDH1. No cases in Group A had IDH1 gene mutation.

Discussion

Our present findings demonstrated that the expression levels of iNOS, 8-OHdG, OGG1 and MTH1 were significantly higher in UCAN and UC samples than in non-inflamed mucosa. In addition, the OGG1 and MTH1 expressions were significantly higher in the UCAN than in the UC cases. It has been reported that increased levels of 8-OHdG induce the activation of MTH1 or OGG1 to minimize 8-OHdG accumulation in mammalian DNA (13,30). Similarly in UC, oxidative stress may upregulate defense systems such as MTH1 and OGG1 to minimize 8-OHdG accumulation.

Liao *et al* reported that OGG1 knockout mice showed a significantly increased risk of adenocarcinoma development in the colon compared to wild-type mice after the induction of dextran sulfate sodium-induced colitis (31). In breast

Table VI. The clinicopathological features and molecular findings of UC-associated neoplasm and mutation analysis for TP53 and K-ras.

Case	Gender	Age	Disease duration	Tumor location	Diagnosis	Clinical stage	p53 IHC	TP53 mutation	K-ras mutation
1	F	63	20	Rectum	Ca	T2	+	codon220, TAT>CAT	wt
2	F	63	16	Descending colon	Ca	T2	-	codon240, AGT>AGC	wt
3	F	71	10	Ascending colon	Ca	T2	-	codon249, AGG>AGT	codon12, GGT>GAT
4	M	43	10	Rectum	Ca	T3	-	codon249, AGG>AGT	wt
5	F	48	24	Rectum	Ca	T3	-	codon285, GAA>GAG	wt
6	M	49	9	Ascending colon	Ca	T2	+	wt	wt
7	F	43	5	NA	Ca	T1	+	wt	wt
8	M	66	33	Descending colon	Ca	T2	+	wt	wt
9	M	38	0.5	Ascending colon	Ca	T1	-	wt	wt
10	F	75	35	Rectum	Ca	T1	-	wt	wt
11	M	48	NA	Rectum	Ca	T1	-	wt	wt
12	M	39	20	Rectum	Ca	T3	-	wt	wt
13	M	62	19	Transverse colon	Ca	T2	-	wt	codon12, GGT>TGT
14	M	68	19	Descending colon	Ca	T2	-	wt	wt
15	M	80	20	Sigmoid colon	HGD	Tis	+	codon197, GTG>ATG	wt
16	M	50	11	Sigmoid colon	HGD	Tis	-	codon197, GTG>ATG	wt
17	M	22	13	Transverse colon	HGD	Tis	+	codon220, TAT>CAT	wt
18	M	34	15	Rectum	HGD	Tis	+	codon312, AGC>AAG	wt
19	F	26	9	Rectum	HGD	Tis	-	wt	wt
20	F	44	11	Rectum	LGD	Tis	+	codon197, GTG>ATG	NA
21	F	68	23	Descending colon	LGD	Tis	+	NA	NA
22	M	43	18	Transverse colon	LGD	Tis	+	wt	Wt
23	F	42	NA	Transverse colon	LGD	Tis	+	wt	Wt

Ca, cancer; HGD, high-grade dysplasia; LGD, low-grade dysplasia; IHC, immunohistochemistry; wt, wild type; NA, not available.

Table VII. The correlations between TP53 mutational status and clinicopathological factors or immunohistochemical features of UC-associated neoplasia (n=22)^a.

Variable	TP53 mutation		P-value
	Positive (%) (n=10)	Negative (%) (n=12)	
Age (year)	46 (25-71)	45.5 (26-77)	0.8948
Gender			
Male (%)	5 (50)	8 (66.6)	0.4285
Female (%)	5 (50)	4 (33.3)	
Disease duration (year)	14 (10-24)	18.5 (0.5-35)	0.9698
iNOS			
Score	8 (6-8)	7 (4-8)	0.1099
8-OHdG			
Labelling index (%)	45.9 (23.4-78.1)	50.7 (10.4-69.1)	0.9212
OGG1			
Score	4 (2-6)	6 (2-6)	0.1028
MTH1			
Score	6 (4-6)	4 (2-6)	0.3515
MUTYH nuclear			
Low (score 0) (%)	8 (80)	10 (83.3)	0.8403
High (score 1) (%)	2 (20)	2 (16.7)	
Cytoplasmic			
Low (score 1) (%)	3 (30)	5 (41.7)	0.5711
High (score 2) (%)	7 (70)	7 (58.3)	
p53 protein			
Positive (%)	5 (50)	5 (41.7)	0.6959
Negative (%)	5 (50)	7 (58.3)	

^aOne case with unsuccessful PCR for TP53 was excluded. In the parentheses, range is described for age, disease duration, iNOS, 8-OHdG, OGG1 and MTH1, whereas percentage is described for other factors. Outside the parentheses, average years are described for age and disease duration, whereas case numbers are described for other factors.

cancer, reduced OGG1 expression was correlated with poor prognosis (17). On the other hand, Kubo *et al* showed 8-OHdG accumulation and cytoplasmic OGG1 overexpression in esophageal squamous cell carcinoma (16). They also showed that cytoplasmic OGG1 expression was correlated with the depth of cancer, lymph node metastasis and the stage of progression. In the present study, the OGG1 cytoplasmic expression was higher in UCAN than UC cases.

It is reported that OGG1 protein is present in the nucleus and mitochondria and that the protein at both of these sites work as DNA repair protein. The dysfunction of each OGG1 leads to high 8-OHdG accumulation and cell death (32). These findings suggested that OGG1 overexpression might represent persisting oxidative stress and cellular response in UCAN and unlike in mouse models, the upregulation of OGG1 function may play some role in the pathogenesis of human UCAN.

Tsuzuki *et al* showed that in an MTH1-deficient mouse model, a greater number of tumors were formed in the lung, liver, and stomach compared to MTH1 wild-type mice (33). In contrast, Song *et al* showed that MTH1 expression was significantly higher in human gastric cancer tissue than

para-cancer tissue, and that MTH1 expression correlated with 8-OHdG accumulation (34). These findings may be explained in part by the notion that cancer cells may require MTH1 to avoid both DNA damage and cell death because an excessive accumulation of 8-OHdG leads to tumor cell death (35,36). In the present study, the MTH1 expression was higher in UCAN than in the UC and non-inflamed mucosa. Our results may support the hypothesis that MTH1 is required for cancer survival for defense against oxidative damage in UCAN. Similar to the previous study, the present UCAN and UC cases exhibited higher cytoplasmic expression and lower nuclear expression of MUTYH compared to the non-inflamed mucosa. Such an abnormal expression pattern of MUTYH may represent dysfunction of MUTYH, as we had suggested (18).

In general, 8-OHdG accumulation leads to the G:C to T:A and A:T to C:G transversion mutations (10). Chaubert *et al* reported that *K-ras* mutations were present in 44% of UCAN patients with a predominance of G:C to T:A transversion (37). However, in our series, despite the accumulation of 8-OHdG, transversion mutations of *K-ras* or *TP53* gene were not frequent.

For example, among our 21 cases of UCAN, *K-ras* mutation was identified in two (9.5%) cases, one of whom showed a G:C to T:A transversion mutation. Although we detected the *TP53* mutations in 10/22 cases (45.5%) of UCAN, only two cases showed transversion mutations.

The frequency of *TP53* mutation was higher in our dysplasia cases compared to the carcinoma cases. Carcinoma may arise from other factors, such as genetic alterations in tumor suppressor genes, oncogenes and genes encoding DNA repair proteins, as well as from loss of genomic stability.

We also observed that the expression levels of OGG1, *MUTYH* and *MTH1* were not correlated with the presence of *TP53* gene mutation or p53 protein accumulation. It thus seems to be less likely that a dysfunction of OGG1, *MUTYH* or *MTH1* caused the excessive accumulation of 8-OHdG and the subsequent transversion mutation of *K-ras* or *TP53* in UCAN. However, the possibility remains that an accumulation of 8-OHdG might cause a transversion mutation of other genes that would participate in the pathogenesis of UCAN. As an additional possibility, dysfunction of OGG1 and *MUTYH* caused by *TP53* mutation might play a role in the tumorigenesis in some populations of UCAN. It has been reported that, since OGG1 and *MUTYH* are transcriptionally regulated by p53, p53 deficiency leads to OGG1 and *MUTYH* dysfunction, resulting in escape from programmed cell death under oxidative stress and the promotion of tumorigenesis (38,39). However, further study is necessary to elucidate the mechanism of *TP53* mutation and its role in the tumorigenesis of UCAN.

A previous study revealed that isocitrate dehydrogenase 1(*IDH1*) mutations were present in 1% of sporadic colorectal carcinomas (22). Hartman *et al* also reported that adenocarcinomas associated with inflammatory bowel disease more frequently demonstrated *IDH1* mutations than conventional intestinal adenocarcinomas did (13% vs. 0%) (23). In addition, *IDH1* mutations were more frequently observed in Crohn's disease-associated neoplasms than UC-associated neoplasms in their study. In the present study, no *IDH1* gene mutation was detected in the neoplasms associated with UC. These findings suggest that *IDH1* mutation may be exceptional in UCAN.

In conclusion, the results of our study demonstrated that iNOS, 8-OHdG, *MTH1* and OGG1 were accumulated in UCAN and the inflamed mucosa of UC patients. We also observed that the expressions of *MTH1* and OGG1 were higher in UCAN than in UC. Our results suggest that both inflamed mucosa and neoplasms of UC are exposed to persisting oxidative damage, which may lead to the increased expressions of *MTH1* and OGG1, and possibly to UC-related carcinogenesis.

Acknowledgements

The authors would like to thank the Research Support Center, Graduate School of Medical Sciences, Kyushu University for their technical support.

References

- Jess T, Rungoe C and Peyrin-Biroulet L: Risk of colorectal cancer in patients with ulcerative colitis: A meta-analysis of population-based cohort studies. *Clin Gastroenterol Hepatol* 10: 639-645, 2012.
- Jess T, LoftusEV Jr, Velayos FS, Harmsen WS, Zinsmeister AR, Smyrk TC, Tremaine WJ, MeltonLJ III, Munkholm P and Sandborn WJ: Incidence and prognosis of colorectal dysplasia in inflammatory bowel disease: A population-based study from Olmsted County, Minnesota. *Inflamm Bowel Dis* 12: 669-676, 2006.
- Brentnall TA, Haggitt RC, Rabinovitch PS, Kimmey MB, Bronner MP, Levine DS, Kowdley KV, Stevens AC, Crispin DA, Emond M and Rubin CE: Risk and natural history of colonic neoplasia in patients with primary sclerosing cholangitis and ulcerative colitis. *Gastroenterology* 110: 331-338, 1996.
- Nuako KW, Ahlquist DA, Mahoney DW, Schaid DJ, Siems DM and Lindor NM: Familial predisposition for colorectal cancer in chronic ulcerative colitis: A case-control study. *Gastroenterology* 115: 1079-1083, 1998.
- Devroede GJ, Taylor WF, Sauer WG, Jackman RJ and Stickler GB: Cancer risk and life expectancy of children with ulcerative colitis. *N Engl J Med* 285: 17-21, 1971.
- D'Inca R, Cardin R, Benazzato L, Angriman I, Martines D and Sturniolo GC: Oxidative DNA damage in the mucosa of ulcerative colitis increases with disease duration and dysplasia. *Inflamm Bowel Dis* 10: 23-27, 2004.
- Kojima M, Morisaki T, Tsukahara Y, Uchiyama A, Matsunari Y, Mibu R and Tanaka M: Nitric oxide synthase expression and nitric oxide production in human colon carcinoma tissue. *J Surg Oncol* 70: 222-229, 1999.
- Choudhari SK, Chaudhary M, Bagde S, Gadbail AR and Joshi V: Nitric oxide and cancer: A review. *World J Surg Oncol* 11: 118, 2013.
- Vannini F, Kashfi K and Nath N: The dual role of iNOS in cancer. *Redox Biol* 6: 334-343, 2015.
- Nakabeppu Y, Tsuchimoto D, Ichinoe A, Ohno M, Ide Y, Hirano S, Yoshimura D, Tominaga Y, Furuichi M and Sakumi K: Biological significance of the defense mechanisms against oxidative damage in nucleic acids caused by reactive oxygen species: From mitochondria to nuclei. *Ann N Y Acad Sci* 1011: 101-111, 2004.
- Shibutani S, Takeshita M and Grollman AP: Insertion of specific bases during DNA synthesis past the oxidation-damaged base 8-oxodG. *Nature* 349: 431-434, 1991.
- Vakakanidis A, Vlachogianni T and Fiotakis C: 8-hydroxy-2'-deoxyguanosine (8-OHdG): A critical biomarker of oxidative stress and carcinogenesis. *J Environ Sci Health C Environ Carcinog Ecotoxicol Rev* 27: 120-139, 2009.
- Kondo S, Toyokuni S, Tanaka T, Hiai H, Onodera H, Kasai H and Imamura M: Overexpression of the hOGG1 gene and high 8-hydroxy-2'-deoxyguanosine (8-OHdG) lyase activity in human colorectal carcinoma: Regulation mechanism of the 8-OHdG level in DNA. *Clin Cancer Res* 6: 1394-1400, 2000.
- Sekiguchi M and Tsuzuki T: Oxidative nucleotide damage: Consequences and prevention. *Oncogene* 21: 8895-8904, 2002.
- García-Quispes WA, Pérez-Machado G, Akdi A, Pastor S, Galofré P, Biarnés F, Castell J, Velázquez A and Marcos R: Association studies of OGG1, XRCC1, XRCC2 and XRCC3 polymorphisms with differentiated thyroid cancer. *Mutat Res* 709-710: 67-72, 2011.
- Kubo N, Morita M, Nakashima Y, Kitao H, Egashira A, Saeki H, Oki E, Kakeji Y, Oda Y and Maehara Y: Oxidative DNA damage in human esophageal cancer: Clinicopathological analysis of 8-hydroxydeoxyguanosine and its repair enzyme. *Dis Esophagus* 27: 285-293, 2014.
- Karihtala P, Kauppila S, Puistola U and Jukkola-Vuorinen A: Absence of the DNA repair enzyme human 8-oxoguanine glycosylase is associated with an aggressive breast cancer phenotype. *Br J Cancer* 106: 344-347, 2012.
- Gushima M, Hirahashi M, Matsumoto T, Fujita K, Fujisawa R, Mizumoto K, Nakabeppu Y, Iida M, Yao T and Tsuneyoshi M: Altered expression of *MUTYH* and an increase in 8-hydroxydeoxyguanosine are early events in ulcerative colitis-associated carcinogenesis. *J Pathol* 219: 77-86, 2009.
- Grasso F, Di Meo S, De Luca G, Pasquini L, Rossi S, Boirivant M, Biffoni M, Bignami M and Di Carlo E: The *MUTYH* base excision repair gene protects against inflammation-associated colorectal carcinogenesis. *Oncotarget* 6: 19671-19684, 2015.
- Scarpa M, Castagliuolo I, Castoro C, Pozza A, Scarpa M, Kotsafti A and Angriman I: Inflammatory colonic carcinogenesis: A review on pathogenesis and immunosurveillance mechanisms in ulcerative colitis. *World J Gastroenterol* 20: 6774-6785, 2014.

21. Yin J, Harpaz N, Tong Y, Huang Y, Laurin J, Greenwald BD, Hontanosas M, Newkirk C and Meltzer SJ: p53 point mutations in dysplastic and cancerous ulcerative colitis lesions. *Gastroenterology* 104: 1633-1639, 1993.
22. Borger DR, Tanabe KK, Fan KC, Lopez HU, Fantin VR, Straley KS, Schenkein DP, Hezel AF, Ancukiewicz M, Liebman HM, *et al*: Frequent mutation of isocitrate dehydrogenase (IDH)1 and IDH2 in cholangiocarcinoma identified through broad-based tumor genotyping. *Oncologist* 17: 72-79, 2012.
23. Hartman DJ, Binion D, Regueiro M, Schraut W, Bahary N, Sun W, Nikiforova M and Pai RK: Isocitrate dehydrogenase-1 is mutated in inflammatory bowel disease-associated intestinal adenocarcinoma with low-grade tubuloglandular histology but not in sporadic intestinal adenocarcinoma. *Am J Surg Pathol* 38: 1147-1156, 2014.
24. Riddell RH, Goldman H, Ransohoff DF, Appelman HD, Fenoglio CM, Haggitt RC, Ahren C, Correa P, Hamilton SR, Morson BC, *et al*: Dysplasia in inflammatory bowel disease: Standardized classification with provisional clinical applications. *Human Pathol* 11: 931-968, 1983.
25. Brennan PA, Palacios-Callender M, Zaki GA, Spedding AV and Langdon JD: Type II nitric oxide synthase (NOS2) expression correlates with lymph node status in oral squamous cell carcinoma. *J Oral Pathol Med* 30: 129-134, 2001.
26. Maehara Y, Tomoda M, Hasuda S, Kabashima A, Tokunaga E, Kakeji Y and Sugimachi K: Prognostic value of p53 protein expression for patients with gastric cancer- a multivariate analysis. *Br J Cancer* 79: 1255-1261, 1999.
27. Feng CW, Wang LD, Jiao LH, Liu B, Zheng S and Xie XJ: Expression of p53, inducible nitric oxide synthase and vascular endothelial growth factor in gastric precancerous and cancerous lesions: Correlation with clinical features. *BMC Cancer* 2: 8, 2002.
28. Oda Y, Sakamoto A, Satio T, Kawachi S, Iwamoto Y and Tsuneyoshi M: Molecular abnormalities of p53, MDM2 and H-ras in synovial sarcoma. *Mod Pathol* 13: 994-1004, 2000.
29. Fujita K, Yamamoto H, Matsumoto T, Hirahashi M, Gushima M, Kishimoto J, Nishiyama K, Taguchi T, Yao T and Oda Y: Sessile serrated adenoma with early neoplastic progression: A clinicopathologic and molecular study. *Am J Surg Pathol* 35: 295-304, 2011.
30. Kajitani K, Yamaguchi H, Dan Y, Furuichi M, Kang D and Nakabeppu Y: MTH1, an oxidized purine nucleoside triphosphatase, suppresses the accumulation of oxidative damage of nucleic acids in the hippocampal microglia during kainate-induced excitotoxicity. *J Neurosci* 26: 1688-1698, 2006.
31. Liao J, Seril DN, Lu GG, Zhang M, Toyokuni S, Yang AL and Yang GY: Increased susceptibility of chronic ulcerative colitis-induced carcinoma development in DNA repair enzyme Ogg1 deficient mice. *Mol Carcinog* 47: 638-646, 2008.
32. Nakabeppu Y: Cellular levels of 8-oxoguanine in either DNA or the nucleotide pool play pivotal roles in carcinogenesis and survival of cancer cells. *Int J Mol Sci* 15: 12543-12557, 2014.
33. Tsuzuki T, Egashira A, Igarashi H, Iwakuma T, Nakatsuru Y, Tominaga Y, Kawate H, Nakao K, Nakamura K, Ide F, *et al*: Spontaneous tumorigenesis in mice defective in the MTH1 gene encoding 8-oxo-dGTPase. *Proc Natl Acad Sci USA* 98: 11456-11461, 2001.
34. Song WJ, Jiang P, Cai JP and Zheng ZQ: Expression of cytoplasmic 8-oxo-Gsn and MTH1 correlates with pathological grading in human gastric cancer. *Asian Pac J Cancer Prev* 16: 6335-6338, 2015.
35. Gad H, Koolmeister T, Jemth AS, Eshtad S, Jacques SA, Ström CE, Svensson LM, Schultz N, Lundbäck T, Einarsdottir BO, *et al*: MTH1 inhibition eradicates cancer by preventing sanitation of the dNTP pool. *Nature* 508, 215-221, 2014.
36. Huber KV, Salah E, Radic B, Gridling M, Elkins JM, Stukalov A, Jemth AS, Göktürk C, Sanjiv K, Strömberg K, *et al*: Stereospecific targeting of MTH1 by (S)-crizotinib as an anticancer strategy. *Nature* 508: 222-227, 2014.
37. Chaubert P, Benhattar J, Saraga E and Costa J: K-ras mutations and p53 alterations in neoplastic and nonneoplastic lesions associated with longstanding ulcerative colitis. *Am J Pathol* 144: 767-775, 1994.
38. Oka S, Leon J, Tsuchimoto D, Sakumi K and Nakabeppu Y: MUTYH, an adenine DNA glycosylase, mediates p53 tumor suppression via PARP-dependent cell death. *Oncogenesis* 4: e142, 2015.
39. Chatterjee A, Mambo E, Osada M, Upadhyay S and Sidransky D: The effect of p53-RNAi and p53 knockout on human 8-oxoguanine DNA glycosylase (hOgg1) activity. *FASEB J* 20: 112-114, 2006.