




DOI: 10.1038/s41467-018-05089-5

OPEN

Author Correction: Next generation histology methods for three-dimensional imaging of fresh and archival human brain tissues

Hei Ming Lai^{1,2}, Alan King Lun Liu ^{1,2}, Harry Ho Man Ng ¹, Marc H. Goldfinger², Tsz Wing Chau ², John DeFelice², Bension S. Tilley², Wai Man Wong¹, Wutian Wu^{1,3,4} & Steve M. Gentleman²

Correction to: *Nature Communications* 9:1066 <https://doi.org/10.1038/s41467-018-03359-w>; published online: 14 March 2018

In the original version of this Article, the concentration of boric acid buffer for the SDS clearing solution was given incorrectly as ‘1 M sodium borate’ and should have read ‘0.2 M boric acid’. Also, the composition of PBST incorrectly read ‘1% Triton X-100 (vol/vol) and 0.1% sodium azide (wt/vol)’ and should have read ‘0.1% Triton X-100 (vol/vol) and 0.01% sodium azide (wt/vol)’. Further, the pH of the OPTIClear solution was not stated, and should have read ‘with a pH between 7 to 8 adjusted with hydrochloric acid’. These errors have been corrected in both the PDF and HTML versions of the Article.

Published online: 10 July 2018



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2018

¹School of Biomedical Sciences, Li Ka Shing Faculty of Medicine, The University of Hong Kong, Hong Kong 852, Hong Kong. ²Neuropathology Unit, Division of Brain Sciences, Imperial College London, Burlington Danes Building, Hammersmith Campus, Du Cane Road, London W12 0NN, UK. ³GMH Institute of CNS Regeneration, Jinan University, 601 Huangpu Avenue West, Guangzhou 510632, China. ⁴Re-Stem Biotechnology Co., Ltd, 2588 Wuzhong Road, Suzhou 215310, China. These authors contributed equally: Hei Ming Lai, Alan King Lun Liu and Harry Ho Man Ng. Correspondence and requests for materials should be addressed to W.W. (email: wtwu@hku.hk) or to S.M.G. (email: s.gentleman@imperial.ac.uk)