

REVIEW

Chemokines in homeostasis and diseases

Keqiang Chen¹, Zhiyao Bao^{1,2}, Peng Tang^{1,3}, Wanghua Gong⁴, Teizo Yoshimura⁵ and Ji Ming Wang¹

For the past twenty years, chemokines have emerged as a family of critical mediators of cell migration during immune surveillance, development, inflammation and cancer progression. Chemokines bind to seven transmembrane G protein-coupled receptors (GPCRs) that are expressed by a wide variety of cell types and cause conformational changes in trimeric G proteins that trigger the intracellular signaling pathways necessary for cell movement and activation. Although chemokines have evolved to benefit the host, inappropriate regulation or utilization of these small proteins may contribute to or even cause diseases. Therefore, understanding the role of chemokines and their GPCRs in the complex physiological and diseased microenvironment is important for the identification of novel therapeutic targets. This review introduces the functional array and signals of multiple chemokine GPCRs in guiding leukocyte trafficking as well as their roles in homeostasis, inflammation, immune responses and cancer.

Cellular and Molecular Immunology (2018) **15**, 324–334; doi:10.1038/cmi.2017.134; published online 29 January 2018

Keywords: chemokines; chemotaxis; diseases; GPCRs; homeostasis; inflammatory cells

INTRODUCTION

Leukocyte accumulation is a hallmark of inflammation, immune responses and cancer progression and is mediated by a variety of small protein mediators named chemokines that bind to G protein-coupled receptors (GPCRs). Chemokine GPCRs are mainly expressed by leukocytes, and their major function was originally proposed to be guiding leukocyte trafficking and homing, activating integrins, generating superoxide anions and releasing of granule contents. These functions constitute the first-line host defense against invading microorganisms and tissue injury.¹ However, over the past decade, chemokines and their GPCRs have also been implicated in many essential pathophysiological processes, including development, angiogenesis, epithelial homeostasis and cancer progression.² This article aims to update the current understanding of the multifaceted functions of chemokines and their GPCRs to better position these molecules in the sphere of diseases for the identification of novel therapeutic targets. The readers are also advised to refer to excellent specific review articles for more in-depth information.^{3–8}

CHEMOATTRACTANT GPCR AND LIGAND FAMILIES

GPCRs that mediate cell chemotaxis are categorized into classical and chemokine subfamilies according to the ligand source.⁹ Classical GPCRs include formyl peptide receptors (FPR1, FPR2 and FPR3), platelet-activating factor receptor (PAFR), activated complement component 5 receptor (C5aR) and leukotriene B₄ receptors (BLT1 and BLT2). Another classical chemoattractant GPCR is ChemR23, known as chemokine-like receptor 1 (CMKLR1). ChemR23 is mainly expressed in myeloid cells, such as human macrophages, immature dendritic cells (DCs)¹⁰ and plasmacytoid DCs (PDCs).¹¹ This receptor is detected in the thymus, bone marrow, spleen, fetal liver and lymphoid organs, suggesting its involvement in diverse pathophysiological processes. Structurally, CMKLR1 is more closely related to chemoattractant receptors, such as FPRs as well as the C3aR and C5aR, rather than to chemokine receptors.¹² The chemotactic ligands for ChemR23 include chemerin, the chemerin-derived peptide, C15 and the lipid resolving E1 peptide, Resolvin E1.^{10,13} Chemerin is produced by cells of high endothelial venules in reactive lymph nodes (LNs) and in dermal blood

¹Cancer and Inflammation Program, Center for Cancer Research, National Cancer Institute at Frederick, Frederick, MD 21702, USA; ²Department of Pulmonary & Critical Care Medicine, Ruijin Hospital, Shanghai Jiaotong University School of Medicine, Shanghai 200025, P. R. China; ³Department of Breast Surgery, Southwest Hospital, Third Military Medical University, Chongqing 400038, China; ⁴Basic Research Program, Leidos Biomedical Research, Inc., Frederick, MD 21702, USA and ⁵Department of Pathology and Experimental Medicine, Graduate School of Medicine, Dentistry and Pharmaceutical Sciences, Okayama University, Okayama 700-8558, Japan

Correspondence: Dr JM Wang, MD, PhD, Cancer and Inflammation Program, Center for Cancer Research, National Cancer Institute at Frederick, 1050 Boyles street, Building 560, Room 31-76, Frederick, MD 21702, USA.

E-mail: wangji@mail.nih.gov

Received: 6 July 2017; Revised: 16 October 2017; Accepted: 18 October 2017

vessels in autoimmune skin lesions. The ChemR23/chemerin pair has received increasing attention due to its involvement in many pathological conditions.¹¹ For example, in hemostasis, the distribution of PDCs is restricted to primary and secondary lymphoid organs. However, in autoimmune diseases (that is, lupus erythematosus disease, psoriasis and rheumatoid arthritis), allergy (that is, contact dermatitis and nasal mucosa polyps) and cancer, PDCs accumulate in non-lymphoid tissues as a result of the ChemR23/chemerin interaction.¹¹ ChemR23/chemerin mediates an important host defense against viral infection, as shown in mouse acute viral pneumonia, in which ChemR23 attenuates lung inflammation and enhances antiviral immunity. As a consequence, ChemR23^{-/-} mice exhibit delayed viral clearance and increased mortality, which are associated with reduced PDC recruitment and production of type I interferon (IFN)¹⁴ (Table 1).

Chemokine GPCRs are composed of four subfamilies based on the conserved N-terminal cysteine residues in their mature ligands, termed C-C chemokine motif receptor (CCR), C-X-C chemokine motif receptor (CXCR), CX3CR and XCR^{15,16} (Table 2). A subset of GPCRs, recently termed atypical chemokine receptors, including ACKR1–ACKR6, have also been identified^{17,18} (Table 3). A prominent feature of many chemoattractants and GPCRs is their promiscuity, that is, some chemoattractants bind more than one GPCR, and conversely, some GPCRs display overlapping ligand specificities with variable affinities and occasionally exhibit diverse functions.¹⁹ This finding may represent a mechanism by which the host is able to mobilize cells with flexibility according to the changing microenvironment.

CHEMOKINES AND GPCRS IN HOMEOSTASIS

Homeostatic chemokines

Under normal conditions, homeostatic chemokines regulate cellular trafficking by directing cells that express corresponding GPCRs to specific locations. For example, in the skin, CCL27 recruits cutaneous leukocyte antigen (CLA)⁺ T cells that express the receptor CCR10.²⁰ In the brain, CXCL12 mediates leukocyte homing via CXCR4²¹; in the small intestine, CCL25 recruits the homing of T cells that express $\alpha 4\beta 7$ integrin and CCR9^{22,23}; in the lungs, CXCL12 recruits stromal cell homing²⁴; in LNs, CCL21 and CCL19 recruit CCR7⁺ T cells, and CXCL13 recruits CXCR5⁺ B and T cells; and in the bone marrow (BM), CXCL12 recruits CXCR4⁺ hematopoietic stem cells (HSCs) and creates an HSC 'niche'.^{25,26} Therefore, inhibition of the CXCL12–CXCR4 interaction 'liberates' HSCs from the BM niche into circulation (Table 4). Accumulating evidence indicates that homeostatic chemokines are critical for tissues and organs to maintain a consistent and dynamic leukocyte composition in preparation for immune responses triggered by insults.

Central nervous system (CNS)

A number of homeostatic chemokines are expressed in the vasculature of the blood–brain barrier, including CXCL12, CCL19, CCL20 and CCL21, to regulate the entry of leukocytes into the CNS for immunosurveillance. CXCL12 is also involved in the development of the CNS, including adult neurogenesis and neuronal survival, with opposing activity to the homeostatic chemokine CXCL14, which regulates synaptic inputs to neural precursors. CX3CL1, another homeostatic chemokine, promotes neuronal cell survival and communication with microglia, a process also partially regulated by CXCL12.²¹

Table 1 Classical chemoattractant GPCRs and ligands

Receptor	Expression	Ligands	Functions
FPR1	Myeloid cells, lymphocytes, tumor cells	Bacteria- and host-derived peptides	Leukocyte chemotaxis and activation, tumor growth, invasion, angiogenesis
FPR2	Myeloid cells, tumor cells	Bacteria- and host-derived peptides	Leukocyte chemotaxis and activation, antitumor defense, tumor invasion
FPR3	Monocytes, dendritic cells, tumor cells	Synthetic and host-derived peptides	Monocyte and DC chemotaxis and activation, tumor invasion
PAFR	Macrophages, granule leucocytes, various tissue cells, tumor cells	PAF	Leukocyte chemotaxis and activation, tumor growth and metastasis; angiogenesis
C5aR	Basophils, dendritic cells, mast cells, various nonimmune cells, tumor cells	C5a	Leukocyte chemotaxis and activation, tumor metastasis; opposing function in angiogenesis
LTB4R (BLT1)	Neutrophils, macrophages, T-lymphocytes, tumor cells	LTB4	Leukocyte chemotaxis and activation, tumor growth
LTB4R (BLT2)	Most human tissues cells, leukocytes, tumor cells	LTB4	Leukocyte chemotaxis and activation, tumor growth and metastasis
CMKLR1 (ChemR23)	Mainly in myeloid cells	Chemerin, C15, RvE1	Migration of macrophages, DCs and PDCs

Abbreviations: CMKLR1, chemokine-like receptor 1; C5aR, activated complement component 5 receptor (C5aR); LTB4R, leukotriene B4 receptor; DC, dendritic cell; FPR, formyl peptide receptor; GPCR, G-protein coupled receptor; PAFR, platelet-activating factor receptor; PDC, plasmacytoid DC; RvE1, Resolvin E1.

Bone marrow

CXCL12 is constitutively produced by BM stromal cells, including vascular endothelial cells and osteoblasts.^{25,28}

Table 2 Chemokine receptors and ligands

Receptor	Ligands
CCR1	CCL3, CCL4, CCL5, CCL7, CCL13, CCL14, CCL15, CCL16, CCL23
CCR2	CCL2, CCL7, CCL8, CCL12, CCL13, CCL16
CCR3	CCL5, CCL7, CCL8, CCL11, CCL13, CCL15, CCL16, CCL24, CCL26, CCL28
CCR4	CCL2, CCL3, CCL5, CCL7, CCL22
CCR5	CCL3, CCL4, CCL5, CCL8, CCL11, CCL14, CCL16
CCR6	CCL20
CCR7	CL19, CCL21
CCR8	CCL1, CCL4, CCL17
CCR9	CCL25
CCR10	CCL27, CCL28
CXCR1	CXCL6, CXCL8
CXCR2	CXCL1, CXCL2, CXCL3, CXCL5, CXCL6, CXCL7, CXCL8
CXCR3	CXCL9, CXCL10, CXCL11
CXCR4	CXCL12
CXCR5	CXCL13
CXCR6	CXCL16
CXCR7	CXCL11, CXCL12
CX3CR1	CX3CL1
XCR1	XCL1, XCL2

Abbreviations: CCL, C-C chemokine motif ligand; CCR, C-C chemokine motif receptor; CXCL, C-X-C chemokine motif ligand; CXCR, C-X-C chemokine motif receptor.

CXCL12 interacts with CXCR4 to retain neutrophils in the BM. *CXCR4* with WHIM-associated mutations encodes a carboxyl terminal truncated receptor with impaired internalization and enhanced activation, resulting in abnormal neutrophil retention in the BM and neutropenia in the blood.^{29–32} Conversely, deletion of *CXCR4* in murine hematopoietic cells causes exacerbated neutrophil mobilization into circulation.^{27,33–37} For example, mice carrying a myeloid-specific deletion of *Cxcr4* (myeloid-specific knock-out (MKO) mice), in which *Cxcr4* in mature neutrophils is deleted in both the BM and blood, display a skewed redistribution of neutrophils from the BM to the blood and spleen.³⁷ Moreover, treatment of humans or mice with an antagonist of *CXCR4* causes rapid neutrophil mobilization into circulation.^{38,39} Similar to MKO mice, mice with a conditional deletion of *Cxcl12* display a significant increase in the number of circulating neutrophils.⁴⁰ Therefore, the *CXCR4* and *CXCL12* axis is critical for the retention of neutrophils in the BM, where granulopoiesis is controlled by granulocyte colony-stimulating factor.²⁵

Secondary lymphoid organs (SLOs)

SLOs include the LNs, spleen and Peyer's patches. In mature SLOs, follicular DCs in B-cell follicles produce *CXCL13*, which promotes homeostatic recruitment of B cells via *CXCR5*. B cells in the marginal zone of the spleen depend on *CXCR7* for physiological positioning. Treatment of mice with *CXCR7* antagonists leads to a decrease of B-cell numbers in the splenic marginal zone.⁴¹ In the T-cell area, fibroblastic reticular cells produce *CCL19*, *CCL21* and *CXCL12*, which localize T cells

Table 3 Atypical chemokine receptors and ligands

Name	Common aliases	Ligands
ACKR1	DARC; Duffy	CCL2, CCL5, CCL11, CCL13, CCL14, CXCL1, CXCL2, CXCL3, CXCL7, CXCL8
ACKR2	D6, CCR9, CCR10	CCL2, CCL3, CCL4, CCL5, CCL7, CCL8, CCL12, CCL13, CCL14, CCL17, CCL22
ACKR3	CXCR7; RDC1	CXCL11, CXCL12
ACKR4	CCRL1; CCX-CKR, CCBP2, CCR11	CCL19, CCL21, CCL25, CXCL13
ACKR5	CKRX, CRAM-A, L-CCR, CRAM-B, HCR, CCR11	CCL5, CCL19, chemerin
ACKR6	Nir1	CCL18

Abbreviations: CCL, C-C chemokine motif ligand; CCR, C-C chemokine motif receptor; CXCL, C-X-C chemokine motif ligand; CXCR, C-X-C chemokine motif receptor.

Table 4 Homeostatic chemokines

Location	Chemokines	Receptors	Function	References
Skin	CCL27	CCR10	Skin homing of T cells	20
Brain	CXCL12	CXCR4	Brain homing of leukocytes	21
Lung	CXCL12	CXCR4	Stromal cell homing	24
Spleen	CCL19, CCL21, CXCL13	CCR7, CXCR5	T cell, B cell and DC homing	27
Small intestine	CCL25	CCR9	Lymphocyte homing	8,9
Secondary lymphoid tissues	CCL21, CCL19, CCL13	CCR7, CXCR5	T cell, B cell and DC homing	22
Bone marrow	CXCL12	CXCR4	Neutrophil retention in BM	25,26

Abbreviations: CCL, C-C chemokine motif ligand; CCR, C-C chemokine motif receptor; CXCL, C-X-C chemokine motif ligand; CXCR, C-X-C chemokine motif receptor; DC, dendritic cell.

and DCs via CCR7 and CXCR4. Thus, even in the absence of an immune response, naive lymphocytes and antigen-presenting cells actively migrate to and remain in SLOs by chemokine and GPCR interactions.⁴

INFLAMMATORY CHEMOKINES

Unlike homeostatic chemokines, inflammatory chemokines are not constitutively expressed at high levels in the absence of tissue damage. Such chemokines are expressed by circulating leukocytes and other cell types only upon activation. Therefore, these molecules are also termed inducible chemokines, the expression of which is often triggered by pro-inflammatory mediators, such as tumor necrosis factor and IFN- γ , and microbial products, including lipopolysaccharide. Inflammatory chemokines are expressed by numerous cell types at almost any tissue location. This expression allows for the relatively transient attraction of inflammatory leukocytes to damaged tissues. Inflammatory chemokines include CCL1–5 and CXCL1–11. In general, CXCL1–8 regulate neutrophil migration and CXCL9–11 regulate activated T-cell migration. Most inflammatory CC chemokines induce chemotaxis of monocytes, macrophages and T cells. Almost all leukocytes express complex patterns of chemokine GPCRs that allow for responses to diverse chemokines that are induced under stress conditions.⁴²

Neutrophil trafficking in inflammation

Neutrophil trafficking in inflammation is guided by multiple chemokine and classical chemoattractant GPCRs that are sequentially expressed by cells in a complex tissue microenvironment. For example, Fprs antecede CXCR2 to rapidly recruit the first wave of neutrophils into the liver of *Listeria*-infected mice in response to bacterial chemoattractant peptides.⁴³ In mouse skin wounds, the leukotriene B4 GPCR, BLT, is critical for neutrophil infiltration, followed by dynamic cell swarming, which requires the participation of the chemokine GPCR CXCR2 and a formyl peptide GPCR Fpr2.⁴⁴ For the normal healing of sterile skin wounds, coordinated neutrophil influx by Fprs and CXCR2 is needed.⁴⁵ Therefore, neutrophil trafficking in inflammatory responses is mediated by chemotactic cues that are established by pathogen-associated molecular patterns (chemotactic PAMPs) and damage-associated molecular patterns (DAMPs) via the cooperation of chemokines as well as classical chemoattractant GPCRs.

Monocytic cell recruitment in inflammation

Monocytes migrate out of the BM by following the CCL2 gradients produced by vascular endothelial cells in response to systemic PAMPs, such as Toll-like receptor (TLR) ligands. In mouse airway inflammation, production of the CCR2 ligand CCL2 and Fpr2 ligand CRAMP is increased in lung tissues. Tightly orchestrated trafficking of monocytic DCs is established by multiple chemoattractant GPCRs, in which CCL2/CCR2 mobilize Ly6C^{high} monocytic DC precursors from the BM into circulation,^{46,47} where cells extravasate into the perivascular regions of the inflamed lung guided by CCL2 and become

immature DCs upon exposure to the PAMPs and DAMPs present in the airway.² Immature DCs subsequently lose functional CCR2, but gain high-level expression of Fpr2, and migrate to peribronchiolar regions in response to the Fpr2 ligand, a host-derived chemotactic DAMP, CRAMP,² which not only forms a chemotactic gradient for DC recruitment but also promotes DC maturation in cooperation with TLR agonistic PAMPs.⁴⁸ A shift of chemoattractant GPCR expression occurs again as DCs mature when Fpr2 is downregulated, with highly elevated expression of the DC-homing chemokine GPCR CCR7, enabling mature DCs to be directed to lymphatic organs. Thus trafficking of monocytic DCs in the inflammatory airway consists of the finely tuned orchestration of chemoattractant GPCRs on the cell surface, initiating from CCR2, with Fpr2 as an intermediate and CCR7 as the final player, which accomplishes the last segment of homing.

CCR2 and CCL2 (MCP-1), as prototype chemokine GPCR and ligand partners, were discovered based on their capacity to mediate monocyte recruitment in inflammation and cancer.⁹ Their participation is represented by a model of acute toxoplasmosis in which mouse resistance depends on the timely appearance of Ly6C^{hi} monocyte-derived effector macrophages via the CCR2/CCL2 interaction. Mice deficient in either the receptor CCR2 or ligand CCL2 were not able to control parasite replication due to a reduction in macrophage-produced interleukin-12 and reactive nitrogen intermediates.^{49,50}

Trafficking of other immune cells

Mast cell precursors migrate from the BM to the periphery via CXCR2, BLT1 and/or CYSLT1, another chemoattractant GPCR, signaling. Eosinophils leave the BM via CCR3, while CCR7 is used by mature tissue-resident DCs to migrate into afferent lymphatic vessels through CCL21 gradients.⁴ Thus, despite the vast number of chemokines and their GPCRs, little redundancy exists in complex pathophysiological processes.

Under normal conditions, CXCR5 mediates the homing of B cells into lymphoid organs. However, in systemic lupus erythematosus (SLE), CXCR4 guides B-cell migration from the dark to the light zone in lymphoid follicle germinal centers in response to locally produced CXCL12. In extrafollicular regions, B cells differentiate into plasma cells to produce anti-dsDNA antibodies. There is an increased CXCR4^{high}CXCR5^{low/-} B-cell population in SLE patients with higher anti-dsDNA titers in circulation. Therefore, dysregulated CXCR4/CXCR5 expression on SLE B cells may skew the translocation of B cells to extrafollicular sites, presumably to allow cells to escape negative selection and apoptosis, leading to more active promotion of the development of SLE.⁵¹ Such a shift in chemokine GPCR levels during disease progression may determine the final destiny of immune cells with the outcome of the disease at stake.

CHEMOKINES AND GPCRS IN HUMAN DISEASES

Chemokines and their GPCRs are associated with an extraordinary array of pathological conditions^{52–55} (Table 5),

including autoimmune disorders^{8,56–61} (for example, psoriasis, rheumatoid arthritis and multiple sclerosis); pulmonary diseases^{62–65} (for example, asthma and chronic obstructive pulmonary disease); transplant rejection^{70,71}; metabolic and vascular diseases^{66–69,78} (for example, obesity, diabetes and atherosclerosis; as well as infectious diseases^{72–77} (for example, HIV, sepsis and inflammatory bowel disease). The active participation of chemokines and GPCRs in disease progression makes these molecules the most prominent targets for drug development.

Table 5 Chemokine GPCRs in inflammation and immunity

Diseases	Chemokine receptors	References
<i>Autoimmune disorder</i>		
Psoriasis	CCR1, CXCR1, CXCR2, CXCR3	56–58
RA	CCR1, CCR2, CXCR1, CXCR2, CXCR3	56–60
MS	CCR1, CCR2, CXCR3	56,57,60,61
<i>Pulmonary diseases</i>		
Asthma	CCR2, CCR3, CCR4, CCR5, CCR6, CCR8, CXCR4, CX3CR1	62,63
COPD	CXCR1, CXCR2	64,65
<i>Metabolic and vascular disease</i>		
Atherosclerosis	CCR2, CX3CR1	66,67
Diabetes	CCR2, CX3CR1	56,68,69
Obesity	CCR2, CCR5, CXCR2	56,68
Transplant rejection	CCR1, CCR2, CCR5, CXCR3	70,71
<i>Infectious diseases</i>		
HIV	CCR5, CXCR4	56,72,73
Sepsis	CXCR1, CXCR2	74,75
IBD	CCR9	56,76,77

Abbreviations: CCR, C-C chemokine motif receptor; COPD, chronic obstructive pulmonary disease; CXCL, C-X-C chemokine motif ligand; CXCR, C-X-C chemokine motif receptor; IBD, inflammatory bowel disease; GPCR, G-protein coupled receptor; HIV, human immunodeficiency virus; MS, multiple sclerosis; RA, rheumatoid arthritis.

Table 6 Chemokine GPCRs in cancer

Chemokine receptors	Cancers	References
CXCR3	Acute lymphoblastic leukemia	85
CXCR4	Breast cancer, ovarian cancer, prostate cancer, pancreatic cancer, melanoma, oesophageal cancer, non-small cell lung cancer, head and neck cancer, bladder cancer, colorectal cancer, osteosarcoma, neuroblastoma, acute lymphoblastic leukemia, chronic lymphocytic leukemia, stomach cancer	85–107
CXCR5	Head and neck cancer	88,108
CCR2	Prostate cancer, glioblastoma, ovarian cancer, breast cancer	109–112
CCR7	Breast cancer, melanoma, non-small cell lung cancer, head and neck cancer, colorectal cancer, chronic lymphocytic leukemia, stomach cancer, non-Hodgkin's lymphoma, T-cell leukemia	113–118
CCR9	Melanoma	119
CCR10	Melanoma	119

Abbreviations: CCR, C-C chemokine motif receptor; CXCR, C-X-C chemokine motif receptor; GPCR, G-protein coupled receptor.

CHEMOKINES AND GPCRS IN CANCER

One of the epoch-making discoveries of chemokines and their GPCRs was the discovery of the ability of these molecules to mediate tumor cell chemotaxis, thus increasing the cell motility required for metastasis. This phenomenon was first reported for the chemokine MCP-1 (CCL2), which mediates organ-specific dissemination of experimental murine lymphoma,^{79,80} an example of tumor cells utilizing the fundamental properties of chemokines for cell progression. This principle was later supported by evidence that chemokine GPCRs mediate metastasis of human tumor cells to distant organs. Since then, chemokines and their GPCRs have been shown to have a key role in the initiation and progression of cancers of multiple organs, including the lungs, colon, liver, breast, cervix, prostate, bladder, ovary, esophagus, skin and lymphatics.^{22,81–84} Many cancer cells express chemokine GPCRs that sense ligands as growth signals and are the driving force for a motile phenotype that enables invasion and metastasis (Table 6). CXCR4 is one of the most frequently identified chemokine GPCRs in cancer and has been implicated in >23 human cancers, including breast,^{86–88} ovary,^{89–91} prostate,^{92,93} pancreas,^{94–96} melanoma,^{97,98} esophagus,⁹⁹ lung,¹⁰⁰ head and neck,¹⁰¹ bladder,¹⁰² colorectum¹⁰³ and stomach cancers¹⁰⁴ as well as osteosarcoma,¹⁰⁵ neuroblastoma,¹⁰⁶ acute lymphoblastic leukemia⁸⁵ and chronic lymphocytic leukemia.¹⁰⁷ Activation of the CXCR4–CXCL12 axis is correlated with angiogenesis,^{90,120–124} proliferation^{124–128} and metastasis of tumors^{129–133} (Table 7). CCR7 is another GPCR that has been frequently implicated in the progression of cancers^{113–118} (Table 8), including breast,¹³⁴ melanoma,^{135,136} lung,¹³⁷ head and neck,¹³⁸ colorectum,¹³⁹ chronic lymphocytic leukemia,¹⁴⁰ stomach,¹⁴¹ non-Hodgkin's lymphoma¹⁴² and T-cell leukemia.¹⁴³ Other chemokine GPCRs, such as CXCR3 in childhood acute lymphoblastic leukemia,⁸⁵ CXCR5 in head and neck cancer¹⁰⁸ and CCR2 in prostate cancer, lymphoma, glioblastoma, ovarian cancer and breast cancer^{109–112} have been described. CCR9 and CCR10¹¹⁹ were also implicated in cancer invasion and melanoma metastasis (Table 6). Notably, some cancers may exploit the function of more than one chemokine GPCR to their advantage, demonstrating the

Table 7 CXCL12–CXCR4 axis in cancer

Function	Mechanism	References
Angiogenesis	Recruiting CXCR4 ⁺ BM-derived monocytes to tumor; stimulating the formation of new tumor blood vessels	85,87,104,134,135
Proliferation	Cross-talking with EGFR in breast cancer	135–139
Metastasis	Promoting EMT by activating MEK/ERK, PI3K/AKT or Wnt/ β -catenin pathway	140–143
Microenvironment	Attracting CXCR4 ⁺ immune cells to tumor sites; stimulating the production of EGF, CCL2, CCL5, IL-1 β and CXCL8 at tumor sites	85

Abbreviations: BM, bone marrow; CCL, C-C chemokine motif ligand; CXCL, C-X-C chemokine motif ligand; CXCR, C-X-C chemokine motif receptor; EGFR, epidermal growth factor receptor; EMT, epithelial–mesenchymal transition; ERK, extracellular signal–regulated kinase; IL, interleukin; MEK, mitogen-activated extracellular signal–regulated kinase; PI3K, phosphoinositide-3 kinase.

Table 8 CCL21–CCR7 axis in cancer

Cancer	Mechanism	References
Colon cancer	Upregulating MMP-9	113,114
B-cell chronic leukemia		
T24 bladder cancer		
Breast cancer	Upregulating VEGF-C	118
Non-small cell lung cancer	Upregulation of VEGF-D	115
Esophageal squamous cell carcinoma	Stimulating MUC1 expression	116
Pancreatic cancer	Activating EMT and Erk/NF- κ B pathway	117

Abbreviations: CCL, C-C chemokine motif ligand; CCR, C-C chemokine motif receptor; EMT, epithelial–mesenchymal transition; Erk, extracellular signal–regulated kinase; MMP, matrix metalloproteinase; MUC1, mucin 1; NF, nuclear factor; VEGF, vascular endothelial growth factor.

delicate evasion of cancer cells from host defense. The bright spot, in this respect, is in the experimental system, as targeting chemokines or their GPCRs has shown beneficial effects on cancer therapy.

THE FINE-TUNED ORCHESTRATION OF CHEMOKINE SYNERGY IN THE INFLAMMATORY RESPONSES

Directional migration and arrest of leukocytes during homeostasis, inflammatory responses and tumor development are delicately controlled by a multitude of chemokines. Immune cell trafficking to inflamed tissues depends on at least three distinct but closely related processes: (1) chemoattractant GPCR expression by migrating leukocytes; (2) gain or loss of GPCR responses regulated by environmental stimuli or homologous and heterologous receptor desensitization; and (3) presence of a dominant ligand gradient in a given anatomical compartment at a given time.^{144–148}

Mounting evidence indicates that many GPCRs, including chemokine GPCRs (that is, CCR2, CCR5, CXCR1, CXCR2 and CXCR4), may function as homodimers or oligomers.¹⁴⁹ Heterodimers between CCR2 and CCR5 or CXCR4,^{150–152} as well as between CXCR1 and CXCR2,¹⁵³ have been reported. The properties of chemokine GPCRs may enable cells to enhance their responsiveness to one or more ligands produced in the vicinity. However, chemokine ligand heterocomplexes may act either as receptor antagonists or exhibit synergy by simultaneously or sequentially activating more than one GPCR.

This synergy is selective for chemokines that are present in the tissue environment, often at lower concentrations. For example, CCL21 and CCL19 may each form heterocomplexes with CXCL13 to activate CCR7 at lower ligand concentrations,¹⁵⁴ CCL22 and CXCL10 complexes more potently activate CXCR3 to enhance migration of CCR4⁺ lymphocytes,¹⁵⁵ CCL19 and CCL21 may each interact with CCL7 to enhance monocyte chemotaxis and prevent CCL7 degradation mediated by atypical chemokine receptor 2;¹⁵⁶ and heterocomplexes of CXCL9 and CXCL12 enhance recruitment of CD8⁺ T cells¹⁵⁷ (Table 9). Therefore, both chemokine GPCRs and their ligands have unique properties, acting singularly or in combinations, to exert functions that should benefit the host defense in complex disease states that call for the optimal and differential mobilization of subsets of leukocytes.

The biological activity of chemokines is also complemented by the pro-inflammatory molecule high-mobility group box 1 protein (HMGB1), which was initially demonstrated to interact with multiple cellular receptors, including TLR4 and receptor for advanced glycation end products. However, HMGB1 attenuates the chemotactic responses of mouse embryonic fibroblasts to the chemokine CXCL12 by blocking the receptor CXCR4 or neutralizing CXCL12, suggesting that the action of HMGB1 requires the expression of cell surface CXCR4 as well as the concomitant presence of CXCL12.^{6,158,159} Interestingly, HMGB1 and CXCL12 form heterocomplexes and exhibit a monocyte chemotactic activity that is more potent than that of CXCL12 alone. Mechanistically, HMGB1 increases binding of CXCL12 to CXCR4 by fixing its N-terminal domain in a conformation that is better suited for receptor activation.^{6,160}

Some chemokine ligands possess sugar moieties that bind glycosaminoglycans (GAGs), which immobilize chemokines on the endothelial surface to ensure the directionality of the chemotactic signals for leukocytes.^{161–163} Chemokines that fail to be immobilized by endothelial GAGs may either be dispersed from the production site or diluted to concentrations below the receptor activating threshold, thereby reducing leukocyte extravasation into tissues. Thus microenvironment GAGs are pivotal to the spatial distribution of chemokine gradients that are necessary for proper leukocyte recruitment.¹⁶⁴

Table 9 Fine tuning of leukocyte trafficking by chemokine complexes

Chemokine heterocomplexes	Functions	References
CCL21 or CCL19+CXCL13	Enhance binding and activating CCR7	145–147
CCL19+CCL21	Enhance the activity of CCR2 ligands and protection from degradation	145–147
CCL22+CCL4	Enhance CCR4 activation	153
CCL22+CXCL10	Enhance CXCL10 binding to CXCR3 or GAGs; stimulate the migration of CCR4-bearing lymphocytes	147
CCL21+CCL19+CCL7	Enhance monocyte migration and prevent CCL7 degradation	147
CXCL9+CXCL12	Enhance recruitment of CD8 ⁺ T cells	155
CXCR4+CXCL12	Enhance cell migration in response to HMGB1	158
HMGB1+CXCL12	HMGB1 promotes CXCL12 binding and activation of CXCR4	156

Abbreviations: CCL, C-C chemokine motif ligand; CXCL, C-X-C chemokine motif ligand; CCR, C-C chemokine motif receptor; CXCR, C-X-C chemokine motif receptor; GAG, glycosaminoglycan; HMGB1, high-mobility group box 1 protein.

PERSPECTIVES

Chemokines and GPCRs are appreciated not only as important mediators for innate immune cell trafficking in response to acute inflammatory insults but also as central cellular fate determinants of adaptive immune responses. There has been rapid progress in the understanding of the biology of chemokines and their GPCRs in the initiation and progression of diseases. The discovery that selected chemokine GPCRs are also expressed on virtually all nonhemopoietic cell types, such as endothelial, epithelial and neuronal and muscular cells, has led to the recognition of the importance of these molecules in more pathophysiological processes, such as angiogenesis, organ development and tumorigenesis.

Despite the expanded knowledge of chemokines and GPCRs generated in past decades, questions remain concerning their involvement in each stage of human disease. Taking cancer as an example, CXCL12–CXCR4 is one of the chemokine/GPCR axes that attract intense attention to its therapeutic potential. Given the recent approval of CXCR4 antagonists for clinical use,¹⁶⁵ it is anticipated that some of these drug leads may soon be tested in cancer patients, particularly those with metastasis. However, owing to the diversity of cancers, it is unlikely that these findings will be able to be generalized. Therefore, more studies are needed to elucidate the whole picture of chemokines and their GPCRs in the growth of primary lesions and establishment of metastatic foci. In this context, it is possible that homeostatic chemokines may directly stimulate the viability and growth of nascent metastatic lesions. Another area in need of rigorous investigation is the biology of atypical chemokine GPCRs, such as D6 and DARC, in tumorigenesis and the metastatic potential of cancer cells. Therefore, recombinant chemokine decoy receptor proteins may have the potential to act as sinks to eliminate excessive chemokine ligands that may stimulate tumor growth and metastasis.

In summary, >20 years after the first discovery of chemokines and their GPCRs, the research field continues to attract a burgeoning army of participants with diverse interests in both basic and clinical research covering almost all aspects of inflammation, immunity, infection and cancer. Thus one should be extremely optimistic about the future benefit to human health derived from more rigorous studies.

CONFLICT OF INTEREST

The authors declare no conflict of interest.

ACKNOWLEDGEMENTS

We thank Ms S Livingstone and Ms C Lamb for secretarial assistance. This project was funded in part by federal funds from the National Cancer Institute, National Institutes of Health, under Contract No. HHSN261200800001E and was supported in part by the Intramural Research Program of the NCI, NIH.

- 1 Le Y, Zhou Y, Iribarren P, Wang J. Chemokines and chemokine receptors: their manifold roles in homeostasis and disease. *Cell Mol Immunol* 2004; **1**: 95–104.
- 2 Chen K, Liu M, Liu Y, Wang C, Yoshimura T, Gong W *et al*. Signal relay by CC chemokine receptor 2 (CCR2) and formylpeptide receptor 2 (Fpr2) in the recruitment of monocyte-derived dendritic cells in allergic airway inflammation. *J Biol Chem* 2013; **288**: 16262–16273.
- 3 Nagarsheth N, Wicha MS, Zou W. Chemokines in the cancer microenvironment and their relevance in cancer immunotherapy. *Nat Rev Immunol* 2017; **17**: 559–572.
- 4 Griffith JW, Sokol CL, Luster AD. Chemokines and chemokine receptors: positioning cells for host defense and immunity. *Annu Rev Immunol* 2014; **32**: 659–702.
- 5 Schulz O, Hammerschmidt SI, Moschovakis GL, Forster R. Chemokines and chemokine receptors in lymphoid tissue dynamics. *Annu Rev Immunol* 2016; **34**: 203–242.
- 6 Cecchinato V, D'Agostino G, Raeli L, Ugucioni M. Chemokine interaction with synergy-inducing molecules: fine tuning modulation of cell trafficking. *J Leukoc Biol* 2016; **99**: 851–855.
- 7 Castan L, Magnan A, Bouchaud G. Chemokine receptors in allergic diseases. *Allergy* 2017; **72**: 682–690.
- 8 Cheng W, Chen G. Chemokines and chemokine receptors in multiple sclerosis. *Mediators Inflamm* 2014; **2014**: 659206.
- 9 Zhou J, Xiang Y, Yoshimura T, Chen K, Gong W, Huang J *et al*. The role of chemoattractant receptors in shaping the tumor microenvironment. *Biomed Res Int* 2014; **2014**: 751392.
- 10 Yoshimura T, Oppenheim JJ. Chemokine-like receptor 1 (CMKLR1) and chemokine (C-C motif) receptor-like 2 (CCRL2); two multifunctional receptors with unusual properties. *Exp Cell Res* 2011; **317**: 674–684.
- 11 Sozzani S, Vermi W, Del Prete A, Facchetti F. Trafficking properties of plasmacytoid dendritic cells in health and disease. *Trends Immunol* 2010; **31**: 270–277.
- 12 Samson M, Edinger AL, Stordeur P, Rucker J, Verhasselt V, Sharron M *et al*. ChemR23, a putative chemoattractant receptor, is expressed in monocyte-derived dendritic cells and macrophages and is a coreceptor for SIV and some primary HIV-1 strains. *Eur J Immunol* 1998; **28**: 1689–1700.

- 13 Gao L, Faibish D, Fredman G, Herrera BS, Chiang N, Serhan CN *et al*. Resolvin E1 and chemokine-like receptor 1 mediate bone preservation. *J Immunol* 2013; **190**: 689–694.
- 14 Bondue B, Vosters O, de Nadai P, Glineur S, De Henau O, Luangsay S *et al*. ChemR23 dampens lung inflammation and enhances anti-viral immunity in a mouse model of acute viral pneumonia. *PLoS Pathog* 2011; **7**: e1002358.
- 15 Huang J, Chen K, Gong W, Dunlop NM, Wang JM. G-protein coupled chemoattractant receptors and cancer. *Front Biosci* 2008; **13**: 3352–3363.
- 16 Fox JC, Nakayama T, Tyler RC, Sander TL, Yoshie O, Volkman BF. Structural and agonist properties of XCL2, the other member of the C-chemokine subfamily. *Cytokine* 2015; **71**: 302–311.
- 17 Bachelier F, Graham GJ, Locati M, Mantovani A, Murphy PM, Nibbs R *et al*. New nomenclature for atypical chemokine receptors. *Nat Immunol* 2014; **15**: 207–208.
- 18 Bachelier F, Ben-Baruch A, Burkhardt AM, Combadiere C, Farber JM, Graham GJ *et al*. International Union of Basic and Clinical Pharmacology. [corrected]. LXXXIX. Update on the extended family of chemokine receptors and introducing a new nomenclature for atypical chemokine receptors. *Pharmacol Rev* 2014; **66**: 1–79.
- 19 Le Y, Cui Y, Iribarren P, Ying G, Wang JM. Manipulating chemoattractant and receptor genes. *In Vivo* 2002; **16**: 1–23.
- 20 Homey B, Alenius H, Muller A, Soto H, Bowman EP, Yuan W *et al*. CCL27-CCR10 interactions regulate T cell-mediated skin inflammation. *Nat Med* 2002; **8**: 157–165.
- 21 Williams JL, Holman DW, Klein RS. Chemokines in the balance: maintenance of homeostasis and protection at CNS barriers. *Front Cell Neurosci* 2014; **8**: 154.
- 22 Zlotnik A, Burkhardt AM, Homey B. Homeostatic chemokine receptors and organ-specific metastasis. *Nat Rev Immunol* 2011; **11**: 597–606.
- 23 Svensson M, Agace WW. Role of CCL25/CCR9 in immune homeostasis and disease. *Expert Rev Clin Immunol* 2006; **2**: 759–773.
- 24 Ruiz EJ, Oeztuerk-Winder F, Ventura JJ. A paracrine network regulates the cross-talk between human lung stem cells and the stroma. *Nat Commun* 2014; **5**: 3175.
- 25 Strydom N, Rankin SM. Regulation of circulating neutrophil numbers under homeostasis and in disease. *J Innate Immun* 2013; **5**: 304–314.
- 26 Eash KJ, Greenbaum AM, Gopalan PK, Link DC. CXCR2 and CXCR4 antagonistically regulate neutrophil trafficking from murine bone marrow. *J Clin Invest* 2010; **120**: 2423–2431.
- 27 Tachibana K, Hirota S, Iizasa H, Yoshida H, Kawabata K, Kataoka Y *et al*. The chemokine receptor CXCR4 is essential for vascularization of the gastrointestinal tract. *Nature* 1998; **393**: 591–594.
- 28 Rankin SM. The bone marrow: a site of neutrophil clearance. *J Leukoc Biol* 2010; **88**: 241–251.
- 29 Gulino AV, Moratto D, Sozzani S, Cavadini P, Otero K, Tassone L *et al*. Altered leukocyte response to CXCL12 in patients with warts hypogammaglobulinemia, infections, myelokathexis (WHIM) syndrome. *Blood* 2004; **104**: 444–452.
- 30 Balabanian K, Lagane B, Pablos JL, Laurent L, Planchenault T, Verola O *et al*. WHIM syndromes with different genetic anomalies are accounted for by impaired CXCR4 desensitization to CXCL12. *Blood* 2005; **105**: 2449–2457.
- 31 Kawai T, Choi U, Whiting-Theobald NL, Linton GF, Brenner S, Sechler JM *et al*. Enhanced function with decreased internalization of carboxy-terminus truncated CXCR4 responsible for WHIM syndrome. *Exp Hematol* 2005; **33**: 460–468.
- 32 Kawai T, Choi U, Cardwell L, DeRavin SS, Naumann N, Whiting-Theobald NL *et al*. WHIM syndrome myelokathexis reproduced in the NOD/SCID mouse xenotransplant model engrafted with healthy human stem cells transduced with C-terminus-truncated CXCR4. *Blood* 2007; **109**: 78–84.
- 33 Zou YR, Kottmann AH, Kuroda M, Taniuchi I, Littman DR. Function of the chemokine receptor CXCR4 in haematopoiesis and in cerebellar development. *Nature* 1998; **393**: 595–599.
- 34 Ma Q, Jones D, Borghesani PR, Segal RA, Nagasawa T, Kishimoto T *et al*. Impaired B-lymphopoiesis, myelopoiesis, and derailed cerebellar neuron migration in CXCR4- and SDF-1-deficient mice. *Proc Natl Acad Sci USA* 1998; **95**: 9448–9453.
- 35 Ma Q, Jones D, Springer TA. The chemokine receptor CXCR4 is required for the retention of B lineage and granulocytic precursors within the bone marrow microenvironment. *Immunity* 1999; **10**: 463–471.
- 36 Kawabata K, Ujikawa M, Egawa T, Kawamoto H, Tachibana K, Iizasa H *et al*. A cell-autonomous requirement for CXCR4 in long-term lymphoid and myeloid reconstitution. *Proc Natl Acad Sci USA* 1999; **96**: 5663–5667.
- 37 Eash KJ, Means JM, White DW, Link DC. CXCR4 is a key regulator of neutrophil release from the bone marrow under basal and stress granulopoiesis conditions. *Blood* 2009; **113**: 4711–4719.
- 38 Liles WC, Broxmeyer HE, Rodger E, Wood B, Hubel K, Cooper S *et al*. Mobilization of hematopoietic progenitor cells in healthy volunteers by AMD3100, a CXCR4 antagonist. *Blood* 2003; **102**: 2728–2730.
- 39 Broxmeyer HE, Orschell CM, Clapp DW, Hangoc G, Cooper S, Plett PA *et al*. Rapid mobilization of murine and human hematopoietic stem and progenitor cells with AMD3100, a CXCR4 antagonist. *J Exp Med* 2005; **201**: 1307–1318.
- 40 Tzeng YS, Li H, Kang YL, Chen WC, Cheng WC, Lai DM. Loss of Cxcl12/Sdf-1 in adult mice decreases the quiescent state of hematopoietic stem/progenitor cells and alters the pattern of hematopoietic regeneration after myelosuppression. *Blood* 2011; **117**: 429–439.
- 41 Wang H, Beaty N, Chen S, Qi CF, Masiuk M, Shin DM *et al*. The CXCR7 chemokine receptor promotes B-cell retention in the splenic marginal zone and serves as a sink for CXCL12. *Blood* 2012; **119**: 465–468.
- 42 Graham GJ. D6 and the atypical chemokine receptor family: novel regulators of immune and inflammatory processes. *Eur J Immunol* 2009; **39**: 342–351.
- 43 Liu M, Chen K, Yoshimura T, Liu Y, Gong W, Wang A *et al*. Formylpeptide receptors are critical for rapid neutrophil mobilization in host defense against *Listeria monocytogenes*. *Sci Rep* 2012; **2**: 786.
- 44 McDonald B, Pittman K, Menezes GB, Hirota SA, Slaba I, Waterhouse CC *et al*. Intravascular danger signals guide neutrophils to sites of sterile inflammation. *Science* 2010; **330**: 362–366.
- 45 Liu M, Chen K, Yoshimura T, Liu Y, Gong W, Le Y *et al*. Formylpeptide receptors mediate rapid neutrophil mobilization to accelerate wound healing. *PLoS ONE* 2014; **9**: e90613.
- 46 Provoost S, Maes T, Joos GF, Tournoy KG. Monocyte-derived dendritic cell recruitment and allergic T(H)2 responses after exposure to diesel particles are CCR2 dependent. *J Allergy Clin Immunol* 2012; **129**: 483–491.
- 47 Serbina NV, Pamer EG. Monocyte emigration from bone marrow during bacterial infection requires signals mediated by chemokine receptor CCR2. *Nat Immunol* 2006; **7**: 311–317.
- 48 Chen K, Xiang Y, Huang J, Gong W, Yoshimura T, Jiang Q *et al*. The formylpeptide receptor 2 (Fpr2) and its endogenous ligand cathelin-related antimicrobial peptide (CRAMP) promote dendritic cell maturation. *J Biol Chem* 2014; **289**: 17553–17563.
- 49 Robben PM, LaRegina M, Kuziel WA, Sibley LD. Recruitment of Gr-1+ monocytes is essential for control of acute toxoplasmosis. *J Exp Med* 2005; **201**: 1761–1769.
- 50 Mordue DG, Sibley LD. A novel population of Gr-1+ activated macrophages induced during acute toxoplasmosis. *J Leukoc Biol* 2003; **74**: 1015–1025.
- 51 Zhao LD, Liang D, Wu XN, Li Y, Niu JW, Zhou C *et al*. Contribution and underlying mechanisms of CXCR4 overexpression in patients with systemic lupus erythematosus. *Cell Mol Immunol* 2017; **14**: 842–849.
- 52 Baggolini M. Chemokines in pathology and medicine. *J Intern Med* 2001; **250**: 91–104.
- 53 Proudfoot AE. Chemokine receptors: multifaceted therapeutic targets. *Nat Rev Immunol* 2002; **2**: 106–115.
- 54 Gerard C, Rollins BJ. Chemokines and disease. *Nat Immunol* 2001; **2**: 108–115.
- 55 Allen SJ, Crown SE, Handel TM. Chemokine: receptor structure, interactions, and antagonism. *Annu Rev Immunol* 2007; **25**: 787–820.
- 56 Kunkel SL, Godessart N. Chemokines in autoimmunity: from pathology to therapeutics. *Autoimmun Rev* 2002; **1**: 313–320.
- 57 Godessart N, Kunkel SL. Chemokines in autoimmune disease. *Curr Opin Immunol* 2001; **13**: 670–675.
- 58 Rottman JB, Smith TL, Ganley KG, Kikuchi T, Krueger JG. Potential role of the chemokine receptors CXCR3, CCR4, and the integrin

- alphaEbeta7 in the pathogenesis of psoriasis vulgaris. *Lab Invest* 2001; **81**: 335–347.
- 59 Bruhl H, Wagner K, Kellner H, Schattenkirchner M, Schlondorff D, Mack M. Surface expression of CC- and CXC-chemokine receptors on leucocyte subsets in inflammatory joint diseases. *Clin Exp Immunol* 2001; **126**: 551–559.
- 60 Szekanecz Z, Kim J, Koch AE. Chemokines and chemokine receptors in rheumatoid arthritis. *Semin Immunol* 2003; **15**: 15–21.
- 61 Sorensen TL, Tani M, Jensen J, Pierce V, Lucchinetti C, Folcik VA et al. Expression of specific chemokines and chemokine receptors in the central nervous system of multiple sclerosis patients. *J Clin Invest* 1999; **103**: 807–815.
- 62 Lukacs NW. Role of chemokines in the pathogenesis of asthma. *Nat Rev Immunol* 2001; **1**: 108–116.
- 63 Mellado M, Martin de Ana A, Gomez L, Martinez C, Rodriguez-Frade JM. Chemokine receptor 2 blockade prevents asthma in a cynomolgus monkey model. *J Pharmacol Exp Ther* 2008; **324**: 769–775.
- 64 Panina P, Mariani M, D'Ambrosio D. Chemokine receptors in chronic obstructive pulmonary disease (COPD). *Curr Drug Targets* 2006; **7**: 669–674.
- 65 Donnelly LE, Barnes PJ. Chemokine receptors as therapeutic targets in chronic obstructive pulmonary disease. *Trends Pharmacol Sci* 2006; **27**: 546–553.
- 66 Charo IF, Taubman MB. Chemokines in the pathogenesis of vascular disease. *Circ Res* 2004; **95**: 858–866.
- 67 Gosling J, Slaymaker S, Gu L, Tseng S, Zlot CH, Young SG et al. MCP-1 deficiency reduces susceptibility to atherosclerosis in mice that overexpress human apolipoprotein B. *J Clin Invest* 1999; **103**: 773–778.
- 68 Awad AS, Kinsey GR, Khutishvili K, Gao T, Bolton WK, Okusa MD. Monocyte/macrophage chemokine receptor CCR2 mediates diabetic renal injury. *Am J Physiol Renal Physiol* 2011; **301**: F1358–F1366.
- 69 Solomon M, Balasa B, Sarvetnick N. CCR2 and CCR5 chemokine receptors differentially influence the development of autoimmune diabetes in the NOD mouse. *Autoimmunity* 2010; **43**: 156–163.
- 70 Nelson PJ, Krensky AM. Chemokines, chemokine receptors, and allograft rejection. *Immunity* 2001; **14**: 377–386.
- 71 Inston NG, Cockwell P. The evolving role of chemokines and their receptors in acute allograft rejection. *Nephrol Dial Transplant* 2002; **17**: 1374–1379.
- 72 Zaitseva M, Peden K, Golding H. HIV coreceptors: role of structure, posttranslational modifications, and internalization in viral-cell fusion and as targets for entry inhibitors. *Biochim Biophys Acta* 2003; **1614**: 51–61.
- 73 Lusso P. HIV and the chemokine system: 10 years later. *EMBO J* 2006; **25**: 447–456.
- 74 Cummings CJ, Martin TR, Frevert CW, Quan JM, Wong VA, Mongovin SM et al. Expression and function of the chemokine receptors CXCR1 and CXCR2 in sepsis. *J Immunol* 1999; **162**: 2341–2346.
- 75 Juffermans NP, Dekkers PE, Peppelenbosch MP, Speelman P, van Deventer SJ, van Der Poll T. Expression of the chemokine receptors CXCR1 and CXCR2 on granulocytes in human endotoxemia and tuberculosis: involvement of the p38 mitogen-activated protein kinase pathway. *J Infect Dis* 2000; **182**: 888–894.
- 76 Eberhardson M, Marits P, Jones M, Jones P, Karlen P, Karlsson M et al. Treatment of inflammatory bowel disease by chemokine receptor-targeted leukapheresis. *Clin Immunol* 2013; **149**: 73–82.
- 77 Zimmermann NP, Vongsa RA, Wendt MK, Dwinell MB. Chemokines and chemokine receptors in mucosal homeostasis at the intestinal epithelial barrier in inflammatory bowel disease. *Inflamm Bowel Dis* 2008; **14**: 1000–1011.
- 78 Xu L, Kitade H, Ni Y, Ota T. Roles of chemokines and chemokine receptors in obesity-associated insulin resistance and nonalcoholic fatty liver disease. *Biomolecules* 2015; **5**: 1563–1579.
- 79 Wang JM, Chertov O, Proost P, Li JJ, Menton P, Xu L et al. Purification and identification of chemokines potentially involved in kidney-specific metastasis by a murine lymphoma variant: induction of migration and NFkappaB activation. *Int J Cancer* 1998; **75**: 900–907.
- 80 Wang JM, Deng X, Gong W, Su S. Chemokines and their role in tumor growth and metastasis. *J Immunol Methods* 1998; **220**: 1–17.
- 81 Balkwill F, Charles KA, Mantovani A. Smoldering and polarized inflammation in the initiation and promotion of malignant disease. *Cancer Cell* 2005; **7**: 211–217.
- 82 Mantovani A. Cancer: inflaming metastasis. *Nature* 2009; **457**: 36–37.
- 83 Ali S, Lazennec G. Chemokines: novel targets for breast cancer metastasis. *Cancer Metastasis Rev* 2007; **26**: 401–420.
- 84 Vindrieux D, Escobar P, Lazennec G. Emerging roles of chemokines in prostate cancer. *Endocr Relat Cancer* 2009; **16**: 663–673.
- 85 Wu S, Gessner R, Taube T, Korte A, von Stackelberg A, Kirchner R et al. Chemokine IL-8 and chemokine receptor CXCR3 and CXCR4 gene expression in childhood acute lymphoblastic leukemia at first relapse. *J Pediatr Hematol Oncol* 2006; **28**: 216–220.
- 86 Zlotnik A, Yoshie O, Nomiya H. The chemokine and chemokine receptor superfamilies and their molecular evolution. *Genome Biol* 2006; **7**: 243.
- 87 Liang Z, Wu T, Lou H, Yu X, Taichman RS, Lau SK et al. Inhibition of breast cancer metastasis by selective synthetic polypeptide against CXCR4. *Cancer Res* 2004; **64**: 4302–4308.
- 88 Li YM, Pan Y, Wei Y, Cheng X, Zhou BP, Tan M et al. Upregulation of CXCR4 is essential for HER2-mediated tumor metastasis. *Cancer Cell* 2004; **6**: 459–469.
- 89 Kajiyama H, Shibata K, Terauchi M, Ino K, Nawa A, Kikkawa F. Involvement of SDF-1alpha/CXCR4 axis in the enhanced peritoneal metastasis of epithelial ovarian carcinoma. *Int J Cancer* 2008; **122**: 91–99.
- 90 Jiang YP, Wu XH, Shi B, Wu WX, Yin GR. Expression of chemokine CXCL12 and its receptor CXCR4 in human epithelial ovarian cancer: an independent prognostic factor for tumor progression. *Gynecol Oncol* 2006; **103**: 226–233.
- 91 Oda Y, Ohishi Y, Basaki Y, Kobayashi H, Hirakawa T, Wake N et al. Prognostic implications of the nuclear localization of Y-box-binding protein-1 and CXCR4 expression in ovarian cancer: their correlation with activated Akt, LRP/MVP and P-glycoprotein expression. *Cancer Sci* 2007; **98**: 1020–1026.
- 92 Akashi T, Koizumi K, Tsuneyama K, Saiki I, Takano Y, Fuse H. Chemokine receptor CXCR4 expression and prognosis in patients with metastatic prostate cancer. *Cancer Sci* 2008; **99**: 539–542.
- 93 Hirata H, Hinoda Y, Kikuno N, Kawamoto K, Dahiya AV, Suehiro Y et al. CXCL12 G801A polymorphism is a risk factor for sporadic prostate cancer susceptibility. *Clin Cancer Res* 2007; **13**: 5056–5062.
- 94 Liang JJ, Zhu S, Bruggeman R, Zaino RJ, Evans DB, Fleming JB et al. High levels of expression of human stromal cell-derived factor-1 are associated with worse prognosis in patients with stage II pancreatic ductal adenocarcinoma. *Cancer Epidemiol Biomarkers Prev* 2010; **19**: 2598–2604.
- 95 Marechal R, Demetter P, Nagy N, Berton A, Decaestecker C, Polus M et al. High expression of CXCR4 may predict poor survival in resected pancreatic adenocarcinoma. *Br J Cancer* 2009; **100**: 1444–1451.
- 96 Marchesi F, Monti P, Leone BE, Zerbi A, Vecchi A, Piemonti L et al. Increased survival, proliferation, and migration in metastatic human pancreatic tumor cells expressing functional CXCR4. *Cancer Res* 2004; **64**: 8420–8427.
- 97 Franco R, Cantile M, Scala S, Catalano E, Cerrone M, Scognamiglio G et al. Histomorphologic parameters and CXCR4 mRNA and protein expression in sentinel node melanoma metastasis are correlated to clinical outcome. *Cancer Biol Ther* 2010; **9**: 423–429.
- 98 Scala S, Giuliano P, Ascierto PA, Ierano C, Franco R, Napolitano M et al. Human melanoma metastases express functional CXCR4. *Clin Cancer Res* 2006; **12**: 2427–2433.
- 99 Kaifi JT, Yekebas EF, Schurr P, Obonyo D, Wachowiak R, Busch P et al. Tumor-cell homing to lymph nodes and bone marrow and CXCR4 expression in esophageal cancer. *J Natl Cancer Inst* 2005; **97**: 1840–1847.
- 100 Liu K, Bao C, Yao N, Miao C, Varlotto J, Sun Q et al. Expression of CXCR4 and non-small cell lung cancer prognosis: a meta-analysis. *Int J Clin Exp Med* 2015; **8**: 7435–7445.
- 101 Katayama A, Ogino T, Bandoh N, Nonaka S, Harabuchi Y. Expression of CXCR4 and its down-regulation by IFN-gamma in head and neck squamous cell carcinoma. *Clin Cancer Res* 2005; **11**: 2937–2946.
- 102 Eisenhardt A, Frey U, Tack M, Roskopf D, Lummen G, Rubben H et al. Expression analysis and potential functional role of the CXCR4 chemokine receptor in bladder cancer. *Eur Urol* 2005; **47**: 111–117.
- 103 Kim J, Mori T, Chen SL, Amersi FF, Martinez SR, Kuo C et al. Chemokine receptor CXCR4 expression in patients with melanoma

- and colorectal cancer liver metastases and the association with disease outcome. *Ann Surg* 2006; **244**: 113–120.
- 104 Yasumoto K, Koizumi K, Kawashima A, Saitoh Y, Arita Y, Shinohara K *et al*. Role of the CXCL12/CXCR4 axis in peritoneal carcinomatosis of gastric cancer. *Cancer Res* 2006; **66**: 2181–2187.
- 105 Oda Y, Yamamoto H, Tamiya S, Matsuda S, Tanaka K, Yokoyama R *et al*. CXCR4 and VEGF expression in the primary site and the metastatic site of human osteosarcoma: analysis within a group of patients, all of whom developed lung metastasis. *Mod Pathol* 2006; **19**: 738–745.
- 106 Russell HV, Hicks J, Okcu MF, Nuchtern JG. CXCR4 expression in neuroblastoma primary tumors is associated with clinical presentation of bone and bone marrow metastases. *J Pediatr Surg* 2004; **39**: 1506–1511.
- 107 O'Hayre M, Salanga CL, Kipps TJ, Messmer D, Dorrestein PC, Handel TM. Elucidating the CXCL12/CXCR4 signaling network in chronic lymphocytic leukemia through phosphoproteomics analysis. *PLoS ONE* 2010; **5**: e11716.
- 108 Muller A, Sonkoly E, Eulert C, Gerber PA, Kubitzka R, Schirlau K *et al*. Chemokine receptors in head and neck cancer: association with metastatic spread and regulation during chemotherapy. *Int J Cancer* 2006; **118**: 2147–2157.
- 109 Lu Y, Chen Q, Corey E, Xie W, Fan J, Mizokami A *et al*. Activation of MCP-1/CCR2 axis promotes prostate cancer growth in bone. *Clin Exp Metastasis* 2009; **26**: 161–169.
- 110 Liang Y, Bollen AW, Gupta N. CC chemokine receptor-2A is frequently overexpressed in glioblastoma. *J Neurooncol* 2008; **86**: 153–163.
- 111 Negus RP, Stamp GW, Relf MG, Burke F, Malik ST, Bernasconi S *et al*. The detection and localization of monocyte chemoattractant protein-1 (MCP-1) in human ovarian cancer. *J Clin Invest* 1995; **95**: 2391–2396.
- 112 Li S, Lu J, Chen Y, Xiong N, Li L, Zhang J *et al*. MCP-1-induced ERK/GSK-3beta/Snail signaling facilitates the epithelial-mesenchymal transition and promotes the migration of MCF-7 human breast carcinoma cells. *Cell Mol Immunol* 2017; **14**: 621–630.
- 113 Li J, Sun R, Tao K, Wang G. The CCL21/CCR7 pathway plays a key role in human colon cancer metastasis through regulation of matrix metalloproteinase-9. *Dig Liver Dis* 2011; **43**: 40–47.
- 114 Mishan MA, Ahmadiankia N, Bahrami AR. CXCR4 and CCR7: two eligible targets in targeted cancer therapy. *Cell Biol Int* 2016; **40**: 955–967.
- 115 Sun L, Zhang Q, Li Y, Tang N, Qiu X. CCL21/CCR7 up-regulate vascular endothelial growth factor-D expression via ERK pathway in human non-small cell lung cancer cells. *Int J Clin Exp Pathol* 2015; **8**: 15729–15738.
- 116 Shi M, Chen D, Yang D, Liu XY. CCL21-CCR7 promotes the lymph node metastasis of esophageal squamous cell carcinoma by up-regulating MUC1. *J Exp Clin Cancer Res* 2015; **34**: 149.
- 117 Zhang L, Wang D, Li Y, Liu Y, Xie X, Wu Y *et al*. CCL21/CCR7 axis contributed to CD133+ pancreatic cancer stem-like cell metastasis via EMT and Erk/NF-kappaB pathway. *PLoS ONE* 2016; **11**: e0158529.
- 118 Tutunea-Fatan E, Majumder M, Xin X, Lala PK. The role of CCL21/CCR7 chemokine axis in breast cancer-induced lymphangiogenesis. *Mol Cancer* 2015; **14**: 35.
- 119 Kuhnelt-Leddihn L, Muller H, Eisendle K, Zelger B, Weinlich G. Overexpression of the chemokine receptors CXCR4, CCR7, CCR9, and CCR10 in human primary cutaneous melanoma: a potential prognostic value for CCR7 and CCR10? *Arch Dermatol Res* 2012; **304**: 185–193.
- 120 Chow MT, Luster AD. Chemokines in cancer. *Cancer Immunol Res* 2014; **2**: 1125–1131.
- 121 Wang J, Wang J, Sun Y, Song W, Nor JE, Wang CY *et al*. Diverse signaling pathways through the SDF-1/CXCR4 chemokine axis in prostate cancer cell lines leads to altered patterns of cytokine secretion and angiogenesis. *Cell Signal* 2005; **17**: 1578–1592.
- 122 Kryczek I, Lange A, Mottram P, Alvarez X, Cheng P, Hogan M *et al*. CXCL12 and vascular endothelial growth factor synergistically induce neoangiogenesis in human ovarian cancers. *Cancer Res* 2005; **65**: 465–472.
- 123 Wang J, Wang J, Dai J, Jung Y, Wei CL, Wang Y *et al*. A glycolytic mechanism regulating an angiogenic switch in prostate cancer. *Cancer Res* 2007; **67**: 149–159.
- 124 Guo ST, Chi MN, Yang RH, Guo XY, Zan LK, Wang CY *et al*. INPP4B is an oncogenic regulator in human colon cancer. *Oncogene* 2016; **35**: 3049–3061.
- 125 Guo F, Wang Y, Liu J, Mok SC, Xue F, Zhang W. CXCL12/CXCR4: a symbiotic bridge linking cancer cells and their stromal neighbors in oncogenic communication networks. *Oncogene* 2016; **35**: 816–826.
- 126 Thomas RM, Kim J, Revelo-Penafiel MP, Angel R, Dawson DW, Lowy AM. The chemokine receptor CXCR4 is expressed in pancreatic intraepithelial neoplasia. *Gut* 2008; **57**: 1555–1560.
- 127 Porcile C, Bajetto A, Barbieri F, Barbero S, Bonavia R, Biglieri M *et al*. Stromal cell-derived factor-1alpha (SDF-1alpha/CXCL12) stimulates ovarian cancer cell growth through the EGF receptor transactivation. *Exp Cell Res* 2005; **308**: 241–253.
- 128 Porcile C, Bajetto A, Barbero S, Pirani P, Schettini G. CXCR4 activation induces epidermal growth factor receptor transactivation in an ovarian cancer cell line. *Ann NY Acad Sci* 2004; **1030**: 162–169.
- 129 Sun X, Wei L, Chen Q, Terek RM. CXCR4/SDF1 mediate hypoxia induced chondrosarcoma cell invasion through ERK signaling and increased MMP1 expression. *Mol Cancer* 2010; **9**: 17.
- 130 Yang P, Wang G, Huo H, Li Q, Zhao Y, Liu Y. SDF-1/CXCR4 signaling up-regulates survivin to regulate human sacral chondrosarcoma cell cycle and epithelial-mesenchymal transition via ERK and PI3K/AKT pathway. *Med Oncol* 2015; **32**: 377.
- 131 Hu TH, Yao Y, Yu S, Han LL, Wang WJ, Guo H *et al*. SDF-1/CXCR4 promotes epithelial-mesenchymal transition and progression of colorectal cancer by activation of the Wnt/beta-catenin signaling pathway. *Cancer Lett* 2014; **354**: 417–426.
- 132 Li X, Ma Q, Xu Q, Liu H, Lei J, Duan W *et al*. SDF-1/CXCR4 signaling induces pancreatic cancer cell invasion and epithelial-mesenchymal transition in vitro through non-canonical activation of Hedgehog pathway. *Cancer Lett* 2012; **322**: 169–176.
- 133 Li X, Li P, Chang Y, Xu Q, Wu Z, Ma Q *et al*. The SDF-1/CXCR4 axis induces epithelial-mesenchymal transition in hepatocellular carcinoma. *Mol Cell Biochem* 2014; **392**: 77–84.
- 134 Boyle ST, Ingman WV, Poltavets V, Faulkner JW, Whitfield RJ, McColl SR *et al*. The chemokine receptor CCR7 promotes mammary tumorigenesis through amplification of stem-like cells. *Oncogene* 2016; **35**: 105–115.
- 135 Emmett MS, Lanati S, Dunn DB, Stone OA, Bates DO. CCR7 mediates directed growth of melanomas towards lymphatics. *Microcirculation* 2011; **18**: 172–182.
- 136 Fang L, Lee VC, Cha E, Zhang H, Hwang ST. CCR7 regulates B16 murine melanoma cell tumorigenesis in skin. *J Leukoc Biol* 2008; **84**: 965–972.
- 137 Takanami I. Overexpression of CCR7 mRNA in nonsmall cell lung cancer: correlation with lymph node metastasis. *Int J Cancer* 2003; **105**: 186–189.
- 138 Wang J, Xi L, Hunt JL, Gooding W, Whiteside TL, Chen Z *et al*. Expression pattern of chemokine receptor 6 (CCR6) and CCR7 in squamous cell carcinoma of the head and neck identifies a novel metastatic phenotype. *Cancer Res* 2004; **64**: 1861–1866.
- 139 Gunther K, Leier J, Henning G, Dimmler A, Weissbach R, Hohenberger W *et al*. Prediction of lymph node metastasis in colorectal carcinoma by expression of chemokine receptor CCR7. *Int J Cancer* 2005; **116**: 726–733.
- 140 Till KJ, Lin K, Zuzel M, Cawley JC. The chemokine receptor CCR7 and alpha4 integrin are important for migration of chronic lymphocytic leukemia cells into lymph nodes. *Blood* 2002; **99**: 2977–2984.
- 141 Mashino K, Sadanaga N, Yamaguchi H, Tanaka F, Ohta M, Shibuta K *et al*. Expression of chemokine receptor CCR7 is associated with lymph node metastasis of gastric carcinoma. *Cancer Res* 2002; **62**: 2937–2941.
- 142 Yang J, Wang S, Zhao G, Sun B. Effect of chemokine receptors CCR7 on disseminated behavior of human T cell lymphoma: clinical and experimental study. *J Exp Clin Cancer Res* 2011; **30**: 51.
- 143 Buonamici S, Trimarchi T, Ruocco MG, Reavie L, Cathelin S, Mar BG *et al*. CCR7 signalling as an essential regulator of CNS infiltration in T-cell leukaemia. *Nature* 2009; **459**: 1000–1004.
- 144 Nguyen MT, Chen A, Lu WJ, Fan W, Li PP, Oh DY *et al*. Regulation of chemokine and chemokine receptor expression by pPARgamma in adipocytes and macrophages. *PLoS ONE* 2012; **7**: e34976.

- 145 Saccani A, Saccani S, Orlando S, Sironi M, Bernasconi S, Ghezzi P *et al*. Redox regulation of chemokine receptor expression. *Proc Natl Acad Sci USA* 2000; **97**: 2761–2766.
- 146 Stein JV, Nombela-Arrieta C. Chemokine control of lymphocyte trafficking: a general overview. *Immunology* 2005; **116**: 1–12.
- 147 Moser B, Wolf M, Walz A, Loetscher P. Chemokines: multiple levels of leukocyte migration control. *Trends Immunol* 2004; **25**: 75–84.
- 148 Olson TS, Ley K. Chemokines and chemokine receptors in leukocyte trafficking. *Am J Physiol Regul Integr Comp Physiol* 2002; **283**: R7–R28.
- 149 Gouwy M, Schiraldi M, Struyf S, Van Damme J, Uguccioni M. Possible mechanisms involved in chemokine synergy fine tuning the inflammatory response. *Immunol Lett* 2012; **145**: 10–14.
- 150 Mellado M, Rodriguez-Frade JM, Vila-Coro AJ, Fernandez S, Martin de Ana A, Jones DR *et al*. Chemokine receptor homo- or heterodimerization activates distinct signaling pathways. *EMBO J* 2001; **20**: 2497–2507.
- 151 El-Asmar L, Springael JY, Ballet S, Andrieu EU, Vassart G, Parmentier M. Evidence for negative binding cooperativity within CCR5-CCR2b heterodimers. *Mol Pharmacol* 2005; **67**: 460–469.
- 152 Contento RL, Molon B, Boullaran C, Pozzan T, Manes S, Marullo S *et al*. CXCR4-CCR5: a couple modulating T cell functions. *Proc Natl Acad Sci USA* 2008; **105**: 10101–10106.
- 153 Martinez Munoz L, Lucas P, Navarro G, Checa AI, Franco R, Martinez AC *et al*. Dynamic regulation of CXCR1 and CXCR2 homo- and heterodimers. *J Immunol* 2009; **183**: 7337–7346.
- 154 Paoletti S, Petkovic V, Sebastiani S, Danelon MG, Uguccioni M, Gerber BO. A rich chemokine environment strongly enhances leukocyte migration and activities. *Blood* 2005; **105**: 3405–3412.
- 155 Sebastiani S, Danelon G, Gerber B, Uguccioni M. CCL22-induced responses are powerfully enhanced by synergy inducing chemokines via CCR4: evidence for the involvement of first beta-strand of chemokine. *Eur J Immunol* 2005; **35**: 746–756.
- 156 Kuscher K, Danelon G, Paoletti S, Stefano L, Schiraldi M, Petkovic V *et al*. Synergy-inducing chemokines enhance CCR2 ligand activities on monocytes. *Eur J Immunol* 2009; **39**: 1118–1128.
- 157 Venetz D, Ponzoni M, Schiraldi M, Ferreri AJ, Bertoni F, Doglioni C *et al*. Perivascular expression of CXCL9 and CXCL12 in primary central nervous system lymphoma: T-cell infiltration and positioning of malignant B cells. *Int J Cancer* 2010; **127**: 2300–2312.
- 158 Kew RR, Penzo M, Habel DM, Marcu KB. The IKKalpha-dependent NF-kappaB p52/RelB noncanonical pathway is essential to sustain a CXCL12 autocrine loop in cells migrating in response to HMGB1. *J Immunol* 2012; **188**: 2380–2386.
- 159 Amara A, Lorthioir O, Valenzuela A, Magerus A, Thelen M, Montes M *et al*. Stromal cell-derived factor-1alpha associates with heparan sulfates through the first beta-strand of the chemokine. *J Biol Chem* 1999; **274**: 23916–23925.
- 160 Drury LJ, Ziarek JJ, Gravel S, Veldkamp CT, Takekoshi T, Hwang ST *et al*. Monomeric and dimeric CXCL12 inhibit metastasis through distinct CXCR4 interactions and signaling pathways. *Proc Natl Acad Sci USA* 2011; **108**: 17655–17660.
- 161 Kuschert GS, Coulin F, Power CA, Proudfoot AE, Hubbard RE, Hoogewerf AJ *et al*. Glycosaminoglycans interact selectively with chemokines and modulate receptor binding and cellular responses. *Biochemistry* 1999; **38**: 12959–12968.
- 162 Rot A. Neutrophil attractant/activation protein-1 (interleukin-8) induces in vitro neutrophil migration by haptotactic mechanism. *Eur J Immunol* 1993; **23**: 303–306.
- 163 Proudfoot AE, Handel TM, Johnson Z, Lau EK, LiWang P, Clark-Lewis I *et al*. Glycosaminoglycan binding and oligomerization are essential for the in vivo activity of certain chemokines. *Proc Natl Acad Sci USA* 2003; **100**: 1885–1890.
- 164 Proudfoot AE, Uguccioni M. Modulation of chemokine responses: synergy and cooperativity. *Front Immunol* 2016; **7**: 183.
- 165 Steinberg M, Silva M. Plerixafor: a chemokine receptor-4 antagonist for mobilization of hematopoietic stem cells for transplantation after high-dose chemotherapy for non-Hodgkin's lymphoma or multiple myeloma. *Clin Ther* 2010; **32**: 821–843.