

Cite this: *Chem. Sci.*, 2016, 7, 6280

Correction: Iminoboronates are efficient intermediates for selective, rapid and reversible *N*-terminal cysteine functionalisation

Hélio Faustino,^a Maria J. S. A. Silva,^a Luís F. Veiros,^b Gonçalo J. L. Bernardes^{cd}
and Pedro M. P. Gois^{*a}

DOI: 10.1039/c6sc90045c

www.rsc.org/chemicalscienceCorrection for 'Iminoboronates are efficient intermediates for selective, rapid and reversible *N*-terminal cysteine functionalisation' by Hélio Faustino *et al.*, *Chem. Sci.*, 2016, 7, 5052–5058.

The authors regret that in the original article there are two instances of incorrect *m/z* values.

In Scheme 9, the *m/z* value of the starting material, calcitonin **22**, is herein corrected from 3471.7 to 3431.9 and in Scheme 10, the *m/z* value of the starting material **23** is herein corrected from 3471.7 to 3547.8.

The Royal Society of Chemistry apologises for these errors and any consequent inconvenience to authors and readers.

^aResearch Institute for Medicines (iMed.Ulisboa), Faculty of Pharmacy, Universidade de Lisboa, Lisbon, Portugal. E-mail: pedrogois@ff.ulisboa.pt

^bCentro de Química Estrutural, Instituto Superior Técnico, Universidade de Lisboa, Av. Rovisco Pais 1, 1049-001 Lisbon, Portugal

^cDepartment of Chemistry, University of Cambridge, Lensfield Road, CB2 1EW, Cambridge, UK

^dInstituto de Medicina Molecular, Faculdade de Medicina, Universidade de Lisboa, Avenida Professor Egas Moniz, 1649-028, Lisboa, Portugal

