

C/EBP γ is dispensable for steady-state and emergency granulopoiesis

The controlled production of granulocytes must be able to fulfill different needs of the organism throughout its lifetime. The maintenance of a stable basal level of mature granulocytes is ensured by steady-state granulopoiesis,¹ whereas in stress situations, such as severe infection, large numbers of neutrophils are required and a program called emergency granulopoiesis is activated.² These two programs are differentially regulated at the transcriptional level,³ and require appropriate expression of cell- and stage-specific transcription factors in order to serve particular demands for granulocyte production. It

has been shown that the CCAAT/enhancer-binding protein (C/EBP) family of transcription factors plays a critical role in these processes,³⁻⁵ however, the function of C/EBP γ in normal and emergency granulopoiesis remains elusive. C/EBP γ is ubiquitously expressed in hematopoietic cells,^{6,7} and despite lacking transactivation domains, it has been reported that C/EBP γ controls gene regulation by dimerizing with other transcription factors.^{8,9} Herein, we generated a C/EBP γ conditional knockout (KO) mouse model in which *Cebpg* is depleted in the hematopoietic system, and demonstrated that the transcription factor C/EBP γ , despite being expressed at high levels in all hematopoietic cells, is dispensable for steady-state and emergency granulopoiesis.

To study the role of C/EBP γ in hematopoiesis we gen-




Figure 1. Steady-state granulopoiesis is not affected by genetic deletion of *Cebpg* gene. (A) BM cellularity in WT and *Cebpg* KO mice. Y axis indicates absolute number of cells per hind leg. (B) Representative flow cytometry plots from WT and *Cebpg* KO BM cells stained for myeloid (Ly6G, CD11b), B-cell (B220), and T-cell (CD3) markers. Upper left quadrant indicates percentage of myeloid cells, upper right the percentage of B cells, and lower right the percentage of T cells. (C) Quantification of myeloid cells (Mye), B cells (B), and T cells (T) based on flow cytometry data illustrated in panel B. Y axes indicate relative percentage (rel., left panel) and absolute number (abs., right panel) of cells. (D) Representative flow cytometry plots from WT and *Cebpg* KO BM cells. Left plots were gated from lineage negative cells and stained for c-kit and Sca-1 markers. Blue box indicates percentage of myeloid progenitor cells (lineage⁻, c-kit⁺, Sca-1⁺), and orange box indicates percentage of LKS (lineage⁻, c-kit⁺, Sca-1⁺) cells. Right plots represent expression of Fc γ R II/III and CD34 in myeloid progenitor cells. Lower left gate indicates percentage of megakaryocyte erythroid progenitor cells (MEP), lower right indicates percentage of common myeloid progenitor cells (CMP), and upper right percentage of granulocyte macrophage progenitor cells (GMP). (E) Quantification of panel D. (C-E) Each group contains 12 mice from three independent experiments. (F) Colony forming assays using WT (white bars) and *Cebpg* KO (black bars) BM cells. 7.5×10^3 cells were plated per well. Y axis indicates total number of colonies per well. X axis indicate the cytokines included in each condition. Two-tailed Student's *t*-tests were used to assess statistical significance. BM: bone marrow; WT: wild-type; KO: knockout; LKS: lineage⁻, c-kit⁺, Sca-1⁺ cells; SCF: stem cell factor; IL: interleukin; TPO: thrombopoietin; FLT-3L: FMS-like tyrosin kinase 3 ligand.

erated a *Cebpg* conditional KO murine model (*Cebpg^{fl/fl}*) (Online Supplementary Figure S4) which we crossed with Vav-iCre transgenic mice, generating *Cebpg^{fl/fl}* Vav-iCre⁻ (wild-type, WT) and *Cebpg^{fl/fl}* Vav-iCre⁺ (KO) animals. We validated our model and demonstrated efficient Cre-mediated recombination resulting in controlled deletion of this transcription factor in the hematopoietic system (Online Supplementary Figure S2). Presence of a tdTomato reporter demonstrated expression of C/EBP γ in all hematopoietic cells (Online Supplementary Figure S2H and data not shown). C/EBP γ lacks transactivation domains and it has been proposed that it acts by heterodimerizing

with other factors, such as C/EBP β and Activating Transcription Factor 4 (ATF4).^{8,9} Using electrophoretic mobility shift assays (EMSA) we demonstrated the presence of C/EBP γ heterodimers in murine WT bone marrow (BM) and spleen (SP) cells; complexes which were not present in *Cebpg* KO cells (Online Supplementary Figure S3). Of note, despite the fact that granulocytes represent a high percentage of the total BM population, our EMSAs do not specifically address whether these complexes are present in the granulocytic lineage cells. In summary, we generated *Cebpg^{fl/fl}* mice which report for C/EBP γ expression in all hematopoietic cells, we demon-




Figure 2. *Cebpg* KO mice respond normally to LPS injection. (A) Schematic representation of the experimental setup. WT and *Cebpg* KO mice were treated intraperitoneally (i.p.) with 1400 μ g/kg LPS at 0 and 48 hours. Analysis was performed 24 hours after last injection. (B) BM cellularity in WT and *Cebpg* KO mice treated with PBS vehicle control (black) or LPS (red). Y axis indicates the number of cells per hind leg. (C) Representative flow cytometry plots from WT and *Cebpg* KO BM cells 24 hours after last PBS or LPS injection. Cells were stained for Gr1 and CD11b cell surface markers. Green boxes indicate percentage of Gr1^{high} CD11b⁺ cells and black boxes indicate percentage of Gr1^{low} CD11b⁺ cells. (D) Quantification of panel C. Graphs indicate relative (rel.) and absolute (abs.) number of Gr1^{high} CD11b⁺ cells (left) and Gr1^{low} CD11b⁺ cells (right). (E) Flow cytometric analysis of blood (BL) from WT and *Cebpg* KO mice 24 hours after last PBS or LPS injection. Green boxes indicate percentage of Gr1^{high} CD11b⁺ cells and black boxes indicate percentage of Gr1^{low} CD11b⁺ cells. (F) Quantification of panel E. Left graph indicates relative number of Gr1^{high} CD11b⁺ cells and right graph indicates relative number of Gr1^{low} CD11b⁺ cells. PBS vehicle control treatment indicated in black and LPS in red. Y axes indicate percentages (%). At least eight animals were included in each group. All data represent mean \pm s.d. from three independent experiments. Two-tailed Student's t-tests were used to assess statistical significance (ns: non-significant, ** P <0.01, **** P <0.0001). BM: bone marrow; WT: wild-type; KO: knockout; LPS: lipopolysaccharide; PBS: phosphate-buffered saline.

strated the presence of C/EBP γ heterodimers in adult hematopoietic cells, and induced a controlled deletion of this transcription factor in the hematopoietic system.

Surprisingly, *Cebpg* KO mice were viable and healthy, and showed no signs of disease during their life-span (mice were observed until the age of 70 weeks). Since granulopoiesis takes place in the BM from multipotent hematopoietic stem cells (HSCs) and C/EBP γ is expressed in long-term (LT)-HSCs,⁶ we investigated whether

C/EBP γ is involved in HSC function. Phenotypically defined LT-HSCs (LKS SLAM; LKS: lineage⁻, c-kit⁺, Sca-1⁺; signaling lymphocyte activation molecule (SLAM): CD48⁺, CD150⁺) were sorted, and gene expression profiling was performed (E-MTAB-6245). Using LIMMA package to assess differential gene expression, we detected 426 probesets identifying dysregulated genes in *Cebpg* KO versus WT LT-HSC ($P < 0.05$ and fold change > 1) (Online Supplementary Figure S4A and Online




Figure 3. *Cebpg* deficient and WT mice respond to *C. albicans* infection in the same fashion. (A) Graphical representation of *C. albicans* infection in WT and *Cebpg* KO mice. Animals were treated once (t=0) intravenously (i.v.) with 5×10^5 CFU/25g body weight. Analysis was performed 24h after infection. (B-C) Representative flow cytometry plots (B) and corresponding quantification (C) from WT and *Cebpg* KO blood (BL) from mice treated with PBS or *C. albicans*. Green boxes indicate percentage of Ly6G^{high} CD11b⁺ cells. Graphic indicates relative number of Ly6G^{high} CD11b⁺ cells. Y axis represents percentage (%). (D) Number of neutrophils per L in peripheral BL determined by Auto Hematology Analyzer. WT and *Cebpg* KO mice were treated with PBS (black) or *C. albicans* (red). (E) BM cellularity in WT and *Cebpg* KO mice treated with PBS (black) or *C. albicans* (red). Y axis indicates the number of cells per hind leg. (F) Representative flow cytometry plots from WT and *Cebpg* KO BM cells. Cells were stained for Ly6G and CD11b cell surface markers. Green boxes indicate percentage of Ly6G^{high} CD11b⁺ cells and black boxes indicate percentage of Ly6G^{low} CD11b⁺ cells. (G) Quantification of panel F. Left graphs indicate relative (rel.) and absolute (abs.) number of Ly6G^{high} CD11b⁺ cells. Right graphs indicate rel. and abs. number of Ly6G^{low} CD11b⁺ cells. Each group contains at least five mice. All data represent mean \pm s.d. from two independent experiments. Two-tailed Student's t-tests were used to assess statistical significance (ns: non-significant, * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$). BM: bone marrow; WT: wild-type; KO: knockout; BL: blood.

Supplementary Table S1 and Table S2). However, gene set enrichment analysis demonstrated scarce differentially regulated pathways in the *Cebpg* KO LT-HSC in comparison to WT LT-HSC (Online Supplementary Table S3). To determine whether the changes in gene expression would have an impact on the functionality of *Cebpg* KO LT-HSC, competitive limiting-dilution repopulating assays were performed, revealing no significant differences in WT and *Cebpg* KO LT-HSC numbers and repopulating abilities (Online Supplementary Figure S4B-D). To corroborate our results *in vitro*, we performed serial replating assays, which demonstrated no differences between WT and *Cebpg* KO cells (Online Supplementary Figure S4E,F). Together, our *in vitro* and *in vivo* assays indicate that *Cebpg* is dispensable for HSC functions. Nevertheless, future studies should determine whether C/EBP γ may play a role in LT-HSC self-renewal.

Next, we investigated whether *Cebpg* ablation affects steady-state granulopoiesis. We demonstrated that the deletion of *Cebpg* did not affect total cell numbers in BM, and that *Cebpg* KO mice presented a similar percentage and numbers of mature cells as their WT littermates (Figure 1A-C). In addition, BM and blood neutrophils presented similar morphology in *Cebpg* KO and WT mice, and no differences in granule content were detected (Online Supplementary Figure S5). Furthermore, flow cytometric analysis showed similar numbers of bone marrow progenitors in both WT and KO mice (Figure 1D,E). Similarly, later stages of granulocytic development⁴ demonstrated no differences between WT and *Cebpg* KO mice (Online Supplementary Figure S6). Since it was previously reported that straight *Cebpg* KO BM cells show reduced colony formation *in vitro*,⁵ we performed colony culture assays using several combinations of hematopoietic growth factors (Figure 1F). Unexpectedly, our model did not recapitulate the changes reported previously,⁵ possibly due to the distinct approaches employed to induce *Cebpg* ablation (straight versus conditional deletion) or the different murine background (C57BL/6-129v versus pure C57BL/6NCr). Taken together these data suggest that in our murine model, C/EBP γ is dispensable for basal granulopoiesis.

Since *Cebpg* deficient mice showed no defects in steady-state conditions, we subsequently examined whether C/EBP γ is required during lipopolysaccharide (LPS)-induced emergency granulopoiesis. We challenged WT and *Cebpg* KO mice with high doses of LPS to mimic a severe systemic infection (Figure 2A), and analyzed typical surrogate hallmarks of emergency granulopoiesis.^{10,11} Strikingly, BM cellularity, and the frequency and absolute numbers of mature Gr1^{high} CD11b⁺ and immature Gr1^{low} CD11b⁺ BM neutrophils after LPS treatment was equal in both *Cebpg* KO and WT controls (Figure 2B-D). Furthermore, we demonstrated that the increase in mature and immature blood neutrophils was similar in WT and *Cebpg* KO mice (Figure 2E,F). In addition, LPS treatment resulted in a general increase in SP cellularity and an increase in both mature and immature granulocytes, which was comparable in WT and *Cebpg* KO mice (Online Supplementary Figure S7). These experiments demonstrate that *Cebpg* deficient mice present a similar LPS-induced emergency granulopoiesis as WT mice, indicating that this transcription factor is dispensable for the response to this type of challenge.

Since we did not observe an effect of *Cebpg* deletion during LPS-induced emergency granulopoiesis, which is dependent on toll-like receptor 4,^{11,12} we proceeded to challenge mice with granulocyte-colony stimulating factor (G-CSF), a more general emergency granulopoiesis

inducer (Online Supplementary Figure S8A). In agreement with published data,^{3,11} we observed that G-CSF alone is sufficient to accurately mimic emergency granulopoiesis. Of note, this chronic G-CSF treatment protocol did not result in significant changes in total BM cellularity, since G-CSF-induced granulopoiesis in the BM is accompanied by the mobilization of granulocytes to blood¹³ (Online Supplementary Figure S8B). We showed similar responses in WT and *Cebpg* KO mice, as demonstrated by a decrease in relative and absolute number of mature Gr1^{high} CD11b⁺ cells and an increase in immature Gr1^{low} CD11b⁺ neutrophils in BM (Online Supplementary Figure S8C,D), whereas in SP we observed an increase in both mature and immature granulocytic populations (Online Supplementary Figure S8E-G). Thus, our results demonstrate that *Cebpg* KO mice respond to G-CSF-induced emergency granulopoiesis in a similar fashion as WT mice.

The presence of C/EBP γ :C/EBP β heterodimers in BM and SP (Online Supplementary Figure S3) suggests that C/EBP γ may modulate C/EBP β function. Since C/EBP β plays a critical role in *C. albicans*-induced emergency granulopoiesis,⁴ we investigated whether ablation of *Cebpg* alters the response to fungal infection (Figure 3A). As expected, we observed an increase in the percentage of mature neutrophils in blood (Figure 3B,C) upon *C. albicans* administration, which was similar in WT and *Cebpg* KO mice. Interestingly, we detected a significant increase in absolute blood neutrophil cell counts in KO mice in comparison to control littermates (~2-fold in KO vs. ~1.5-fold in WT) (Figure 3D). This difference was not limited to neutrophils because higher cell counts were measured for the remaining leukocyte subsets as well (*data not shown*). Since higher egress of the BM cells could explain higher absolute cell counts in blood, we compared the reduction in BM cellularity between WT and KO mice following *C. albicans* treatment. However, no significant differences were observed (Figure 3E). Further, we showed that both WT and *Cebpg* KO mice presented typical hallmarks of emergency granulopoiesis (Figure 3F,G). Since it was reported that C/EBP β is crucial for the amplification of immature hematopoietic cells after induction of systemic candidemia,⁴ we next assessed early granulocytic developmental stages in control mice treated with phosphate-buffered saline (PBS) and *C. albicans* infected mice. Although small differences were observed in the relative number of LKS upon treatment, detailed analysis of myeloid progenitors and distinct granulocytic differentiation stages revealed identical dynamics in relative and absolute cell counts in WT and *Cebpg* KO mice (Online Supplementary Figure S9 and Figure S10A,B). We then assessed whether the minor differences mentioned above are functionally relevant. However, we observed no significant differences in the microbiological outcome and survival rate in *Cebpg* KO mice compared to WT animals (Online Supplementary Figure S10C,D). These data suggest that C/EBP γ is not required for efficient emergency granulopoiesis induced by systemic candidemia.

Taken together, our data reveal that C/EBP γ is dispensable for steady-state and emergency granulopoiesis. This is in sharp contrast to the results obtained with a previously reported *Cebpg* KO murine model generated by Kaisho *et al.*,¹⁴ in which mice exhibited a high mortality rate within 48h after birth. However, *Cebpg* ablation in this model was not limited to the hematopoietic system, and it was suggested that the early neonatal death could be caused by lung lesions.¹⁴ On the contrary, in our conditional KO mice, C/EBP γ expression is abolished

mainly in hematopoietic cells, which makes these animals optimal for studying C/EBP γ function in the hematopoietic system.

In summary, the data presented herein demonstrate that specific deletion of *Cebpg* in the murine hematopoietic system did not alter hematopoietic stem and progenitor cell properties, their ability to commit to the myeloid lineage, and produce granulocytes in normal and stress conditions. As a whole, these surprising observations point to a transcription factor redundancy responsible for controlled production of granulocytes in steady-state and emergency granulopoiesis.

Miroslava Kardosova,^{1,2} Polina Zjablovskaja,^{1,3,4} Petr Danek,¹ Pavla Angelisova,¹ Lorena Lobo de Figueiredo-Pontes,⁵ Robert S. Welner,⁶ Tomas Brdicka,¹ Sanghoon Lee,⁷ Daniel G. Tenen^{8,9} and Meritxell Alberich-Jorda^{1,3,4}*

¹Institute of Molecular Genetics of the Czech Academy of Sciences, Prague, Czech Republic; ²Department of Cell Biology, Charles University, Faculty of Science, Prague, Czech Republic; ³CLIP-Childhood Leukaemia Investigation, Prague, Czech Republic; ⁴Department of Paediatric Haematology and Oncology, Charles University, Second Faculty of Medicine, Prague, Czech Republic; ⁵Hematology/Oncology Division of the Department of Internal Medicine, Medical School of Ribeirao Preto, University of São Paulo, Brazil; ⁶Hematology/Oncology Department, University of Alabama at Birmingham, AL, USA; ⁷Department of Systems Biology, University of Texas MD Anderson Cancer Center, Houston, TX, USA; ⁸Cancer Science Institute, National University of Singapore, Singapore and ⁹Harvard Stem Cell Institute, Harvard Medical School, Boston, MA, USA

*DGT and MA-J contributed equally to this work

Acknowledgements: the authors would like to thank the Flow Cytometry Core Facility, IMG AS CR, Prague, Czech Republic for their support with the flow cytometry and sorting presented herein. We would like to thank Dr. P. Johnson and Dr. J. Salotti for advice in EMSA assays and Dr. V. Korinek for his technical support.

Funding: the study was supported by a GACR grant (18-08577S) and institutional funding from the IMG AS CR (RVO 68378050) to MA-J, a FAPESP grant (2015/21866-1) to LLF-P, and by a GA UK fellowship (project No. 341015) from Charles University in Prague to MK. DGT was supported by a STaR Investigator Award, an RCE Core grant, a Tier 3 RNA Biology Center grant MOE2014-T3-1-006 from the NRF and MOE, Singapore; and NIH/NCI grant P01 CA66996. This work used instruments provided by C4Sys infrastructure and it was also supported by the Czech Centre for Phenogenomics (CCP, project no. LM2015040), and OP RDI CZ.1.05/2.1.00/19.0395 (Higher quality and capacity for transgenic models).

Correspondence: alberich@img.cas.cz
doi:10.3324/haematol.2017.173781

Information on authorship, contributions, and financial & other disclosures was provided by the authors and is available with the online version of this article at www.haematologica.org.

References

- Bugl S, Wirths S, Radsak MP, et al. Steady-state neutrophil homeostasis is dependent on TLR4/TRIF signaling. *Blood*. 2013;121(5):723-733.
- Manz MG, Boettcher S. Emergency granulopoiesis. *Nat Rev Immunol*. 2014;14(5):302-314.
- Hirai H, Zhang P, Dayaram T, et al. C/EBPbeta is required for 'emergency' granulopoiesis. *Nat Immunol*. 2006;7(7):732-739.
- Satake S, Hirai H, Hayashi Y, et al. C/EBPbeta is involved in the amplification of early granulocyte precursors during candidemia-induced "emergency" granulopoiesis. *J Immunol*. 2012;189(9):4546-4555.
- Zhang DE, Zhang P, Wang ND, Hetherington CJ, Darlington CJ, Tenen DG. Absence of granulocyte colony-stimulating factor signaling and neutrophil development in CCAAT enhancer binding protein alpha-deficient mice. *Proc Natl Acad Sci U S A*. 1997;94(2):569-574.
- Alberich-Jorda M, Wouters B, Balastik M, et al. C/EBPgamma deregulation results in differentiation arrest in acute myeloid leukemia. *J Clin Invest*. 2012;122(12):4490-4504.
- Roman C, Platero JS, Shuman J, Calame K. Ig/EBP-1: a ubiquitously expressed immunoglobulin enhancer binding protein that is similar to C/EBP and heterodimerizes with C/EBP. *Genes Dev*. 1990;4(8):1404-1415.
- Huggins CJ, Malik R, Lee S, et al. C/EBPgamma suppresses senescence and inflammatory gene expression by heterodimerizing with C/EBPbeta. *Mol Cell Biol*. 2013;33(16):3242-3258.
- Huggins CJ, Mayekar MK, Martin N, et al. C/EBPgamma as a critical regulator of cellular stress response networks through heterodimerization with ATF4. *Mol Cell Biol*. 2015;36(5):693-713.
- Boettcher S, Gerosa RC, Radpour R, et al. Endothelial cells translate pathogen signals into G-CSF-driven emergency granulopoiesis. *Blood*. 2014;124(9):1393-1403.
- Boettcher S, Ziegler P, Schmid MA, et al. Cutting edge: LPS-induced emergency myelopoiesis depends on TLR4-expressing non-hematopoietic cells. *J Immunol*. 2012;188(12):5824-5828.
- Poltorak A, He X, Smirnova I, et al. Defective LPS signaling in C3H/HeJ and C57BL/10ScCr mice: mutations in Tlr4 gene. *Science*. 1998;282(5396):2085-2088.
- Lieschke GJ, Grail D, Hodgson G, et al. Mice lacking granulocyte colony-stimulating factor have chronic neutropenia, granulocyte and macrophage progenitor cell deficiency, and impaired neutrophil mobilization. *Blood*. 1994;84(6):1737-1746.
- Kaisho T, Tsutsui H, Tanaka T, et al. Impairment of natural killer cytotoxic activity and interferon gamma production in CCAAT/enhancer binding protein gamma-deficient mice. *J Exp Med*. 1999;190(11):1573-1582.