



Published in final edited form as:

*J Am Chem Soc.* 2017 March 01; 139(8): 2960–2963. doi:10.1021/jacs.7b01114.

## Polycyclic Aromatic Hydrocarbons via Iron(III)-Catalyzed Carbonyl-Olefin Metathesis

Christopher C. McAtee, Paul S. Riehl, and Corinna S. Schindler


University of Michigan, Department of Chemistry, Willard Henry Dow Laboratory, 930 North University Ave., Ann Arbor, MI 48109, US

### Abstract

Polycyclic aromatic hydrocarbons are important structural motifs in organic chemistry, pharmaceutical chemistry and materials science. The development of a new synthetic strategy toward these compounds is described based on the design principle of iron(III)-catalyzed carbonyl-olefin metathesis reactions. This approach is characterized by its operational simplicity, high functional group compatibility, and regioselectivity while relying on  $\text{FeCl}_3$  as an environmentally benign, earth-abundant metal catalyst. Experimental evidence for oxetanes as reactive intermediates in the catalytic carbonyl-olefin ring-closing metathesis has been obtained.

### Graphical abstract

Authors are required to submit a graphic entry for the Table of Contents (TOC) that, in conjunction with the manuscript title, should give the reader a representative idea of one of the following: A key structure, reaction, equation, concept, or theorem, etc., that is discussed in the manuscript. Consult the journal's Instructions for Authors for TOC graphic specifications.



Polycyclic aromatic compounds (PACs),<sup>1</sup> including phenanthrenes, pyrenes and chrysenes, are important structural motifs that exhibit desirable optical,<sup>2</sup> electronic,<sup>3</sup> and chelating<sup>4</sup> properties. Consequently, diverse fields of research such as materials science,<sup>4</sup> natural product synthesis,<sup>6</sup> asymmetric catalysis,<sup>7</sup> and molecular recognition<sup>8</sup> rely on efficient strategies to access condensed polyaromatic compounds. Established procedures toward these motifs include McMurry coupling reactions<sup>9, 10</sup> that are mediated by low-valent titanium reagents (Fig. 1A II) or oxidative photocyclization strategies<sup>11</sup> of stilbene

<sup>\*</sup>Corresponding Author: corinnas@umich.edu.

#### Supporting Information

Experimental data as well as <sup>1</sup>H and <sup>13</sup>C NMR spectra for all new compounds prepared in the course of these studies are provided in the Supporting Information to this manuscript. The material is available free of charge via the Internet at <http://pubs.acs.org>.


derivatives. These classical approaches<sup>12</sup> have been hampered by the need for stoichiometric reagents, harsh reaction conditions, or competing substrate dimerization. Complementary approaches have been developed to overcome these challenges that are based on Diels-Alder cycloaddition reactions,<sup>13</sup> radical cyclizations,<sup>14</sup> and metal-mediated cycloisomerizations.<sup>15</sup> Additionally, rhodium- and ruthenium-catalyzed procedures have been reported that rely on bis(N-tosylhydrazone)<sup>16</sup> **2** as substrate (Fig. 1A I) and olefin-metathesis reactions of bis(alkenes)<sup>17</sup> **4** (Fig. 1A III). We have recently reported the development of an efficient iron(III)-catalyzed carbonyl-olefin metathesis reaction<sup>18</sup> that proceeds under mild reaction conditions and ambient temperature. Our synthetic strategy for ring-closing metathesis enables the direct coupling of carbonyl and olefin functional groups upon activation by a Lewis acid catalyst to forge the desired alkene bonds. Based on this design principle, we report the development of a new strategy for the synthesis of electronically and sterically diverse PACs. This strategy is compatible with both ketones and aldehydes, proceeding via intermediate oxetanes **6** to provide the corresponding metathesis products in good to excellent yields (Fig. 1B). While several Lewis acids were previously found capable of promoting carbonyl-olefin metathesis reactions,<sup>18</sup> a fine-tuned combination of Lewis acidity<sup>19</sup> and oxophilicity<sup>20</sup> proved essential to give high yields of product. Indeed, when biaryl ketone **8** was reacted with numerous Lewis acids (e.g. TiCl<sub>4</sub>, SnCl<sub>4</sub>, FeCl<sub>2</sub>, Cu(OTf)<sub>2</sub>) no formation or only trace amounts of the metathesis product **9** was observed (entries 1–4, Table 1). Stronger Lewis acids, GaCl<sub>3</sub> and AlCl<sub>3</sub>,<sup>18</sup> were able to promote the desired transformation in 88% and 93% yield, respectively with complete conversion of starting material **8** (entries 7 and 8, Table 1). Notably, substoichiometric BF<sub>3</sub>·Et<sub>2</sub>O led to the formation of **9** in only modest yield and conversion (entry 6, Table 1).<sup>21</sup> Ultimately, 5 mol% FeCl<sub>3</sub> in either dichloroethane or toluene was identified as an optimal set of reaction conditions, resulting in quantitative formation of the product **9** in 97% and 99% yield, respectively (entries 9 and 11, Table 1). More dilute reaction conditions led to slightly lower yields of **9** (entry 10, Table 1). When the reaction was conducted in ethereal solvents (1,4-dioxane), or polar aprotic solvents (DMF), no formation of phenanthrene **9** was observed—presumably due to competing Lewis basicity of these solvents (entries 12 and 13, Table 1). Moreover, the Brønsted acids, anhydrous HCl<sup>22</sup> and *p*TsOH in dichloroethane, did not form phenanthrene **9** and resulted in quantitative reisolation of starting material (entries 16 and 17, Table 1).

We next sought to investigate the ability of biaryl substrates with various olefin subunits (**11–19**) to undergo the metathesis reaction (Table 2). While both electron-rich and electron-poor styrenes (entries 1–6, Table 2) proved to be efficient substrates resulting in high yields of **9**, all but styrene **11** and prenylated **17** required elevated temperatures of 50 °C to proceed to full conversion. Notably, no difference in reactivity between *E*- and *Z*-isomers was observed; both *para*-methyl styrenes **12** and **13** formed metathesis product **9** in yields up to 89% which indicates an indiscriminate reaction pathway of the carbonyl-olefin metathesis reaction. Although the formation of the respective benzaldehydes was observed as the corresponding metathesis byproducts in the course of the reaction, they did not impede reaction progress. Moreover, substrates **11–16** bearing styrenyl moieties proved superior to their prenylated analog **17**, which resulted in the formation of **9** in only 79% yield (entries 1–7, Table 2). In comparison, no reaction was observed when terminal alkene **19** was

subjected to the optimized reaction conditions (entry 9, Table 2). Conversion of biaryl **18** bearing a crotyl moiety under the reaction conditions resulted in low yields (18%) of the desired product. The hampered yields of the non-styrenyl substrates **17** and **18** were found to be caused by a competing carbonyl-ene reaction pathway which led to the formation of **20** and **21** in 21% and 47% yield, respectively, when subjected to the optimized reaction conditions (Fig. 2). These findings contrast distinctly with previous results obtained in our lab<sup>18</sup> in the iron(III)-catalyzed carbonyl-olefin metathesis reaction of aliphatic aryl ketones, in which prenylated substrates proved superior to the analogous styrenes.

The conditions developed for the iron(III)-catalyzed carbonyl-olefin metathesis reaction proved efficient for a range of sterically and electronically differentiated ketones and aldehydes (entries 1–9, Table 3). Although aldehydes have previously been found unreactive in catalytic carbonyl-olefin ring-closing metathesis reactions,<sup>18</sup> **22b** was found to yield the desired metathesis product **23b** in 75% under the optimized conditions.

In addition to methyl ketone **22a** and aldehyde **22b**, substrates bearing sterically demanding isopropyl (**22c**) and *tert*-butyl (**22d**) moieties formed the alkylated phenanthrenes in 79% and 55%, respectively, although the latter required elevated temperatures for efficient conversion (entries 3 and 4, Table 3). Phenyl and naphthyl substituted carbonyl substrates (**22e** and **22f**) were able to undergo metathesis in efficient yields (entries 5 and 6, Table 3). Importantly, biaryl enone **22g** led to the corresponding polycycle **23g** incorporating an exocyclic alkene as a functional handle in 50% yield, albeit at elevated temperatures (entry 7, Table 3). Additionally,  $\beta$ -ketoester **22h** resulted in the formation of metathesis product **23h** in satisfactory yield (72%), while electron-deficient trifluoromethyl ketone **22i** also proved viable as a substrate converting to 9-trifluoromethyl phenanthrene **23i** in 52% (entries 8 and 9, Table 3). Various PAC frameworks were accessible utilizing the optimal reaction conditions (Table 4). Upon subjection to metathesis conditions, the desired PACs were obtained with benzaldehyde as the corresponding byproduct. Electron-deficient phenanthrenes bearing halogen, trifluoromethyl, nitro, or nitrile substitution were formed in yields greater than 85% (**27**, **29**, **45**, **46**, **55** and **56**, Table 4). Similarly, electron-rich substrates incorporating methoxy or benzyl ether functionalities underwent the desired transformation in excellent yields (**30**, **31**, **32**, **38**, **42**, Table 4). However, diminished yields of 75% and 57% were observed for substrates bearing *ortho*-methoxy substitution (**34** and **37**, Table 4). Dioxoles **40** and **44** were formed in 99% and 68% yield, respectively, under the optimized reaction conditions. Moreover, sulfur-containing heterocycles proved viable substrates for metathesis and resulted in the formation of thiophene **39** and benzothiophenes **35** and **41** in good yields. Alternative strategies to these structural motifs are currently hampered by harsh reaction conditions and competing reaction pathways resulting in low overall yields.<sup>23</sup> Unprotected phenols as well as aldehydes readily underwent metathesis resulting in the formation of phenanthrene **28** or aldehyde **50** in 74% and 90% yield, respectively. Furthermore, extended PACs are accessible employing this metathesis strategy. Specifically, methylchrysene **25** is generated in 80% yield, while benzo(c)phenanthrene **36** is accessible in 89% yield from the respective biaryl aldehyde (Table 4). Notably, dibenz[a,h]anthracene **59** is afforded in excellent yield via biscarbonyl-olefin metathesis (eq 1).



(1)

Interestingly, when the prenylated analog of **22i** was converted under the optimized reaction conditions, no formation of the desired carbonyl-olefin metathesis product **23i** was observed. Oxetane **6** was identified as the major product (45% yield, Table 4). This result supports our hypothesis that iron(III)-catalyzed carbonyl-olefin metathesis reactions do proceed via oxetanes as reactive intermediates.<sup>18</sup>

The development of a new approach toward the synthesis of polyaromatic hydrocarbons is reported relying on the design principle of an iron(III)-catalyzed carbonyl-olefin metathesis reaction. This strategy is characterized by its operational simplicity, mild reaction conditions, as well as chemo- and regioselectivity. Analysis of the two reaction partners (olefin and carbonyl) revealed that the respective olefin moieties can readily couple to a variety of differentiated aryl-ketones or aryl aldehydes to garner the corresponding functionalized PACs as metathesis products. Isolation of aryl oxetane **6** supports the notion that this new strategy for the synthesis of polyaromatic hydrocarbons does indeed proceed via oxetanes as reactive intermediates.<sup>18</sup>

## Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

## Acknowledgments

We thank the Petroleum Research Fund (PRF#54688-DN11), the University of Michigan Office of Research and the NIH/National Institute of General Medical Sciences (GM118644) for financial support. P.S.R. thanks Eli Lilly for a summer predoctoral fellowship. We thank Dr. Jeff W. Kampf and Ren Wiscons for X-ray crystallographic studies.


## References

1. a) Harvey RG. Polycyclic Aromatic Hydrocarbons. Wiley-VCH; New York: 1997. b) Gingras M. Chem Soc Rev. 2013; 42:968. [PubMed: 23151799] c) Gingras M, Felix G, Peresutti R. Chem Soc Rev. 2013; 42:1007. [PubMed: 23151610] d) Gingras M. Chem Soc Rev. 2013; 42:1051. [PubMed: 23151680] e) Floyd AJ, Dyke SF, Ward SE. Chem Rev. 1976; 76:509. For additional examples, see: e) Uchida K, Ito S, Nakano M, Abe M, Kubo T. J Am Chem Soc. 2016; 138:2399. [PubMed: 26824692] f) Ji F, Li X, Wu W, Jiang H. J Org Chem. 2014; 79:11246. [PubMed: 25353393]
2. Wigglesworth TJ, Sud D, Norsten TB, Lekhi VS, Branda NR. J Am Chem Soc. 2005; 127:7272. [PubMed: 15898750]
3. Furche F, Ahlrichs R, Wachsmann C, Weber E, Sobanski A, Vögtle F, Grimme S. J Am Chem Soc. 2000; 122:1717.
4. Xu Y, Zhang YX, Sugiyama H, Umamo T, Osuga H, Tanaka K. J Am Chem Soc. 2004; 126:6566. [PubMed: 15161280]


5. a) Lovinger AJ, Nuckolls C, Katz TJ. *J Am Chem Soc.* 1998; 120:264. b) Zöphel L, Enkelmann V, Müllen K. *Org Lett.* 2013; 15:804. [PubMed: 23387523]
6. Kovacs A, Vasas A, Hohmann J. *Phytochemistry.* 2008; 69:1084. [PubMed: 18243254]
7. For representative examples, see: a) Knowles RR, Lin S, Jacobsen EN. *J Am Chem Soc.* 2010; 132:5030. [PubMed: 20369901] b) Narcis MJ, Takenaka N. *Eur J Org Chem.* 2014; 1:21.
8. Dreher SD, Katz TJ, Lam KC, Rheingold AL. *J Org Chem.* 2000; 65:815.
9. McMurry JE, Lectka T, Rico JG. *J Org Chem.* 1989; 54:3748.
10. Dubois F, Gingras M. *Tetrahedron Lett.* 1998; 39:5039.
11. a) Flammang-Barbieux M, Nasielski J, Martin RH. *Tetrahedron Lett.* 1967; 8:743. b) Liu L, Yang B, Katz TJ, Poindexter MK. *J Org Chem.* 1991; 56:3769. c) Martin RH. *Angew Chem Int Ed.* 1974; 13:649.
12. For homolytic aromatic substitution strategy, see Harrowven DC, Guy IL, Nanson L. *Angew Chem Int Ed.* 2006; 45:2242.
13. a) Carreño MC, González-López M, Urbano A. *Chem Comm.* 2005; 5:611. b) Liu L, Katz TJ. *Tetrahedron Lett.* 1990; 31:3983. c) Katz TJ, Liu L, Willmore ND, Fox JM, Rheingold AL, Shi S, Nuckolls C, Rickman BH. *J Am Chem Soc.* 1997; 119:10054. d) Fox JM, Goldberg NR, Katz TJ. *J Org Chem.* 1998; 63:7456. [PubMed: 11672398] e) Dreher SD, Weix DJ, Katz TJ. *J Org Chem.* 1999; 64:3671. [PubMed: 11674496]
14. a) Harrowven DC, Guy IL, Nanson L. *Angew Chem Int Ed.* 2006; 45:2242. b) Harrowven DC, Nunn MIT, Fenwick DR. *Tetrahedron Lett.* 2002; 43:7345.
15. For examples, see: a) Fürstner A, Mamane V. *J Org Chem.* 2002; 67:6264. [PubMed: 12182677] b) Mamane V, Hannen P, Fürstner A. *Chem Eur J.* 2004; 10:4556. [PubMed: 15378635] c) Komeyama K, Igawa R, Takaki K. *Chem Comm.* 2010; 46:1748. [PubMed: 20177637] d) Chernyak N, Gevorgyan V. *J Am Chem Soc.* 2008; 130:5636. [PubMed: 18357988] e) Chernyak N, Gevorgyan V. *Adv Synth Catal.* 2009; 351:1101. [PubMed: 20160971]
16. Xia Y, Liu Z, Xiao Q, Qu P, Ge R, Zhang Y, Wang J. *Angew Chem Int Ed.* 2012; 51:5714.
17. a) Iuliano A, Piccioli P, Fabbri D. *Org Lett.* 2004; 6:3711. [PubMed: 15469330] b) Donohoe TJ, Orr AJ, Bingham M. *Angew Chem Int Ed.* 2006; 45:2664.
18. This work was first reported as Ludwig JR, Gianino JB, Schindler C. Metal-catalyzed carbonyl-olefin metathesis. Abstracts of Papers, 250th ACS National Meeting & Exposition; Boston, MA, United States. August 16th–20th, 2015; Proceedings of the American Chemical Society; Boston, MA. August 16th–20th; Washington, DC: American Chemical Society; 2015. 1328933ORGN-447b) Ludwig JR, Zimmerman PM, Gianino JB, Schindler CS. *Nature.* 2016; 533:374. [PubMed: 27120158]
19. For metal-mediated carbonyl-olefin metathesis reactions, see: a) Schopov I, Jossifov C. *Makromol Chem Rapid Commun.* 1983; 4:659. b) Fu GC, Grubbs RH. *J Am Chem Soc.* 1993; 115:3800. For carbonyl-olefin metathesis reactions proceeding via oxetan photoadducts, see: c) Jones G II, Schwartz SB, Marton MT. *J Chem Soc Chem Comm.* 1973; 11:374. d) Jones G II, Acquadro MA, Carmody MA. *J Chem Soc Chem Comm.* 1975; 6:206. e) Carless HAJ, Trivedi HS. *J Chem Soc Chem Commun.* 1979; 8:382. f) D'Auria M, Racioppi R, Viggiani L. *Photochem Photobiol Sci.* 2010; 9:1134. [PubMed: 20563353] g) Pérez-Ruiz R, Gil S, Miranda MA. *J Org Chem.* 2005; 70:1376. [PubMed: 15704973] h) Pérez-Ruiz R, Miranda MA, Alle R, Meerholz K, Griesbeck AG. *Photochem Photobiol Sci.* 2006; 5:51. [PubMed: 16395427] i) Valiulin RA, Arisco TM, Kutateladze AG. *J Org Chem.* 2011; 76:1319. [PubMed: 21268619] j) Valiulin RA, Arisco TM, Kutateladze AG. *J Org Chem.* 2013; 78:2012. [PubMed: 23106813] For Brønsted and Lewis acid mediated carbonyl-olefin metathesis reactions, see: k) Soicke A, Slavov N, Neudörfel J-M, Schmalz H-G. *Synlett.* 2011; 17:2487. l) van Schaik HP, Vijn RJ, Bickelhaupt F. *Angew Chem Int Ed.* 1994; 33:1611. m) Bah J, Franzén J, Naidu VR. *Eur J Org Chem.* 2015; 8:1834. n) Jossifov C, Kalinova R, Demonceau A. *Chim Oggi.* 2008; 26:85. For catalytic carbonyl-olefin metathesis reactions proceeding via (3+2)/retro-(3+2)-cycloaddition, see: o) Griffith AK, Vanos CM, Lambert TH. *J Am Chem Soc.* 2012; 134:18581. [PubMed: 23126620] p) Hong X, Liang Y, Griffith AK, Lambert TH, Houk KN. *Chem Sci.* 2014; 5:471.
20. a) Satchell DPN, Satchell RS. *Chem Rev.* 1969; 69:251. b) Jensen WB. *Chem Rev.* 1978; 78:1.
21. Kepp KP. *Inorg Chem.* 2016; 55:9461. [PubMed: 27580183]

22. In the carbonyl-olefin metathesis reaction leading to cyclopentenes and cyclohexenes, catalytic amounts of  $\text{BF}_3 \cdot \text{Et}_2\text{O}$  formed the metathesis products in 71% (86% conversion).
23. Arnáiz FJ. *J Chem Educ.* 1995; 72:1139.
24. a) Sanz R, Fernández Y, Castroviejo MP, Pérez A, Fañanás FJ. *J Org Chem.* 2006; 71:6291. [PubMed: 16872224] b) Che R, Wu Z, Li Z, Xiang H, Zhou X. *Chem Eur J.* 2014; 20:7258. [PubMed: 24829172] c) Qiao Z, Ge N, Jiang X. *Chem Comm.* 2015; 51:10295. [PubMed: 26022891]

**A. Literature Precedent:** Selected strategies for the synthesis of PACs.




**B. This work:** Fe-catalyzed carbonyl-olefin metathesis for the synthesis of PACs.



**Figure 1.**

A. Select strategies to access PACs. B. Carbonyl-olefin metathesis approach reported.




**Figure 2.**

Competing metathesis and carbonyl-ene reactions.

**Condition:** biaryl (0.13 mmol), FeCl<sub>3</sub> (5 mol%) in dichloroethane (0.1M), rt, 1h;<sup>a</sup>)yield determined by <sup>1</sup>H NMR analysis with 1,3,5-trimethoxybenzene as internal standard.




Table 1

Reaction optimization for synthesis of **9**.

entry	Lewis acid	solvent	yield <b>9</b> (%)	conversion (%)
1	TiCl <sub>4</sub>	DCE	3	7
2	SnCl <sub>4</sub>	DCE	0	6
3	FeCl <sub>2</sub>	DCE	0	2
4	Cu(OTf) <sub>2</sub>	DCE	0	0
5	ZnCl <sub>2</sub>	DCE	22	26
6	BF <sub>3</sub> ·Et <sub>2</sub> O	DCE	31	35
7	AlCl <sub>3</sub>	DCE	93	100
8	GaCl <sub>3</sub>	DCE	88	100
9	FeCl <sub>3</sub>	DCE	97	100
10	FeCl <sub>3</sub>	DCE (0.01M)	95	100
11	FeCl <sub>3</sub>	toluene	99	100
12	FeCl <sub>3</sub>	DMF	0	0
13	FeCl <sub>3</sub>	1,4-dioxane(0.1M)	0	6
14	HCl	DCE	0	0
15	<i>p</i> TsOH	DCE	0	0

**Conditions:** biaryl **8** (0.13 mmol), Lewis or Brønsted acid (5 mol%) in solvent listed (0.1–0.01M), rt, 1h; yield determined by <sup>1</sup>H NMR analysis with 1,3,5-trimethoxy-benzene as internal standard.

Table 2


Alkene evaluation for formation of **9**.










entry	alkene	yield (%)	entry	alkene	yield (%)
1		90	5		80
2		82	6		84
3		89	7		79
4		86	8		11
			9		n.r.

**Conditions:** biaryl (0.13 mmol), FeCl<sub>3</sub> (5 mol%) in toluene (0.1 M);<sup>a</sup> mixture of *E/Z* (2:1) isomers;<sup>b</sup> reaction heated to 50°C.

Table 3

Evaluation of carbonyl substituents.




entry	substrate	yield (%)	entry	substrate	yield (%)
1	 <b>22a</b>	99	6	 <b>22f</b>	53
2	 <b>22b</b>	75	7	 <b>22g<sup>a</sup></b>	50
3	 <b>22c</b>	79	8	 <b>22h</b>	72
4	 <b>22d</b>	55 <sup>a</sup>	9	 <b>22i<sup>a</sup></b>	52
5	 <b>22e</b>	67			

**Conditions:** biaryl (0.13 mmol), FeCl<sub>3</sub> (5 mol%), in dichloroethane (0.1M), rt, 1–12 h;

<sup>a</sup> reaction heated to 50°C;

Table 4

Scope of the iron(III)-catalyzed carbonyl-olefin metathesis reaction for the synthesis of PACs.



**Conditions:** biaryl (0.13 mmol), FeCl<sub>3</sub> (5 mol%), in DCE (0.1M), rt, 1–12 h;

<sup>a</sup> reaction heated to 50°C;

<sup>b</sup> reaction was run with 20 mol% catalyst loading;

<sup>c</sup> starting material is bis-prenylated biaryl ketone (see Supporting Information For details);

<sup>d</sup> substrate is the prenylated analog of **22i**; reaction was run in toluene as solvent;

<sup>e</sup> starting material is reisolated;

<sup>f</sup> substrate decomposition was observed at the elevated reaction temperatures;

<sup>g</sup> low solubility in organic solvents.