

RESEARCH ARTICLE

Open Access



Antioxidant and anticancer activities of *Trigonella foenum-graecum*, *Cassia acutifolia* and *Rhazya stricta*

Bayan Al-Dabbagh^{1*}, Ismail A. Elhaty¹, Ala'a Al Hrou², Reem Al Sakkaf¹, Raafat El-Awady⁴, S. Salman Ashraf¹ and Amr Amin^{2,3*}

Abstract

Background: Here, we determined in vitro antioxidant activity, total phenols and flavonoids and evaluated antiproliferative activity of three medicinal plant extracts: *Trigonella foenum-graecum* (Fenugreek), *Cassia acutifolia* (Senna) and *Rhazya stricta* (Harmal).

Methods: The leaves of the three medicinal plants were extracted with 70% ethanol. Antioxidant activities of the extracts were determined by using DPPH (1,1-diphenyl-2-picrylhydrazyl) assay. Total flavonoid and phenolic contents were determined using colorimetric assays. MTT assay was used to estimate the antiproliferative activities of the extracts against human hepatoma (HepG2) cancer cell line. In addition, the effects of *R. stricta* extract on cell cycle, colony formation, and wound healing of HepG2 cells and tube formation of HUVEC cells were assessed.

Results: Percentage inhibition of DPPH scavenging activity were dose-dependent and ranged between (89.9% ± 0.51) and (28.6% ± 0.07). Phenolic contents ranged between (11.5 ± 0.013) and (9.7 ± 0.008) mg GAE/g while flavonoid content ranged between (20.8 ± 0.40) and (0.12 ± 0.001) mg QE/g. Antiproliferative results of the extracts were found to be consistent with their antioxidant activity. Among the extracts evaluated, that of *R. stricta* showed the best antioxidant, antiproliferative and antimetastatic activities at low concentration. It also inhibited the colony-formation capacity of HepG2 cells and exhibited antiangiogenic activity. Cell cycle analysis showed significant arrest of cells at G2/M phase 12 and 48 h after treatment and significant arrest at G1/S phase after 24 h of treatment. Consistent data were observed in western blot analysis of protein levels of Cdc2 and its cyclin partners.

Conclusions: These findings introduce *R. stricta* as a potentially useful anti-metastatic agent and a novel potential anti-tumour agent for hepatocellular carcinoma (HCC) treatment.

Keywords: Antioxidants, Anticancer, Medicinal plants, Traditional medicine, *Rhazya stricta*

Background

Free radicals are mainly produced by oxidation processes and they have an important role in the processes of food spoilage and chemical materials degradation. They also contribute to human disorders such as aging-associated diseases, cardiovascular diseases, cancer and inflammatory diseases [1, 2]. Free radicals may also cause a depletion of the immune system antioxidants, a change in the

gene expression and may induce the synthesis of abnormal proteins. About 5% or more of the inhaled oxygen (O₂) is converted to reactive oxygen species (ROS) such as O₂⁻, H₂O₂, and OH radicals. ROS represents the major type of free radicals in any biological system. They are produced through the mitochondrial electrons transport chain [2, 3].

Antioxidants are used to neutralize the effects of free radicals. Thus, they protect humans against infection and degenerative diseases. Antioxidants can be classified into two major categories, natural and synthetic. Synthetic antioxidants include butylated hydroxy anisole (BHA), butylated hydroxyl toluenes (BHT), tertiary butylated hydroquinone and gallic acid esters. These antioxidants effectively inhibit

* Correspondence: bayan.al-dabbagh@uaeu.ac.ae; a.amin@uaeu.ac.ae

¹Department of Chemistry, College of Science, UAE University, PO Box 15551, Al Ain, UAE

²Department of Biology, College of Science, UAE University, PO Box 15551, Al Ain, UAE

Full list of author information is available at the end of the article



oxidation, may serve as chelating agents such as ethylene diamine tetra acetic acid (EDTA), and can bind metals reducing their contribution to the process [4]. However, antioxidants are thought to cause or promote negative health effects such as mutagenesis and carcinogenesis in humans [5]. Therefore, there is a strong trend to replace the synthetic with naturally occurring antioxidants that can prevent free radical-related diseases [4, 6].

Natural antioxidants help in controlling the formation of free radicals and activated oxygen species or they can inhibit their reaction with biological structures [7]. These antioxidants include antioxidative enzymes, such as superoxide dismutase, catalase, and glutathione peroxidase, and small nonenzymatic antioxidant molecules, such as glutathione and vitamins C and E [8]. Many herbs and spices (rosemary, thyme, oregano, sage, basil, pepper, clove, cinnamon, nutmeg, and saffron), and plant extracts (tea, grapeseed, and lemon balm) contain antioxidant components [9–11].

In this study, three hydroalcoholic extracts of traditionally medicinal plants used in the United Arab Emirates (UAE) were evaluated for their antioxidant activities, phenol and flavonoid contents. These plants are, *Trigonella foenum-graecum*, locally known as “Helba”, *Cassia acutifolia*, commonly known as “Holoul” and *Rhazya stricta* which is locally called “Harmal”.

T. foenum-graecum seeds are traditionally used as herbal medicine for their carminative, tonic, aphrodisiac and anticancer effects [12–14]. The leaves of *C. acutifolia* is frequently used in folk medicine as a purgative for a long time [15]. The extracts of *R. stricta* leaves are traditionally used for the treatment of various disorders such as diabetes, sore throat, helminthiasis, inflammatory conditions and rheumatism [16].

Available treatment for hepatocellular carcinoma are mainly limited to invasive hepatectomy or chemotherapy. However, the attention has shifted in recent years to natural-based products for candidate anticancer therapeutics. In the present study, the antiproliferative effects of *Trigonella foenum-graecum*, *Cassia acutifolia*, and *Rhazya stricta* on hepatoma cell line HepG2 were investigated. The use of HepG2 cells to test the cytotoxic effects of a wide range of drugs has been well documented, due to their wide availability, well-differentiation, and drug metabolizing activity [17].

Despite playing a key role in cellular processes, free radicals pose a threat to cells by damaging DNA, proteins, and cellular membranes, leading to onset of many diseases including cancer [18, 19]. Thus, by decreasing free radicals and oxidative stress, antioxidants play a role in ameliorating DNA damage, reducing the rate of abnormal cell division, and decreasing mutagenesis [20]. Therefore, many antioxidant-rich plants possess anticancer activity [21–23].

Vascular endothelial growth factor (VEGF) has been recognized to be involved in several stages of angiogenesis

in malignant diseases by its multi-functional effects in activating and integrating signalling pathway networks [24]. VEGF signalling blockade reduces new vessel growth and leads to endothelial cell apoptosis. Therefore, using tyrosine kinase inhibitors or VEGF/VEGF receptor (VEGFR) antibodies to inhibit crucial angiogenic steps is a practical therapeutic strategy when treating neovascularisation diseases [25]. A potent angiogenesis inhibitor known as E7820, has been shown to reduce integrin $\alpha 2$ mRNA expression and inhibit basic fibroblast growth factor/VEGF-induced HUVEC proliferation and tube formation [26, 27]. Integrin $\alpha 2\beta 1/\alpha 1\beta 1$ expression is reportedly regulated by VEGF and an inhibitory antibody against $\alpha 2\beta 1/\alpha 1\beta 1$ has been shown to inhibit angiogenesis and tumour growth in VEGF-overexpressing tumour cells [28, 29]. Therefore, we investigated here the effect of *R. stricta* leaves extract on angiogenesis utilizing HUVEC tube formation assay; as it showed the most promising antiproliferative activity. In addition, this work was set to determine the in vitro antioxidant activity, total phenols and flavonoids, anticancer activities of tested plants with special interest in *R. stricta*.

Methods

Chemicals

All solvents were analytical grade. Agilent Cary 60 UV-Vis Spectrophotometer was used in all spectrophotometric measurements. Ascorbic acid, ferric chloride, aluminium chloride, potassium acetate, quercetin, DPPH reagent, Folin-Ciocalteu reagent, gallic acid, sodium carbonate, methanol and ethanol were obtained from Sigma Chemical Co. (St. Louis, MO, USA). Millipore deionized water was used throughout. Thiazolyl Blue Tetrazolium Bromide (Sigma Aldrich, USA), Dimethyl Sulfoxide (Sigma Aldrich, USA).

Plant samples

Dried leaves of *C. acutifolia*, *R. stricta* and *T. foenum-graecum* were purchased from the local market. The taxonomic authentication of all the plants was carried out by Dr. Fatima Al-Ansari at the Biology Department, College of Science, United Arab Emirates University. Voucher specimens were deposited at the herbarium of the Biology Department (voucher reference numbers: BA2018–1, BA2018–2, BA2018–3).

Preparation of plant extracts

The leaves of the medicinal plants were crushed separately in a grinder. A sample of 10 g of each plant was extracted with 150 mL of 70% ethanol and 30% water as it showed the best extraction yield [30]. The crushed plants were macerated for 48 h at 4 °C. The resulting mixture was then filtered under vacuum and concentrated under reduced pressure in a rotary evaporator at 40 °C. The extracts were further dried using a TELSTAR

CRYODOS freeze dryer machine then kept at $-20\text{ }^{\circ}\text{C}$ for further analysis. A solution of 30 mg/mL of each plant was prepared in 50% ethanol for the following tests.

Determination of total polyphenol content

The total phenolic content (TPC) was determined by using the Folin-Ciocalteu reagent [31]. A 10% solution was prepared from the stock solution (30 mg/mL) using 50% ethanol. 100 μl of this solution was mixed with 200 μl of the Folin-Ciocalteu reagent and 2 mL of de-ionized water then incubated at room temperature for 3 min. A sample of 20% aqueous sodium carbonate (*w/w*, 1 mL) was then added to the mixture. The total polyphenols were determined after 1 h of incubation at room temperature. A negative control sample was also prepared using the same procedure. The absorbance of the resulting blue colour was measured at 765 nm. Results were expressed in mg gallic acid equivalents (GAE) per g dry weight of plant material using an equation obtained from gallic acid calibration curve. The samples were analyzed in triplicate.

Free-radical scavenging activity

The antioxidant activity of the extracts was assessed based on their ability to scavenge the stable 1,1-diphenyl-2-picrylhydrazyl (DPPH) radical as described previously [32]. Various concentrations of the three extracts in methanol were prepared (0.15 to 1.5 mg/mL). A methanolic solution of DPPH (3.8 mL, 60 $\mu\text{g/mL}$) was rapidly mixed with the plant extract (200 μl , 30 mg/mL) in a test tube, with methanol serving as the blank sample and a control was also assayed simultaneously. The contents of the tubes were swirled then allowed to stand for 30 min at room temperature in the dark. The absorbance was measured at 517 nm in a spectrophotometer. The scavenging ability of the plant extract was calculated using this equation: DPPH Scavenging activity (%) = [(Abs control - Abs sample) / (Abs control)] \times 100, where Abs control is the absorbance of DPPH + methanol; Abs sample is the absorbance of DPPH radical + sample (sample or standard). The EC_{50} value ($\mu\text{g/mL}$), the effective concentration at which DPPH radicals are scavenged by 50%, was determined graphically. The total antioxidant activity was expressed as ascorbic acid equivalent/g dry extract. The assay was done in triplicates.

Determination of total flavonoids

The total flavonoids content in the extracts was determined using the aluminium chloride colorimetric method [33]. A known concentration (600 $\mu\text{g/mL}$) of each extract in methanol was prepared. A 500 μl of the extracts were mixed separately with 0.1 mL of 10% (*w/v*) aluminium chloride solution, 0.1 mL of 1 M potassium acetate solution, 1.5 mL of methanol and 2.8 mL of

distilled water. The solutions were thoroughly mixed and incubated at room temperature for 30 min. The absorbance of the reaction mixture was measured at 415 nm using a spectrophotometer. The total flavonoids content was determined using a standard curve with quercetin (1 to 25 $\mu\text{g/mL}$) as the standard. The mean of three readings was used and expressed as mg of quercetin equivalents (QE)/ g of the dry extract.

Cell culture

A human hepatocellular carcinoma (HCC)-derived cell line (HepG2) was cultured in RPMI 1640 medium containing 1% antibiotic cocktail and supplemented with 10% fetal bovine serum. Cells were incubated at $37\text{ }^{\circ}\text{C}$ in 5% CO_2 humidified incubator. Cells were passaged every 2–3 days using 0.25% trypsin-EDTA.

Cytotoxicity assay

HepG2 were seeded at a density of 5000 cells/well in a 96-well plate, and were allowed to attach overnight. Thereafter, cells were treated with various concentrations of the plants extracts for 24 h. To assess the cytotoxic effect of the three plants extracts, MTT (3-[4,5-dimethylthiazol-2-yl]-2,5-diphenyltriazolium bromide) assay was carried out. Briefly, cells treated with the plant extracts were exposed to tetrazolium MTT at a concentration of 5 mg/mL. Viable active cells reduced yellow MTT salt to insoluble purple formazan, which was dissolved using DMSO. The absorbance of the coloured solution was measured at a wavelength of 570 nm using Epoch microplate spectrophotometer (BioTek). The obtained absorbance at 570 nm of both control and treated cells was used to calculate percentage of cell viability. Assuming 100% viability in control cells, percentage of treated cells viability will be calculated accordingly:

Percent of viable cells

$$= (\text{Abs. of treated cells} / \text{Abs. of control cells}) \times 100$$

Assessment of morphological changes

HepG2 were seeded at a density of 0.25×10^6 cells/ well in a 6-well plate, and were allowed to attach overnight. After which, cells were treated without (0 μg , control) or with increasing concentrations of *R. stricta* (10, 20, 30, 50, 70 μg) for 24 h. The morphology of the cells was assessed after being fixed and stained with 0.5% crystal violet using bright-field microscopy (200 \times magnification, scale = 200 μm).

Colony formation assay

To assess the effects of *R. stricta* on cell survival, the colony formation assay was carried out in vitro. Briefly, HepG2 cells were seeded at a density of 1000 cells/ well

in a 6-well plate, and were incubated for 24 h to allow attachment. The second day, the cells were treated without (0 µg, control) or with increasing concentrations (10, 20, 30 µg) of the extract for 24 h. After which, the media was replaced with fresh complete growth media without the extract, and cells were left to incubate until visible colonies were formed; while changing the media every 3–4 days. The experiment was carried out in triplicates. Colonies were fixed with absolute methanol, then stained with 0.5% crystal violet. Results are represented as the percentage of the well area that is covered by colonies (colony area percentage). Analysis has been carried out using ImageJ plugin ColonyArea [34]. In addition, an absorption-based method was carried out to validate the earlier results, by which the absorption of the crystal violet dye in each well is measured after being dissolved. Briefly, the samples that had been analysed using ImageJ were subjected to 10% acetic acid solution, then were placed on an orbital shaker for 15 min. After which, 100 µL of each triplicate sample was transferred to a 96-well plate (in triplicates), and absorbance was measured using Epoch microplate spectrophotometer (BioTek).

Wound-healing assay

To assess the ability of HepG2 to migrate after the treatment with *R. stricta*, wound-healing assay was carried out in vitro. Cells were seeded at a density of 0.5×10^6 /well in 6-well plate, and were allowed to attach overnight. A scratch in the cell monolayer was made using a sterile plastic pipetting tip, and then the monolayer was washed with PBS. The cells were treated without (0 µg) or with 20, 30 µg of the extract. Images were taken at 0, 24, 48, 72 h using bright-field microscopy (40 x magnification). Analysis was carried out using ImageJ, percent of open area was calculated according to the following formula [$T_x = T_{24}, T_{48},$ or T_{72} (at time 24, 48, or 72 h, respectively)]: Percent of open area = (open area at T_x / open area at T_0) X 100.

Experiment was carried out in triplicates, data is representative of 3 random regions in each triplicate of each sample.

Cell cycle analysis

Effect of *R. stricta* extract on cell cycle progression of HepG2 cells was analysed as previously described [35]. Cells were treated without or with 30 µg of *R. stricta* extract at different time intervals (6 – 48 h), collected by trypsinization, washed twice with PBS, fixed in 70% ethanol, treated with RNase, stained with propidium iodide and then cell cycle distribution was analysed in BD Accuri C6 cytometer and software (BD Biosciences, USA).

Western blotting

HepG2 were seeded at a density of 1×10^6 in 60 mm dish and allowed to attach overnight. Cells were treated without or with 30 µg of extract for 6, 12, 24, 48 h. Cells

were lysed and total protein was quantified using BCA. 20 µg of total protein was separated on SDS-PAGE, transferred onto nitrocellulose membranes that were blocked using 5% BSA TBST. Primary antibodies against Cdc2 (1:000; cell signalling), p-Cdc2 (1:1000; cell signalling), Cyclin B1 (1:1000; cell signalling), Cyclin A1 (1:000; Abcam) were used. GAPDH (1:15000; Abcam) was used as loading control. Proteins were detected using LI-COR C-DiGit Chemiluminescence Western Blot Scanner.

Matrigel capillary tube formation

96-well plate was coated with Matrigel matrix (Corning, NY, USA) at 50 µl/well and allowed to polymerize for 60 min at 37 °C. HUVEC cells were then seeded on the Matrigel at a concentration of 2×10^4 cells/well without (0 µg) or with (10, 20, 30 µg) *R. stricta* extract. After incubation for 18 h, tubules were imaged using an inverted microscope and analyzed with ImageJ software.

Statistical analysis

All data were expressed as mean ± standard deviation (SD) of three independent experiments. Correlation analysis of antioxidants versus the total phenolic and flavonoid contents were carried out using the regression analysis, with GraphPad Prism 6.0 and Microsoft Excel 2016. $P < 0.05$ was considered to indicate a significant difference.

Results and discussion

Medicinal plants have been of great interest as a source of natural antioxidants used for health promotion. The therapeutic activity of plants is mostly due to their biologically active polyphenolic substances, mostly flavonoids and phenolic acids. These substances exhibit antioxidant, anti-lipoxygenase and anticancer activities. The present study elaborates on the antioxidant activity, polyphenolic and flavonoid contents of three folk plants from the UAE; *T. foenum-graecum*, *C. acutifolia* and *R. stricta*. The antiproliferative effect of such plants was studied against human cancer cells HepG2 in an attempt to find a correlation with the antioxidant activity of those extracts that are based on their phenolic and flavonoid contents.

Plant extraction


Different solvents have been used in the literature for the preparation of plant extracts [36]. In this study, we used 70% ethanol as the extraction solvent. The amorphous solid of the leave extracts under investigation was obtained by complete evaporation of ethanol/water. The yield of each extract was calculated as w/w percent yield. The yields of *T. foenum-graecum*, *C. acutifolia* and *R. stricta* extracts were 25, 23 and 30% respectively.

Total polyphenol content of the extracts

Polyphenols are aromatic secondary plant metabolites and are widely spread throughout plants. They have been associated with colour, sensory qualities, and nutritional and antioxidant properties of food [37]. It is reported that there is a strong relationship between total polyphenol contents and antioxidant activity. The hydroxyl groups in phenols have a strong scavenging ability for free radicals. Therefore, the total polyphenol contents of plants may directly contribute to their antioxidant activity [32, 38]. The Folin-Ciocalteu reagent is commonly used in the literature to determine phenolic compounds. This reagent reacts with phenolic compounds and gives a blue colour complex that absorbs radiation and allows quantification [39]. The total phenolic content for the ethanolic extracts of *C. acutifolia*, *R. stricta* and *T. foenum-graecum* was determined by the Folin-Ciocalteu method using gallic acid as a standard. The calibration curve showed linearity for gallic acid in the range of 0.5–26 $\mu\text{g/mL}$, with a correlation coefficient (R^2) of 0.984. *R. stricta* contained the highest total polyphenols (11.5 ± 0.013 mg GAE/g extract), followed by *C. acutifolia* (10.8 ± 0.025 mg GAE/g extract) and *T. foenum-graecum* (9.7 ± 0.008 mg GAE/g extract) (Fig. 1). Belguith-Hadriche et al. [40] reported the total phenolic content of various extracts of *T. foenum-graecum* ranged between 9.42 ± 0.50 in hexane and 78.1 ± 0.90 mg GAE/g dry weight extracts in methanol. A study conducted in Saudi Arabia showed the total phenolic content of *R. stricta* extracts ranged between 62.5 ± 0.2 and 66.63 ± 0.03 mg GAE/g extract [41].

The DPPH radical scavenging activity


The antioxidant activity of each plant was also assessed based on its ability to reduce the stable DPPH radical according to the method reported by Lim [32]. The DPPH



radical (DPPH \cdot) is a stable radical and has the ability to accept an electron or hydrogen radical and form a stable diamagnetic molecule producing a colour change from blue to yellow [42]. The colour change of DPPH has been widely used to measure the radical scavenging activity because of its stability, simplicity, and reproducibility [43]. The free radical scavenging capacity of the ethanolic extracts of the three plants were assayed based on the remaining amount (%) of DPPH \cdot as a function of time (30 min). The total antioxidant activity was expressed as ascorbic acid equivalent/g dry extract.

The calibration curve of ascorbic acid showed linearity in the range of 5–20 $\mu\text{g/mL}$, with a correlation coefficient (R^2) of 0.994. The percentage inhibitions of DPPH scavenging activity in all the extracts were dose-dependent (Fig. 2). The DPPH scavenging activity by the *T. foenum-graecum* extract was ($89.7\% \pm 1.54$) at 1.5 mg/mL and ($28.6\% \pm 2.07$) at 0.15 mg/mL. DPPH of *C. acutifolia* was ($86.3\% \pm 0.64$) at 1.5 mg/mL and ($30.0\% \pm 1.37$) at 0.15 mg/mL while that of *R. stricta* was ($89.9\% \pm 0.51$) at 1.5 mg/mL and ($28.7\% \pm 1.27$) at 0.15 mg/mL. The results showed that there is no clear difference between the obtained DPPH scavenging activities of the three extracts. These results are almost in agreement with their total polyphenol contents. Consequently, the antioxidant activity of these plants might be related to their contents of phenolic compounds.

In addition, the DPPH radical scavenging ability of the extracts was evaluated as EC_{50} ($\mu\text{g/mL}$) value. The smallest EC_{50} values indicates the best free radical scavenging activity. The highest scavenging activity was exerted by *R. stricta* ($EC_{50} = 241.8$ $\mu\text{g/mL}$) which contained the highest amount of total polyphenol content (11.5 $\mu\text{g/mL}$), followed by *C. acutifolia* ($EC_{50} = 244.8$, TPC = 10.8 $\mu\text{g/mL}$). The lowest radical scavenging activity was exhibited by the *T. foenum-graecum* extract ($EC_{50} = 245.9$, TPC = 9.7 $\mu\text{g/mL}$). The relationship between total phenol content and free radical scavenging activity (using EC_{50}) was also studied using linear regression analysis (Fig. 3a). The results showed a






Fig. 3 Linear correlations between the amount of total phenols and DPPH· radical scavenging activity (a) and between the flavonoid content and DPPH· radical scavenging activity (b)

significant negative correlation ($R^2 = -0.856$, P -value < 0.05) between EC_{50} (DPPH· scavenging) and total phenolic content suggesting that the presence of the phenolic compounds contributed significantly to the antioxidant activity of the tested plants. These results are consistent with previous works that showed a linear correlation between the total phenolic content and the reducing antioxidant capacity of some plant extracts [32, 38, 44].

Total flavonoid contents

Flavonoids are a class of secondary plant phenolics. Flavonoids and their derivatives have a wide range of biological actions including anticancer activity. The anticancer activity of flavonoids is attributed to their potent antioxidant effects which include metal chelation and free-radical scavenging activities [45]. Flavonoids present in herbs were found to significantly contribute to their antioxidant properties [44]. The flavonoid content was obtained using aluminium chloride assay which based on the formation of a complex between the aluminium ion, Al (III), and the carbonyl and hydroxyl groups of flavones and flavonols that produce a yellow colour [46]. Flavonoid content was calculated from the regression equation of quercetin calibration curve and was expressed as quercetin equivalents. The calibration curve showed linearity in the range of 1–25 µg/mL, with a correlation coefficient (R^2) of 0.999.

Figure 4 shows the flavonoids contents in all the extracts. Using the standard curve generated by quercetin, the total flavonoids content of the *T. foenum-graecum* extract was $(14.6 \pm 0.21$ mg QE/g), whereas, the *C. actitifolia* extract was $(20.8 \pm 0.40$ mg QE/g) and finally that of *R. stricta* was $(9.2 \pm 0.22$ mg QE/g).

The relationship between the total flavonoids content and the free radical scavenging activity (using EC_{50}) was studied using linear regression analysis (Fig. 3b). The results showed a positive correlation ($R^2 = 0.460$, P -value < 0.05) between EC_{50} (DPPH scavenging) and total flavonoids content. The obtained correlation was moderate

suggesting that other compounds maybe participating in the radical scavenging activity of these plant extracts.

Effects of extracts on cell viability in HepG2 cells

HCC remains among the leading cause of cancer-related death worldwide [47, 48]. Although therapeutic approaches for advanced HCC are limited to the use of multikinase inhibitors, such as sorafenib, only modest survival benefits have clinically been reported. Thus, identifying new compounds with promise antitumor activity against HCC is exceedingly needed [48, 49]. Natural products, especially from plants, are often better tolerated than their synthetic analogs used in cancer treatments [50]. They contain a wide spectrum of bioactive secondary metabolites that are the foundation of the recently introduced notion of broad-spectrum integrative approach for cancer prevention and treatment [51].

Effects of tested extracts were investigated against human hepatoma (HepG2) cancer cell line. A dose-dependent reduction in cell viability was reported in cells treated with all tested extracts (Fig. 5). The IC_{50} values of extracts ranged from 30 µg/mL to 200 µg/mL. Treatment with *R. stricta* significantly enhanced the mortality of cancer cells at the lowest concentration (30 µg/mL). *T. foenum-graecum* was





Fig. 4 Flavonoids content of the tested extracts



however less potent with IC₅₀ of 200 µg/mL where 50% of HepG2 cancer cells were eradicated at 200 µg/mL. A fenugreek-enriched diet decreased colon tumour incidence and hepatic lipid peroxidation in liver cancer-induced rats in addition to increasing the endogenous antioxidant activities in liver [52]. Li et al. [53] showed that diosgenin, fenugreek's main active ingredient, down regulated the expression of various STAT3-regulated genes, inhibited proliferation and potentiated the apoptotic effects of paclitaxel and doxorubicin, suggesting that diosgenin could be a novel and potential treatment option for HCC and other cancers. Therefore, the role of fenugreek extract and its active principals as supplements in diet-based preventive/therapeutic strategies to improve health care continues to be a fast growing field of research. The correlation between the antioxidant activities (free radical scavenging activity) and the anticancer activities (cell viability) of *C. acutifolia*,

R. stricta and *T. foenum-graecum* extracts was studied using linear regression analysis. The results showed a significant positive correlation ($R^2 = -0.933, 0.997$ and 0.797 , P -value < 0.05) for *C. acutifolia*, *R. stricta* and *T. foenum-graecum* extracts respectively.

At a concentration range of (100–200 µg/mL) *Cassia acutifolia*'s extract was similarly cytotoxic to HepG2 cells. Alkaloids extracted from Senna species reduced cell viability in a concentration-dependent manner of different tumour cell lines including HepG2 [49]. Senna alkaloids showed important antiproliferative activity on HepG2 cells that was mediated by ERK inactivation and down-regulation of cyclin D1 expression. Similarly, extracts of different cassia species were able to inhibit growth of colorectal (DLD1), among other, human cancer cell lines [54]. Thus, Senna's extract may represent a potential new antitumor and/or adjuvant treatment

against liver and colorectal cancer and further investigations should be conducted to unravel its molecular mechanism. *R. stricta*'s extract was by far the most effective cytotoxic agent tested at the present study at as little as 30 µg/mL. The δ-tocopherol and alkaloid fraction of *R. stricta*'s leaves extract have been shown to delay many angiogenic and inflammatory activities and to inhibit cell viability of HepG2, hence potentially useful against cancer [55, 56]. As *R. stricta* extract showed the most promising antiproliferative activity, its antiproliferative effects were tested against other GI cell line. The present study also showed that *R. stricta* extract inhibited cell proliferation in colorectal cancer cells (HCT116), in a dose-dependent manner (Additional file 1: Figure S1). To get insight into the molecular effects of *R. stricta* on HepG2 cells, its effect on cell cycle progression was analysed. *R. stricta* treatment induced arrest of cells in the G2/M phase (12 and 48 h time points) and in G1/S phase (24 h time point) (Fig. 5d and e). This cell cycle arrest reflects the cytotoxic effects of *R. stricta* on HepG2 cells where the cells are arrested at these phases to repair the toxic lesions induced by *R. stricta* or to be removed by death pathways such as apoptosis. The significant increase of the fraction of cells in G2/M after long term treatment (48 h) may indicate permanent arrest of cells in this phase (quiescence). These findings are consistent with western blot data, where cell cycle promoting proteins cdc2 and its cyclin partners (cyclin A1 and cyclin B1) are down regulated post *R. stricta* treatment (Fig. 5f). Cdc2 activity is regulated through two mechanisms, first by binding to its cyclin partners [57], and second by being phosphorylated at Tyr15, and subsequently inhibited [58, 59]. Cdc2/cyclin complexes contribute to cell cycle progression through phosphorylation of target proteins [60]. As shown, Cdc2, and its partners cyclin A and cyclin B were downregulated in a time-dependent manner (Fig. 5f). In addition, treatment with *R. stricta* extract resulted in a slight increase in Tyr15 phosphorylated Cdc2 (Fig. 5f).

***R. stricta* extract inhibited the colony formation of HepG2 cells**

This assay measures the ability of tumour cells to survive and grow to form colonies after treatment with cytotoxic agents. To assess clonogenicity, cells were plated onto 6-well plates and incubated with 0.0, 10, 20, and 30 µg of *R. stricta* extract and treated cultures were maintained in culture for an additional 10 days to allow formation of colonies [61, 62]. The results from that assay showed that *R. stricta* extract inhibited the colony formation of HepG2 cells in a dose-dependent manner (0, 10, 20, 30 µg). ImageJ analysis results were consistent with results obtained from the absorption-based method where *R. stricta* extract also inhibited the survival of HepG2 cells in a dose-dependent manner (Fig. 6a and b).

***R. stricta* extract inhibits the migration ability of HepG2 cells**

Many cancer patients die of tumour mobilization which often lead to metastatic foci in distant part/s of the body. Therefore, attempts to prevent or slowdown progression and metastasis of cancer cells are crucial [63]. Earlier study showed that *R. stricta* extract exerts potent anti-cancer ability against human breast cancer [64]. To date, however, limited information is available regarding *R. stricta* extract effect on cell migration in any cancer type. The potential role of *R. stricta* extract on HepG2 cells' migration was assessed using wound healing assay. Cell migration was inhibited in a dose-dependent manner (0, 20 and 30 µg) starting from 24 to 72 h post *R. stricta* extract treatments compared to the control (Fig. 6c).

***R. stricta* extract inhibits endothelial cell tube formation**

Cell migration and invasion are crucial steps in various physiological processes such as morphogenesis, angiogenesis, wound healing and inflammation [65]. Invasion and migration are critical events for tumour progression and tumour recurrence [66, 67]. HUVEC cells had greater numbers of branching tube networks after 18 h of incubation with no *R. stricta* extract (Fig. 7a). This tube branching was attenuated by *R. stricta* extract treatments in a dose-dependent manner (Fig. 7b and c). Colony formation results demonstrate that colony numbers in all dose groups were decreased. These results suggest that *R. stricta* extract reduced cell migration and colony formation capacity in HepG2 cancer cells. *R. stricta* extract may, therefore, be introduced as a novel agent to treat/prevent HCC and as a single agent or in combination with other drugs. It would also be of interest to identify *R. stricta* extract bioactive molecules and assess their effects against cancer, particularly HCC. Further studies are currently underway to identify and characterize *R. stricta* extract bioactive ingredients and to unravel their molecular mechanism against cancer.

The obtained antiproliferative results of the three extracts against HepG2 cancer cell line were found to be consistent with their antioxidant activity, free radical scavenging ability and phenolic content. Out of the three extracts, *R. stricta* showed considerable antiproliferative activity at low concentration and was the most potent among the tested plants with $IC_{50} = 30$ µg/mL. In addition, *R. stricta* showed the highest total polyphenols (11.5 ± 0.013 mg GAE/g extract). It has also the most percentage inhibitions of DPPH scavenging activity ($89.9\% \pm 0.51$) at 1.5 mg/ml and ($28.7\% \pm 1.27$) at 0.15 mg/ml. This indicates that the antiproliferative effect of the three extracts, in particular *R. stricta*, may be attributed to their antioxidant polyphenolic and flavonoid contents. Phenolic compounds have been found to counteract cancer either by means of




Fig. 6 *Rhazya stricta* extract inhibits colony forming ability and wound healing of HepG2 cells in a dose-dependent manner. **a** Representative images of HepG2 colonies after treatment with increasing concentrations of *R. stricta* extract. **b** Percent of area occupied by colonies in treated and non-treated wells (representative of triplicate samples; * $P < 0.05$, ** $P < 0.005$, *** $P < 0.0001$) and absorbance of each treated and non-treated wells (representative of 3 biological triplicates, each in triplicate; * $P < 0.05$, ** $P < 0.001$, *** $P < 0.0001$). **c** Representative images and quantification (three regions of three biological triplicates; * $P < 0.05$, ** $P < 0.001$, *** $P < 0.0001$) of wound-healing assay results of HepG2 cells treated without or with 20 and 30 µg of *R. stricta* extract




Fig. 7 *Rhazya stricta* extract inhibits tube formation in HUVECs on Matrigel in a dose-dependent manner. Photographs of tube formation in HUVECs on Matrigel after incubation with or without *R. stricta* extract at 18 h. Cells were treated with *R. stricta* extract at a series of concentrations (10 µg; **b**, 20 µg; **c**, 30 µg; **d**) or DMSO vehicle (control; **a**) for 18 h

antioxidant effect or by inhibiting the formation of carcinogenic metabolites that damage the vital biomolecules [68].

Finally, the investigated UAE plants have shown a strong reducing antioxidant capacity and free radical scavenging ability. These findings validate the use of these plants in folk medicine for the treatment of certain diseases. Oxidative cell damage events are frequently correlated with the oxidative stress. We show here that both of those properties are present in the ethanolic extract of *R. stricta*, which has significant antiproliferative activity and a great antioxidant activity. The results reported here are very promising indicators for the potential application of this plant for preventive and therapeutic purposes.

Conclusions

In conclusion, natural antioxidants have many important applications in health promotion, food preservation, food flavouring and cosmetics. They are preferred over synthetic antioxidants because they are safer for consumption and more environmentally friendly. The present study investigates the antioxidant activity and polyphenolic content of three medicinal plants from the UAE. We found extracts rich in antioxidants and in polyphenols, which merit further investigations. Of the extracts evaluated, that of *R. stricta* (Harmal) showed the greatest antioxidant, antiangiogenic and antiproliferative activities, a discovery that makes this species a promising source of anticancer agent development especially for colon and liver cancers, and

hence worthy of further investigation. Isolation of active compounds and exploring their mode of action against tumours by using in vivo experimental models would make an important future study.

Additional file

Additional file 1: Figure S1. Assessment of the cytotoxic effects of *Trigonella foenum-graecum* (Helba), *Cassia acutifolia* (Holoul), and *Rhazya stricta* (Harmal) extracts on HCT116 in vitro. MTT assay results of 9cHCT116 cells viability after treatment with increasing concentrations of Helba (a) Holoul (b) and Harmal (c) for 24 h. * $P < 0.05$, ** $P < 0.005$, *** $P < 0.0001$. (DOCX 99 kb)

Acknowledgments

This work was supported by a start-up grant (Grant 31S 215), Division of Research and Graduate Studies, UAEU and an individual research grant (Grant 31S 188), College of Science, UAEU to the PI: Dr. Bayan Al-Dabbagh and by Zayed Bin Sultan Centre for Health Sciences (ZCHS) grant number 31R050 for Dr. Amr Amin.

Funding

This work was supported by a start-up grant (Grant 31S 215), Division of Research and Graduate Studies, UAEU and an individual research grant (Grant 31S 188), College of Science, UAEU to the PI: Dr. Bayan Al-Dabbagh and by Zayed Bin Sultan Centre for Health Sciences (ZCHS) grant number 31R050 for Dr. Amr Amin.

Availability of data and materials

"The datasets used and/or analyzed during the current study are available from the corresponding authors upon reasonable request."

Authors' contributions

BA and AA designed the research. BA, IAE, AAH, RA and RE performed the experiments. SSA participated in the interpretation of the results. BA and AA analyzed the data and wrote the paper. All authors read and approved the final manuscript.

Ethics approval and consent to participate

Not applicable

Consent for publication

Not applicable

Competing interests

The authors declare that they have no competing interests.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Author details

¹Department of Chemistry, College of Science, UAE University, PO Box 15551, Al Ain, UAE. ²Department of Biology, College of Science, UAE University, PO Box 15551, Al Ain, UAE. ³Zoology Department, Cairo University, Giza, Egypt. ⁴Department of Pharmacy Practice and Pharmacotherapeutics, Sharjah Institute for Medical Research and College of Pharmacy, University of Sharjah, Sharjah, UAE.

Received: 28 March 2018 Accepted: 11 July 2018

Published online: 22 August 2018

References

- Aruoma OI. Free radicals, antioxidants and international nutrition. *Asia Pac J Clin Nutr.* 1999;8:53–63.
- Gupta VK, Sharma SK. Plants as natural antioxidants. *Nat Prod Rad.* 2006;5:326–34.
- Magder S. Reactive oxygen species: toxic molecules or spark of life? *Crit Care.* 2006;10:208.
- Brewer MS. Natural antioxidants: sources, compounds, mechanisms of action, and potential applications. *Compr Rev Food Sci Food Saf.* 2011;10:221–47.
- Lu LY, Ou N, Lu Q-B. Antioxidant induces DNA damage, cell death and mutagenicity in human lung and skin normal cells. *Sci Rep.* 2013;3:3169.
- Soobrattee MA, Neergheen VS, Luximon-Ramma A, Aruoma OI, Bahorun T. Phenolics as potential antioxidant therapeutic agents: mechanism and actions. *Mutat Res.* 2005;579:200–13.
- Chaudière J, Ferrari-Iliou R. Intracellular antioxidants: from chemical to biochemical mechanisms. *Food Chem Toxicol.* 1999;37:949–62.
- Fridovich I. Fundamental aspects of reactive oxygen species, or what's the matter with oxygen? *Ann N Y Acad Sci.* 1999;893:13–8.
- Hinneburg I, Damien Dorman HJ, Hiltunen R. Antioxidant activities of extracts from selected culinary herbs and spices. *Food Chem.* 2006;97:122–9.
- Amin A, Hamza AA, Bajbouj K, Ashraf SS, Daoud S. Saffron: a potential candidate for a novel anticancer drug against hepatocellular carcinoma. *Hepatology.* 2011;54:857–67.
- Hamza AA, Ahmed MM, Elwey HM, Amin A. Melissa officinalis protects against doxorubicin-induced cardiotoxicity in rats and potentiates its anticancer activity on MCF-7 cells. *PLoS One.* 2016;11:e0167049.
- Xue WL, Li XS, Zhang J, Liu YH, Wang ZL, Zhang RJ. Effect of *Trigonella foenum-graecum* (fenugreek) extract on blood glucose, blood lipid and hemorheological properties in streptozotocin-induced diabetic rats. *Asia Pac J Clin Nutr.* 2007;16(Suppl 1):422–6.
- Amin A, Alkaabi A, Al-Falasi S, Daoud SA. Chemopreventive activities of *Trigonella foenum-graecum* (fenugreek) against breast cancer. *Cell Biol Int.* 2005;29:687–94.
- El Bairi K, Ouzir M, Agnieszka N, Khalki L. Anticancer potential of *Trigonella foenum-graecum*: cellular and molecular targets. *Biomed Pharmacother.* 2017;90:479–91.
- Seethapathy GS, Ganesh D, Kumar JUS, Senthilkumar U, Newmaster SG, Ragupathy S, et al. Assessing product adulteration in natural health products for laxative yielding plants, Cassia, Senna, and Chamaecrista, in southern India using DNA barcoding. *Int J Legal Med.* 2015;129:693–700.
- Elkady AI. Crude alkaloid extract of *Rhazya stricta* inhibits cell growth and sensitizes human lung cancer cells to cisplatin through induction of apoptosis. *Genet Mol Biol.* 2013;36:12–21.
- Faqi AS. A comprehensive guide to toxicology in preclinical drug development: Academic Press; 2013.
- Diplock AT, Charuleux JL, Crozier-Willi G, Kok FJ, Rice-Evans C, Roberfroid M, et al. Functional food science and defence against reactive oxidative species. *Br J Nutr.* 1998;80:577–5112.
- Valko M, Leibfritz D, Moncol J, Cronin MT, Mazur M, Telser J. Free radicals and antioxidants in normal physiological functions and human disease. *Int J Biochem Cell Biol.* 2007;39:44–84.
- Shigenaga MK, Ames BN. Oxidants and mitogenesis as causes of mutation and cancer: the influence of diet. *Basic Life Sci.* 1993;61:419–36.
- Greenwell M, Rahman P. Medicinal plants: their use in anticancer treatment. *Int J Pharm Sci Res.* 2015;6:4103.
- Alam AHMK, Hossain ASMS, Khan MA, Kabir SR, Reza MA, Rahman MM, et al. The Antioxidative Fraction of White Mulberry Induces Apoptosis through Regulation of p53 and NFκB in EAC Cells. *PLoS ONE.* 2016;11:e0167536.
- Islam S, Nasrin S, Khan MA, Hossain ASMS, Islam F, Khandokhar P, et al. Evaluation of antioxidant and anticancer properties of the seed extracts of *Syzygium fruticosum* Roxb. growing in Rajshahi, Bangladesh. *BMC Complement Altern Med.* 2013;13:142.
- Carmeliet P. Angiogenesis in life, disease and medicine. *Nature.* 2005;438:932.
- Bar J, Goss GD. Tumor vasculature as a therapeutic target in non-small cell lung cancer. *J Thorac Oncol.* 2012;7:609–20.
- Keizer RJ, Funahashi Y, Semba T, Wanders J, Beijnen J, Schellens J, et al. Evaluation of α2-integrin expression as a biomarker for tumor growth inhibition for the investigational integrin inhibitor E7820 in preclinical and clinical studies. *AAPS J.* 2011;13:230–9.
- Mita M, Kelly KR, Mita A, Ricart AD, Romero O, Tolcher A, et al. Phase I study of E7820, an oral inhibitor of integrin α2 expression with antiangiogenic properties, in patients with advanced malignancies. *Clin Cancer Res.* 2011;17:193–200.
- Senger DR, Perruzzi CA, Streit M, Koteliansky VE, de Fougères AR, Detmar M. The α1β1 and α2β1 integrins provide critical support for vascular endothelial growth factor signaling, endothelial cell migration, and tumor angiogenesis. *Am J Pathol.* 2002;160:195–204.
- Senger DR, Claffey KP, Benes JE, Perruzzi CA, Sergiou AP, Detmar M. Angiogenesis promoted by vascular endothelial growth factor: regulation through α1β1 and α2β1 integrins. *Proc Natl Acad Sci U S A.* 1997;94:13612–7.
- Amzad Hossain M, Shah MD. A study on the total phenols content and antioxidant activity of essential oil and different solvent extracts of endemic plant *Merremia borneensis*. *Arab J Chem.* 2015;8:66–71.
- Singleton VL, Orthofer R, Lamuela-Raventós RM. Analysis of total phenols and other oxidation substrates and antioxidants by means of folin-ciocalteu reagent: *Methods Enzymol Academic Press*; 1999. p. 152–78.
- Lim YY, Quah EPL. Antioxidant properties of different cultivars of *Portulaca oleracea*. *Food Chem.* 2007;103:734–40.
- Chang C-C, Yang M-H, Wen H-M, Chern J-C. Estimation of total flavonoid content in propolis by two complementary colorimetric methods. *J Food Drug Anal.* 2002;10:178–82.
- Guzmán C, Bagga M, Kaur A, Westermarck J, Abankwa D. ColonyArea: an ImageJ plugin to automatically quantify Colony formation in Clonogenic assays. *PLoS One.* 2014;9:e92444.
- Saleh E, El-Awady R, Anis N. Predictive markers for the response to 5-fluorouracil therapy in cancer cells: constant-field gel electrophoresis as a tool for prediction of response to 5-fluorouracil-based chemotherapy. *Oncol Lett.* 2013;5:321–7.
- Khan MA, Rahman AA, Islam S, Khandokhar P, Parvin S, Islam MB, et al. A comparative study on the antioxidant activity of methanolic extracts from different parts of *Morus alba* L. (Moraceae). *BMC Res Notes.* 2013;6:24.
- Robbins RJ. Phenolic acids in foods: an overview of analytical methodology. *J Agric Food Chem.* 2003;51:2866–87.
- Wojdyło A, Oszmiański J, Czemerys R. Antioxidant activity and phenolic compounds in 32 selected herbs. *Food Chem.* 2007;105:940–9.
- Pontis JA, LAMAd C, SJRd S, Flach A. Color, phenolic and flavonoid content, and antioxidant activity of honey from Roraima, Brazil. *Food Sci Technol (Campinas).* 2014;34:69–73.
- Belguith-Hadrieh O, Bouaziz M, Jamoussi K, Simmonds MS, El Feki A, Makni-Ayedi F. Comparative study on hypocholesterolemic and antioxidant activities of various extracts of fenugreek seeds. *Food Chem.* 2013;138:1448–53.
- Bukhari NA, Al-Otaibi RA, Ibrahim MM. Phytochemical and taxonomic evaluation of *Rhazya stricta* in Saudi Arabia. *Saudi J Biol Sci.* 2017;24:1513–21.
- Robards K, Prenzler PD, Tucker G, Swatsitang P, Glover W. Phenolic compounds and their role in oxidative processes in fruits. *Food Chem.* 1999;66:401–36.
- Kitts DD, Wijewickreme AN, Hu C. Antioxidant properties of a north American ginseng extract. *Mol Cell Biochem.* 2000;203:1–10.
- Shan B, Cai YZ, Sun M, Corke H. Antioxidant capacity of 26 spice extracts and characterization of their phenolic constituents. *J Agric Food Chem.* 2005;53:7749–59.

45. Amin A, Mousa M. Merits of anti-cancer plants from the Arabian gulf region. *Cancer Ther.* 2007;5:55–66.
46. Popova M, Bankova V, Butovska D, Petkov V, Nikolova-Damyanova B, Sabatini AG, et al. Validated methods for the quantification of biologically active constituents of poplar-type propolis. *Phytochem Anal.* 2004;15:235–40.
47. Amin A, Hamza AA, Daoud S, Khazanehdari K, Hrouf AA, Baig B, et al. Saffron-based Crocin prevents early lesions of liver Cancer: in vivo, in vitro and network analyses. *Recent Pat Anticancer Drug Discov.* 2016;11:121–33.
48. Wu C-T, Tsai Y-T, Lai J-N. Demographic and medication characteristics of traditional Chinese medicine users among colorectal cancer survivors: a nationwide database study in Taiwan. *J Tradit Complement Med.* 2016;7:188–94.
49. Pereira RM, Ferreira-Silva GA, Pivatto M, Santos Lde A, Bolzani Vda S, Chagas de Paula DA, et al. Alkaloids derived from flowers of *Senna spectabilis*, (–)-cassine and (–)-spectaline, have antiproliferative activity on HepG2 cells for inducing cell cycle arrest in G1/S transition through ERK inactivation and downregulation of cyclin D1 expression. *Toxicol In Vitro.* 2016;31:86–92.
50. Newman DJ, Cragg GM. Natural products as sources of new drugs from 1981 to 2014. *J Nat Prod.* 2016;79:629–61.
51. Block KI, Gyllenhaal C, Lowe L, Amedei A, Amin ARMR, Amin A, et al. Designing a broad-spectrum integrative approach for cancer prevention and treatment. *Semin Cancer Biol.* 2015;35(Supplement):S276–304.
52. Devasena T, Venugopal MP. Fenugreek seeds modulate 1,2-dimethylhydrazine-induced hepatic oxidative stress during colon carcinogenesis. *Ital J Biochem.* 2007;56:28–34.
53. Li F, Fernandez PP, Rajendran P, Hui KM, Sethi G. Diosgenin: a steroidal saponin, inhibits STAT3 signaling pathway leading to suppression of proliferation and chemosensitization of human hepatocellular carcinoma cells. *Cancer Lett.* 2010;292:197–207.
54. Chandra P, Pandey R, Kumar B, Srivastva M, Pandey P, Sarkar J, et al. Quantification of multianalyte by UPLC–QqQLIT–MS/MS and in-vitro anti-proliferative screening in *Cassia* species. *Ind Crop Prod.* 2015;76:1133–41.
55. Nehdi IA, Sbihi HM, Tan CP, Al-Resayes SI. Seed oil from *Harmal* (*Rhazya stricta* Decne) grown in Riyadh (Saudi Arabia): a potential source of δ -tocopherol. *J Saudi Chem Soc.* 2016;20:107–13.
56. El Gendy MAM, Ali BH, Michail K, Siraki AG, El-Kadi AOS. Induction of quinone oxidoreductase 1 enzyme by *Rhazya stricta* through Nrf2-dependent mechanism. *J Ethnopharmacol.* 2012;144:416–24.
57. Brown NR, Noble ME, Endicott JA, Johnson LN. The structural basis for specificity of substrate and recruitment peptides for cyclin-dependent kinases. *Nat Cell Biol.* 1999;1:438–43.
58. McGowan CH, Russell P. Human Wee1 kinase inhibits cell division by phosphorylating p34cdc2 exclusively on Tyr15. *EMBO J.* 1993;12:75.
59. Wells NJ, Watanabe N, Tokusumi T, Jiang W, Verdecia MA, Hunter T. The C-terminal domain of the Cdc2 inhibitory kinase Myt1 interacts with Cdc2 complexes and is required for inhibition of G (2)/M progression. *J Cell Sci.* 1999;112:3361–71.
60. Enserink JM, Kolodner RD. An overview of Cdk1-controlled targets and processes. *Cell Div.* 2010;5:11–52.
61. Gach K, Grądzka I, Wasyk I, Męczyńska-Wielgosz S, Iwanańko T, Szymański J, et al. Anticancer activity and radiosensitization effect of methyleneisoxazolidin-5-ones in hepatocellular carcinoma HepG2 cells. *Chem Biol Interact.* 2016;248:68–73.
62. Rafehi H, Orłowski C, Georgiadis GT, Ververis K, El-Osta A, Karagiannis TC. Clonogenic assay: adherent cells. *J Vis Exp.* 2011;49:2573.
63. Jiang WG, Sanders AJ, Katoh M, Ungefroren H, Gieseler F, Prince M, et al. Molecular, biological and clinical perspectives. *Semin Cancer Biol.* 2015; 35(Supplement):S244–75.
64. Baeshen NA, Elkady AI, Abuzinadah OA, Mutwakil MH. Potential anticancer activity of the medicinal herb, *Rhazya stricta*, against human breast cancer. *Afr J Biotechnol.* 2012;11:8960–72.
65. Bozzuto G, Ruggieri P, Molinari A. Molecular aspects of tumor cell migration and invasion. *Ann Ist Super Sanita.* 2010;46:66–80.
66. Alur I, Dodurga Y, Secme M, Elmas L, Bagci G, Goksin I, et al. Anti-tumor effects of bemiparin in HepG2 and MIA PaCa-2 cells. *Gene.* 2016;585:241–6.
67. Lavictoire SJ, Parolin DA, Klimowicz AC, Kelly JF, Lorimer IA. Interaction of Hsp90 with the nascent form of the mutant epidermal growth factor receptor EGFRvIII. *J Biol Chem.* 2003;278:5292–9.
68. Hocman G. Chemoprevention of cancer: phenolic antioxidants (BHT, BHA). *Int J BioChemPhys.* 1988;20:639–51.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

