

RESEARCH ARTICLE

# Assessing dietary intakes from household budget surveys: A national analysis in Bangladesh

Dimitra Karageorgou<sup>1</sup>, Fumiaki Imamura<sup>2</sup>, Jianyi Zhang<sup>1</sup>, Peilin Shi<sup>1</sup>, Dariush Mozaffarian<sup>1</sup>, Renata Micha<sup>1\*</sup>

<sup>1</sup> Friedman School of Nutrition Science and Policy, Tufts University, Boston, MA, United States of America, <sup>2</sup> MRC Epidemiology Unit, University of Cambridge, Cambridge, United Kingdom

\* [renata.micha@tufts.edu](mailto:renata.micha@tufts.edu)



## Abstract

### Background

Accurate national information on dietary intakes, including heterogeneity among individuals, is critical to inform health implications and policy priorities. In low- and middle-income countries, household expenditure surveys constitute the major source of food data, but with uncertain validity for individual-level intakes.

### Objective

To investigate how individualized dietary consumption estimated from household survey data compared with individual-level 24-hr dietary recalls (24hR); and to assess potential heterogeneity by method for individualizing household intakes, dietary indicator, and individual characteristics (age, sex, education, religion, household income).

### Methods

We evaluated data from the 2011–2012 Bangladesh Household Integrated Survey (BIHS), which included household-level consumption data (5,503 households) and individual-level dietary data based on 24hR from these households (22,173 participants). Household and 24hR estimates were standardized and harmonized for 33 dietary indicators, including 9 food groups, total energy, 8 macronutrients, and 15 micronutrients. Individual consumption was estimated from household data using two approaches, the Adult Male Equivalent (AME) and per capita (PC) approach. For each dietary indicator, differences in household vs. individual mean estimates were evaluated overall and by strata of individual characteristics, using Spearman's correlations and univariate and multivariate linear regression models.

### Results

Individualized household estimates overestimated individual intakes from 24hR for all dietary factors using either estimation method ( $P < 0.001$  for each), except for starchy

## OPEN ACCESS

**Citation:** Karageorgou D, Imamura F, Zhang J, Shi P, Mozaffarian D, Micha R (2018) Assessing dietary intakes from household budget surveys: A national analysis in Bangladesh. PLoS ONE 13(8): e0202831. <https://doi.org/10.1371/journal.pone.0202831>

**Editor:** Brian Luckett, Tulane University, UNITED STATES

**Received:** January 19, 2018

**Accepted:** August 9, 2018

**Published:** August 27, 2018

**Copyright:** © 2018 Karageorgou et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Data Availability Statement:** All relevant data are within the paper and its Supporting Information files.

**Funding:** This research was supported by the World Bank and Gates Foundation. The funding agencies did not contribute to design or conduct of the study; collection, management, analysis, or interpretation of the data; preparation, review, or approval of the manuscript; or decision to submit the manuscript for publication.

**Competing interests:** Ms Karageorgou and Drs Zhang, Shi, Mozaffarian and Micha report grants from the Gates Foundation during the conduct of the study. Drs. Mozaffarian and Micha report fees from the World Bank during the conduct of the study. Dr. Imamura reports funding from the United Kingdom Medical Research Council Epidemiology Unit Core support (MC\_UU\_12015/5) during the conduct of this study. Dr. Micha is PI of a research grant from Unilever on an investigator-initiated project to assess the effects of omega-6 fatty acid biomarkers on diabetes and heart disease; reports personal fees from Bunge; all outside the submitted work. Dr. Mozaffarian reports personal fees from Boston Heart Diagnostics, Haas Avocado board, Astra Zeneca, GOED, DSM, Life Sciences Research Organization, and UpToDate, outside the submitted work. We affirm that this does not alter our adherence to PLOS ONE policies on sharing data and materials.

vegetables (AME:  $P = 0.15$ ; PC:  $P = 0.85$ ). For foods, overestimation ranged from 4% for seafood to about 240% for fruits, and for nutrients from 11% for carbohydrates and polyunsaturated fats to 55% for vitamin C, with similar overestimation for the AME and the PC method. By strata, overestimation was modestly higher in men vs. women, in children (0–10y) vs. adolescents (11–19y) and adults (20–44y,  $\geq 45$ y), among adults of higher ( $\geq 6$ y) vs. lower ( $< 6$ y) education, in Muslims vs. other religions (Christians, Hindus), and for the lowest vs. all other income groups. This overestimation was notably higher in young children (0–5y) vs. all other age groups and in the lowest vs. all other income groups. Underestimation was rarely observed, for example for milk intake (-56%) in young children (0–5y). The PC approach did not capture heterogeneity in validity of estimation of different dietary factors by age, mainly in children (0–5y, 6–10y). Spearman's correlations between individualized household estimates and 24hR data were higher for the AME (0.30–0.70) than PC (0.20–0.50) approach. Findings were similar with and without multivariate regression, with proportions of variance ( $R^2$ ) in 24hR intakes explained by the AME being generally greater than PC estimates, yet still low to modest.

## Conclusions

In this national survey, established methods for estimating individual level intakes from household surveys produce overestimation of intakes of nearly all dietary indicators, with significant variation depending on the dietary factor and modest variation depending on individual characteristics. These findings suggest a need for new methods to estimate individual-level consumption from household survey estimates.

## Introduction

The major global health and economic impacts of food insecurity and undernutrition have been recognized, now joined by tremendous diet-induced burdens of non-communicable diseases (NCDs) [1–5]. In nearly every region of the world, suboptimal diet is a leading modifiable risk factor for mortality and morbidity, exceeding the burdens attributable to most other global health challenges [6–8]. Even modest dietary changes are associated with improvements in maternal and child undernutrition and micronutrient deficiencies [9–13], as well as meaningful reductions in NCDs [14–17]. Based on the crucial role of nutrition in health, a better understanding of patterns and distributions of dietary habits globally is critical to inform and establish dietary priorities and improvement goals [6, 18].

For most countries around the world [18, 19], particularly low- and middle-income countries (LMIC), limited survey data are available on individual-level dietary intakes. Because of this, methods have been developed to utilize all available individual-level dietary data worldwide, together with estimates of national food-supply availability (food balance sheets) from the UN Food and Agriculture Organization (FAO) [20–22], to estimate individual-level dietary intakes in every country globally [6, 19, 23–27]. Yet, the utility of another major potential source of dietary information, household-level consumption and expenditure surveys (HCES), is not well established. Household surveys have the advantage of being done regularly (typically every 3–5 years in most countries) and including large samples, providing potentially relevant data to augment existing estimates of individual-level dietary intakes. However, such surveys are designed mainly to evaluate financial and living conditions of households, rather than specifically for nutrition or food habits; and also collect data on overall household food

consumption, not individual intakes [28–30]. Other potential limitations of household surveys include short recall reference periods, a limited number of foods assessed, absence of data on within-household distributions of intakes, and insufficient accounting for food waste, for foods acquired for purposes other than household consumption (e.g., for storage, guests, livestock), for foods consumed away from home, and for cooking effects on food weight and nutrient content [30–33]. As such, their potential validity for estimating individual-level dietary intake distributions, as well as potential variability in this validity according to individual characteristics such as age, sex, education, or income, is not established.

Two main methods have been proposed for estimating individualized dietary intakes from household data, including the adult male equivalent (AME) and the per capita (PC) approach [20, 34–42]. The PC approach assumes that each person within the household has equal access to and intake of food, while the AME assumes that each person's intake is proportional to their age and sex-specific caloric requirements. However, the validity of such approaches in predicting individual dietary intakes, including for total caloric/energy intake, major foods, macro- and micronutrients and by various population subgroups, remains uncertain. Few prior studies have used and tested the PC [39] or AME approach [34, 35, 40, 41]. These have focused on total energy and a limited number of macro- or micro-nutrients (e.g., protein, fat, fiber, iron) [34, 40, 41] or specific foods associated with malnutrition [35, 39]; have included only specific population groups, such as women of reproductive age and young children (up to 5y) [34, 35] rather than the general population [39–41]; and have used heterogeneous sources of household-level dietary data, including acquisition [35, 39], consumption [34], or computed from individual intakes [40, 41]. Multiple other foods, macronutrients and micronutrients linked to the double burden of malnutrition, as well as potential differences by individual characteristics or method of estimation (PC, AME) have not been evaluated. To address these gaps in knowledge, we compared estimates of individualized dietary intakes from household data to 24-hr recall (24hR) dietary estimates among the same individuals in the nationally representative Bangladesh Integrated Household Survey (BIHS). We evaluated validity overall as well as according to method of estimation (AME, PC), dietary factor, and key individual characteristics.

## Methods

### Dietary survey

We utilized the BIHS 2011–2012 [43], a comprehensive nationally representative survey from a LMIC that includes both household survey and individual-level 24hR dietary estimates from the same individuals. BIHS data are publicly available [43], and household-level dietary data further meet the International Household Survey Network (IHSN) reliability and relevance assessment criteria (**Table A in S1 File**) [30]. BIHS used a two-stage stratified sampling design and covered a total of 6,503 households including 27,285 individuals (47.6% men, 0–120 years, mean age 26.6 (SD: 19.9) years) [43, 44]. Of those, 5,503 households were representative of the rural Bangladesh and 2,040 of southwest Bangladesh as part of the Feed the Future (FTF) global hunger and food security initiative; 1,040 households contributed to the representativeness of both national and FTF zone samples [44]. For the present analysis, and in line with local experts, we used the BIHS national sample, which included 5,503 households and 23,135 individuals. Of these, we excluded 962 individuals with missing 24hR data that were not home at the time of the interview, did not report any consumption of foods or beverages for unknown reasons, or were exclusively breastfed babies. Excluded individuals did not differ in key sociodemographics compared to the overall sample, yet as expected, they were younger ( $17.0 \pm 19.6$  years), since 26% were exclusively breastfed babies. The final analytical sample consisted of 5,503 households and 22,173 individuals (**Table 1**).

**Table 1. Characteristics of the 2011–2012 Bangladesh Integrated Household Survey (BIHS)<sup>1</sup>.**

	BIHS sample for analysis
Representativeness	National of rural Bangladesh
Sample size	5,503 households 22,173 individuals
Sex (n, %)	
Men	10,502 (47.4%)
Women	11,671 (52.6%)
Age (years)	
Range	0–120
Mean (SD)	26.8±19.8
Age groups (n, %)	
0–5	2,807 (12.7%)
6–10	3,078 (13.9%)
11–19	3,885 (17.5%)
20–44	7,728 (34.8%)
≥45	4,675 (21.1%)
Education (n, %) <sup>2</sup>	
<6 years	8,934 (72.0%)
≥6 years	3,469 (28.0%)
Religion (n, %) <sup>3</sup>	
Muslims	19,735 (89%)
Other	2,438 (11%)
Household income (n, %)	
1 <sup>st</sup> quintile	4,441 (20.0%); 71% of low education
2 <sup>nd</sup> quintile	4,450 (20.1%); 79% of low education
3 <sup>rd</sup> quintile	4,429 (20.0%); 78% of low education
4 <sup>th</sup> quintile	4,420 (19.9%); 72% of low education
5 <sup>th</sup> quintile	4,433 (20.0%); 61% of low education
Sampling design	Two-stage stratification; selection of PSUs and selection of households within each PSU (2001 population census–adjusted for 2011 population census)
Seasonality	December 15 –March 15, 2012 (national sample) “Monga” periods are not covered [82] <sup>4</sup>
Training of the researchers	Bangladeshi consulting firm with expertise in complex surveys and data analysis. IFPRI researchers and the consulting firm experts trained experienced enumerators, researchers, and editors to edit the completed questionnaires during the survey.
Food groups included in the food list <sup>5</sup>	Cereal (17), Pulses (9), Edible oil (7), Vegetables (43), Leafy vegetables (31), Meat/ Eggs and milk (16), Fruits (31), Fish (65), Spices (20), Drinks and beverages (5), Other foods (6), Other foods prepared outside home (37)

<sup>1</sup> The reported descriptive statistics are unweighted.

<sup>2</sup> Estimated only for the adult population (≥20 years old).

<sup>3</sup> Religion was reported only for household head. We assumed that other household members were of the same religion. Other religions refer to Christians and Hindus.

<sup>4</sup> “Monga” is most commonly defined as a famine-like state that occurs twice per year; a severe period from mid-September to mid-November, and a less severe from mid-March to mid-May. It affects mainly northwestern Bangladesh.

<sup>5</sup> The household consumption questionnaire included a list of selected food items, organized in food groups. IFPRI, International Food Policy Research Institute; PSU, Primary sampling units

<https://doi.org/10.1371/journal.pone.0202831.t001>

24hR data were collected in-person by trained interviewers using an open-ended recall. The household member responsible for preparing the meals (women 98.8%, 36.2±12.3 years) reported the foods (single-ingredient, mixed dishes) consumed during the previous day in the household from any source, including own cooking, purchased foods, and gifts. Information was collected on the total “as consumed” weight of each food item, the disaggregated ingredients and corresponding raw weights in mixed dishes, and on how much of these food items was consumed by each household member, stored as leftovers, thrown away, or given to guests, others, and animals/livestock.

Household-level consumption was assessed by a 7-day 287-food item questionnaire, including 37 food items for food consumed away from home. Foods consumed and their quantity for the household (raw weight for single-ingredient foods; cooked weight for mixed dishes) were reported by the same household member as for the 24hR. Dietary data collection was performed from December 2011 to March 2012. Details on the BIHS administration and questionnaires can be found elsewhere [43].

### Dietary dataset harmonization

Dietary data were harmonized within and between the 24hR and household datasets. This process involved 7 key steps (**Appendix A in S1 File**): 1. *dataset retrieval*, involving identification and retrieval of relevant dietary and sociodemographic BIHS datasets and variables; 2. *unique food item identification and description*, identifying the unique food items (single-ingredient or disaggregated ingredient) across the diet assessment methods by matching their available food description, further accounting for food consumed away from home; 3. *food matching*, matching food items to available food composition data [45, 46] for nutrient profiling (if nutrient composition was available for the overall recipe/mixed dish then that was preferred) further accounting for alterations in nutrient content during cooking (use of retention factors [45, 47, 48]) [49, 50]; 4. *unit standardization*, accounting for non-edible portions and cooking alterations (use of yield factors [45, 47, 51, 52]), converting and reporting in standardized “as consumed” metrics, i.e., g/day for foods and macronutrients (other than cholesterol), and mg/day or µg/day for micronutrients; 5. *food classification*, classifying unique food items (including disaggregated ingredients) to food groups (e.g., fruits, vegetables) using previously established methods [19, 53, 54]; 6. *individualization of household consumption*, where household food and nutrient consumption was individualized by the AME [20] and PC [55] approach (**Appendix B in S1 File**); and, 7. *final dataset preparation*, merging and creating a complete dataset including individual-level dietary and sociodemographic information. Local experts provided advice on each of those steps, particularly for steps 2, 3, and 5.

The AME method [20] was our primary approach for individualizing household consumption [34, 35, 37]. This method assumes that the intra-household food distribution is proportional to the individual’s share of total household energy requirements, and as such household members do not receive an equal share of the food available for consumption. The energy requirements of household members of different age, sex, and status (pregnant/ lactating women) were expressed in proportion to an adult male’s energy requirements (**Appendix B in S1 File**). In secondary analysis, we used the PC approach [56] to estimate the per capita consumption, assuming that the available food in the household is equally distributed among household members.

### Selection of key dietary targets

We included dietary indicators that captured an individual’s intake from all sources. Among the array of dietary factors that could be assessed, we identified 48 potential dietary indicators

(18 food groups, 11 macronutrients, 18 micronutrients, and total energy) based on evidence for etiological effects on a) major chronic diseases (e.g., type 2 diabetes, stroke, heart disease) and related risk factors (e.g., blood lipids, blood pressure, obesity) [7, 8, 57–62], or b) deficiency-related health conditions and mortality (e.g., anemia, blindness, maternal mortality) [1, 10, 12, 13]. Among these 48 factors, the final selection of dietary indicators was based on observed intake levels (foods) and available food composition data (nutrients) in this survey. For example, if intake for a selected food group was low, it was combined into a broader category (e.g., whole grains and refined grains were combined into total grains due to low whole grain intake), or omitted if very rarely consumed (e.g., sugar-sweetened beverages). Nutrients were not analyzed if food composition data were not available (iodine), missing for >70% of foods (trans fats, omega-6 fats, vitamin B<sub>12</sub>, selenium) or missing for major dietary sources (omega-3 contents for seafood).

In sum, 33 dietary indicators were included in the final analysis (**Table B in S1 File**), including 9 food groups (fruits, non-starchy vegetables, starchy vegetables, legumes, total grains, meat/eggs, seafood, whole-fat milk, fats/oils), total energy, 8 macronutrients (protein, carbohydrates, total fats, saturated fatty acids (SFA), monounsaturated fatty acids (MUFA), polyunsaturated fatty acids (PUFA), cholesterol, fiber), and 15 micronutrients (vitamins A, B<sub>1</sub> (thiamine), B<sub>2</sub> (riboflavin), B<sub>3</sub> (niacin), B<sub>6</sub>, B<sub>9</sub> (folate), C, D, and E, sodium, potassium, magnesium, calcium, iron, zinc).

### Statistical analysis

Average dietary consumption was estimated and compared for the individual 24hR intakes and individualized household estimates overall and by population strata, including by age (<5, 6–10, 11–19, 20–44, and ≥45 years), sex (men, women), education (<6 years of education, ≥6 years of education), religion (Muslims, other religions), and monthly household income (quintiles). To assess how well individualized household estimates ranked participants in comparison to 24hR estimates, Spearman’s correlations were used.

To assess differences in dietary means and also the proportion of variance explained, the relation between 24hR and AME estimates was assessed by using univariate and multivariable random-intercept linear regression analysis which accounted for household clusters [19].

$$24hR\ estimate_{ij} = \beta_0 + \beta \times AME\ estimate_{ij} \quad (univariate\ model)$$

$$24hR\ estimates_{ij} = \beta_0 + \beta \times AME\ estimate_{ij} + \beta' \times covariates_{ij} \quad (multivariable\ model)$$

where 24hR estimate<sub>ij</sub> represents intake estimates for individual *i* and household *j*; β<sub>0</sub> represents the intercept; β (slope) represents the difference in the 24hR mean for a 1-unit difference in the AME mean; AME estimate<sub>ij</sub> represents individualized intake estimates for individual *i* and household *j*; covariates<sub>ij</sub> were covariates specific to individual *i* and household *j*; β' represents a set of regression coefficients for differences in the 24hR mean for a 1-unit difference in covariates; and random effects were modeled for β<sub>0</sub> and fixed effects were modeled for βs. Analyses were repeated for the PC approach. For the multivariable models, we selected covariates which would be available in household surveys, including basic demographics, such as age and sex, in the minimally adjusted model, and additionally education, religion, household income, respondent’s characteristics (age, sex, education), and household characteristics (household size, number of children, and wastage percentage) in the fully adjusted model.

For the aim to assess potential heterogeneity, regression analyses (univariate, multivariate) were performed by sex, age, sex and age, education, religion, and household income. There was no correlation between education and income, as assessed by Cohen’s kappa coefficient

( $\kappa = -0.02$ ), which justified stratified analysis by each variable separately. Stratification by all demographic factors was not performed because of low sample size and unstable estimates in some strata. Missing covariate values for education ( $n = 17$ , 0.0008%) were imputed with a single regression imputation as the missing values were very few, assuming education was missing at random [63], and using age, sex, and household size as predictors; predictors had no missing values.

Analyses were performed using STATA 14 (College Station, TX: StataCorp LP). Results from statistical tests were considered significant with two-sided alpha = 0.05.

## Results

### Findings for overall population

Participants were 47.4% men, less educated (72% with <6 years of education), mainly Muslims (89.0%), and had mean (SD) monthly household income of 6,701 (9,339) BDT (79.4 (110.6) USD/month) (Table 1). Their diet was characterized by staples consisting primarily of total grains (e.g., rice, flour), starchy vegetables (mainly potatoes), non-starchy vegetables, and seafood (mainly fish) (Table 2).

Considering relative rankings (Spearman's correlations) of individuals, correlations between individualized household estimates and 24hR data were generally higher for AME compared with PC estimates (Table 2). For AME estimates, correlations were generally modest (ranging from about 0.30 to 0.70) while for PC estimates correlations were lower (ranging from about 0.20 to 0.50).

For all dietary indicators except starchy vegetables, mean individualized household estimates significantly exceeded individual intakes ( $P < 0.001$  each) (Table 2). The degree of overestimation was generally very similar between the AME and PC approaches (Fig 1). Among different foods, the overestimation was smallest for seafood (AME: 4%, PC: 4%) and total grains (8%, 7%), and greatest for non-starchy vegetables (56%, 54%) and fruits (242%, 239%). Total energy was overestimated by about 12%. Among macro- and micronutrients, the degree of overestimation ranged from about 11–55% and was highest for vitamins A (50%, 49%) and C (55%, 54%) and lowest for carbohydrates (12%, 11%), and protein (13%, 12%). The shape of the dietary factor distribution, including narrowness, was generally similar between the AME and 24hR estimates (Figure A in S1 File). The magnitude of the narrowness, as assessed by the standard deviations, was also quite similar between the two types of estimates (Table 2). The AME estimates were generally characterized by less variable distributions.

In unadjusted linear regression analyses, proportions of the variance ( $R^2$ ) in 24hR intakes explained by individualized household estimates (AME) were generally modest to low (Table 3, Table C in S1 File). For foods, for example, modest values were seen for total grains ( $R^2 = 0.48$ ), milk (0.24) and fats/oils (0.23), and lower values for fruits (0.06), legumes (0.13) and meat/eggs (0.13). For total energy and nutrients, variation explained was highest for niacin (0.50), vitamin B<sub>6</sub> (0.49), energy (0.46) and carbohydrates (0.46), and lowest for sodium (0.08) and vitamin C (0.11). Proportions explained by the PC estimates were generally smaller. Adjustment for sex and age improved the variation explained for all dietary factors, mainly for energy, protein, carbohydrates, fiber, potassium and magnesium. Additional adjustments did not appreciably change observed relations.

### Findings by sex

Overall dietary consumption patterns were broadly similar by sex, although men often had higher mean intakes across foods, especially for milk, meat/eggs, legumes, and total grains, and certain nutrients, mainly B vitamins, cholesterol, fatty acids and calcium (Tables D and E

Table 2. Comparison of individualized household consumption and individual dietary estimates by dietary factor in the 2011–2012 BIHS.

Dietary Factor, unit <sup>1</sup>	Observations (n) <sup>2</sup>	Consumption (mean, SD) <sup>3</sup>			Correlation (rho coefficient) <sup>4</sup>		Difference between 24hR and AME (mean, SD) <sup>5</sup>		Difference between 24hR and PC (mean, SD) <sup>5</sup>	
		AME <sup>3</sup>	PC <sup>3</sup>	24hR <sup>3</sup>	24hR-AME	24hR-PC	Absolute <sup>6</sup>	Percent	Absolute <sup>7</sup>	Percent
<b>Food groups</b>										
Fruits, g/d	22,146	30.8 (53.1)	30.5 (50.5)	9.0 (35.7)	0.29	0.30	21.8 (56.3)	242	21.5 (54.0)	239
Non-starchy vegetables, g/d	22,173	260.3 (166.8)	258.4 (150.6)	167.3 (147.2)	0.46	0.32	92.9 (174.1)	56	91.0 (176.6)	54
Starchy vegetables, g/d	21,919	105.6 (69.4)	104.8 (62.6)	104.7 (97.8)	0.41	0.31	0.9 (95.4)	1	0.1 (99.0)	0
Legumes, g/d	19,885	22.5 (32.3)	22.3 (30.9)	20.1 (50.2)	0.39	0.39	2.4 (48.9)	12	2.2 (49.1)	11
Total grains, g/d	13,310	1473.7 (536.7)	1463.4 (399.9)	1367.2 (563.9)	0.69	0.37	106.5 (430.3)	8	96.1 (561.1)	7
Meat/Eggs, g/d	15,019	17.2 (20.5)	17.1 (19.5)	11.8 (26.3)	0.36	0.37	5.4 (26.9)	46	5.2 (26.7)	44
Seafood, g/d	22,173	25.7 (24.7)	25.4 (23.1)	24.6 (29.9)	0.50	0.44	1.1 (28.6)	4	0.9 (29.2)	4
Milk, whole fat, g/d	21,201	26.3 (54.1)	26.2 (52.0)	17.7 (60.7)	0.51	0.52	8.6 (58.2)	49	8.4 (54.8)	47
Fats/Oils, g/d	22,162	20.2 (13.2)	20.1 (12.0)	17.8 (16.0)	0.56	0.43	2.4 (15.2)	13	2.3 (15.7)	13
<b>Energy &amp; Macronutrients</b>										
Energy, kcal/d	22,173	2322.4 (871.1)	2305.4 (654.5)	2064.6 (818.7)	0.69	0.36	257.8 (679.5)	12	240.7 (847.0)	12
Protein, g/d	22,173	56.5 (23.6)	56.1 (18.9)	50.0 (22.3)	0.67	0.39	6.5 (19.8)	13	6.1 (23.2)	12
Carbohydrates, g/d	22,173	444.1 (164.2)	440.8 (121.9)	397.7 (158.5)	0.68	0.35	46.3 (129.4)	12	43.1 (163.8)	11
Total fat, g/d	22,173	30.0 (18.6)	29.8 (16.7)	25.6 (19.7)	0.61	0.50	4.4 (18.7)	17	4.1 (19.3)	16
SFA, g/d	22,173	6.5 (5.4)	6.5 (5.0)	5.3 (4.8)	0.62	0.52	1.2 (5.5)	23	1.2 (5.4)	23
MUFA, g/d	22,173	9.1 (5.6)	9.0 (5.0)	7.8 (6.0)	0.61	0.49	1.3 (5.5)	17	1.3 (5.7)	17
PUFA, g/d	22,173	13.7 (9.0)	13.6 (8.1)	12.2 (10.3)	0.67	0.55	1.5 (9.3)	12	1.4 (9.7)	11
Cholesterol, mg/d	22,172	43.4 (46.6)	43.0 (44.0)	33.4 (58.8)	0.47	0.46	9.9 (59.7)	30	9.6 (58.9)	29
Fiber, g/d	22,173	31.4 (12.7)	31.1 (9.9)	26.0 (11.4)	0.65	0.34	5.4 (10.6)	21	5.1 (12.5)	20
<b>Vitamins</b>										
Vitamin A, µg RAE/d	22,173	322.8 (356.2)	320.5 (336.9)	214.8 (404.8)	0.41	0.38	107.9 (439.4)	50	105.6 (438.7)	49
Vitamin D, µg/d	22,173	1.3 (1.9)	1.3 (1.8)	1.1 (2.7)	0.47	0.46	0.2 (2.5)	18	0.2 (2.6)	18
Vitamin E, mg/d	22,173	5.4 (3.2)	5.3 (2.8)	4.5 (3.4)	0.67	0.55	0.9 (3.1)	20	0.8 (3.2)	18
Thiamine, mg/d	22,173	0.9 (0.4)	0.9 (0.3)	0.8 (0.4)	0.63	0.36	0.2 (0.3)	25	0.2 (0.4)	25
Riboflavin, mg/d	22,173	0.6 (0.3)	0.6 (0.2)	0.5 (0.2)	0.60	0.41	0.1 (0.3)	20	0.1 (0.3)	20
Niacin, mg/d	22,173	16.0 (6.4)	15.9 (5.0)	14.1 (6.2)	0.73	0.45	1.9 (4.8)	13	1.8 (6.0)	13
Vitamin B6, mg/d	22,173	1.4 (0.7)	1.4 (0.6)	1.2 (0.7)	0.69	0.46	0.2 (0.5)	17	0.2 (0.6)	17
Folate, µg/d	22,173	157.3 (82.1)	156.1 (71.0)	120.5 (77.2)	0.53	0.33	36.8 (84.0)	31	35.6 (87.1)	30
Vitamin C, mg/d	22,173	65.4 (45.3)	65.0 (41.3)	42.2 (40.7)	0.41	0.27	23.3 (49.7)	55	22.8 (49.7)	54
<b>Minerals</b>										
Calcium, mg/d	22,173	343.4 (207.8)	340.8 (185.1)	274.2 (222.7)	0.58	0.43	69.2 (222.3)	25	66.6 (225.4)	24
Iron, mg/d	22,173	12.0 (5.7)	11.9 (4.8)	9.9 (5.3)	0.63	0.38	2.1 (5.1)	21	2.0 (5.6)	20
Sodium, mg/d	22,173	5855.2 (3145.8)	5812.9 (2739.4)	4225.0 (3073.7)	0.39	0.19	1630.2 (3737.0)	39	1587.9 (3790.8)	38
Potassium, mg/d	22,173	1745.6 (784.3)	1732.6 (647.3)	1394.6 (638.9)	0.63	0.36	351.0 (670.1)	25	338.0 (735.4)	24
Magnesium, mg/d	22,173	376.9 (146.7)	374.1 (112.8)	320.6 (133.4)	0.67	0.34	56.3 (119.0)	18	53.5 (143.3)	17

(Continued)



Table 2. (Continued)

Dietary Factor, unit <sup>1</sup>	Observations (n) <sup>2</sup>	Consumption (mean, SD) <sup>3</sup>			Correlation (rho coefficient) <sup>4</sup>		Difference between 24hR and AME (mean, SD) <sup>5</sup>		Difference between 24hR and PC (mean, SD) <sup>5</sup>	
		AME <sup>3</sup>	PC <sup>3</sup>	24hR <sup>3</sup>	24hR-AME	24hR-PC	Absolute <sup>6</sup>	Percent	Absolute <sup>7</sup>	Percent
Zinc, mg/d	22,173	9.8 (3.8)	9.7 (2.9)	8.6 (3.5)	0.68	0.36	1.2 (3.0)	14	1.1 (3.7)	13

<sup>1</sup> Dietary factors presented had adequate data/information for the present analysis (see Selection of dietary targets).

<sup>2</sup> Sample sizes differ because we performed paired analysis for each dietary factor, i.e., for each analysis we used only the individuals with available intake data for both diet assessments.

<sup>3</sup> Bangladesh Integrated Household Survey (BIHS) 2011–2012 provided household-level dietary data from a 7-day household consumption questionnaire and individual-level data from 24-hour recalls (24hR). Household consumption was individualized by applying a) the Adult Male Equivalent (AME) method [20] as proposed by FAO [56], assuming moderate physical activity, and b) the per capita (PC) approach assuming equal distribution among household members (Appendix B in S1 File). Individual intake was estimated from 24hR.

<sup>4</sup> Spearman correlation coefficients (rho). All correlations were significant (P<0.001).

<sup>5</sup> Absolute differences correspond to AME-24hR and PC-24hR respectively, and percentage differences correspond to absolute difference/24hR\*100.

<sup>6</sup> Differences between means were significant for all dietary factors (paired t-test, P<0.001), with the exception of starchy vegetables (p = 0.15).

<sup>7</sup> Differences between means were significant for all dietary factors (paired t-test, P<0.001), with the exception of starchy vegetables (p = 0.85).

MUFA, Monounsaturated fats; PUFA, Polyunsaturated fats; SFA, Saturated fat

<https://doi.org/10.1371/journal.pone.0202831.t002>


in S1 File). The overestimation of AME household estimates was modestly higher in men than women for most factors (Fig 1) except for meats/eggs, milk, and dietary cholesterol. In contrast, the PC estimates often produced higher overestimation in women than men, highest for milk (men: 32%, women: 66%), fruits (226%, 255%), meats/eggs (31%, 58%), and cholesterol (19%, 39%). In men vs. women, the variance explained for each dietary factor by individualized household estimates was similar to that seen for the overall population (Table T in S1 File).

### Findings by age

Overall dietary consumption patterns were similar by age groups (Tables F–J in S1 File), except that younger adults (20–44 years) generally had higher consumption levels compared to other ages; legume, vitamin A, and vitamin D consumption was higher at older ages, and milk and fruit consumption were lower at older ages. Variability was evident in the relationship between the household estimates and individual intakes by age, especially among children (≤10 years) (Fig 1, Tables F–J in S1 File). For 0–5 year-olds, underestimation was seen for milk (-56%) with AME estimates, while milk was overestimated in all other ages (e.g., 114% in 20–44 year-olds); intakes of most other foods were overestimated to a greater extent in 0–5 year-olds compared with older children and adults. Notably, overestimation in children was substantially higher with the PC method that did not capture heterogeneity in validity of estimation of dietary factors by age (Fig 1). Variance explained by household estimates for each dietary factor by age was generally lower than that observed for the overall population; across age groups it was higher among younger children (0–5y) vs. all other ages (Table U in S1 File).

### Findings by education, religion and household income

Stratification by education revealed similar overall dietary consumption patterns, although adults with higher education (≥6 years) had generally higher intakes, especially for milk, cholesterol, meat/eggs, and fruits (Tables K and L in S1 File). The overestimation of AME estimates was modestly higher in individuals of higher vs. lower education (<6 years). PC



**Fig 1. National mean individualized household estimates compared with 24-hour recall intakes as the reference measure of individual-level consumption, overall and by sex and age for selected dietary factors in the 2011–2012 BIHS.** Mean individualized dietary consumption estimated from household survey data by the Adult Male Equivalent (AME) and per capita (PC) approach (Appendix B in [S1 File](#)) and individual-level 24-hr dietary recall (24hr) intakes are presented for the overall population (all), by sex (men, women), and by age (0-5y, 6-10y, 11-19y, 20-44y, 45+y). Intakes are presented in g/d for foods, and in mg/d (except for vitamin A,  $\mu$ g/d) for nutrients.

<https://doi.org/10.1371/journal.pone.0202831.g001>

estimates were quite similar by education, but diverged considerably from those of the overall population; notably for starchy vegetables (individuals with higher vs. lower education: -12%, -12%), total grains (-8%, -6%), and seafood (-7%, -9%) that were underestimated. Variance explained with the AME method tended to be higher among adults of lower vs. higher education ([Table V in S1 File](#)). The opposite was observed with the PC approach, but this was reversed with sex and age adjustment.

Table 3. Relation between individualized household intake estimates as predictors of individual dietary intakes in the 2011–2012 BIHS<sup>1</sup>.

Dietary Factor, unit	AME			PC		
	Intercept (SE)	$\beta$ (SE)	R <sup>2</sup>	Intercept (SE)	$\beta$ (SE)	R <sup>2</sup>
<b>Food groups</b>						
Fruits, g/d						
Unadjusted	5.4 (0.5)	0.1 (0.02)	0.06	3.7 (0.5)	0.2 (0.02)	0.06
Sex and age	7.0 (0.7)	0.1 (0.02)	0.06	3.9 (0.7)	0.2 (0.02)	0.06
Multivariate	10.7 (7.0)	0.1 (0.02)	0.07	7.2 (7.0)	0.2 (0.02)	0.06
Non-starchy vegetables, g/d						
Unadjusted	29.4 (2.5)	0.5 (0.01)	0.15	96.6 (3.7)	0.3 (0.01)	0.09
Sex and age	23.8 (2.8)	0.3 (0.01)	0.19	8.0 (3.8)	0.2 (0.01)	0.18
Multivariate	14.4 (16.6)	0.3 (0.01)	0.20	-9.8 (17.0)	0.3 (0.01)	0.19
Legumes, g/d						
Unadjusted	6.1 (0.7)	0.6 (0.04)	0.13	7.8 (0.8)	0.5 (0.04)	0.12
Sex and age	2.4 (0.8)	0.5 (0.04)	0.13	-2.3 (0.9)	0.5 (0.04)	0.13
Multivariate	9.7 (11.9)	0.5 (0.04)	0.13	3.9 (11.8)	0.5 (0.04)	0.13
Total grains, g/d						
Unadjusted	86.9 (13.1)	0.9 (0.01)	0.48	626.3 (17.2)	0.5 (0.01)	0.12
Sex and age	137.1 (14.1)	0.5 (0.01)	0.57	77.2 (16.8)	0.4 (0.01)	0.53
Multivariate	207.4 (51.0)	0.6 (0.01)	0.60	42.2 (57.8)	0.4 (0.01)	0.55
Seafood, g/d						
Unadjusted	6.8 (0.5)	0.7 (0.02)	0.22	11.0 (0.7)	0.5 (0.03)	0.17
Sex and age	3.8 (0.5)	0.5 (0.02)	0.23	-1.8 (0.7)	0.5 (0.03)	0.22
Multivariate	3.2 (4.0)	0.5 (0.02)	0.23	-3.9 (4.0)	0.5 (0.03)	0.22
<b>Energy &amp; Macronutrients</b>						
Energy, kcal/d						
Unadjusted	269.5 (15.8)	0.8 (0.01)	0.46	1052.8 (26.7)	0.4 (0.01)	0.13
Sex and age	413.2 (16.5)	0.4 (0.01)	0.55	308.4 (25.0)	0.3 (0.01)	0.51
Multivariate	498.6 (70.5)	0.5 (0.01)	0.57	237.0 (75.7)	0.4 (0.01)	0.53
Protein, g/d						
Unadjusted	7.0 (0.4)	0.8 (0.01)	0.39	25.8 (0.7)	0.4 (0.01)	0.14
Sex and age	10.0 (0.4)	0.4 (0.01)	0.47	5.9 (0.7)	0.3 (0.01)	0.43
Multivariate	11.8 (2.4)	0.5 (0.01)	0.49	4.0 (2.5)	0.4 (0.01)	0.45
Carbohydrates, g/d						
Unadjusted	44.8 (3.0)	0.8 (0.01)	0.46	204.1 (4.9)	0.4 (0.01)	0.12
Sex and age	71.4 (3.2)	0.4 (0.01)	0.55	56.9 (4.7)	0.3 (0.01)	0.51
Multivariate	80.3 (12.7)	0.5 (0.01)	0.58	35.3 (13.8)	0.4 (0.01)	0.54
MUFA, g/d						
Unadjusted	2.0 (0.1)	0.6 (0.01)	0.30	2.9 (0.2)	0.5 (0.03)	0.23
Sex and age	2.1 (0.1)	0.5 (0.02)	0.30	0.2 (0.2)	0.5 (0.03)	0.29
Multivariate	3.6 (0.7)	0.5 (0.02)	0.31	1.2 (0.7)	0.5 (0.03)	0.30
PUFA, g/d						
Unadjusted	2.3 (0.2)	0.7 (0.01)	0.29	4.7 (0.3)	0.6 (0.02)	0.21
Sex and age	2.1 (0.2)	0.6 (0.02)	0.30	-0.6 (0.3)	0.5 (0.02)	0.28
Multivariate	5.5 (1.4)	0.6 (0.02)	0.31	2.3 (1.4)	0.5 (0.02)	0.30
Fiber, g/d						
Unadjusted	2.9 (0.2)	0.7 (0.01)	0.38	14.7 (0.4)	0.4 (0.01)	0.11
Sex and age	4.6 (0.2)	0.4 (0.01)	0.48	3.9 (0.4)	0.3 (0.01)	0.45
Multivariate	5.0 (1.2)	0.4 (0.01)	0.50	2.5 (1.3)	0.3 (0.01)	0.47

(Continued)

Table 3. (Continued)

Dietary Factor, unit	AME			PC		
	Intercept (SE)	$\beta$ (SE)	R <sup>2</sup>	Intercept (SE)	$\beta$ (SE)	R <sup>2</sup>
<b>Vitamins</b>						
Vitamin A, $\mu\text{g}$ RAE/d						
Unadjusted	64.3 (7.1)	0.5 (0.03)	0.11	98.4 (7.9)	0.4 (0.03)	0.10
Sex and age	37.0 (7.2)	0.4 (0.03)	0.12	-5.3 (8.6)	0.4 (0.03)	0.11
Multivariate	17.1 (61.0)	0.4 (0.03)	0.12	-36.3 (60.5)	0.4 (0.03)	0.11
Thiamine, mg/d						
Unadjusted	0.1 (0.01)	0.7 (0.01)	0.34	0.4 (0.01)	0.4 (0.01)	0.12
Sex and age	0.2 (0.01)	0.4 (0.01)	0.42	0.1 (0.01)	0.3 (0.01)	0.39
Multivariate	0.2 (0.04)	0.4 (0.01)	0.43	0.1 (0.04)	0.3 (0.01)	0.40
Riboflavin, mg/d						
Unadjusted	0.2 (0.01)	0.5 (0.01)	0.29	0.3 (0.01)	0.4 (0.01)	0.16
Sex and age	0.2 (0.01)	0.3 (0.01)	0.33	0.1 (0.01)	0.3 (0.01)	0.31
Multivariate	0.2 (0.02)	0.3 (0.01)	0.34	0.1 (0.02)	0.4 (0.02)	0.33
Niacin, mg/d						
Unadjusted	1.1 (0.1)	0.8 (0.01)	0.50	5.8 (0.2)	0.5 (0.01)	0.19
Sex and age	1.5 (0.1)	0.5 (0.01)	0.56	0.1 (0.2)	0.4 (0.01)	0.51
Multivariate	1.8 (0.5)	0.6 (0.01)	0.58	-0.8 (0.5)	0.5 (0.01)	0.53
<b>Minerals</b>						
Calcium, mg/d						
Unadjusted	82.6 (4.7)	0.5 (0.02)	0.22	118.4 (7.6)	0.5 (0.02)	0.16
Sex and age	88.9 (5.3)	0.4 (0.02)	0.23	23.3 (7.9)	0.4 (0.02)	0.22
Multivariate	84.4 (30.8)	0.4 (0.02)	0.23	-0.5 (31.0)	0.4 (0.03)	0.22
Iron, mg/d						
Unadjusted	1.5 (0.1)	0.7 (0.01)	0.32	5.1 (0.2)	0.4 (0.02)	0.15
Sex and age	1.9 (0.1)	0.4 (0.01)	0.38	0.9 (0.2)	0.3 (0.01)	0.35
Multivariate	1.8 (0.6)	0.4 (0.01)	0.39	0.1 (0.6)	0.4 (0.02)	0.37
Sodium, mg/d						
Unadjusted	1472.9 (53.7)	0.5 (0.01)	0.08	3284.4 (90.3)	0.2 (0.02)	0.02
Sex and age	1530.3 (55.6)	0.2 (0.01)	0.14	1361.0 (91.0)	0.1 (0.02)	0.13
Multivariate	2794.9 (570.6)	0.2 (0.01)	0.14	2574.0 (579.3)	0.1 (0.02)	0.14
Zinc, mg/d						
Unadjusted	1.1 (0.1)	0.8 (0.01)	0.43	4.6 (0.1)	0.4 (0.01)	0.12
Sex and age	1.7 (0.1)	0.4 (0.01)	0.52	1.3 (0.1)	0.3 (0.01)	0.49
Multivariate	1.9 (0.3)	0.5 (0.01)	0.54	0.9 (0.4)	0.3 (0.01)	0.51

All p-values are <0.001.

<sup>1</sup> Results are presented for selected dietary factors and for overall population in the Bangladesh Integrated Household Survey (BIHS) 2011–2012. Results for all 33 dietary factors are presented in the Supplement (Table C in S1 File).

On the basis of linear regression models with 24-hour diet recall (24hR) intakes as the dependent variable and individualized Adult Male Equivalent (AME) or per capita (PC) consumption estimates from household questionnaire as the independent variable. The sex and age model was categorized as follows: age,  $\leq 5$ , 6–10, 11–19, 20–44, and  $\geq 45$  years; sex, men and women. The multivariate model was adjusted for the following covariates: age ( $\leq 5$ , 6–10, 11–19, 20–44, and  $\geq 45$  years), sex (men, women), education ( $< 6$  years of education,  $\geq 6$  years of education), religion (Muslims, other), household income (quintiles), respondent's age (continuous), sex (men, women) and education ( $< 6$  years of education,  $\geq 6$  years of education), household size, number of children within household, and food wastage percentage (using 24hR data, we calculated for each household, the percent of food wastage -sum of food waste, and food given to guests, others and animals- to total consumed food (mean: 11.6%, SD: 13.6)). *B*s represent the change in the individual intake (24hR) for every unit increase in the respective mean of household estimates. SEs for the intercept and  $\beta$ s are presented. R<sup>2</sup> represents the coefficient of determination for the overall model.

MUFA, Monounsaturated fats; PUFA, Polyunsaturated fats; SFA, Saturated fat

<https://doi.org/10.1371/journal.pone.0202831.t003>

Intakes were only modestly higher among other religions (Christians, Hindus) vs. Muslims (Tables M and N in S1 File). The overestimation by both individualized household estimates was similar and generally higher in Muslims vs. other religions, except for seafood intakes (6%, 53%, respectively) (Table W in S1 File). Higher variance was explained in other religions vs. Muslims.

Consumption patterns were similar by household income, with higher intakes generally seen in the highest income group (Tables O-S in S1 File). Variability was unremarkable between household estimates and overestimation was higher for the lowest vs. all other income groups. Proportions of variance explained in 24hR intakes were generally modestly higher in the middle (2<sup>nd</sup>, 3<sup>rd</sup> quintile) income groups (Table X in S1 File).

## Discussion

This investigation of household-level data and 24-hour dietary recall information from the same households in rural Bangladesh showed that individualized household estimates significantly exceeded individual intakes for nearly all dietary factors assessed, including by 12% for total energy, 0–242% for major food groups, 11%–30% for macronutrients and 13%–55% for micronutrients. The degree of overestimation varied by both sex and estimation method, with larger overestimation by AME in men and by PC in women; and also for young children (0–5 years), where milk intake was underestimated and intakes of other dietary factors were greatly overestimated. For all dietary factors, low to modest variation in intakes was explained by individualized household estimates, higher for the AME than the PC approach. These novel findings suggest that current methods to utilize household-level survey data are problematic for estimating individual dietary intakes.

Smallest to modest overestimation (<10–15%) was observed for key staple foods, including starchy vegetables, seafood, total grains, legumes and fats/oils; total energy; macronutrients; and specific vitamins and minerals, including niacin and zinc. This small to modest overestimation was similar between the AME and PC approach, and across all strata, except for 0–5 year-old children and individuals of low income in whom all dietary indicators were greatly overestimated. Interestingly, overestimation was higher among adults of higher vs. lower education, but given this population is mostly less educated, these results should be interpreted within that context. These findings suggest that established methods for estimating individual intakes from household surveys could be used to approximate specific dietary indicators, such as most frequently consumed foods, energy and macronutrients. Yet, intakes were still overestimated with further differences noted by individual characteristics and estimation method, and thus future applications of these methods should acknowledge and potentially try to account for this likely limitation.

In contrast, largest overestimation ( $\geq 20\%$ ) was seen for foods with high seasonal variation and increased variability between individuals, such as fruits and non-starchy vegetables; and less commonly consumed foods with fewer questions assessing their consumption in the household questionnaire, such as milk and meats/eggs. Almost all vitamins and minerals assessed were substantially overestimated, particularly those found in the above food sources, such as vitamin A, vitamin C, folate, and calcium. Overestimation was also larger for sodium consistent with increased variability in salt use -irrespectively of total energy intake levels-between individuals [64]. Similar findings were generally observed by individual characteristics (other than younger children) and estimation method. Our findings suggest that household estimates may not reasonably estimate dietary intakes of foods that may be under-represented in the household questionnaire, as well as foods and most vitamins and minerals that are generally characterized by high individual variability. These findings are consistent

with methods used to generate such data, which were developed to capture household consumption rather than actual individual intake, and do not account for food wastage, food preparation alterations in weight and nutrient content, or food eaten away from home [65].

Notably, in really young (0–5 years) children all dietary factors were substantially overestimated, except for milk that was greatly underestimated. Household-level data are challenging for accurately estimating dietary intakes in infants and toddlers [34, 35, 40]. It is quite usual for children to leave a substantial proportion of leftover food [66], and this potential misconception between what is offered vs. what is consumed could contribute to an overall overestimation of household consumption. Furthermore, food consumption by different household members is not necessarily proportional to their energy requirements, particularly for young children and/or for specific foods. Our results showed that 24hR milk intake was highest in very young children compared to all other ages. Yet with the AME method, 0–5 year-olds were assigned the smallest proportion of the household milk consumption (relative to their energy requirements), while with the PC method they were treated as any other individual (assuming equal consumption). These findings confirm that household estimates are not appropriate for estimating dietary intakes of young children, whose dietary patterns vary from the rest of the population.

Both AME and PC approaches appear to be quite problematic for estimating individual intakes from household-level data, with key differences noted by individual characteristics. The AME improved estimation in women, children (0–5 and 6–10 years) and the PC in men, adolescents and adults, and individuals of low or high education, with no substantial differences in the overall population estimates or by religion or income. Women and children ( $\leq 10$  years) in particular are two of the top interest population groups for several international organizations and priority guidelines [67–70]. These vulnerable populations are more likely to develop nutrient deficiencies, especially in LMICs [71], while there is increasing evidence [72–75] of obesity and NCD originating in early development stages and as a potential consequence of maternal and childhood malnutrition. Our results recognize the usefulness of the AME over the PC method for these populations, a finding consistent with the method used to derive PC estimates. The PC method is by design crude and assumes equal consumption within household members, thus limiting its ability to capture variations in individual intake. For all dietary factors variation explained by the AME estimates was consistently higher than the PC estimates, though it was still low to moderate. Potential improvements in the AME estimations -further dependent on data availability- could include the use of caloric equations that account for each individual's anthropometrics and actual physical activity levels, to enable more accurate redistribution of household consumption.

Household surveys suggest several appropriate uses for its household consumption estimates, including constructing food balance sheets, providing food security indicators and poverty measurements, and informing food nutrition interventions [30]; yet, their use for assessing dietary quality or examining diet-disease burden relations [42, 76–78] can be problematic, further supported by the present findings. Given that several LMICs rely solely on household surveys for their food consumption estimates [38], these results highlight the need to further adapt existing individualization methods or develop new ones for better approximating individual-level intakes from household-level data. They also highlight the need to investigate the reasons behind observed overestimations, particularly for dietary factors and population groups with largest discrepancies.

The present findings support and greatly expand on prior reports which compared individualized household estimates to individual dietary data for energy and selected macro- and micronutrients such as protein, vitamin C, and iron, but not foods, or did not evaluate differences by individualization approach, and key population subgroups, such as sex, age,

education, and income [34–36, 39–41, 79]. In Uganda, AME underestimated the energy adjusted intakes of key nutrients (e.g., vitamin C, folate, calcium) related to deficiencies in women (15–49y) and young children (2–5y) compared to 24hR estimates [34]. In Cameroon, AME estimates overestimated 24hR intakes for key foods assessed (vegetable oil, sugar, bouillon cube) in women (15–49y), and either over- (vegetable oil, bouillon cube) or under- (wheat flour, sugar) estimated intakes in children (1–5y) [35]. These studies used different surveys as sources for individual- and household-level data, not always comparable (i.e., both nationally representative); the sample was limited to households with only women and/or children of certain age; AME nutrient estimates were compared with energy-adjusted individual intakes; or only a few nutrients or single food items (for purposes of fortification) were assessed.

In prior analysis using the same BIHS survey, individual intakes from the 24hR were summed back to the household level rather than using actual household-level data [40]; subsequent application of AME approach efficiently redistributed energy, iron, zinc, vitamin A, and calcium among household members aged 4 years and above. In similar analyses -using computed household dietary data- in Ethiopia and Bangladesh, AME estimates compared well with individual intakes of energy, iron, and protein in adults and children, but not in women of reproductive age and infants (<2y) where substantial overestimation was seen [41]. These analyses do not test the validity of the AME approach in individualizing actual household data, which is particularly important when household questionnaires are the only source of dietary data.

Our investigation has several strengths. We systematically quantified differences between estimated individual dietary intake from household-level data and individual 24-hr recalls for multiple dietary indicators and different estimation methods, evaluating both rankings within the population, differences in means, and variation explained. We further assessed heterogeneity in this validity according to several key individual characteristics, and for a range of population subgroups, including children, women, and men. We included a wide range of nutrients related to both chronic diseases, and deficiencies, undernourishment, and child-maternal outcomes [10, 12, 13]. We adjusted our analyses for food wastage, as reported in the 24hR, and accounted for food consumed away in all estimates. To maximize comparability between diet assessment methods, we followed a series of standardized steps to harmonize description, classification, and quantification of food and nutrient intakes, including application of nutrient retention factors and yield factors, highlighting key crucial preparatory methodological steps in individualizing household consumption.

Despite comprehensive approaches to harmonize dietary data, there are inherent limitations in the dietary collection methods. We could not assess certain foods or nutrients, such as sugar-sweetened beverages, iodine, omega-3 and omega-6 fats. In this survey, as in other LMIC settings [38, 40, 41, 80], the main person responsible for cooking reported 24hR intakes for all other household members; though this is standard for children, it is generally not recommended for other adults, as it could lead to systematic reporting biases [81]. Only one 24hR was administered, which may have affected the accuracy of the estimated individual intakes. Yet, single 24hR can provide valid estimates of the absolute mean “usual” intake of a population subgroup, as assessed in the present analysis. Energy requirements for the AME approach were based on standard FAO equations that may be less sensitive in capturing individual variation. Seasonality (monga period in rural Bangladesh) was not covered [82], and our findings may differ for certain foods with substantial seasonal variation. Our results are based on a rural low-income population and their generalizability to urban or middle-income populations may be limited. Conversely, rural low-income populations globally are more likely to be lacking individual-level surveys and, thus, most relevant to assess in the present analysis.

Future studies are needed to replicate the validity of and extend our approaches in different settings and populations over time.

In conclusion, household estimates substantially overestimated individual intakes in a national survey in rural Bangladesh with significant heterogeneity according to sex, age, education, and income. Methodology constructed in the present analysis showed that current methods for estimating individual intakes from household-level data are problematic, yet it confirmed usefulness of the AME vs. the PC approach in better approximating dietary intakes for key populations, mainly children and women. These findings will facilitate future use of household consumption estimates by scientists and policy makers to more accurately estimate dietary intakes, when household questionnaires are the only source of dietary data. Leveraging national household surveys already in place to routinely collect individual-level dietary data, even in a reasonably powered subset of the population, would be of great value to LMIC settings [83]. Relative to its importance as a global risk factor for health, disparities, and sustainability, national investment in the routine collection of individual-level dietary data should be prioritized for accurate diet assessment, burden analyses and policy implementation.

## Supporting information

**S1 File. Supplementary material. Appendix A.** Dietary dataset preparation.

**Appendix B.** Methods for individualizing consumption data from household surveys.

**Table A.** Reliability and relevance assessment of the 2011–2012 BIHS by the International Household Survey Network criteria.

**Table B.** Definitions and units of dietary factors used in the 2011–2012 BIHS.

**Table C.** Relation between individualized household intake estimates as predictors of individual dietary intakes in the 2011–2012 BIHS.

**Table D.** Comparison of individualized household consumption and individual dietary intake estimates by dietary factor in men in the 2011–2012 BIHS.

**Table E.** Comparison of individualized household consumption and individual dietary intake estimates by dietary factor in women in the 2011–2012 BIHS.

**Table F.** Comparison of individualized household consumption and individual dietary intake estimates by dietary factor in children 5 years and under in the 2011–2012 BIHS.

**Table G.** Comparison of individualized household consumption and individual dietary intake estimates by dietary factor in children 6–10 years old in the 2011–2012 BIHS.

**Table H.** Comparison of individualized household consumption and individual dietary intake estimates by dietary factor in adolescents 11–19 years old in the 2011–2012 BIHS.

**Table I.** Comparison of individualized household consumption and individual dietary intake estimates by dietary factor in adults 20–44 years old in the 2011–2012 BIHS.

**Table J.** Comparison of individualized household consumption and individual dietary intake estimates by dietary factor in adults over 45 years of age in the 2011–2012 BIHS.

**Table K.** Comparison of individualized household consumption and individual dietary intake estimates by dietary factor in adults ( $\geq 20$  years old) of low educational level ( $< 6$  years) in the 2011–2012 BIHS.

**Table L.** Comparison of individualized household consumption and individual dietary intake estimates by dietary factor in adults ( $\geq 20$  years old) of medium and high educational level ( $\geq 6$  years) in the 2011–2012 BIHS.

**Table M.** Comparison of individualized household consumption and individual dietary intake estimates by dietary factor in Muslims<sup>1</sup> in the 2011–2012 BIHS.

**Table N.** Comparison of individualized household consumption and individual dietary intake



estimates by dietary factor among other religions<sup>1</sup> in the 2011–2012 BIHS.

**Table O.** Comparison of individualized household consumption and individual dietary intake estimates by dietary factor among individuals in the first quintile of household income in the 2011–2012 BIHS.

**Table P.** Comparison of individualized household consumption and individual dietary intake estimates by dietary factor among individuals in the second quintile of household income in the 2011–2012 BIHS.

**Table Q.** Comparison of individualized household consumption and individual dietary intake estimates by dietary factor among individuals in the third quintile of household income in the 2011–2012 BIHS.

**Table R.** Comparison of individualized household consumption and individual dietary intake estimates by dietary factor among individuals in the fourth quintile of household income in the 2011–2012 BIHS.

**Table S.** Comparison of individualized household consumption and individual dietary intake estimates by dietary factor among individuals in the fifth quintile of household income in the 2011–2012 BIHS.

**Table T.** Relation between individualized household intake estimates as predictors of individual dietary by sex in the 2011–2012 BIHS.

**Table U.** Relation between individualized household intake estimates as predictors of individual dietary intakes by age in the 2011–2012 BIHS.

**Table V.** Relation between individualized household intake estimates as predictors of individual dietary intakes by education in the 2011–2012 BIHS.

**Table W.** Relation between individualized household intake estimates as predictors of individual dietary intakes by religion in the 2011–2012 BIHS.

**Table X.** Relation between individualized household intake estimates as predictors of individual dietary intakes by household income in the 2011–2012 BIHS.

**Figure A.** Distribution of individualized household estimates and 24-hour recall intakes for selected dietary factors in the overall population in the 2011–2012 BIHS.  
(DOCX)

## Acknowledgments

**Funding:** This research was supported by the World Bank and Gates Foundation. The funding agencies did not contribute to design or conduct of the study; collection, management, analysis, or interpretation of the data; preparation, review, or approval of the manuscript; or decision to submit the manuscript for publication.

### Author conflict of interest disclosures

Ms Karageorgou and Drs Zhang, Shi, Mozaffarian and Micha report grants from the Gates Foundation during the conduct of the study. Drs. Mozaffarian and Micha report fees from the World Bank during the conduct of the study. Dr. Imamura reports funding from the United Kingdom Medical Research Council Epidemiology Unit Core support (MC\_UU\_12015/5) during the conduct of this study. Dr. Micha is PI of a research grant from Unilever on an investigator-initiated project to assess the effects of omega-6 fatty acid biomarkers on diabetes and heart disease; reports personal fees from Bunge; all outside the submitted work. Dr. Mozaffarian reports personal fees from Boston Heart Diagnostics, Haas Avocado board, Astra Zeneca, GOED, DSM, Life Sciences Research Organization, and UpToDate, outside the submitted work. These do not alter our adherence to PLOS ONE policies on sharing data and materials.

**Non author contributions:** We thank Wahid Quabili, research analyst in the International Food Policy Research Institute, and Atonu Rabbani, Associate Professor at Department of

Economics, University of Dhaka, for assistance with the BIHS dataset and dietary dataset preparation.

## Author Contributions

**Conceptualization:** Renata Micha.

**Formal analysis:** Dimitra Karageorgou, Fumiaki Imamura, Jianyi Zhang, Peilin Shi.

**Funding acquisition:** Dariush Mozaffarian, Renata Micha.

**Supervision:** Renata Micha.

**Writing – original draft:** Dimitra Karageorgou, Renata Micha.

**Writing – review & editing:** Dimitra Karageorgou, Dariush Mozaffarian, Renata Micha.

## References

1. UNICEF, WHO, The World Bank Group: Joint child malnutrition estimates. Levels and trends in child malnutrition: Key findings of the 2017 edition 2017.
2. Shamah-Levy T, Mundo-Rosas V, Morales-Ruan C, Cuevas-Nasu L, Mendez-Gomez-Humaran I, Perez-Escamilla R. Food insecurity and maternal-child nutritional status in Mexico: cross-sectional analysis of the National Health and Nutrition Survey 2012. *BMJ open*. 2017; 7(7):e014371. <https://doi.org/10.1136/bmjopen-2016-014371> PMID: 28760785.
3. World Food Programme. Food insecurity and undernutrition in the urban slums of Bangladesh. 2015.
4. Kaur J, Lamb MM, Ogden CL. The Association between Food Insecurity and Obesity in Children-The National Health and Nutrition Examination Survey. *Journal of the Academy of Nutrition and Dietetics*. 2015; 115(5):751–8. <https://doi.org/10.1016/j.jand.2015.01.003> PMID: 25737437.
5. Popkin BM, Adair LS, Ng SW. Global nutrition transition and the pandemic of obesity in developing countries. *Nutrition reviews*. 2012; 70(1):3–21. <https://doi.org/10.1111/j.1753-4887.2011.00456.x> PMID: 22221213; PubMed Central PMCID: PMC3257829.
6. Micha R, Khatibzadeh S, Shi P, Andrews KG, Engell RE, Mozaffarian D, et al. Global, regional and national consumption of major food groups in 1990 and 2010: a systematic analysis including 266 country-specific nutrition surveys worldwide. *BMJ open*. 2015; 5(9):e008705. <https://doi.org/10.1136/bmjopen-2015-008705> PMID: 26408285; PubMed Central PMCID: PMC4593162.
7. Micha R, Penalvo JL, Cudhea F, Imamura F, Rehm CD, Mozaffarian D. Association Between Dietary Factors and Mortality From Heart Disease, Stroke, and Type 2 Diabetes in the United States. *JAMA*. 2017; 317(9):912–24. Epub 2017/03/08. <https://doi.org/10.1001/jama.2017.0947> PMID: 28267855; PubMed Central PMCID: PMC5852674.
8. Micha R, Shulkin ML, Penalvo JL, Khatibzadeh S, Singh GM, Rao M, et al. Etiologic effects and optimal intakes of foods and nutrients for risk of cardiovascular diseases and diabetes: Systematic reviews and meta-analyses from the Nutrition and Chronic Diseases Expert Group (NutriCoDE). *PloS one*. 2017; 12(4):e0175149. <https://doi.org/10.1371/journal.pone.0175149> PMID: 28448503; PubMed Central PMCID: PMC5407851.
9. Bhutta ZA, Das JK, Rizvi A, Gaffey MF, Walker N, Horton S, et al. Evidence-based interventions for improvement of maternal and child nutrition: what can be done and at what cost? *Lancet*. 2013; 382(9890):452–77. [https://doi.org/10.1016/S0140-6736\(13\)60996-4](https://doi.org/10.1016/S0140-6736(13)60996-4) PMID: 23746776.
10. Black RE, Victora CG, Walker SP, Bhutta ZA, Christian P, de Onis M, et al. Maternal and child undernutrition and overweight in low-income and middle-income countries. *Lancet*. 2013; 382(9890):427–51. [https://doi.org/10.1016/S0140-6736\(13\)60937-X](https://doi.org/10.1016/S0140-6736(13)60937-X) PMID: 23746772.
11. Ruel MT, Alderman H, Maternal, Child Nutrition Study G. Nutrition-sensitive interventions and programmes: how can they help to accelerate progress in improving maternal and child nutrition? *Lancet*. 2013; 382(9891):536–51. [https://doi.org/10.1016/S0140-6736\(13\)60843-0](https://doi.org/10.1016/S0140-6736(13)60843-0) PMID: 23746780.
12. Stammers AL, Lowe NM, Medina MW, Patel S, Dykes F, Perez-Rodrigo C, et al. The relationship between zinc intake and growth in children aged 1–8 years: a systematic review and meta-analysis. *European journal of clinical nutrition*. 2015; 69(2):147–53. <https://doi.org/10.1038/ejcn.2014.204> PMID: 25335444.
13. Low MS, Speedy J, Styles CE, De-Regil LM, Pasricha SR. Daily iron supplementation for improving anaemia, iron status and health in menstruating women. *The Cochrane database of systematic reviews*. 2016; 4:CD009747. <https://doi.org/10.1002/14651858.CD009747.pub2> PMID: 27087396.

14. Lloyd-Jones DM, Hong Y, Labarthe D, Mozaffarian D, Appel LJ, Van Horn L, et al. Defining and setting national goals for cardiovascular health promotion and disease reduction: the American Heart Association's strategic Impact Goal through 2020 and beyond. *Circulation*. 2010; 121(4):586–613. <https://doi.org/10.1161/CIRCULATIONAHA.109.192703> PMID: 20089546.
15. Mozaffarian D, Capewell S. United Nations' dietary policies to prevent cardiovascular disease. *Bmj*. 2011; 343:d5747. <https://doi.org/10.1136/bmj.d5747> PMID: 21917832; PubMed Central PMCID: PMC3230082.
16. Mozaffarian D, Hao T, Rimm EB, Willett WC, Hu FB. Changes in diet and lifestyle and long-term weight gain in women and men. *The New England journal of medicine*. 2011; 364(25):2392–404. <https://doi.org/10.1056/NEJMoa1014296> PMID: 21696306; PubMed Central PMCID: PMC3151731.
17. Mozaffarian D, Appel LJ, Van Horn L. Components of a cardioprotective diet: new insights. *Circulation*. 2011; 123(24):2870–91. <https://doi.org/10.1161/CIRCULATIONAHA.110.968735> PMID: 21690503.
18. Khatibzadeh S, Saheb Kashaf M, Micha R, Fahimi S, Shi P, Elmadfa I, et al. A global database of food and nutrient consumption. *Bulletin of the World Health Organization*. 2016; 94(12):931–4. <https://doi.org/10.2471/BLT.15.156323> PMID: 27994286; PubMed Central PMCID: PMC5153920.
19. Del Gobbo LC, Khatibzadeh S, Imamura F, Micha R, Shi P, Smith M, et al. Assessing global dietary habits: a comparison of national estimates from the FAO and the Global Dietary Database. *The American journal of clinical nutrition*. 2015; 101(5):1038–46. <https://doi.org/10.3945/ajcn.114.087403> PMID: 25788002; PubMed Central PMCID: PMC4409685.
20. Weisell R, Dop MC. The adult male equivalent concept and its application to Household Consumption and Expenditures Surveys (HCES). *Food and nutrition bulletin*. 2012; 33(3 Suppl):S157–62. <https://doi.org/10.1177/15648265120333S203> PMID: 23193766.
21. Khoury CK, Bjorkman AD, Dempewolf H, Ramirez-Villegas J, Guarino L, Jarvis A, et al. Increasing homogeneity in global food supplies and the implications for food security. *Proceedings of the National Academy of Sciences of the United States of America*. 2014; 111(11):4001–6. <https://doi.org/10.1073/pnas.1313490111> PMID: 24591623; PubMed Central PMCID: PMC3964121.
22. Kastner T, Rivas MJ, Koch W, Nonhebel S. Global changes in diets and the consequences for land requirements for food. *Proceedings of the National Academy of Sciences of the United States of America*. 2012; 109(18):6868–72. <https://doi.org/10.1073/pnas.1117054109> PMID: 22509032; PubMed Central PMCID: PMC3345016.
23. Global Nutrition and Policy Consortium. Home of the Global Dietary Database. Available from: <http://www.globaldietarydatabase.org/>.
24. Imamura F, Micha R, Khatibzadeh S, Fahimi S, Shi P, Powles J, et al. Dietary quality among men and women in 187 countries in 1990 and 2010: a systematic assessment. *The Lancet Global health*. 2015; 3(3):e132–42. [https://doi.org/10.1016/S2214-109X\(14\)70381-X](https://doi.org/10.1016/S2214-109X(14)70381-X) PMID: 25701991; PubMed Central PMCID: PMC4342410.
25. Singh GM, Micha R, Khatibzadeh S, Lim S, Ezzati M, Mozaffarian D, et al. Estimated Global, Regional, and National Disease Burdens Related to Sugar-Sweetened Beverage Consumption in 2010. *Circulation*. 2015; 132(8):639–66. <https://doi.org/10.1161/CIRCULATIONAHA.114.010636> PMID: 26124185; PubMed Central PMCID: PMC4550496.
26. Micha R, Khatibzadeh S, Shi P, Fahimi S, Lim S, Andrews KG, et al. Global, regional, and national consumption levels of dietary fats and oils in 1990 and 2010: a systematic analysis including 266 country-specific nutrition surveys. *Bmj*. 2014; 348:g2272. <https://doi.org/10.1136/bmj.g2272> PMID: 24736206; PubMed Central PMCID: PMC3987052.
27. Mozaffarian D, Fahimi S, Singh GM, Micha R, Khatibzadeh S, Engell RE, et al. Global sodium consumption and death from cardiovascular causes. *The New England journal of medicine*. 2014; 371(7):624–34. <https://doi.org/10.1056/NEJMoa1304127> PMID: 25119608.
28. Sekula W, Nelson M, Figurska K, Oltarzewski M, Weisell R, Szponar L. Comparison between household budget survey and 24-hour recall data in a nationally representative sample of Polish households. *Public health nutrition*. 2005; 8(4):430–9. PMID: 15975190.
29. Sichieri R, Pereira RA, Martins A, Vasconcellos A, Trichopoulou A. Rationale, design, and analysis of combined Brazilian household budget survey and food intake individual data. *BMC public health*. 2008; 8:89. <https://doi.org/10.1186/1471-2458-8-89> PMID: 18366647; PubMed Central PMCID: PMC2288604.
30. International Household Survey Network. Assessment of the reliability and relevance of the food data collected in National Household Consumption and Expenditure Surveys. 2014 Contract No.: IHSN Working Paper No. 008.
31. D'Souza A. How Well Do Household-Level Data Characterize Food Security? Evidence from the Bangladesh Integrated Household Survey. 2014.

32. Friel S, Nelson M, McCormack K, Kelleher C, Thriskos P. Methodological issues using household budget survey expenditure data for individual food availability estimation: Irish experience in the DAFNE pan-European project. *DAta Food NETworking. Public health nutrition*. 2001; 4(5B):1143–7. PMID: [11924938](#).
33. Paterakis SE, Nelson M. A comparison between the National Food Survey and the Family Expenditure Survey food expenditure data. *Public health nutrition*. 2003; 6(6):571–80. PMID: [14690038](#).
34. Jariseta ZR, Dary O, Fiedler JL, Franklin N. Comparison of estimates of the nutrient density of the diet of women and children in Uganda by Household Consumption and Expenditures Surveys (HCES) and 24-hour recall. *Food and nutrition bulletin*. 2012; 33(3 Suppl):S199–207. <https://doi.org/10.1177/15648265120333S208> PMID: [23193771](#).
35. Engle-Stone R, Brown KH. Comparison of a Household Consumption and Expenditures Survey with Nationally Representative Food Frequency Questionnaire and 24-hour Dietary Recall Data for Assessing Consumption of Fortifiable Foods by Women and Young Children in Cameroon. *Food and nutrition bulletin*. 2015; 36(2):211–30. <https://doi.org/10.1177/0379572115587272> PMID: [26121703](#).
36. Dary O, Jariseta ZR. Validation of dietary applications of Household Consumption and Expenditures Surveys (HCES) against a 24-hour recall method in Uganda. *Food and nutrition bulletin*. 2012; 33(3 Suppl):S190–8. <https://doi.org/10.1177/15648265120333S207> PMID: [23193770](#).
37. Bermudez OI, Lividini K, Smitz MF, Fiedler JL. Estimating micronutrient intakes from Household Consumption and Expenditures Surveys (HCES): an example from Bangladesh. *Food and nutrition bulletin*. 2012; 33(3 Suppl):S208–13. <https://doi.org/10.1177/15648265120333S209> PMID: [23193772](#).
38. Fiedler JL, Lividini K, Bermudez OI, Smitz MF. Household Consumption and Expenditures Surveys (HCES): a primer for food and nutrition analysts in low- and middle-income countries. *Food and nutrition bulletin*. 2012; 33(3 Suppl):S170–84. <https://doi.org/10.1177/15648265120333S205> PMID: [23193768](#).
39. Louzada MLdC, Levy RB, Martins APB, Claro RM, Steele EM, Verly E Jr, et al. Validating the usage of household food acquisition surveys to assess the consumption of ultra-processed foods: Evidence from Brazil. *Food Policy*. 2017; 72:112–20. <https://doi.org/10.1016/j.foodpol.2017.08.017>.
40. Sununtnasuk C, Fiedler JL. Can household-based food consumption surveys be used to make inferences about nutrient intakes and inadequacies? A Bangladesh case study. *Food Policy*. 2017; 72:121–31. <https://doi.org/10.1016/j.foodpol.2017.08.018>.
41. Coates J, Rogers BL, Blau A, Lauer J, Roba A. Filling a dietary data gap? Validation of the adult male equivalent method of estimating individual nutrient intakes from household-level data in Ethiopia and Bangladesh. *Food Policy*. 2017; 72:27–42. <https://doi.org/10.1016/j.foodpol.2017.08.010>.
42. Claro RM, Levy RB, Bandoni DH, Mondini L. Per capita versus adult-equivalent estimates of calorie availability in household budget surveys. *Cadernos de saude publica*. 2010; 26(11):2188–95. PMID: [21180992](#).
43. Ahmed A. Bangladesh Integrated Household Survey (BIHS) 2011–2012 International Food Policy Research Institute, Dataverse2013 [cited 2015]. Available from: <https://dataverse.harvard.edu/dataset.xhtml?persistentId=hdl:1902.1/21266>.
44. International Food Policy Research Institute. The status of food security in the Feed the Future Zone and other regions of Bangladesh: Results from the 2011–2012 Bangladesh Integrated Household Survey. 2013.
45. Shaheen N, Rahim ATM, Mohiduzzaman M, Parvin Banu C, Bari ML, Basak Tukun A, et al. Food composition table for Bangladesh. 1 ed: Institute of Nutrition and Food Science, Centre for Advanced Research in Sciences, University of Dhaka; 2013.
46. USDA National Nutrient Database for Standard Reference, Release 28. Version Current [Internet]. 2015. Available from: <https://ndb.nal.usda.gov/ndb/>.
47. Boggar A. Tables on weight yield of food and retention factors of food constituents for the calculation of nutrient composition of cooked foods (dishes). 2002.
48. U.S. Department of Agriculture. USDA Table of Nutrient Retention Factors. Release 6. 2007.
49. FAO/INFOODS. Guidelines for Food Matching. Version 1.2. 2012. FAO, Rome.
50. Charrondiere UR, Rittenschober D, Nowak V, Stadlmayr B, Wijesinha-Bettoni R, Haytowitz D. Improving food composition data quality: Three new FAO/INFOODS guidelines on conversions, data evaluation and food matching. *Food chemistry*. 2016; 193:75–81. <https://doi.org/10.1016/j.foodchem.2014.11.055> PMID: [26433290](#).
51. Islam R. Consumption of Unsafe Foods: Evidence from Heavy Metal, Mineral and Trace Element Contamination. Department of Soil Science, Bangladesh Agricultural University, 2013.
52. Department of Agriculture U.S. Food yields. Summarized by different stages of preparation. *Agriculture Handbook*. 1975; 102.

53. European Food Safety Authority. The food classification and description system FoodEx 2 (draft-revision 1). Parma, Italy: 2011.
54. European Food Safety Authority. Guidance on the EU Menu methodology. *EFSA Journal*. 2014; 12(12). <https://doi.org/10.2903/j.efsa.2014.3944>
55. FAO. Food Balance Sheets. A handbook. 2001. FAO. Rome.
56. FAO. Human energy requirements. Report of a Joint FAO/WHO/UNU Expert Consultation. Rome: 2001.
57. Afshin A, Micha R, Khatibzadeh S, Mozaffarian D. Consumption of nuts and legumes and risk of incident ischemic heart disease, stroke, and diabetes: a systematic review and meta-analysis. *The American journal of clinical nutrition*. 2014; 100(1):278–88. <https://doi.org/10.3945/ajcn.113.076901> PMID: [24898241](https://pubmed.ncbi.nlm.nih.gov/24898241/); PubMed Central PMCID: [PMC4144102](https://pubmed.ncbi.nlm.nih.gov/PMC4144102/).
58. Aune D, Norat T, Romundstad P, Vatten LJ. Whole grain and refined grain consumption and the risk of type 2 diabetes: a systematic review and dose-response meta-analysis of cohort studies. *European journal of epidemiology*. 2013; 28(11):845–58. <https://doi.org/10.1007/s10654-013-9852-5> PMID: [24158434](https://pubmed.ncbi.nlm.nih.gov/24158434/).
59. He FJ, Nowson CA, Lucas M, MacGregor GA. Increased consumption of fruit and vegetables is related to a reduced risk of coronary heart disease: meta-analysis of cohort studies. *Journal of human hypertension*. 2007; 21(9):717–28. <https://doi.org/10.1038/sj.jhh.1002212> PMID: [17443205](https://pubmed.ncbi.nlm.nih.gov/17443205/).
60. He K, Song Y, Daviglius ML, Liu K, Van Horn L, Dyer AR, et al. Accumulated evidence on fish consumption and coronary heart disease mortality: a meta-analysis of cohort studies. *Circulation*. 2004; 109(22):2705–11. <https://doi.org/10.1161/01.CIR.0000132503.19410.6B> PMID: [15184295](https://pubmed.ncbi.nlm.nih.gov/15184295/).
61. Mellen PB, Walsh TF, Herrington DM. Whole grain intake and cardiovascular disease: a meta-analysis. *Nutrition, metabolism, and cardiovascular diseases: NMCD*. 2008; 18(4):283–90. <https://doi.org/10.1016/j.numecd.2006.12.008> PMID: [17449231](https://pubmed.ncbi.nlm.nih.gov/17449231/).
62. Micha R, Wallace SK, Mozaffarian D. Red and processed meat consumption and risk of incident coronary heart disease, stroke, and diabetes mellitus: a systematic review and meta-analysis. *Circulation*. 2010; 121(21):2271–83. <https://doi.org/10.1161/CIRCULATIONAHA.109.924977> PMID: [20479151](https://pubmed.ncbi.nlm.nih.gov/20479151/); PubMed Central PMCID: [PMC2885952](https://pubmed.ncbi.nlm.nih.gov/PMC2885952/).
63. He Y. Missing data analysis using multiple imputation: getting to the heart of the matter. *Circulation Cardiovascular quality and outcomes*. 2010; 3(1):98–105. <https://doi.org/10.1161/CIRCOUTCOMES.109.875658> PMID: [20123676](https://pubmed.ncbi.nlm.nih.gov/20123676/); PubMed Central PMCID: [PMC2818781](https://pubmed.ncbi.nlm.nih.gov/PMC2818781/).
64. McLean RM. Measuring population sodium intake: a review of methods. *Nutrients*. 2014; 6(11):4651–62. <https://doi.org/10.3390/nu6114651> PMID: [25353661](https://pubmed.ncbi.nlm.nih.gov/25353661/); PubMed Central PMCID: [PMC4245554](https://pubmed.ncbi.nlm.nih.gov/PMC4245554/).
65. Jacobs K, Sumner D. The food balance sheets of the Food and Agriculture Organization: a review of potential ways to broaden the appropriate uses of the data. 2002. Review sponsored by the FAO.
66. Foster E, Hawkins A, Barton KL, Stamp E, Matthews JN, Adamson AJ. Development of food photographs for use with children aged 18 months to 16 years: Comparison against weighed food diaries—The Young Person's Food Atlas (UK). *PloS one*. 2017; 12(2):e0169084. <https://doi.org/10.1371/journal.pone.0169084> PMID: [28199319](https://pubmed.ncbi.nlm.nih.gov/28199319/); PubMed Central PMCID: [PMC5310878](https://pubmed.ncbi.nlm.nih.gov/PMC5310878/).
67. FAO. Introducing the Minimum Dietary Diversity—Women (MDD-W) Global Dietary Diversity Indicator for Women Washington DC: 2014.
68. Kothari M, Abderrahim N. Nutrition Update 2010. Calverton, Maryland, USA: ICF Macro: 2010.
69. United Nations. The Millennium Development Goals Report. 2015. New York.
70. United Nations. United Nations Millennium Declaration. September 2000.
71. World Health Organization. Available from: [www.who.int](http://www.who.int).
72. Elshenawy S, Simmons R. Maternal obesity and prenatal programming. *Molecular and cellular endocrinology*. 2016; 435:2–6. Epub 2016/07/10. <https://doi.org/10.1016/j.mce.2016.07.002> PMID: [27392495](https://pubmed.ncbi.nlm.nih.gov/27392495/).
73. Godfrey KM, Gluckman PD, Hanson MA. Developmental origins of metabolic disease: life course and intergenerational perspectives. *Trends in endocrinology and metabolism: TEM*. 2010; 21(4):199–205. Epub 2010/01/19. <https://doi.org/10.1016/j.tem.2009.12.008> PMID: [20080045](https://pubmed.ncbi.nlm.nih.gov/20080045/).
74. Christian P, Stewart CP. Maternal micronutrient deficiency, fetal development, and the risk of chronic disease. *The Journal of nutrition*. 2010; 140(3):437–45. Epub 2010/01/15. <https://doi.org/10.3945/jn.109.116327> PMID: [20071652](https://pubmed.ncbi.nlm.nih.gov/20071652/).
75. Pereira TJ, Moyce BL, Kereliuk SM, Dolinsky VW. Influence of maternal overnutrition and gestational diabetes on the programming of metabolic health outcomes in the offspring: experimental evidence. *Biochemistry and cell biology = Biochimie et biologie cellulaire*. 2015; 93(5):438–51. Epub 2015/02/13. <https://doi.org/10.1139/bcb-2014-0141> PMID: [25673017](https://pubmed.ncbi.nlm.nih.gov/25673017/).

76. Murphy S, Ruel M, Carriquiry A. Should Household Consumption and Expenditures Surveys (HCES) be used for nutritional assessment and planning? *Food and nutrition bulletin*. 2012; 33(3 Suppl):S235–41. <https://doi.org/10.1177/15648265120333S213> PMID: 23193776.
77. Naska A, Berg MA, Cuadrado C, Freisling H, Gedrich K, Gregoric M, et al. Food balance sheet and household budget survey dietary data and mortality patterns in Europe. *The British journal of nutrition*. 2009; 102(1):166–71. <https://doi.org/10.1017/S000711450809466X> PMID: 18986595.
78. Claro RM, Jaime PC, Lock K, Fisberg RM, Monteiro CA. Discrepancies among ecological, household, and individual data on fruits and vegetables consumption in Brazil. *Cadernos de saude publica*. 2010; 26(11):2168–76. PMID: 21180990.
79. Berti PR. Intra-household distribution of food: a review of the literature and discussion of the implications for food fortification programs. *Food and nutrition bulletin*. 2012; 33(3 Suppl):S163–9. <https://doi.org/10.1177/15648265120333S204> PMID: 23193767.
80. Beegle K, De Weerd J, Friedman J, Gibson J. Methods of household consumption measurement through surveys: Experimental results from Tanzania. *Journal of Development Economics*. 2012; 98(1):3–18. <https://doi.org/10.1016/j.jdeveco.2011.11.001>.
81. Livingstone MB, Robson PJ. Measurement of dietary intake in children. *The Proceedings of the Nutrition Society*. 2000; 59(2):279–93. PMID: 10946797.
82. Ansari MNA, Atkins PJ. Understanding the munga in northwest Bangladesh: household perceptions and perceptual connotations *International Research Journal of Social Sciences*. 2014; 3(8):22–9.
83. Micha R, Coates J, Leclercq C, Charrondiere UR, Mozaffarian D. Global Dietary Surveillance: Data Gaps and Challenges. *Food and nutrition bulletin*. 2018; 39(2):175–205. Epub 2018/02/27. <https://doi.org/10.1177/0379572117752986> PMID: 29478333.