

Published in final edited form as:

Neuron. 2006 October 19; 52(2): 271–280. doi:10.1016/j.neuron.2006.07.023.

Definition of estrogen receptor pathway critical for estrogen positive feedback to gonadotropin-releasing hormone (GnRH) neurons and fertility

Tim M. Wintermantel^{#1}, Rebecca E. Campbell^{#2}, Robert Porteous^{#2}, Dagmar Bock¹, Hermann-Josef Gröne¹, Martin G. Todman², Kenneth S. Korach³, Erich Greiner¹, Cristian A. Pérez⁴, Günther Schütz¹, and Allan E. Herbison²

¹Molecular Biology of the Cell I, German Cancer Research Center, Im Neuenheimer Feld 280, D-69120 Heidelberg, Germany ²Centre for Neuroendocrinology and Department of Physiology, University of Otago School of Medical Sciences, Dunedin 9001, New Zealand ³Receptor Biology Section, Laboratory of Reproductive and Developmental Toxicology, NIEHS, Research Triangle Park, USA ⁴Laboratory of Molecular Genetics, The Rockefeller University, 1230 York Ave., New York, New York 10021, USA

These authors contributed equally to this work.

Summary

The mechanisms through which estrogen regulates gonadotropin-releasing hormone (GnRH) neurons to control mammalian ovulation are unknown. We found that estrogen positive feedback to generate the preovulatory gonadotropin surge was normal in estrogen receptor β knockout (ER β) mutant mice but absent in ER α mutant mice. An ER α -selective compound was sufficient to generate positive feedback in wild-type mice. As GnRH neurons do not express ER α , estrogen positive feedback upon GnRH neurons must be indirect in nature. To establish the cell type responsible, we generated a neuron-specific ER α mutant mouse line. These mice failed to exhibit estrogen positive feedback demonstrating that neurons expressing ER α are critical. We then used GnRH neuron-specific Cre-dependent Pseudorabies virus tracing to show that the ER α -expressing neurons modulating GnRH neurons are located within rostral periventricular regions of the hypothalamus. These studies demonstrate ovulation is driven by estrogen actions upon ER α -expressing neuronal afferents to GnRH neurons.

Introduction

The gonadotropin releasing hormone (GnRH) neurons represent the key output cells of the neuronal network controlling fertility in all mammalian species. The GnRH neurons and associated cells that comprise the “GnRH neuronal network” are responsible for integrating

Correspondence to: Allan E. Herbison, Centre for Neuroendocrinology, Department of Physiology, University of Otago School of Medical Sciences, P.O. Box 913, Dunedin, New Zealand, *Telephone* -64-3-479 7312, *Fax* -64-3-479 7323, allan.herbison@stonebow.otago.ac.nz or to: Günther Schütz, Department of Molecular Biology of the Cell I, German Cancer Research Center, Im Neuenheimer Feld 280, D-69120 Heidelberg, Germany, *Telephone* -49-6221-42 34 11, *Fax* -49-6221-42 34 70, g.schuetz@dkfz.de.

multiple internal homeostatic and external environmental signals to ensure appropriate levels of fertility for the individual (Levine, 1997; Herbison, 2006). Arguably, the most important of these signals is that of estrogen, secreted by the gonads to achieve feedback regulation of gonadotropin secretion (Herbison, 1998; Petersen et al., 2003). Throughout most of the menstrual cycle, estrogen suppresses gonadotropin secretion but, at mid-cycle, switches to have a potent stimulatory or “positive feedback” action to evoke the luteinizing hormone (LH) surge that triggers ovulation. Although feedback effects of estrogen are known to occur at the pituitary gland (Shupnik, 1996), actions of estrogen within the brain are accepted as being critical for the generation of the GnRH surge that drives the preovulatory LH surge in all mammals, including primates (Karsch et al., 1997; Herbison, 1998).

Even though estrogen positive feedback is central to mammalian fertility, the underlying mechanism remains poorly understood. Since GnRH neurons express ER β , but not ER α , it is possible that estrogen acts directly upon them to generate the GnRH surge (Herbison and Pape, 2001; Petersen et al., 2003). However, several lines of evidence indicate that effects of estrogens may be transmitted to GnRH neurons in an indirect manner by ER α - and/or ER β -expressing neurons, glia or endothelial cells (Rage et al., 1997; Smith and Jennes, 2001; Prevot, 2002; Petersen et al., 2003). The detailed investigation of this mechanism has been hampered by the scattered distribution of the GnRH neurons that makes them difficult to investigate technically. Thus, at present, neither the estrogen receptor subtype (ER α vs. ER β), nor the critical cell types involved in estrogen positive feedback, have been defined and much controversy surrounds this critical issue. A genetic approach to define which of these two estradiol receptors is crucial for GnRH neuronal activation to induce ovulation, and to characterize its role in specific cells might provide valuable insights. Using mice in which ER α or ER β have been inactivated, we first demonstrate here that ER α , but not ER β , is required for estrogen positive feedback to GnRH neurons. Using an ER α -selective ligand in wild-type mice we show that ER α is not only necessary but sufficient to generate estrogen positive feedback. Second, exploiting a novel neuron-specific mutation of the ER α gene we were able to identify neurons, as opposed to other cell types, as critical targets for estradiol action. This clearly establishes that neurons expressing ER α are required for estrogen to activate GnRH neurons. Finally, through use of a GnRH neuron-specific Pseudorabies virus (PRV) tracing approach we have been able to define the location of ER α -expressing neurons projecting to GnRH neurons. These data demonstrate a key role for ER α in mammalian estrogen positive feedback and provide definitive evidence for the “indirect model” of estrogen action whereby estrogen regulates ER α -expressing neuronal afferents to the GnRH neurons to bring about the preovulatory GnRH/LH surge.

Results

Estrogen positive feedback on LH secretion and GnRH neuron activation is absent in ER α mutant and normal in ER β mutant female mice

The stimulatory effects of estradiol positive feedback were evaluated using a protocol that enabled the activation status of GnRH neurons to be assessed alongside changes in plasma LH levels. Ovariectomized wild-type C57BL6/J mice given an estradiol capsule followed by an injection of estradiol benzoate were found to exhibit an LH surge centered around

19:00h, on lights out (Fig.1A). In the ER α mutant mice, there was no evidence of an LH surge in any of the mice (n=8) in response to the same estrogen regimen with mean LH levels at 19:00h of 1.6 ± 0.3 ng/ml compared with 5.2 ± 1.0 ng/ml in wild-type control animals (n=7, $p<0.05$; Fig.1B). In contrast, clear evidence of an LH surge was observed in each of the ovariectomized, estrogen-treated ER β mutant mice (mean LH levels of 8.6 ± 1.7 ng/ml compared with 8.9 ± 1.5 ng/ml in controls, n= 7 each group) (Fig.2A).

Approximately 40% of the GnRH neuronal population express the immediate early gene c-Fos at the time of GnRH surge and this is believed to be an accurate indicator of those GnRH neurons activated by estrogen to generate the GnRH surge (Hoffman et al., 1993). In addition, there is increasing evidence that neurons within the anteroventral periventricular nucleus (AVPV) of the rostral hypothalamus may be an important target for estrogen in bringing about the GnRH surge (Herbison, 1998; Simerly, 2002). Dual labeling immunocytochemistry (Fig.1C,D) revealed that approximately 40% of GnRH neurons in wild-type mice expressed c-Fos at the time of the estrogen-induced LH surge (Fig.1C,E). The GnRH neurons expressing c-Fos were located preferentially within the preoptic area with dual-labelled GnRH neurons only rarely detected in more rostral areas such as the medial septum (data not shown). In contrast, no GnRH neurons were detected to express c-Fos in any brain region of ER α mutant mice (Fig.1D,E). The distribution and number of GnRH neurons detected in wild-type and ER α mutant females were not different (data not shown). In addition, we found that the numbers of singly-labelled c-Fos-expressing cells located within the AVPV were significantly greater in wild-type mice (Fig.1F) compared with ER α mutant females ($p<0.05$; Fig.1G,H).

Single and dual-label immunocytochemistry experiments in ER β mutant females revealed no differences in the number of dual-labelled c-Fos-GnRH neurons compared with wild-type controls (Fig.2B) or in the number of c-Fos-expressing cells detected in the AVPV (Fig.2C). The distribution and number of GnRH neurons in wild-type and ER β mutant females were not different (data not shown).

An ER α -specific ligand is sufficient to generate positive feedback in wild-type mice

The results with the ER α and β mutant mice indicated that ER α was necessary for estrogen positive feedback. To evaluate whether ER α -pathways were sufficient for positive feedback, wild-type mice were given the ER α -selective compound 16 α -LE2 (Hegele-Hartung et al., 2004) as the second estradiol injection alongside control mice receiving the normal estrogen protocol. The second estradiol injection is critical for evoking positive feedback as mice given vehicle control at this time point never exhibit an LH surge. Mice treated with 16 α -LE2 (n=5) exhibited an LH surge at 19:00h (LH, 5.9 ± 1.7 ng/ml compared with normal estrogen-treated controls 6.5 ± 2.3) and had $49\pm 7\%$ of rostral preoptic area GnRH neurons expressing c-Fos (compared with controls that had $60\pm 11\%$).

Together, these observations demonstrate that cells expressing ER α are both necessary and sufficient for estrogen positive feedback actions upon GnRH neurons. As GnRH neurons do not express ER α , these results demonstrate that ER α -expressing neuronal, glial or other cell types must mediate estrogen positive feedback actions upon GnRH neurons.

Neuron-specific ER α mutant mice are infertile

To examine the hypothesis that estrogen acts through ER α -expressing neurons, rather than other cell types, to activate GnRH neurons, neuron-specific ER α mutant mice were developed by breeding ER α flox mice with a CamKiCre transgenic mouse line. Immunocytochemical analyses of ER α expression in the resulting mice showed that ER α immunoreactivity was absent from the hypothalamus (Fig.3 AB) and all other brain regions (Suppl. Fig. 2) of ER $\alpha^{fl/fl};CamKiCre$ mice. In contrast, ER α protein was detected in the pituitary of all genotypes including female ER $\alpha^{fl/fl};CamKiCre$ mice (Fig.3 AB insets). Female ER $\alpha^{fl/fl};CamKiCre$ mice were found to be infertile and, from an age of 5-6 weeks, to exhibit striking abnormalities in their reproductive organs (Fig.4). In the mutants, the uterus was grossly enlarged and filled with liquid (Fig.4B). The endometrium, however, was severely atrophic and lacked all glandular structures. Whereas the endometrial stroma was loose and vascularized in control mice (Fig.4C), it appeared condensed and showed granulocyte infiltration in the mutants (Fig.4D). Histologically, the ovaries of ER $\alpha^{fl/fl};CamKiCre$ mice showed signs of gonadotropin hyperactivation; in mutant mice, ovaries contained a large number of antral follicles compared with wildtype animals (Fig. 4E,F). In the ovarian hilus, theca cells appeared hypertrophic and luteinized in mutant animals, indicating inappropriate stimulation of the ovaries. Furthermore, no corpora lutea were observed in the ovaries of mutants (Fig.4F) suggesting a failure of ovulation. Basal LH levels were not different between diestrous controls (0.91 ± 0.11 ng/ml) and ER $\alpha^{fl/fl};CamKiCre$ (1.23 ± 0.08 ng/ml) female mice.

Neuron-specific ER α mutant mice are unable to generate estrogen positive feedback

Ovariectomized, estrogen-treated ER $\alpha^{fl/fl};CamKiCre$ mice ($n=6$) failed to exhibit an LH surge with mean LH levels at 19:00h of 0.9 ± 0.2 ng/ml compared with 5.8 ± 2.1 ng/ml in littermate controls ($n=5$; Fig.3C). Immunocytochemical experiments did not detect c-Fos in any GnRH neurons in ER $\alpha^{fl/fl};CamKiCre$ mice following estrogen treatment. This compared with c-Fos in $41\pm 8\%$ of GnRH neurons in littermate controls (Fig.3D).

As c-Fos protein is detectable for 5-6h in GnRH neurons following the GnRH/Lh surge (Lee et al., 1990) we additionally evaluated c-Fos expression in GnRH neurons at midday of the expected day of the surge and at 1am on the following day to evaluate whether the timing of positive feedback had been altered in ER $\alpha^{fl/fl};CamKiCre$ mice. We found a complete absence of c-Fos in GnRH neurons at these time points. As we find no evidence for c-Fos in GnRH neurons of ER $\alpha^{fl/fl};CamKiCre$ at 13:00h, 19:00h or 1am the next day, it seems unlikely that the GnRH neurons receive either an early or delayed positive feedback signal in ER $\alpha^{fl/fl};CamKiCre$.

The distribution and number of GnRH neurons in ER $\alpha^{fl/fl};CamKiCre$ brains was identical to that of controls (data not shown). Whereas numbers of c-Fos-positive cells in a control brain region were not different between control littermates (5.1 ± 2.2 cells/unit area/section) and ER $\alpha^{fl/fl};CamKiCre$ mice (8.3 ± 1.6 cells/unit area/section), significantly more singly-labelled c-Fos cells were detected within the AVPV of littermate controls compared to ER $\alpha^{fl/fl};CamKiCre$ mice ($p<0.05$; Fig.3E). These data closely resemble that of the global

ER α mutant mouse and indicate that ER α -expressing neurons are critical for estrogen positive feedback to occur.

The ER α -expressing primary afferents to GnRH neurons are located within periventricular regions of the rostral hypothalamus

The studies above show that ER α -expressing neurons are critical for estrogen positive feedback to GnRH neurons. To identify the locations of these cells within the “GnRH neuronal network”, we have used a Cre-dependent PRV tracing strategy in GnRH-Cre mice (DeFalco et al., 2001; Yoon et al., 2005). In this approach, the Ba2001 PRV is activated only after infecting a Cre-expressing GnRH neuron *in vivo*. This allows for the now unconditional PRV to replicate and to pass in a retrograde manner to the primary afferents of the GnRH neuron and, subsequently, their own afferents in a time-dependent manner. The retrograde chain of infection can be followed by evaluating GFP expression as the unconditional Ba2001 PRV also expresses GFP in each cell it infects.

Five founder GnRH-Cre lines were generated. Analysis of Cre expression by dual-label immunocytochemistry in adult female mice revealed that one of these lines expressed Cre in a highly selective manner within 97 \pm 2% of all GnRH neurons (n=6; Fig.5A). Crossing this line with the ROSA-26 indicator mice (Soriano, 1999) revealed Cre-dependent recombination in 97 \pm 1% of GnRH neurons (n=4; not shown).

To evaluate the ER α -expressing afferents to the GnRH neuron population that are activated at the time of the GnRH surge (i.e. those in the rostral preoptic area), a single 500nl injection of Ba2001 PRV was given into the rostral preoptic area of adult female mice. Mice examined at 6, 12, 24 and 30h after injection of Ba2001 (n=8) displayed no evidence of GFP expression in any region of the brain. However, 48h later (n=4), we found that between 5 and 9 GnRH neurons located around the injection site expressed GFP (Fig.5B,C) indicating that Cre-mediated recombination had occurred in these cells. No other GFP-expressing cells were detected within the brain at this time point. Seventy-two hours following PRV injection, GFP-expressing cells (representing afferent neurons to rostral preoptic GnRH neurons, Fig.5D) were identified in multiple brain regions. Twenty-four hours later (96h post-injection), we found a much larger distribution and number of GFP-expressing cells. As 24h is required for PRV to move from one order of neurons to the next (Card and Enquist, 1995; Horvath et al., 2002), the neurons identified at 72h post-injection are almost certainly primary afferents to GnRH neurons.

At 72h post-injection (n=6), GFP-expressing neurons were identified in multiple areas of the forebrain and brainstem (Fig.5G). The regions displaying the largest numbers of retrogradely-labeled neurons were the hypothalamic periventricular nucleus (PeN; mean of 36 cells/mouse in a 1:3 series of brain sections), AVPV (23 cells), median preoptic nucleus (MnPO; 14 cells) and dorsal raphe (16 cells). Other brain regions that consistently exhibited GFP-expressing neurons at lower densities were the lateral septum, arcuate nucleus (ARN) and nucleus tractus solitarius (Fig.5G).

Dual-labelling immunocytochemistry (n=6) for GFP and ER α (Fig.5D,E) identified ER α -expressing afferents to be located in four brain regions; the AVPV, MnPO, PeN and ARN

(Fig.5F). The percentage of GFP-positive neurons expressing ER α was 53 \pm 12%, 34 \pm 21%, 24 \pm 11% in the AVPV, MnPO and PeN respectively, and 56 \pm 19%, for the ARN although we only ever observed 1-2 GFP-expressing cells in the ARN (Fig.5G). To ensure the neuronal identity of PRV-infected cells expressing ER α , immunocytochemical labeling for GFP and ER α was undertaken in combination with Nissl staining. All GFP/ER α -expressing cells were also positive for Nissl (Suppl. Fig. 3). Injections of Ba2001 PRV into the rostral preoptic area of wild-type, non-Cre-expressing mice, or into the striatum of GnRH-Cre mice, resulted in a complete absence of GFP-expressing neurons. These findings demonstrate that ER α -expressing primary afferents to GnRH neurons have their cell bodies located principally within rostral periventricular nuclei of the hypothalamus.

The findings of the PRV study demonstrated that significant estrogen-receptive primary afferents to the GnRH neurons existed in the PeN and MnPO in addition to the AVPV. As such, we returned to the estrogen positive feedback c-Fos studies and examined c-Fos expression in the MnPO and PeN of ER α and ER β mutant mice. The numbers of c-Fos cells in the PeN of mice treated with estrogen were significantly reduced in ER α mutants (22 \pm 3 cells, p <0.05) compared with wild-type mice (40 \pm 6 cells per section) but were unchanged in ER β mutant females. The numbers of c-Fos-expressing cells in the MnPO were not significantly different in either ER α - or ER β -mutant mice compared with controls. These data indicate that a sub-population of cells within the PeN, as well as the AVPV, are activated by estrogen positive feedback.

Discussion

We have used here a series of ER mutant mouse models and an ER-selective ligand to demonstrate the critical receptor isoform and cell type necessary and sufficient for estrogen to initiate the GnRH/LH surge and ovulation. We show that global ER β mutant mice exhibit normal patterns of estrogen-induced GnRH activation and LH surge secretion. In contrast, global ER α mutant mice fail to exhibit GnRH neuron activation or LH surge secretion in response to estrogen. As GnRH neurons do not express ER α (Herbison and Pape, 2001), this indicated that positive feedback effects of estrogen must occur in an indirect manner. To define the critical ER α -expressing cell-type in the pathway, neuron-specific ER α mutant mice were generated by crossing a new floxed exon 3 ER α mouse line with CamKII α -Cre mice. CamKII α is expressed selectively by forebrain neurons (Ouimet et al., 1984; Burgin et al., 1990) and the CamKII α -Cre mouse line used here has been shown previously to efficiently delete loxP-flanked target sequences in neurons of the CNS, leading to neuron-specific gene ablation (Casanova et al., 2001; Marsicano et al., 2003). Mice harbouring a neuron-specific ER α deletion were found to be infertile and lack estrogen positive feedback demonstrating that ER α -expressing neurons are critical. To define the location of these neurons we have used a novel conditional Cre-Pseudorabies virus approach to demonstrate that the ER α -expressing primary afferents to GnRH neurons are located primarily within the periventricular nuclei of the rostral hypothalamus. Together, these observations provide conclusive evidence that critical positive feedback actions of estrogen upon GnRH neurons are mediated by ER α -expressing neuronal afferents within the GnRH neuronal network.

It is now recognized that estrogen can modulate the activity of neuronal networks through multiple mechanisms including both slow genomic and rapid actions (McEwen and Alves, 1999). In the case of estrogen positive feedback, however, it seems clear that classical genomic mechanisms underlie the estrogen activation of GnRH neurons. In all species examined, positive feedback only occurs after several hours of estrogen exposure (Legan et al., 1975; Bronson, 1981; Xia et al., 1992; Evans et al., 1997). Furthermore, once the animal has been exposed to estrogen for a sufficient period, it can be removed without having any negative effect upon subsequent GnRH neuron activation. Thus, rapid and immediate actions of estrogen are not likely to be critical for GnRH neuron activation and positive feedback relies upon a classical genomic mechanism. The down-stream genes regulated by estrogen to bring about GnRH neuron activation are not yet established although the progesterone receptor is one candidate known to be regulated potently by estrogen (Levine, 1997; Shughrue et al., 1997). The finding of ER β mRNA and protein in GnRH neurons of the mouse (Skynner et al., 1999a), the rat (Hrabovszky et al., 2001) and, more recently, the sheep (Skinner and Dufourny, 2005), raised the possibility that estrogen might act directly upon GnRH neurons via this isoform to generate the GnRH/LH surge (Herbison and Pape, 2001). However, in terms of both LH secretion and immediate early gene expression in GnRH neurons, we found that estrogen positive feedback is normal in the global ER β mutant mouse. This strongly suggests that ER β -regulated signaling within GnRH neurons is not critical for estrogen's positive feedback effects upon these cells.

The global ER α mutant mice are infertile and exhibit an anovulatory ovarian phenotype with polycystic follicles and an absence of corpora lutea (Couse et al., 1999; Couse and Korach, 1999). We provide evidence here that the failure of estrogen positive feedback on the GnRH neuron network is very likely to underlie the anovulatory phenotype of the global ER α mutant mice. Estrogen administration to ovariectomized global ER α mutant mice failed to generate an LH surge or evidence of GnRH neuron activation. This provided direct evidence of the critical importance of ER α -expressing cells in positive feedback. In keeping with this observation, the ER α -selective compound 16 α -LE2 (Hegele-Hartung et al., 2004) was found to be capable of eliciting normal positive feedback in wild-type mice. This indicates that ER α activation is both necessary and sufficient for positive feedback to occur. As the GnRH neurons do not themselves express ER α (Herbison and Pape, 2001), these observations in ER α mutant mice suggested that other ER α -expressing cell types within the GnRH neuronal network were critical for positive feedback to occur. Several different modes of indirect estrogen input to the GnRH neurons have been proposed and include effects mediated by vascular endothelial cells, tanycytes, glial cells, and interneurons (Rage et al., 1997; Herbison, 1998; Smith and Jennes, 2001; Prevot, 2002; Petersen et al., 2003). Using a Cre-loxP strategy that generates a neuron-specific ER α knockout, we have been able to demonstrate that, of these possibilities, it is ER α -expressing neurons within the network that are critical for estrogen positive feedback. Other cell types expressing ER α may play a role in positive feedback but are insufficient, by themselves, to activate the GnRH neurons to generate the GnRH/LH surge. Thus, the present genetic dissection of the estrogen positive feedback mechanism identifies clearly both the critical ER isoform and cell type involved.

The absence of estrogen positive feedback in neuron-specific ER α mutant mice is entirely compatible with their infertile reproductive phenotype. Ovarian histology shows an absence

of corpora lutea and abundance of antral follicles supporting the failure of the ovulatory mechanism. In contrast to the global ER α mutant line, there was no evidence in neuron-specific ER α mutant mice for a polycystic hemorrhagic ovarian phenotype generated by excessive gonadotrophin secretion (Couse et al., 1999). Indeed, basal LH levels are not elevated in neuron-specific ER α mutant mice suggesting that pituitary estrogen negative feedback is probably sufficient to restrain gonadotrophin secretion. Another prominent feature of the neuron-specific ER α mutant mice is that of dilated, fluid-filled uteri reminiscent of that found on proestrus. Unlike in the global ER α mutant mouse (Couse and Korach, 1999; Dupont et al., 2000), the uteri of neuron-specific ER α mutant mice remain sensitive to estrogen and may be exposed to a continual high level of estrogen originating from the large numbers of developing follicles. Thus, the reproductive phenotype of the neuron-specific ER α mutant mouse is compatible with that of a mouse in which estrogen negative feedback is competent but positive feedback is absent. This would enable relatively normal basal gonadotrophin secretion, sufficient to promote follicular growth and estrogen production, to exist in association with a complete failure of ovulation.

Studies in the rat have implicated the AVPV as an estrogen-sensitive brain region involved in the positive feedback mechanism (Wiegand and Terasawa, 1982; Herbison, 1998; Simerly, 2002). The present results provide definitive evidence for this hypothesis and also suggest that the locations of estrogen-sensitive neurons involved in estrogen positive feedback include the preoptic PeN immediately caudal to the AVPV. To determine the locations of ER α -expressing afferents to GnRH neurons we generated a GnRH-Cre transgenic mouse for use with a novel Cre-dependent PRV retrograde tracing strategy (DeFalco et al., 2001). By establishing a time-course of viral infection and restricting our PRV injection sites to the rostral preoptic area, where GnRH neurons expressing c-Fos at the time of positive feedback reside, we traced out the primary afferents to the sub-population of GnRH neurons activated by estrogen to create the preovulatory GnRH surge. Primary afferent inputs to GnRH neurons were found to originate from a variety of predominantly mid-line hypothalamic and brainstem locations. However, the sub-populations of ER α -expressing afferents were found to be tightly clustered within the rostral preoptic area. In addition to a strong projection from ER α -expressing neurons in the AVPV, we also identified populations of ER α -primary afferents originating from the PeN and, nearby the GnRH somata, in the MnPO. Our c-Fos studies support the functional relevance of both AVPV and PeN neurons in estrogen positive feedback as a strong correlation existed between levels of c-Fos in these brain regions and GnRH neuron activation and the occurrence of an LH surge. A similar observation has been made in the rat where c-Fos is expressed by AVPV/PeN neurons in response to estrogen positive feedback (Le et al., 1999).

Brought together, the evidence for the AVPV and PeN as sites of primary afferents mediating estrogen positive feedback to rodent GnRH neurons is now very strong; (1) lesions of the AVPV/ventral PeN abolish positive feedback (Wiegand and Terasawa, 1982), (2) implantation of anti-estrogens in the AVPV/PeN abolish positive feedback (Petersen and Barraclough, 1989), (3) AVPV and PeN neurons provide inputs to GnRH neurons (present study and (Horvath et al., 1993; Gu and Simerly, 1997; Simonian et al., 1999)), (4) neurons expressing ER α are critical for estrogen positive feedback (present study), (5) AVPV and PeN neurons express ER α and are activated by estrogen positive feedback (present study

and (Le et al., 1999)), and (6) AVPV/PeN neurons expressing ER α project directly to GnRH neurons (present study).

Together, these studies clarify the mechanism of estrogen positive feedback to GnRH neurons by providing definitive evidence that it is ER α , and not ER β , that is critical and that a neuron-specific ablation of ER α renders mice infertile with a complete ablation of estrogen positive feedback. Thus, estrogen acts indirectly upon GnRH neurons to bring about positive feedback driving ovulation and the ER α -expressing neurons mediating this pathway are located in periventricular regions. Future studies involving the genetic ablation of ER α in specific neuronal phenotypes will now be required to define the hierarchy of periventricular neurons mediating estrogen feedback to GnRH neurons.

Experimental Procedures

Generation of an ER α flox (ER α fl) mouse line

A conditional allele of the mouse *Esr1* allele was generated. The targeting construct (pER α flox.tkneo.DTA) (Suppl. Fig. 1A, top panel) was based on a 9 kb *Bam*HI fragment representing the genomic sequence of exon 3 and surrounding introns of the mouse *Esr1* gene (Suppl. Fig. 1A, second panel). This fragment, obtained from the RPCI21 mouse genomic library (Vente et al., 1999), was modified using homologous recombination in *E. Coli* (Zhang et al., 1998) to carry a loxP site 5' to exon 3, a PGKtkneo cassette flanked by flr sites and one loxP site in the 3' direction of exon 3. E14 embryonic stem cells were transfected with the linearised targeting construct and selected for construct integration (Tronche et al., 1999). G418-resistant clones were characterised by Southern blot using external genomic probes from the *Estra* locus (Suppl. Fig. 1B). Clones that had undergone homologous recombination were transiently transfected with the expression plasmid pCAGGS-Flpe (Schaff et al., 2001) and selected with 1 μ M Gancyclovir to isolate subclones that have lost the selection cassette after Flp recombination. This was verified by Southern blot analysis (Suppl. Fig. 1C). From the resulting embryonic stem cell clones, chimeric mice were generated by blastocyst injection and uterine transfer. By breeding these chimeras to C57BL/6 mice, the ER α fl mouse line was established. To generate a neuron-specific ER α knockout mouse, ER α fl mice were bred to CamKII α -iCre BAC transgenic mice (referred to as CamKiCre mice) (Casanova et al., 2001). CamKII α is expressed around birth in almost all forebrain neurons but not glial cells (Ouimet et al., 1984; Burgin et al., 1990). This resulted in the generation of ER α ^{fl/fl};CamKiCre mice with heterozygous ER α ^{+fl}; CamKiCre mice, and ER α ^{fl/fl} mice serving as controls.

Generation of a GnRH-cre transgenic mouse

The GnRH-Cre mouse was generated in the same manner as reported previously (Skynner et al., 1999b) using a ~12kb transgene incorporating all the introns and exons of the GnRH gene, 5.5 kb of upstream (5') and 3.5 kb of downstream (3') flanking sequence, and a SmaI site engineered by site-directed mutagenesis in exon II between sequences encoding amino acids 2 and 3 of the GnRH decapeptide. An expression cassette consisting of (in 5' to 3' order) a synthetic intron, nuclear-localizing signal, *CRE recombinase* and a polyadenylation (polyA) sequence was then inserted at the SmaI site to produce the transgene. Transgenic

mice were produced by pronuclear injection and identified by PCR analysis of genomic DNA isolated from tail biopsies.

Animals and estradiol-induced LH surge protocol

All animal experimental procedures were approved by the University of Otago Animal Ethics Committee and the Regierungspräsidium Karlsruhe. Animals were housed in groups of 3-4/cage with food and water freely available and a 12:12 lighting schedule (lights on 07:00h, off 19:00h). Genotyping of ER α mutant and ER β mutant mice was undertaken as described previously (Abraham et al., 2003). Adult (>60 days of age) wild-type and ER mutant female C57BL6/J mice were anesthetized with Halothane, ovariectomized and an estradiol-filled Silastic capsule implanted subcutaneously. Estradiol capsules were made according to the protocol of Bronson (Bronson, 1981), and involved filling Silastic tubing (1.0 mm internal, 2.1mm external diameter; Dow Corning, Michigan) with Silastic medical-grade adhesive (Dow Corning) containing 0.1mg 17- β -estradiol (Sigma, Missouri)/ml adhesive. Each mouse was given an ~ 1cm length of Silastic tubing (1 μ g estradiol/20g body weight). To prevent infection in brain-specific ER α knockout mice (see below), ER $\alpha^{fl/fl};CamKiCre$ mice and controls were given daily s.c. injections of 5 μ l/g Baytril 0.1 % (Bayer, Leverkusen, Germany) for five consecutive days, starting two days pre-ovariectomy (OVX). Six days after OVX, mice received a subcutaneous injection of estradiol benzoate (1 μ g/20g body weight, Intervet, Castle Hill, Australia) at 09:00h. On the following day, animals were killed with an i.p. overdose of pentobarbitone and trunk blood collected between 17:00 and 21:00h for LH radioimmunoassay. The sensitivity of the radioimmunoassay was 0.4 ng/ml and had an intra-assay coefficient of variation of 12.1%.

Five separate LH surge experiments were undertaken. In the first, wild-type mice were treated as above and groups of 4-5 mice killed at 17:00h, 18:00h, 19:00h, 20:00h and 21:00 to ascertain the profile of the LH surge. As this showed that 19:00h represented to time of peak LH secretion, further experiments were undertaken by killing mice at 19:00h. In the second experiment, ER α mutant and wild-type littermates (N=7-8 each group) were treated as above, anesthetized with pentobarbitone, and a 0.3ml blood sample obtained from the right atrium before perfusion of the mouse through the left ventricle with 15ml 4% paraformaldehyde fixative solution. Brains were then removed and post-fixed for 90 min at room temperature in 4% paraformaldehyde before being placed in a 30% sucrose/Tris-buffered saline (TBS) solution overnight. The third experiment involved the same procedure but using ER β mutant mice and wild-type littermates (n=7 each group). The fourth experiment used wild-type adult female mice given the estrogen-replacement protocol but with half of the mice receiving the ER α -selective agonist 3,17-dihydroxy-19-nor-17 α -pregna-1,3,5 (10)-triene-21,16 α -lactone (16 α -LE2; 0.8 μ g/20g body weight, s.c.; kind gift of Schering AG, Berlin, Germany) instead of the estradiol benzoate injection. At this dose, 16 α -LE2 is highly selective for ER α *in vivo* (Hegele-Hartung et al., 2004). The fifth experiment involved neuron-specific ER α knockout mice (n=4; ER $\alpha^{fl/fl};CamKiCre$ genotype) and control littermates (n=4-5; ER $\alpha^{+/fl};CamKiCre$, ER $\alpha^{fl/fl}$ and ER $\alpha^{+/fl}$ genotypes) treated as above. To evaluate possible altered timing of positive feedback and the LH surge in these mice, animals were perfused at 13:00h, 19:00h and 01:00h with the

addition that pituitaries were collected alongside the brains. In each experiment, individual control and ER mutant mice were perfused in an alternate manner.

Immunohistochemistry

Single labeling ER α immunohistochemistry was performed using either the monoclonal rat H222 antibody (1.3 μ g/ml, gift of Abbott Laboratories, IL) for free-floating brain sections or a polyclonal rabbit antibody (1 μ g/ml, MC-20, SC-542, Santa Cruz Biotechnology, CA) for the pituitary. Brains sections were cut in the coronal plane at 30 μ m on a freezing microtome and processed as free-floating sections using biotinylated anti-rat IgGs followed by the VECTASTAIN ABC system (Vector Laboratories) and nickel-diaminobenzidine as the chromagen. Fixed pituitaries were dehydrated, embedded in paraffin wax, and cut into 6- μ m-sections for slide-mounted immunohistochemistry using biotinylated anti-rabbit IgGs and the VECTASTAIN ABC system with diaminobenzidine.

Dual-labeling c-Fos and GnRH immunocytochemistry was undertaken as described previously (Herbison et al., 1995). Briefly, the rostral forebrain was cut into three sets of 30 μ m-thick coronal sections and one set immunostained for c-Fos (Li et al., 1999) using a polyclonal rabbit anti-cFos antibody (1:10,000, SC52, Santa Cruz Biotechnology Inc., Santa Cruz, CA) followed by biotinylated anti-rabbit IgGs (1:200; Vector, Burlingame, CA) and Vector Elite avidin-peroxidase (1:100), and revealed using nickel-diaminobenzidine hydrochloride. A second sequential staining using a rabbit anti-GnRH antibody (1:40,000; LR1, gift of R. Benoit, Montreal) and peroxidase-labelled anti-rabbit immunoglobulins (1:400; Vector Labs) revealed by diaminobenzidine hydrochloride alone was used to visualize the GnRH neurons. The removal of either primary antibodies resulted in a complete absence of the respective immunoreactivity.

Analysis was undertaken by counting the numbers of single-labelled (brown cytoplasm only) and dual-labelled (brown cytoplasm and black nucleus) GnRH neurons. The distribution and number of dual labelled GnRH neurons were counted in the regions of the medial septum, rostral preoptic area and anterior hypothalamic area represented by plates 22-24, 25-27 and 28-31 respectively, of the Franklin and Paxinos brain atlas (Franklin and Paxinos, 1997). In addition, the number of c-Fos-labelled cells identified within the AVPV, PeN and MePN were counted. Cell counts in the AVPV were determined by placing a right-angle triangle (160 μ m base x 400 μ m side) over the AVPV (plate 29) and counting all immunoreactive nuclei bilaterally on two sections from each mouse. The same procedure was undertaken for the MnPO by placing a square (300 μ m x 300 μ m) over the midline MnPO (plate 27) and counting the numbers of c-Fos-immunoreactive cells in 2 sections from each mouse. Counts in the PeN were undertaken in the same way by placing a rectangle (60 μ m x 800 μ m) next to the third ventricle and the level of plate 31. Values for each mouse were used to determine mean counts and these used to generate means+SEM values for each group. All data were analysed by non-parametric Mann-Whitney U tests with a $p < 0.05$ considered significant.

Conditional Pseudorabies virus tracing

The conditional PRV strain, Ba2001, was generated as described previously (DeFalco et al., 2001) (3.8×10^8 pfu/ml) and stored at -80C until use. Intracerebral stereotaxic injections of

Ba2001 were made in Avertin-anaesthetized adult female GnRH-Cre mice. A Hamilton syringe was used to deliver 500nl of Ba2001 into the rostral preoptic area (coordinates: 0.5mm bregma, 0mm lateral, -5.1mm dorsal-ventral) at a rate of 20nl/min. The needle was left in place for 5 minutes following injection. Control injections were made into the preoptic area of wild-type mice and into the striatum of GnRH-Cre mice. Mice were killed at various time points after injection and brain tissue was prepared for immunohistochemical processing as described above. Thirty micrometer thick sections underwent dual-labelling for either GFP (to detect PRV infected cells) and GnRH, or GFP and ER α . The following antibodies were used: chicken anti-GFP (1:2,500; Chemicon), rabbit anti-GFP (1:5,000, Molecular Probes), rabbit anti-GnRH antibody (1:40,000; LR1, gift of R. Benoit, Montreal), and rabbit anti-ER α (1:10,000, Upstate). GFP/GnRH dual labelling was visualized using anti-rabbit IgGs followed by streptavidin 568 (1:200; Molecular Probes) and anti-chicken FITC (1:200; Jackson Immunolabs). GFP/ER α dual labelling was visualized with biotinylated anti-rabbit IgGs followed by streptavidin 568 (1:200; Molecular Probes), and anti-chicken FITC (1:200; Jackson Immunolabs). Every section throughout the brain of 72h post-injection female mice (n=6) was analysed for GFP/GnRH dual labelling using an Olympus BX51 epifluorescence microscope. Dual labelling for GFP/ER α was analysed in every third section throughout the forebrain (n=5). GFP/ER α /Nissl staining was undertaken by including a final incubation step in Neurotrace 435/455 blue fluorescent Nissl stain (1:100, Molecular Probes).

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgements

This research was supported by the Wellcome Trust, Royal Society of New Zealand Marsden Fund, “Deutsche Forschungsgemeinschaft” through Collaborative Research Centres 405, 488 and 636., FOR 302, GRK 791/1.02, GRK 484, and Sachbeihilfe Schu 51/7-2, by the “Fonds der Chemischen Industrie”, the European Community through grant QLG1-CT-2001-01574, the Bundesministerium für Bildung und Forschung (BMBF) through NGFN grants FZK 01GS01117 and KGCV1/01GS0416, German-Polish cooperation project 01GZ0310 and project number 0313074C (systems biology), and by the Alexander von Humboldt-Stiftung through the Max-Planck-Forschungspreis für Internationale Kooperation, and by the Division of Intramural Research, NIEHS/NIH. CAP is a Research Fellow of the Sjogren’s Syndrome Foundation. Joachim Elzer is thanked for help with generation of the ER α mutant mice, and Katja Prella, Schering AG Berlin, for support of the 16aLE2 study.

References

- Abraham IM, Han K, Todman MG, Korach KS, Herbison AE. Estrogen receptor β mediates rapid estrogen actions on gonadotropin-releasing hormone neurons *in vivo*. *J Neurosci*. 2003; 23:5771–5777. [PubMed: 12843281]
- Bronson FH. The regulation of luteinizing hormone secretion by estrogen: relationships among negative feedback, surge potential, and male stimulation in juvenile, peripubertal, and adult female mice. *Endocrinology*. 1981; 108:506–516. [PubMed: 7449740]
- Burgin KE, Waxham MN, Rickling S, Westgate SA, Mobley WC, Kelly PT. In situ hybridization histochemistry of Ca²⁺/calmodulin-dependent protein kinase in developing rat brain. *J Neurosci*. 1990; 10:1788–1798. [PubMed: 2162385]
- Card JP, Enquist LW. Neurovirulence of pseudorabies virus. *Crit Rev Neurobiol*. 1995; 9:137–162. [PubMed: 8581980]

- Casanova E, Fehsenfeld S, Mantamadiotis T, Lemberger T, Greiner E, Stewart AF, Schutz G. A CamKIIalpha iCre BAC allows brain-specific gene inactivation. *Genesis*. 2001; 31:37–42. [PubMed: 11668676]
- Couse JF, Bunch DO, Lindzey J, Schomberg DW, Korach KS. Prevention of the polycystic ovarian phenotype and characterization of ovulatory capacity in the estrogen receptor-alpha knockout mouse. *Endocrinology*. 1999; 140:5855–5865. [PubMed: 10579351]
- Couse JF, Korach KS. Estrogen receptor null mice: what have we learned and where will they lead us? *Endocr Rev*. 1999; 20:358–417. [PubMed: 10368776]
- DeFalco J, Tomishima M, Liu H, Zhao C, Cai X, Marth JD, Enquist L, Friedman JM. Virus-assisted mapping of neural inputs to a feeding center in the hypothalamus. *Science*. 2001; 291:2608–2613. [PubMed: 11283374]
- Dupont S, Krust A, Gansmuller A, Dierich A, Chambon P, Mark M. Effect of single and compound knockouts of estrogen receptors alpha (ERalpha) and beta (ERbeta) on mouse reproductive phenotypes. *Development*. 2000; 127:4277–4291. [PubMed: 10976058]
- Evans NP, Dahl GE, Padmanabhan V, Thrun LA, Karsch FJ. Estradiol requirements for induction and maintenance of the gonadotropin-releasing hormone surge: implications for neuroendocrine processing of the estradiol signal. *Endocrinology*. 1997; 138:5408–5414. [PubMed: 9389526]
- Franklin KBJ, Paxinos G. *The Mouse Brain in Stereotaxic Coordinates*. San Diego: Academic Press; 1997.
- Gu GB, Simerly RB. Projections of the sexually dimorphic anteroventral periventricular nucleus in the female rat. *J Comp Neurol*. 1997; 384:142–164. [PubMed: 9214545]
- Hegele-Hartung C, Siebel P, Peters O, Kosemund D, Muller G, Hillisch A, Walter A, Kraetzschmar J, Fritzscheier KH. Impact of isotype-selective estrogen receptor agonists on ovarian function. *Proc Natl Acad Sci U S A*. 2004; 101:5129–5134. [PubMed: 15037755]
- Herbison AE. Multimodal influence of estrogen upon gonadotropin-releasing hormone neurons. *Endocr Rev*. 1998; 19:302–330. [PubMed: 9626556]
- Herbison AE. *Physiology of the GnRH neuronal network* Knobil and Neill's *Physiology of Reproduction*. Neill JD, editor San Diego: Academic Press; 2006. 1415–1482.
- Herbison AE, King IS, Tan KKC, Dye S. Increased fos expression in preoptic calcitonin gene-related peptide (CGRP) neurones following mating but not the luteinizing hormone surge in female rats. *J Neuroendocrinol*. 1995; 7:377–385. [PubMed: 7550284]
- Herbison AE, Pape JR. New evidence for estrogen receptors in gonadotropin-releasing hormone neurons. *Front Neuroendocrinol*. 2001; 22:292–308. [PubMed: 11587554]
- Hoffman GE, Smith MS, Verbalis JG. c-Fos and related immediate early gene products as markers of activity in neuroendocrine systems. *Front Neuroendocrinol*. 1993; 14:173–213. [PubMed: 8349003]
- Horvath S, Kis Z, Boldogkoi Z, Nogradi A, Toldi J. Oestrogen-dependent tracing in the rat CNS after pseudorabies virus infection. *Eur J Neurosci*. 2002; 15:937–943. [PubMed: 11906536]
- Horvath TL, Naftolin F, Leranath C. Luteinizing hormone-releasing hormone and gamma-aminobutyric acid neurons in the medial preoptic area are synaptic targets of dopamine axons originating in anterior periventricular areas. *Journal of Neuroendocrinology*. 1993; 5:71–79. [PubMed: 8097941]
- Hrabovszky E, Steinhäuser A, Barabás K, Shughrue PJ, Petersen SL, Merchenthaler I, Liposits Z. Estrogen receptor- β immunoreactivity in luteinizing hormone-releasing hormone neurons of the rat brain. *Endocrinology*. 2001; 142:3261–3264. [PubMed: 11416051]
- Karsch FJ, Bowen JM, Caraty A, Evans NP, Moenter SM. Gonadotropin-releasing hormone requirements for ovulation. *Biol Reprod*. 1997; 56:303–309. [PubMed: 9116125]
- Le WW, Berghorn KA, Rassnick S, Hoffman GE. Periventricular preoptic area neurons coactivated with luteinizing hormone (LH)-releasing hormone (LHRH) neurons at the time of the LH surge are LHRH afferents. *Endocrinology*. 1999; 140:510–519. [PubMed: 9886864]
- Lee WS, Smith MS, Hoffman GE. Luteinizing hormone-releasing hormone neurons express Fos protein during the proestrous surge of luteinizing hormone. *Proc Natl Acad Sci U S A*. 1990; 87:5163–5167. [PubMed: 2114646]
- Legan SJ, Coon GA, Karsch FJ. Role of estrogen as initiator of daily LH surges in the ovariectomized rat. *Endocrinology*. 1975; 96:50–56. [PubMed: 1109905]

- Levine JE. New concepts of the neuroendocrine regulation of gonadotropin surges in rats. *Biology of Reproduction*. 1997; 56:293–302. [PubMed: 9116124]
- Li C, Chen P, Smith MS. Identification of neuronal input to the arcuate nucleus (ARH) activated during lactation: implications in the activation of neuropeptide Y neurons. *Brain Res*. 1999; 824:267–276. [PubMed: 10196458]
- Marsicano G, Goodenough S, Monory K, Hermann H, Eder M, Cannich A, Azad SC, Cascio MG, Gutierrez SO, van der Stelt M, et al. CB1 cannabinoid receptors and on-demand defense against excitotoxicity. *Science*. 2003; 302:84–88. [PubMed: 14526074]
- McEwen BS, Alves SE. Estrogen actions in the central nervous system. *Endocr Revs*. 1999; 20:279–307. [PubMed: 10368772]
- Quimet CC, McGuinness TL, Greengard P. Immunocytochemical localization of calcium/calmodulin-dependent protein kinase II in rat brain. *Proc Natl Acad Sci U S A*. 1984; 81:5604–5608. [PubMed: 6591208]
- Petersen SL, Barraclough CA. Suppression of spontaneous LH surges in estrogen-treated ovariectomized rats by microimplants of antiestrogens into the preoptic brain. *Brain Research*. 1989; 484:279–289. [PubMed: 2713688]
- Petersen SL, Ottem EN, Carpenter CD. Direct and indirect regulation of gonadotropin-releasing hormone neurons by estradiol. *Biol Reprod*. 2003; 69:1771–1778. [PubMed: 12890720]
- Prevot V. Glial-neuronal-endothelial interactions are involved in the control of GnRH secretion. *J Neuroendocrinol*. 2002; 14:247–255. [PubMed: 11999726]
- Rage F, Lee BJ, Ma YJ, Ojeda SR. Estradiol enhances prostaglandin E2 receptor gene expression in luteinizing hormone-releasing hormone (LHRH) neurons and facilitates the LHRH response to PGE2 by activating a glia-to-neuron signaling pathway. *J Neurosci*. 1997; 17:9145–9156. [PubMed: 9364061]
- Schaft J, Ashery-Padan R, van der Hoeven F, Gruss P, Stewart AF. Efficient FLP recombination in mouse ES cells and oocytes. *Genesis*. 2001; 31:6–10. [PubMed: 11668672]
- Shughrue PJ, Lane MV, Merchenthaler I. Regulation of progesterone receptor messenger ribonucleic acid in the rat medial preoptic nucleus by estrogenic and antiestrogenic compounds: an in situ hybridization study. *Endocrinology*. 1997; 138:5476–84. [PubMed: 9389534]
- Shupnik MA. Gonadal hormone feedback on pituitary gonadotropin genes. *Trends in Endocrinology and Metabolism*. 1996; 7:272–276. [PubMed: 18406759]
- Simerly RB. Wired for reproduction: organization and development of sexually dimorphic circuits in the mammalian forebrain. *Annu Rev Neurosci*. 2002; 25:507–536. [PubMed: 12052919]
- Simonian SX, Spratt DP, Herbison AE. Identification and characterization of estrogen receptor α -containing neurons projecting to the vicinity of the gonadotropin-releasing hormone perikarya in the rostral preoptic area of the rat. *J Comp Neurol*. 1999; 411:346–358. [PubMed: 10404258]
- Skinner DC, Dufourny L. Oestrogen receptor beta-immunoreactive neurones in the ovine hypothalamus: distribution and colocalisation with gonadotropin-releasing hormone. *J Neuroendocrinol*. 2005; 17:29–39. [PubMed: 15720473]
- Skytner MJ, Sim JS, Herbison AE. Detection of estrogen receptor α and β messenger ribonucleic acids in adult gonadotropin-releasing hormone neurons. *Endocrinology*. 1999a; 140:5195–5201. [PubMed: 10537149]
- Skytner MJ, Slater R, Sim JA, Allen ND, Herbison AE. Promoter transgenics reveal multiple gonadotropin-releasing hormone-1-expressing cell populations of different embryological origin in mouse brain. *J Neurosci*. 1999b; 19:5955–5966. [PubMed: 10407034]
- Smith MJ, Jennes L. Neural signals that regulate GnRH neurones directly during the oestrous cycle. *Reproduction*. 2001; 122:1–10. [PubMed: 11425324]
- Soriano P. Generalized lacZ expression with the ROSA26 Cre reporter strain. *Nat Genet*. 1999; 21:70–71. [PubMed: 9916792]
- Tronche F, Kellendonk C, Kretz O, Gass P, Anlag K, Orban PC, Bock R, Klein R, Schutz G. Disruption of the glucocorticoid receptor gene in the nervous system results in reduced anxiety. *Nat Genet*. 1999; 23:99–103. [PubMed: 10471508]
- Vente A, Korn B, Zehetner G, Poustka A, Lehrach H. Distribution and early development of microarray technology in Europe. *Nat Genet*. 1999; 22:22. [PubMed: 10319856]

- Wiegand SJ, Terasawa E. Discrete lesions reveal functional heterogeneity of suprachiasmatic structures in regulation of gonadotropin secretion in the female rat. *Neuroendocrinology*. 1982; 34:395–404. [PubMed: 6808412]
- Xia L, Van Vugt D, Alston EJ, Luckhaus J, Ferin M. A surge of gonadotropin-releasing hormone accompanies the estradiol-induced gonadotropin surge in the rhesus monkey. *Endocrinology*. 1992; 131:2812–2820. [PubMed: 1446619]
- Yoon H, Enquist LW, Dulac C. Olfactory inputs to hypothalamic neurons controlling reproduction and fertility. *Cell*. 2005; 123:669–682. [PubMed: 16290037]
- Zhang Y, Buchholz F, Muyrers JP, Stewart AF. A new logic for DNA engineering using recombination in *Escherichia coli*. *Nat Genet*. 1998; 20:123–128. [PubMed: 9771703]




Figure 1. Absence of estrogen positive feedback in ER α mutant mice.

A. Depicts the profile of the LH surge in wild-type ovariectomized mice treated with estrogen. $n=4-5$ at each time point. **B.** Mean (\pm SEM) LH levels in wildtype-littermates ($n=7$) and ER α mutant mice ($n=8$), ovariectomized, treated with estrogen and killed at 19:00h, * $p<0.05$. **C-E,** Dual-labelling c-Fos (black nuclei) and GnRH (brown cytoplasmic staining) immunocytochemistry in wild-type (C) and ER α mutant (D) mice. Whereas approximately 40% of GnRH neurons express c-Fos (C, arrowheads and E) in wild-type mice, no GnRH neurons express c-Fos in ER α mutant mice (D). **F-H,** Low-power photomicrographs

showing the location of many c-Fos-expressing cells within the AVPV of wild-type mice (F) compared with few in ER α mutant mice (G). The mean (+SEM) number of c-Fos cells per unit area in the AVPV is shown in H. $p < 0.05$. Scale bars represent 30 μ m in C and 100 μ m in F.




Figure 2. Positive feedback actions of estrogen appear normal in ER β mutant mice.

A. Mean (+SEM) LH levels in wildtype-littermates (n=7) and ER β mutant mice (n=7), ovariectomized, treated with estrogen and killed at 19:00h. **B.** The percentage of GnRH neurons (+SEM) found to express c-Fos in wildtype-littermates (n=7) and ER α mutant mice (n=8), ovariectomized, treated with estrogen and killed at 19:00h. **C.** The number of c-Fos-expressing neurons detected within the AVPV of wildtype-littermates (n=7) and ER α mutant mice (n=8), ovariectomized, treated with estrogen and killed at 19:00h.




Figure 3. Absence of estrogen positive feedback in neuron-specific ER α mutant mice. **A,B.** show ER α expression in the rostral hypothalamus within the anteroventral periventricular nucleus (AVPV) of **(A)** control (ER $\alpha^{fl/fl}$) and **(B)** neuron-specific ER α mutant (ER $\alpha^{fl/fl};CamKiCre$) mice. Scale bar is 100 μ m, 3V= third ventricle, OC = optic chiasm. **Insets** show equivalent ER α staining in the anterior pituitary of **(A)** control and **(B)** neuron-specific ER α mutant mice. **C.** Mean (+SEM) LH levels in control-littermates (n=5) and ER $\alpha^{fl/fl};CamKiCre$ mice (n=4), ovariectomized, treated with estrogen and killed at 19:00h. * p<0.05 **D.** Approximately 40% of GnRH neurons express c-Fos in control mice ovariectomized, treated with estrogen and killed at 19:00h whereas none are found in ER $\alpha^{fl/fl};CamKiCre$ mice. **E.** The number of c-Fos-expressing neurons detected within the

AVPV of controls and $ER\alpha^{fl/fl};CamKiCre$ mice, ovariectomized, treated with estrogen and killed at 19:00h. * $p < 0.05$.




Figure 4. Ovarian and uterine phenotype of neuron-specific ER α mutant mice.

A,B. reproductive tract of control ER $\alpha^{fl/fl}$ (**A**) and mutant ER $\alpha^{fl/fl};CamKiCre$ (**B**) mice showing fluid-filled uteri in mutants. **C,D** H&E staining of uteri from control ER $\alpha^{fl/fl}$ (**C**) and mutant ER $\alpha^{fl/fl};CamKiCre$ (**D**) mice shows atrophy and lack of glandular structures in the mutant. **E,F.** H&E staining of ovaries from control ER $\alpha^{fl/fl}$ (**E**) and mutant ER $\alpha^{fl/fl};CamKiCre$ (**F**) mice shows increased numbers of antral follicles and lack of corpora lutea in the mutant.




Figure 5. Identification of ER α -expressing primary afferents to GnRH neurons using GnRH-specific viral mediated tracing.

A. Restricted expression of Cre (black nuclei) in GnRH neurons (brown cytoplasm) of transgenic GnRH-Cre mice. **B,C.** A rostral preoptic area GnRH neuron adjacent to the injection site (arrowheads) exhibiting GFP immunoreactivity 48h following injection of Ba2001. B shows GFP immunoreactivity and C is GnRH immunostaining. **D.** Four neurons (arrows) in the AVPV exhibiting GFP immunoreactivity 72h after Ba2001 injection into the rostral preoptic area. **E.** Dual-labelling for GFP (green) and ER α (red) reveals that 2 of these cells (arrows) express ER α . **F.** Schematic brain maps demonstrating GFP-immunoreactive neurons (open stars) and GFP+ER α immunoreactive (filled stars) cells detected in individual 30 μ m-thick coronal brain sections at three levels in the hypothalamus. Abbreviations, aca, anterior commissure; ARN, arcuate nucleus; AVPV, anteroventral periventricular nucleus; f, fornix; MnPO, median preoptic nucleus; ME, median eminence, PeN, periventricular nucleus; 3V, third ventricle. Scale bar is 15 μ m in A,C,D. **G.** Summary of locations of primary afferents to GnRH neurons following stereotaxic injection of Ba2001 PRV into the rPOA of GnRH-Cre mice. Various colors in boxes adjacent to listed brain areas indicate ranges of mean numbers of GFP-positive neurons identified in female mice (n=6). Brain regions in which labelled cells were detected in <5 of the 6 mice are indicated as yellow boxes (variable). Right-most column indicates the mean percentage (\pm SEM) of GFP-positive neurons expressing ER α immunoreactivity. Abbreviations; ARN, arcuate nucleus; AVPV, anteroventral periventricular nucleus; DBB, diagonal band of Broca; DMH, dorsomedial hypothalamus; DR, dorsal raphe; LC, locus coereulus; LH, lateral

hypothalamus; LS, lateral septum; MnPO, median preoptic nucleus; MPN, medial preoptic nucleus; MS, medial septum; NTS, nucleus tractus solitarius; PAG, periaqueductal grey; PeN, periventricular nucleus; PVN, paraventricular nucleus; SFO, subfornical organ; VMN, ventromedial nucleus.