

## Invited Mini Review

BLT2, a leukotriene B<sub>4</sub> receptor 2, as a novel prognostic biomarker of triple-negative breast cancerJaeIn Park<sup>1,#</sup>, Jae-Hyun Jang<sup>1,#</sup>, Geun-Soo Park<sup>1</sup>, Yunro Chung<sup>2</sup>, Hye Jin You<sup>3,\*</sup> & Jae-Hong Kim<sup>1,\*</sup><sup>1</sup>Department of Biotechnology, College of Life Sciences and Biotechnology, Korea University, Seoul 02841, Korea, <sup>2</sup>Department of Biomedical Informatics, Arizona State University, Scottsdale, AZ 85259, USA, <sup>3</sup>Translational Research Branch, Division of Translational Science, National Cancer Center, Goyang 10408, Korea

Triple-negative breast cancer (TNBC) is considered to be a notorious type of cancer due to its aggressive metastatic potential and poor prognosis. Recent evidence suggests that BLT2, a low-affinity LTB<sub>4</sub> receptor is critically associated with the phenotypes of TNBC cells, including invasion, metastasis, and survival. Furthermore, in a group of 545 breast cancer patients with metastasis, we observed that the high-BLT2 subgroup had a lower disease-free-survival rate than the low-BLT2 subgroup. Thus, we theorized that anti-BLT2 strategies could facilitate the development of new therapies used for TNBC. This review focuses on recent discoveries regarding BLT2 and its roles in as a novel prognostic biomarker in TNBC. [BMB Reports 2018; 51(8): 373-377]

## INTRODUCTION

Breast cancer is the most common cancer diagnosis in women and the second leading cause of cancer-related death in women worldwide. Several different subtypes of breast cancers have been identified. Triple-negative breast cancer (TNBC), being among the most devastating types of breast cancer, is a disease with high morbidity and mortality rates. TNBC patients account for only 15-20% of all breast cancer patients (1), and patients with TNBC have an increased likelihood of distant recurrence and death compared to patients with other types of breast cancer. TNBC patients also tend to develop visceral metastases early during their disease. A recent study reported the five-year survival rate for TNBC as

approximately 30% lower than other types of breast cancer (2). TNBC is very heterogeneous and lacks expression of the estrogen receptor (ER), progesterone receptor (PR), and epidermal growth factor receptor 2 (HER2). The absence of ER/PR/HER2 reduces the therapeutic effects of many targeted anti-cancer drugs, cytotoxic drugs, and radiation; as a result, patients with TNBC are only managed with standard chemotherapy (3). Therefore, the identification of new therapeutic targets for TNBC is urgently needed.

Leukotriene B<sub>4</sub> (LTB<sub>4</sub>) is synthesized from arachidonic acid via the 5-lipoxygenase pathway, and it mediates intracellular functions via G protein-coupled receptors (GPCRs): LTB<sub>4</sub> receptor 1 (BLT1) and LTB<sub>4</sub> receptor 2 (BLT2) (4). BLT2 is a low-affinity LTB<sub>4</sub> receptor that can be stimulated through various ligands, such as 12(S)-hydroxy-5(Z),8(Z),10(E),14(Z)-eicosatetraenoic acid (12(S)-HETE), and 12(S)-hydroxy-5(Z),8(E),10(E)-heptadecatrienoic acid (12-HHT), as well as LTB<sub>4</sub> (5). Recent studies have suggested that BLT2 is closely associated with various aspects of survival, invasion, and metastasis in TNBC (6). This review focuses on the recently discovered roles of BLT2 in TNBC progression.

## ROLE OF BLT2 IN THE SURVIVAL AND PROLIFERATION OF TNBC CELLS

TNBC cells are known to be insensitive to most specific target treatments, including targeted small-molecule drugs (7). Therefore, conventional chemotherapeutic agents, such as Taxol and anthracyclines, are still the preferred treatments for TNBC patients. However, these drugs are not targeted and hair loss, diarrhea, nausea, and vomiting are among their many undesirable side effects. Many TNBC studies focus on survival and proliferation mechanisms in order to identify more effective therapeutic treatments (8). One of the mechanisms for TNBC progression is the generation of reactive oxygen species (ROS). ROS are tumorigenic by virtue of their ability to increase cell proliferation, survival, and DNA damage induction, leading to genetic lesions that cause tumorigenicity and sustain subsequent tumor progression (9, 10). Due to their stimulating actions on TNBC proliferation, up-regulation of ROS generation plays an important role in aggressive cancer

\*Corresponding authors. Jae-Hong Kim, Tel: +82-2-3290-3452; Fax: +82-31-920-2232; E-mail: jhongkim@korea.ac.kr; Hye Jin You, Tel: +82-31-920-2206; Fax: +82-31-920-2232; E-mail: hjyou@ncc.re.kr

<sup>#</sup>These authors contributed equally to this work.

<https://doi.org/10.5483/BMBRep.2018.51.8.127>

Received 20 May 2018

**Keywords:** BLT2, Leukotriene B<sub>4</sub> receptor-2, TNBC, Triple-negative breast cancer

phenotypes (11). In TNBC, tumor-associated ROS are produced from nicotinamide adenine dinucleotide phosphate-oxidase (NOX) complexes. Recent studies have showed that the NOX1 cascade attenuates the transcriptional activity of tumor suppressor and tumor cell apoptosis processes (12). Our previous study showed that BLT2 is critically associated with the generation of ROS via NOX1, supporting the proliferation and survival of human TNBC MDA-MB-468 cells (13). In summary, BLT2 is suggested to play critical role in TNBC survival and proliferation (Fig. 1).

### ROLE OF BLT2 IN THE INTERACTION BETWEEN TNBC CELLS AND IMMUNE CELLS


The interactions between cancer and immune cells have been suggested to be important components of tumor and TNBC progression (14, 15). Cell adhesion molecules are involved in cell to cell interaction, which is an especially important component in the cancer-immune system (16, 17). Intercellular adhesion molecule 1 (ICAM-1) is a member of the immunoglobulin superfamily and is expressed on many cell types, such as leukocytes, endothelial cells, and cancer cells (18, 19). In these cell types, ICAM-1 can be up-regulated in response to pro-inflammatory cytokines or stimuli (20). Moreover, studies have shown that ICAM-1 promotes TNBC progression,

aggressiveness, and shorter recurrence-free survival (21-24). Our recent study demonstrated that BLT2 regulates LPS-induced ICAM-1 expression in MDA-MB-231 cells via BLT2-ERK-NF- $\kappa$ B-linked cascades (Fig. 1). Therefore, up-regulation of BLT2 in TNBC potentially increases TNBC adhesion to monocytes (25). Our findings suggest that BLT2 can influence the microenvironment of TNBC, suggesting another critical role for BLT2 in TNBC progression.

### ROLE OF BLT2 IN THE INVASION AND METASTASIS OF TNBC CELLS

The metastatic phenotype is a major cause of death in cancer patients. Various mechanisms are involved in the metastasis of TNBC, including the acquisition of invasive characteristics in the tumor microenvironment (26, 27). Moreover, this particular cancer type shows enhanced metastatic potential. Patients with TNBC are known to have an increased risk of metastasis and distant recurrence compared to patients with other types of breast cancer (28). Given that metastasis can significantly contribute to cancer severity, finding new therapeutic targets for these types of metastatic, aggressive breast cancer is critical for treating patients.

Recently, pro-inflammatory cytokine interleukins (IL)-6 and IL-8 have been suggested to play roles in the invasion and metastasis of breast cancer (29). IL-6 has reportedly been correlated with TNBC progression, and clinical studies have shown increased IL-8 levels in aggressive breast tumor tissue when compared with normal breast tissue (30, 31). Due to the importance of pro-inflammatory cytokines in TNBC invasion, in-depth study is needed to validate a better target for inflammatory signaling. Our recent studies have shown that IL-6 and IL-8 are produced through a BLT2-dependent pathway (6, 32) (Fig. 1). Kim et al. found that BLT2 expression was increased in human TNBC cell lines, human TNBC MDA-MB-231, and MDA-MB-435 cells when compared with its expression in a human breast cancer cell line, MCF-10A. Therefore, we examined whether BLT2 can reduce the invasiveness and secretion of IL-8. MDA-MB-231 and MDA-MB-435 cells were treated with a specific BLT2 antagonist LY255283 or transfected with BLT2 small interfering RNA (siRNA). In these experiments, we demonstrated that IL-8 secretion, invasiveness in TNBC cells, and lung invasion in a TNBC xenograft mouse model are regulated in a BLT2-dependent manner. Park et al. also demonstrated that IL-6/IL-8 is downstream of the BLT2-dependent pathway in LPS-treated TNBC cell lines and an MDA-MB-231 xenograft mouse model. The activity of NF- $\kappa$ B is up-regulated in TNBC, and the NF- $\kappa$ B signaling pathway plays a crucial role in regulating the invasiveness of TNBC. In TNBC, NF- $\kappa$ B signaling cascades, which are related to invasion, are regulated by the BLT2-NOX-1 pathway, leading to the production of IL-6 and IL-8 in TNBC (Fig. 1). Together, these observations suggest that BLT2 plays a critical role in TNBC cell invasion and metastasis




**Fig. 1.** Comprehensive overview of the BLT2 signaling pathways. See text for details.



via IL-6/IL-8 up-regulation.

### SIGNIFICANCE OF BLT2 IN LYMPH NODE-POSITIVE BREAST CANCER PATIENTS AND HER2-NEGATIVE BREAST CANCER PATIENTS

We analyzed BLT2 with *Breastmark* by dichotomizing gene expression data around a media into 'high' and 'low' expression, and with 50% of patient samples with lymph node metastasis in each group, in order to investigate whether BLT2 contributes to breast cancer patient survival (33, 34). Fig. 2 shows Kaplan-Meier curves for the high- and low-BLT2 subgroups; the high-BLT2 subgroup exhibited a lower disease-free-survival (DFS) rate than the low-BLT2 subgroup.



**Fig. 2.** The Kaplan-Meier curves of the overall survival rates of patients with lymph node metastasis stratified by BLT2 expression (n = 545; HR = 1.341; 95% CI = (1.036, 1.735), P = 0.03).



**Fig. 3.** Overall survival of breast cancer patients with and without up-regulated BLT2 expression at the mRNA level and a pie chart of the patients' HER2 status. Absolute patient numbers for each subgroup are indicated.

Additionally, we fit a univariate COX proportional hazards model in which those with higher BLT2 expression are 1.34-fold more likely to have disease progression than those with lower BLT2 expression (Hazard ratio (HR) = 1.34; 95% CI = (1.036, 1.735); P = 0.025584). Patients with a basal subtype in the high-BLT2 expression group also had significantly reduced DFS (data not shown). These results suggest that BLT2 plays a significant role in the lymph node metastasis of breast cancer patients. Another dataset showed the effect of BLT2 up-regulation, at the mRNA level, on overall survival rates (Fig. 3). In the Metabric dataset (StudyID brca\_metabric) from the cBioportal for Cancer Genomics (35), the overall survival of 56 cases with up-regulated BLT2, at mRNA level, was significantly lower than in the remaining whole sequenced cases (Fig. 3) (P = 0.041). Furthermore, 52 of 56 cases with up-regulated BLT2 were identified as HER2-negative (Fig. 3). In general, most patients with metastatic breast cancer have HER2-negative breast cancer (Reference PMID: 28942029 DOI: 10.1016/j.ctrv.2017.09.001), implying that BLT2 plays a role in the breast cancer malignancy of a subset of patients with metastasis and HER2 negativity. These results suggest that BLT2 is associated with malignancy in breast cancer, such as in TNBC.

### CONCLUSION AND FUTURE PERSPECTIVES

In this review, we discussed the role of BLT2 in TNBC progression. TNBC is commonly associated with a poor prognosis, which is unpredictable and measured by considering standard clinicopathological values, such as lymph node status or size at initial presentation (36). The studies reviewed in this article demonstrate that these characteristics are likely to be mediated, at least in part, by BLT2. As shown in Fig. 1, BLT2 activation through its ligands activates a number of signaling pathways that support the malignant phenotypes of TNBC cells. Our findings reveal a previously unrecognized role BLT2 plays in the progression of TNBC. Our study enhances the

understanding of the molecular mechanism of TNBC and provides potential targets for developing new therapies to be used in the treatment of TNBC patients.

## ACKNOWLEDGEMENTS

This research was supported by Bio & Medical Technology Development Program Grants (2017M3A9D8063317) and a Mid-Career Researcher Program grant (2017R1A2B4002203) provided by the National Research (NRF) funded by the Ministry of Science, Information and Communication Technologies (ICT), and Future Planning, Republic of Korea. Additionally, this research was supported by a grant from the Korea Healthcare Technology R&D Project, Ministry of Health & Welfare, Republic of Korea (A101032), National Research Foundation of Korea Grant funded by the Korea Government (MSIP, South Korea) (No.2015R1A2A2A01003829) and was also supported by a Korea University grant and the BK21 Plus Program (School of Life Sciences and Biotechnology, Korea University).

## CONFLICTS OF INTEREST

The authors have no conflicting interests.

## REFERENCES

1. Anders CK, Winer EP, Ford JM et al (2010) Poly(ADP-Ribose) polymerase inhibition: "targeted" therapy for triple-negative breast cancer. *Clin Cancer Res* 16, 4702-4710
2. Carey L, Winer E, Viale G, Cameron D and Gianni L (2010) Triple-negative breast cancer: disease entity or title of convenience? *Nat Rev Clin Oncol* 7, 683-692
3. Kassam F, Enright K, Dent R et al (2009) Survival outcomes for patients with metastatic triple-negative breast cancer: implications for clinical practice and trial design. *Clin Breast Cancer* 9, 29-33
4. Dinarello CA (2006) The paradox of pro-inflammatory cytokines in cancer. *Cancer Metastasis Rev* 25, 307-313
5. Tager AM and Luster AD (2003) BLT1 and BLT2: the leukotriene B(4) receptors. *Prostaglandins Leukot Essent Fatty Acids* 69, 123-134
6. Park GS and Kim JH (2015) Myeloid differentiation primary response gene 88-leukotriene B4 receptor 2 cascade mediates lipopolysaccharide-potentiated invasiveness of breast cancer cells. *Oncotarget* 6, 5749-5759
7. Maiello MR, D'Alessio A, Bevilacqua S, Gallo M, Normanno N and De Luca A (2015) EGFR and MEK Blockade in Triple Negative Breast Cancer Cells. *J Cell Biochem* 116, 2778-2785
8. Lonne GK, Masoumi KC, Lennartsson J and Larsson C (2009) Protein kinase Cdelta supports survival of MDA-MB-231 breast cancer cells by suppressing the ERK1/2 pathway. *J Biol Chem* 284, 33456-33465
9. Bogucki DJ and Domaradzki JA (2005) Numerical study of light scattering by a boundary-layer flow. *Appl Opt* 44, 5286-5291
10. Weng MS, Chang JH, Hung WY, Yang YC and Chien MH (2018) The interplay of reactive oxygen species and the epidermal growth factor receptor in tumor progression and drug resistance. *J Exp Clin Cancer Res* 37, 61
11. Pelicano H, Zhang W, Liu J et al (2014) Mitochondrial dysfunction in some triple-negative breast cancer cell lines: role of mTOR pathway and therapeutic potential. *Breast Cancer Res* 16, 434
12. Puca R, Nardinocchi L, Starace G et al (2010) Nox1 is involved in p53 deacetylation and suppression of its transcriptional activity and apoptosis. *Free Radic Biol Med* 48, 1338-1346
13. Choi JA, Lee JW, Kim H et al (2010) Pro-survival of estrogen receptor-negative breast cancer cells is regulated by a BLT2-reactive oxygen species-linked signaling pathway. *Carcinogenesis* 31, 543-551
14. Blot E, Chen W, Vasse M et al (2003) Cooperation between monocytes and breast cancer cells promotes factors involved in cancer aggressiveness. *Br J Cancer* 88, 1207-1212
15. Chittechath M, Dhillon MK, Lim JY et al (2014) Molecular profiling reveals a tumor-promoting phenotype of monocytes and macrophages in human cancer progression. *Immunity* 41, 815-829
16. Nakasone ES, Hurvitz SA and McCann KE (2018) Harnessing the immune system in the battle against breast cancer. *Drugs Context* 7, 212520
17. Brown EJ (1997) Adhesive interactions in the immune system. *Trends Cell Biol* 7, 289-295
18. Ohene-Abuakwa Y and Pignatelli M (2000) Adhesion molecules as diagnostic tools in tumor pathology. *Int J Surg Pathol* 8, 191-200
19. Zhong L, Simard MJ and Huot J (2018) Endothelial microRNAs regulating the NF-kappaB pathway and cell adhesion molecules during inflammation. *FASEB J*, fj201701536R
20. Hubbard AK and Rothlein R (2000) Intercellular adhesion molecule-1 (ICAM-1) expression and cell signaling cascades. *Free Radic Biol Med* 28, 1379-1386
21. Evani SJ, Prabhu RG, Gnanaruban V, Finol EA and Ramasubramanian AK (2013) Monocytes mediate metastatic breast tumor cell adhesion to endothelium under flow. *FASEB J* 27, 3017-3029
22. Rosette C, Roth RB, Oeth P et al (2005) Role of ICAM1 in invasion of human breast cancer cells. *Carcinogenesis* 26, 943-950
23. Strell C, Lang K, Niggemann B, Zaenker KS and Entschladen F (2010) Neutrophil granulocytes promote the migratory activity of MDA-MB-468 human breast carcinoma cells via ICAM-1. *Exp Cell Res* 316, 138-148
24. Schroder C, Witzel I, Muller V et al (2011) Prognostic value of intercellular adhesion molecule (ICAM)-1 expression in breast cancer. *J Cancer Res Clin Oncol* 137, 1193-1201
25. Park GS and Kim JH (2015) LPS Up-Regulates ICAM-1 expression in breast cancer cells by stimulating a MyD88-BLT2-ERK-Linked cascade, which promotes adhesion to monocytes. *Mol Cells* 38, 821-828
26. Neophytou C, Boutsikos P and Papageorgis P (2018)

- Molecular mechanisms and emerging therapeutic targets of triple-negative breast cancer metastasis. *Front Oncol* 8, 31
27. Christofori G (2006) New signals from the invasive front. *Nature* 441, 444-450
  28. Liang Z, Bian X and Shim H (2016) Downregulation of microRNA-206 promotes invasion and angiogenesis of triple negative breast cancer. *Biochem Biophys Res Commun* 477, 461-466
  29. Ma Y, Ren Y, Dai ZJ, Wu CJ, Ji YH and Xu J (2017) IL-6, IL-8 and TNF-alpha levels correlate with disease stage in breast cancer patients. *Adv Clin Exp Med* 26, 421-426
  30. Hartman ZC, Poage GM, den Hollander P et al (2013) Growth of triple-negative breast cancer cells relies upon coordinate autocrine expression of the proinflammatory cytokines IL-6 and IL-8. *Cancer Res* 73, 3470-3480
  31. De Larco JE, Wuertz BR, Rosner KA et al (2001) A potential role for interleukin-8 in the metastatic phenotype of breast carcinoma cells. *Am J Pathol* 158, 639-646
  32. Kim H, Choi JA, Park GS and Kim JH (2012) BLT2 up-regulates interleukin-8 production and promotes the invasiveness of breast cancer cells. *PLoS One* 7, e49186
  33. Madden SF, Clarke C, Gaule P et al (2013) BreastMark: an integrated approach to mining publicly available transcriptomic datasets relating to breast cancer outcome. *Breast Cancer Res* 15, R52
  34. Connolly C, Madden SF, Buggy DJ and Gallagher HC (2017) Expression of anaesthetic and analgesic drug target genes in excised breast tumour tissue: Association with clinical disease recurrence or metastasis. *PLoS One* 12, e0177105
  35. Gao J, Aksoy BA, Dogrusoz U et al (2013) Integrative analysis of complex cancer genomics and clinical profiles using the cBioPortal. *Sci Signal* 6, p11
  36. Lavigne AC, Castells M, Mermet J, Kocanova S, Dalvai M and Bystricky K (2014) Increased macroH2A1.1 expression correlates with poor survival of triple-negative breast cancer patients. *PLoS One* 9, e98930