

Regulation of Cyclin D1, DNA Topoisomerase I, and Proliferating Cell Nuclear Antigen Promoters During the Cell Cycle

HU-HUI LEE,* WEN-HSUI CHIANG,*† SHIAN-HUEY CHIANG,*‡ YIN-CHANG LIU,†
JAULANG HWANG,* AND SUN-YU NG*¹

**Institute of Molecular Biology, Academia Sinica, NanKang, Taipei 11529, Taiwan, Republic of China*

†*Institute of Life Science, National Tsing Hua University, Hsinchu 30043, Taiwan, Republic of China*

‡*Institute of Biochemistry, National Yang-Ming Medical College, Shih-Pai 11221,
Taiwan, Republic of China*

Cyclin D1, DNA topoisomerase I, and proliferating cell nuclear antigen (PCNA) are three important cell cycle regulatory proteins. Recently, their promoters have been isolated, thus facilitating molecular analysis of transcriptional control mechanisms of these genes. Transcription of these three promoters in stable K562 transfectants during different cell cycle phases was analyzed after cell cycle synchronization. About 1 kb of 5' flanking region from either cyclin D1 or DNA topoisomerase I gene is sufficient to confer G₁- or S-phase-specific transcription activity to chloramphenicol acetyltransferase (CAT) reporter genes, respectively. In contrast, 2.8 kb of 5' flanking sequences from the PCNA gene led to constitutive transcription, but the inclusion of a segment of the PCNA gene first intron, which contains evolutionarily conserved sequences, could enhance transcription in G₁/S-enriched nuclei. This PCNA intron region contains a binding site recognized by the transcription factor E2F. To test whether this site is functional, we cotransfected PCNA-CAT genes with E2F-1 and DP-1 expression plasmids. Expression of the E2F-1/DP-1 heterodimer activated the CAT gene with the PCNA intron. Therefore, this intron region, involved in transcriptional activation at the cell cycle G₁/S boundary, is also E2F inducible.

Cell cycle PCNA gene Transcription factor E2F Transcription regulation

THE cellular processes of DNA replication and cell division, carried out during the S- and M-phases of the cell cycle, involve a large number of proteins. Expression of those genes encoding many of these proteins is regulated in a cell cycle-dependent manner. Studies of molecular mechanisms involved in cell cycle transcriptional control have been limited by the small number of promoters that have so far been isolated. We investigated the transcriptional regulation of three important cell cycle proteins: cyclin D1, DNA topoisomerase I, and proliferating cell nuclear antigen (PCNA).

Cyclin D1 is one of the mammalian homologs of yeast G₁ cyclins (reviewed in Sherr, 1993). Although the cell cycle function(s) of this protein has not been well defined, it is believed to be important for cell cycle progression (Baldin et al., 1993; Quelle et al., 1993). Cyclin D1 protein interacts with cyclin-dependent kinases (Xiong et al., 1992b) and the retinoblastoma protein (Dowdy et al., 1993). Expression of mouse cyclin D1 is stimulated by colony-stimulating factor 1 (CSF-1) in macrophages (reviewed in Sherr, 1991). Overexpression of cyclin D1 is observed in many tumors,

Received June 8, 1994; revision accepted July 21, 1994.

¹Address correspondence to Sun-Yu Ng at his present address: Institute of Molecular Medical Sciences, 460 Page Mill Road, Palo Alto, CA 94306.

due to chromosomal translocation or gene amplification, thus linking it to the *bcl-1* proto-oncogene (Motokura and Arnold, 1993). Recently, the human cyclin D1 promoter has been isolated (Inaba et al., 1992; Xiong et al., 1992a; Herber et al., 1994). Here we show that a reporter gene under the control of this promoter is transcribed actively during G₁.

Both DNA topoisomerases are DNA replication enzymes, and are probably involved in the relaxation of supercoils generated during the replication process (Wang, 1991). In yeast, at least 17 replication enzyme genes, including a topoisomerase gene, are coordinately activated at START by the Mlu I cell cycle box (MCB) binding factor (MBF) complex (reviewed in Andrews and Herskowitz, 1990; Johnston, 1992). The existence of a similar subset of cell cycle-regulated mammalian genes is likely (McKinney and Heintz, 1991). The recent isolation of human type I and type II DNA topoisomerase promoters will facilitate such studies (Kunze et al., 1991; Hochhauser et al., 1992). In this article, we show that the DNA topoisomerase I promoter is transcriptionally most active during S-phase.

PCNA, as the DNA polymerase delta auxiliary protein, is also required for DNA replication (Bravo et al., 1987; Prelich and Stillman, 1988; Waga and Stillman, 1994). In addition, PCNA is associated with cyclin-dependent kinases (Zhang et al., 1993). The yeast PCNA gene is regulated by MBF at START (reviewed in Andrews and Herskowitz, 1990; Johnston, 1992). Previously, the mammalian PCNA gene was thought to be growth dependent but not cell cycle regulated (reviewed in Baserga, 1991). Recently, the human and rodent PCNA genes were isolated (Travali et al., 1989; Yamaguchi et al., 1991; Liu, 1992). We found evolutionarily conserved sequences in both the promoter regions and within the first intron. However, previous analyses of the PCNA promoter were focused on the 5' flanking region (Charollais et al., 1992; Labrie et al., 1993), with the exception of negative regions in intron 1 (Alder et al., 1992) and intron 4 (Ottavio et al., 1990). We constructed two PCNA reporter genes, both containing 2.8 kb of 5' flanking sequences, but only one with 413 bp of conserved intron sequences. We show that intron 1 sequences can enhance transcription in G₁/S-enriched nuclei. This conserved PCNA intron region contains a binding site recognized by the transcription factor E2F, which is known to regulate expression of S-phase genes at the G₁/S cell cycle transition (Nevins, 1992; Helin and Harlow, 1993; La Thangue,

1994). To investigate whether this E2F site is functional, we cotransfected E2F-1 (Helin et al., 1992; Kaelin et al., 1992; Shan et al., 1992) and DP-1 (Girling et al., 1993; Helin et al., 1993b) expression plasmids with the two PCNA reporter genes and found that only the PCNA intron reporter gene was activated.

MATERIALS AND METHODS

Plasmid Constructions

Cyclin D1-CAT plasmid was constructed from 1.3-kb *Pvu* II fragment of pD1-G0650 (Xiong et al., 1992a) inserted into the *Sma* I site of pUMSV-0CAT (Salier and Kurachi, 1989). DNA topoisomerase I-CAT plasmid was constructed from 990-bp *Eco*RI-*Xho*I fragment of pTOPOI (Kunze et al., 1991) inserted into the *Sma* I site of pUMSV-0CAT. PCNA-HN-CAT and PCNA-HH-CAT plasmids were constructed from 2.8-kb *Hinc* II-*Nru* I or 3.6-kb *Hinc* II fragment of p3BS2 (Travali et al., 1989) inserted into the *Sma* I site of pUMSV0CAT. A 2.6-kb *Xba* I-*Eag* I fragment was excised from PCNA-HH-CAT to yield PCNA-EH-CAT. A 320-bp *Hinc* II-*Nae* I fragment was inserted into PCNA-EH-CAT to produce PCNA-EN-CAT. A 74-bp *Eco* 47 III-*Apa* I fragment was excised from PCNA-EH-CAT to generate PCNA-EHΔ60/133-CAT. A 319-bp *Eco* 47 III-*Pfl* M I fragment was removed from PCNA-EH-CAT to generate PCNA-EHΔ60/378-CAT. An oligonucleotide containing the PCNA intron E2F sequence (CGCGTTTGTGGCTTTG-GCGCGAAAAAAGAGGGGAC) was inserted between *Asc* I and *Ppu* MI sites of PCNA-EHΔ60/378-CAT to yield PCNA-E2F-CAT.

Cell Culture

Human K562 erythroleukemia cells (ATCC CCL 243) were cultured in RPMI 1640 medium supplemented with 10% calf serum. Human Saos-2 osteosarcoma cells (ATCC HTB 85) were cultured in McCoy's 5a medium supplemented with 15% fetal bovine serum (FBS).

Stable DNA Transfection by Electroporation

Electroporation of K562 cells was performed as described (Spandidos et al., 1987). Stable K562 transfectants were selected with 300 μg/ml hygromycin. A mixed population of stable transfectants was expanded for cell cycle synchronization.

Synchronization of K562 Cells by Drug Treatment

Mitotic K562 cells were first synchronized with nocodazole as described (Liu et al., 1994) and were later G₂-arrested with mimosine. S-phase K562 cells were obtained by thymidine block.

Counterflow Centrifugal Elutriation and Flow Cytometry

Counterflow centrifugal elutriation of K562 cells was performed as described (Kauffman et al., 1990) with Beckman JE 5.0 rotor. We routinely used 12–15 l cultures ($3 \sim 5 \times 10^9$ cells). The overall cell cycle distribution of a population of approximately 5×10^5 cells was determined as described (Vindelov and Christensen, 1990). Fluorescence was measured using a Becton Dickinson FACScan and was analyzed on a Hewlett Packard computer with CellFIT software.

Nuclear Transcription Reactions

Nuclear run on transcription was performed as described (Greenberg and Ziff, 1984) with the following modifications. K562 cells were harvested at 1500 rpm for 5 min, then washed with cold PBS three times. The cell pellet (5×10^7 cells) was resuspended in 4 ml 0.1% NP40 lysis buffer [10 mM Tris-HCl, pH 7.5, 10 mM NaCl, 3 mM MgCl₂, 0.1% (v/v) NP40] by pipetting up-and-down three to five times, followed by centrifugation. The nuclear pellet was washed once with 4 ml storage buffer (50 mM Tris-HCl, pH 8.0, 0.1 mM EDTA, 5 mM MgCl₂, 40% glycerol), and then resuspended in 200 μ l storage buffer for storage in liquid N₂. For run-on transcription assay, the nuclei were thawed and mixed with 25 μ l 10 \times reaction buffer (100 mM Tris-HCl, pH 8.0, 10 mM EDTA, 3 mM MgCl₂, 25 mM DTT, 700 mM KCl), 5 μ l of 25 mM each of ATP, CTP, and GTP, 20 μ l of [α -³²P]UTP (200 μ Ci), and 1 μ l RNase inhibitor to react at 30°C for 30 min. To terminate transcription reaction, 1 μ l 250 mM CaCl₂ and 20 μ l of 1 unit/ μ l DNase I were added and incubated at 30°C for 10 min. To remove proteins, 30 μ l 10 \times SET buffer (5% SDS, 50 mM EDTA, 100 mM Tris-HCl, pH 7.4) and 2 μ l proteinase K (10 mg/ml) were added to react at 42°C for 30 min. The guanidinium thiocyanate phenol chloroform procedure was used to isolate RNA as described (Chomczynski and Sacchi, 1987). After ethanol precipitation, ³²P-labeled RNA was dissolved in 100 μ l of TES (10 mM N-Tris[hydroxymethyl]-methyl 2-aminoethane sulphonic acid, pH 7.4,

0.2% SDS, and 10 mM EDTA) and then 1 μ l was used for scintillation counting. The value of counting should be more than 1×10^5 CPM. NaOH was added to give a final concentration of 0.2 M to denature ³²P-labeled RNA for 10 min on ice. The solution was neutralized by the addition of HEPES to a final concentration of 0.5 M and NaCl concentration was adjusted to 0.7 M. In a given experiment, each set of filters was hybridized with the same number of CPM of ³²P-labeled RNA in 1 ml of TES buffer containing 0.2% SDS at 65°C for 36 h. After hybridization, the filters were washed with several changes of 2 \times SSC (1 \times SSC = 0.15 M NaCl, 0.0125 M Na citrate, pH 7.0) twice for 1 h at 65°C and then incubated at 37°C in 2 \times SSC with 10 μ g/ml RNase A for 30 min. The filters were then washed again in 2 \times SSC at 37°C for 1 h, air dried, and exposed in a PhosphoImager cassette (Molecular Dynamics). For binding to nitrocellulose, plasmid DNA was linearized by restriction enzyme digestion and the DNA was denatured by incubation with 0.2 M NaOH for 30 min at room temperature, followed by neutralization with 10 vol of 6 \times SSC. The DNA was loaded onto nitrocellulose using the Schleicher and Schuell slot blot apparatus; 20 μ g DNA was applied per slot.



Transient DNA Transfections and CAT Assays

Human Saos-2 osteosarcoma cells were transfected by calcium phosphate-mediated precipitation, with a total of 25 μ g of DNA per 100-mm dish, including internal control plasmid p β J3, which contains the human β -actin promoter (Ng et al., 1989)-directed β -galactosidase gene. Cell extracts were prepared, and CAT assays were performed as previously described (Liu et al., 1993).

RESULTS

Transcription of Cyclin D1 and DNA Topoisomerase I Genes in Cell Cycle Synchronized Nuclei

Cyclin D1-CAT plasmid was constructed as described in the Materials and Methods section. We transfected this plasmid, together with pSV2-*hph* (Margolskee et al., 1988), into human K562 cells by electroporation. Hygromycin-resistant stable transfectants were assayed for CAT activities and expanded into 1-l suspension culture. To examine transcription of the reporter gene during the cell cycle, we used a two-step procedure for synchronization of cells in mitosis and G₂ by nocodazole-






or mimosine-arrest, respectively (Liu et al., 1994). Thymidine block was used to obtain S-phase-arrested cells. Flow cytometric analysis of DNA content showed that drug-arrested cells were about 85% synchronized (Fig. 1A). About 2×10^8 cells were harvested for nuclear transcription reaction, and labeled RNA was hybridized to CAT, cyclin D1, and glyceraldehyde 3-phosphate dehydrogenase (GAPDH) DNA probes (Fig. 1B). PhosphoImager quantitation of nascent transcripts showed that, relative to the constitutive level of GAPDH transcription, transcription of both endogenous cyclin D1 and transfected cyclin D1-CAT genes is active during G₂ only (Fig. 1C).

DNA topoisomerase I-CAT plasmid was constructed as described in the Materials and Methods section. Transfection, selection, cell cycle synchronization, and nuclear transcription were carried out as described earlier for cyclin D1-CAT. Flow cytometric analysis of DNA content showed that these cells were synchronized from 74% to 91% (Fig. 2A). RNA labeled in nuclear transcription reactions were hybridized to CAT, DNA topoisomerase I, and GAPDH DNA probes (Fig. 2B), and quantitation of nascent transcripts showed that, relative to the constitutive level of GAPDH transcription, transcription of both endogenous DNA topoisomerase I and transfected DNA topoisomerase I-CAT genes is most active during S-phase (Fig. 2C).

Transcription of PCNA Gene in Cell Cycle Synchronized Nuclei

Two PCNA-CAT plasmids were constructed as described in the Materials and Methods section, one containing 413 bp of conserved intron sequences (PCNA-HH-CAT) and the other containing an SV40 intron only (PCNA-HN-CAT). Transfection, selection, cell cycle synchronization, and nuclear transcription were carried out as described earlier for cyclin D1-CAT. Flow cytomet-

FIG. 1. K562 cyclin D1-CAT stable transfectant cell cycle synchronization and transcription analysis. (A) Flow cytometry analysis of mimosine-arrested cells (top), thymidine-blocked cells (middle), and nocodazole-inhibited cells (bottom). The distribution of DNA contents was determined by FACScan. Percentage of G₁, S-phase, and G₂ + M cells in each synchronized cell fraction is shown in each inset. (B) Hybridization of nascent nuclear transcripts from mimosine-arrested (Mms), thymidine-blocked (TdR), or nocodazole-inhibited (Ncdz) cells to cyclin D1, CAT, and GAPDH DNA probes. (C) Quantitation of nascent cyclin D1 and CAT transcripts normalized with nascent GAPDH transcripts, respectively, in mimosine-arrested (Mms), thymidine-blocked (TdR), or nocodazole-inhibited (Ncdz) cells.









ric analysis of DNA content showed that these cells were synchronized from 74% to 96% (Fig. 3A,B). RNA labeled in nuclear transcription reactions was hybridized to CAT, PCNA, and GAPDH DNA probes, and quantitation of nascent transcripts showed that, relative to the constitutive level of GAPDH transcription, transcription of the endogenous PCNA and transfected intron-containing CAT genes during S-phase is slightly higher than the intronless CAT gene (Fig. 3E,F).

Due to possible problems of drug-induced alterations of gene expression (reviewed in Pardee and Keyomarsi, 1992), we repeated nuclear transcription analyses of cyclin D1-CAT and PCNA-CAT genes with nuclei fractionated by counterflow centrifugal elutriation (Kauffman et al., 1990). Cell populations were fractionated on the basis of sedimentation properties; therefore, counterflow centrifugal elutriation currently is the best method to obtain physiologically unperturbed cell cycle phase-specific subpopulations. In fact, we chose to work with human K562 cells because it is the cell line of choice for counterflow centrifugal elutriation (Kauffman et al., 1990). Results from a nuclear transcription experiment with elutriation-synchronized cyclin D1-CAT cells confirm our previous conclusion that this promoter is G₁ inducible (Fig. 4).

To examine whether transcription of PCNA would be transiently activated during G₁/S in the cell cycle, we fractionated K562 PCNA-HH-CAT cells into eight subpopulations, labeled as F1 to F8. Unlike drug-synchronized cell populations, counterflow centrifugal elutriation can fractionate K562 cells to yield a G₁/S subpopulation. Flow cytometric analysis of DNA content showed that F5 cells were 50% G₁ and 41% S-phase (Fig. 5A). Nuclei were prepared from F5 and five other cell fractions for nuclear transcription reactions: labeled RNA was hybridized to CAT, PCNA, and

FIG. 2. K562 DNA topoisomerase I-CAT stable transfectant cell cycle synchronization and transcription analysis. (A) Flow cytometry analysis of mimosine-arrested cells (top), thymidine-blocked cells (middle), and nocodazole-inhibited cells (bottom). The distribution of DNA contents was determined by FACSscan. Percentage of G₁, S-phase, and G₂ + M cells in each synchronized cell fraction is shown in each inset. (B) Hybridization of nascent nuclear transcripts from mimosine-arrested (Mms), thymidine-blocked (TdR), or nocodazole-inhibited (Ncdz) cells to DNA topoisomerase I, CAT, and GAPDH DNA probes. (C) Quantitation of nascent DNA topoisomerase I and CAT transcripts normalized with nascent GAPDH transcripts, respectively, in mimosine-arrested (Mms), thymidine-blocked (TdR), or nocodazole inhibited (Ncdz) cells.



GAPDH DNA probes (Fig. 5B), and quantitation of nascent transcripts showed that, relative to the constitutive level of GAPDH transcription, transcription of both endogenous PCNA and transfected intron-containing CAT genes is clearly more active in F5 nuclei (Fig. 5C). In contrast, transcription reactions carried out with nuclei derived from counterflow centrifugal elutriation-fractionated K562 PCNA-HN-CAT cells showed that the endogenous PCNA gene, but not the transfected intronless CAT gene, is transcriptionally more active in G₁/S (F6) nuclei (Fig. 6). Therefore, element(s) within the PCNA gene first intron is (are) essential for transcription activation of the G₁/S transition.

PCNA Gene First Intron Contained Conserved E2F Sites

When the three PCNA intron 1 sequences were compared, we detected a conserved motif TTTTCGCGCCAAA in human, TTTTCCGCCAAA in rat, and TTTCCGCGCCAAA in mouse (Fig. 7), which resembles the E2F sites of adenovirus E2 and mouse dihydrofolate reductase (DHFR) genes, TTTTCGCGCTTAAA and ATTTTCGCGCCAAA, respectively. Furthermore, 18 bp downstream of this intron site, there is yet another conserved sequence (GCGGGAAAA in all three genes) related to the E2F site in the human thymidine kinase promoter (Kim and Lee, 1992). E2F was previously identified as an early adenovirus gene transcription factor. It is now shown to be involved in cell cycle control as a functional target of the tumor suppressor protein pRb that is encoded by the retinoblastoma susceptibility gene (Nevins, 1992; Johnson et al., 1993). The disruption of E2F-pRb interaction is apparently effected by pRb phosphorylation at the G₁/S transition (reviewed in Cobrinik et al., 1992; Hamel et al., 1992), and/or cyclin D-pRb complex formation (Dowdy et al., 1993; Ewen et al., 1993).

Activation of PCNA Intron Reporter Genes by E2F-1/DP-1 Heterodimer

Recent cloning of the E2F-1 cDNA has facilitated analysis of E2F-mediated cell cycle gene regulation (Helin et al., 1992; Kaelin et al., 1992; Shan et al., 1992). Subsequently, another cDNA, named DP-1 for DRTF-polypeptide-1, which encodes a second E2F site binding protein, was isolated (Girling et al., 1993; Helin et al., 1993b). E2F-1 and DP-1 can heterodimerize in vitro as well as in vivo, leading to cooperative transactivation of the E2F site (Bandara et al., 1993; Helin et al., 1993b; Krek et al., 1993). The PCNA intron E2F sites can be recognized by E2F-1 recombinant protein in vitro (our unpublished observation). To test whether this E2F-1/DP-1 heterodimer can regulate PCNA gene expression, we transfected E2F-1 and DP-1 expression plasmids, together with one of the following reporter genes: PCNA-HH-CAT, PCNA-HN-CAT, and DHFR-CAT (Fig. 8B). Mouse DHFR promoter contains an E2F binding site and can be activated by E2F-1 expression (Slansky et al., 1993). After β -gal normalization, both PCNA-HH-CAT and DHFR-CAT activities were stimulated either four- or seven-fold by E2F-1 and DP-1 expression, respectively. In contrast, the PCNA-HN-CAT gene was not activated by E2F-1 and DP-1 expression.

The PCNA-HH-CAT gene contains over 1500 bp of 5' flanking region for which the nucleotide sequence has not been reported. To better define *cis*-acting element(s) required for activation of the PCNA-HH-CAT gene, first we deleted over 2 kb of 5' flanking sequences to remove this unknown region (Fig. 8A). Cotransfection of this PCNA-EH-CAT gene containing 413 bp intron sequences or a similar PCNA-EN-CAT construct, which contains an additional 295 bp of intron sequences (Fig. 8A), with E2F-1 and DP-1 expression plasmids resulted in activation of both reporter genes (Fig. 8B). Next we made internal deletions of the PCNA first intron: PCNA-EH Δ 60/133-CAT is

FACING PAGE

FIG. 3. K562 PCNA-HH-CAT and PCNA-HN-CAT stable transfectant cell cycle synchronization and transcription analysis. (A) Flow cytometry analysis of mimosine-arrested K562 PCNA-HH-CAT cells (top), thymidine-blocked cells (middle), and nocodazole-inhibited cells (bottom). The distribution of DNA contents was determined by FACScan. Percentage of G₁, S-phase, and G₂ + M cells in each synchronized cell fraction is shown in each inset. (B) Flow cytometry analysis of mimosine-arrested K562 PCNA-HN-CAT cells (top), thymidine-blocked cells (middle), and nocodazole-inhibited cells (bottom). The distribution of DNA contents was determined by FACScan. Percentage of G₁, S-phase, and G₂ + M cells in each synchronized cell fraction is shown in each inset. (C) Hybridization of K562 PCNA-HH-CAT nascent nuclear transcripts to PCNA, CAT, and GAPDH DNA probes. (D) Hybridization of K562 PCNA-HN-CAT nascent nuclear transcripts to PCNA, CAT, and GAPDH DNA probes. (E) Quantitation of K562 PCNA-HH-CAT nascent PCNA and CAT transcripts normalized with nascent GAPDH transcripts, respectively. (F) Quantitation of K562 PCNA-HN-CAT nascent PCNA and CAT transcripts normalized with nascent GAPDH transcripts, respectively.






FIG. 4. K562 cyclin D1-CAT stable transfectant cell cycle synchronization and transcription analysis. (A) Flow cytometry analysis of DNA contents of counterflow centrifugal elutriated cells (F1-F15) as determined by FACScan. Percentage of G₁, S-phase, and G₂ + M cells in each synchronized cell fraction is shown in each inset. (B) Hybridization of nascent nuclear transcripts to cyclin D1, CAT, and GAPDH DNA probes. (C) Quantitation of nascent cyclin D1 and CAT transcripts normalized with nascent GAPDH transcripts, respectively.






FIG. 5. K562 PCNA-HH-CAT stable transfectant cell cycle synchronization and transcription analysis. (A) Flow cytometry analysis of DNA contents of counterflow centrifugal elutriated cells (F1-F8) as determined by FACScan. Percentage of G₁, S-phase, and G₂ + M cells in each synchronized cell fraction is shown in each inset. (B) Hybridization of nascent nuclear transcripts to PCNA, CAT, and GAPDH DNA probes. (C) Quantitation of nascent PCNA and CAT transcripts normalized with nascent GAPDH transcripts, respectively.




FIG. 6. K562 PCNA-HN-CAT stable transfectant cell cycle synchronization and transcription analysis. (A) Flow cytometry analysis of DNA contents of counterflow centrifugal elutriated cells (F1-F9) as determined by FACScan. Percentage of G₁, S-phase, and G₂ + M cells in each synchronized cell fraction is shown in each inset. (B) Hybridization of nascent nuclear transcripts to PCNA, CAT, and GAPDH DNA probes. (C) Quantitation of nascent PCNA and CAT transcripts normalized with nascent GAPDH transcripts, respectively.




FIG. 7. Conserved first intron sequences in the PCNA genes. Alignment of human, mouse, and rat PCNA intron sequences with the GCG pileup program and displayed with PrettyBox. Shown are human intron sequence from base 75 to 600 (Travali et al., 1989), mouse intron sequence from base 71 to 581 (Yamaguchi et al., 1991), and rat intron sequence from base 66 to 579 (Liu, 1992). The positions of E2F sites are indicated. The 35-bp negative regulatory element (Alder et al., 1992) is between base positions 48 and 82 in the human sequence. The 3' junctions of the deletion constructs PCNA-EH Δ 60/133-CAT and PCNA-EH Δ 60/378-CAT are indicated.

missing a known negative region in this intron (Alder et al., 1992), and PCNA-EH Δ 60/378-CAT lacks both this negative region and E2F binding sites (Fig. 8A). As expected, PCNA-EH Δ 60/378-CAT is not inducible by E2F-1 and DP-1, whereas PCNA-EH Δ 60/133-CAT still is (Fig. 8B). Finally, we reinserted the E2 and DHFR-like E2F site (TTTTCGCGCCAAA) back into the PCNA-EH Δ 60/378-CAT plasmid downstream of the transcription initiation site (Fig. 8A). This E2F binding site, originally derived from the first intron, is able to confer inducibility by E2F-1 and DP-1 to this PCNA-E2F-CAT gene (Fig. 8B). We conclude that E2F-1- and DP-1-responsive sequences in the PCNA intron are likely the E2F binding sites.

Cyclin D1 promoter (Herber et al., 1994; Philipp et al., 1994) contains putative E2F binding sites (TTTCGGGCA and TTTGGCGCC) but expression of E2F-1 and DP-1 stimulated the cyclin D1-CAT reporter gene less than twofold (data not

shown). Therefore, although cyclin D1 and PCNA can associate with each other (Xiong et al., 1992b), they are independently regulated at the transcriptional level.

The DNA topoisomerase I promoter also contains two putative E2F binding sites (TTTGCC-CCG and CGGCGGAAA), but this reporter gene was inhibited by E2F-1 and DP-1 (data not shown). Further work will be required to determine whether these sites are recognized by the E2F complex in vivo, or if their flanking sequences contain antagonistic motif(s).

DISCUSSION

Characterization of Cell Cycle-Regulated Promoters

We have examined the promoters of three cell cycle regulatory proteins. The human cyclin D1 promoter activity is restricted to G₁; the human




FIG. 8. Regulation of PCNA promoter activities by CMV E2F-1 and DP-1 expression plasmids. (A) The structure of PCNA reporter genes is shown with the position of E2F site as indicated. (B) The ability of human E2F-1 and DP-1, transfected into human Saos-2 cells, to activate transcription of PCNA-HN-CAT, PCNA-HH-CAT, PCNA-EN-CAT, PCNA-EH-CAT, PCNA-EHΔ60/133-CAT, PCNA-EHΔ60/378-CAT, PCNA-E2F-CAT, or DHFR(-850/+16)pCAT (Schmidt et al., 1990) reporter genes was determined. CAT activities, derived from duplicate transfections, were normalized to the β-galactosidase activities produced by an internal control plasmid, pβJ3. Shown is the stimulation of promoter-driven CAT activity in response to CMV E2F-1 and CMV DP-1 (Helin et al., 1993b) divided by the activity obtained with the CMV vector alone. Activation of PCNA-HH-CAT, PCNA-EN-CAT, PCNA-EH-CAT, PCNA-EHΔ60/133-CAT, and PCNA-E2F-CAT by E2F-1 and DP-1 was observed reproducibly in at least three independent transfections.

DNA topoisomerase I promoter is active throughout S-phase; the human PCNA promoter is transiently activated at the G₁/S transition. Specific cell cycle-regulated elements of these three promoters have not been defined in our present study, but we have mapped the location of such elements within a region of 1 kb for human cyclin D1 and

DNA topoisomerase I promoters, and within a 733-bp region of the PCNA first exon and intron. Further mapping of *cis*-acting elements by promoter deletions, combined with transcription analysis in transfected nuclei, after cell cycle synchronization, should yield definitive information about cell cycle control elements for each promoter.

In previous studies, activation of cell cycle-regulated promoters was only demonstrated in G₀-arrested cells stimulated with serum. These promoters included the human PCNA promoter (Chang et al., 1990; Ottavio et al., 1990), human DNA polymerase α promoter (Pearson et al., 1991), the human cdc2 promoter (Dalton, 1992), the mouse DHFR promoter (Slansky et al., 1993), the mouse B-*myb* promoter (Lam and Watson, 1993), and the human cyclin D1 promoter (Herber et al., 1994). We made the following modification to the analysis of cell cycle-regulated promoters in cell cycle-synchronized cells. First, we choose to use human K562 erythroleukemia cells instead of HeLa cells. Both HeLa and K562 grow in suspension cultures, but the latter is the cell line of choice for cell cycle synchronization by counterflow centrifugal elutriation (Kauffman et al., 1990). HeLa cells also express endogenous HPV-18 E7 protein, which can disrupt interaction between pRb protein and transcription factors (Pagano et al., 1992), including E2Fs (Flemington et al., 1993; Hagemeyer et al., 1993; Helin et al., 1993a; Lees et al., 1993), ATF-2 (Kim et al., 1992), MYC (Rustgi et al., 1991), and Sp-1 (Udvardi et al., 1993). Secondly, we use nuclear transcription reactions to analyze reporter gene expression. Our results thus give a more direct measurement of transcription rates than RNA analyses or reporter enzyme assays.

Regulation of the PCNA Gene by E2F

The E2F binding site complex consists of five polypeptides (Huber et al., 1993), four of which have been cloned: E2F-1 (Helin et al., 1992; Kaelin et al., 1992; Shan et al., 1992), DP-1 (Girling et al., 1993; Helin et al., 1993b), E2F-2 and E2F-3 (Ivey-Hoyle et al., 1993; Lees et al., 1993). The E2F-1 and DP-1 expression plasmids (Helin et al., 1992; Kaelin et al., 1992; Helin et al., 1993b) can activate the E2F site-containing PCNA, DHFR, DNA polymerase α , and cdc2 promoters (our unpublished observation). In contrast, cyclin D1 and DNA topoisomerase I promoters are not activated by E2F-1 and DP-1. The identification of transcriptional regulators for the latter promoters will be facilitated by further characterization of *cis*-acting elements important for cell cycle activation.

The PCNA E2F site, unlike other E2F sites (Nevins, 1992), is localized in the first intron. This intron contains about 500 bp of evolutionarily conserved DNA sequences. Basal activity of the PCNA-EH-CAT gene is about sevenfold lower than the PCNA-EH Δ 60/133-CAT gene, perhaps

due to the negative regulatory region (Alder et al., 1992). The latter region and E2F site are within 100 bp of each other, thus their binding proteins may interact, and possibly may contribute to the transient activation of this gene.

The use of a common transcription factor represents the simplest mechanism for coordinating S-phase gene expression. This mechanism is used by yeast for coordinating the transcription of many S-phase genes. DNA topoisomerase I is activated during S-phase, but unlike the PCNA promoter, it is not E2F inducible. At least one of the two E2F site-related sequences (TTTGCCCCG) can bind E2F, but transactivation may be inhibited by either *trans*-acting or *cis*-acting mechanisms. At least two additional E2F-1-related cDNAs have been cloned (Lees et al., 1993), and whether distinct subsets of E2F sites can be activated by the expression of individual E2F/DRTF polypeptide or specific combinations thereof is being tested. Another possibility is that transcription of the DNA topoisomerase I gene during S-phase is regulated by other cell cycle transcription factor(s) for which binding site(s) reside within the 1-kb 5' flanking region (Heiland et al., 1993).

In summary, we carried out cell cycle transcription analysis of three promoters. Only one of them, the PCNA intron-containing promoter, was shown to be transiently activated at G₁/S and also E2F inducible. The above results, taken together, identified the human PCNA gene as a bona fide cell cycle- and E2F-regulated gene. Many promoters (e.g., *c-myc*) also contain an E2F site, and presumably can be activated by E2F expression. However, *c-myc* is an early G₁ gene and E2F is thought to be associated with pRb during most of G₁, and this complex acts as a transcriptional repressor (Flemington et al., 1993; Hagemeyer et al., 1993; Helin et al., 1993a). Therefore, the presence of E2F site(s) may not infer cell cycle regulation by E2F. The latter process, the coordinate activation of S-phase genes, by E2F and its associated proteins, may be controlled by new E2F synthesis (Slansky et al., 1993; Shan et al., 1994), heterodimerization between E2F-1 and DP-1 or related polypeptide(s) (Bandara et al., 1993; Helin et al., 1993b; Krek et al., 1993), phosphorylation of pRb and inhibition of pRb-E2F interaction, and/or disruption of p107-E2F interaction by unknown mechanisms (Schwarz et al., 1993; Zamanian and La Thangue, 1993; Zhu et al., 1993). A better understanding of E2F regulatory mechanisms will be facilitated by the identification of other E2F-regulated promoters as we have shown here for the PCNA gene.

ACKNOWLEDGEMENTS

We thank Andrew Arnold for cyclin D1/*PRADI* cDNA, Renato Baserga for the human PCNA gene and cDNA, David Beach and Yue Xiong for the human cyclin D1 genomic clone, Kristian Helin and Ed Harlow for CMV E2F-1 and CMV DP-1, Bill Kaelin for CMV RBAP-1, Rowena Girling and Nic La Thangue for SV DP-1, Chi-Hon Lee for DNA topoisomerase I-

CAT, Gary Merrill for DHFR-CAT, and Rick Westerman for PrettyBox. We also thank Heinz-Peter Nasheuer and Teresa S.-F. Wang for helping with counterflow centrifugal elutriation of K562 cells, Chiou-Fen Chuang and Mei-Ling Liu for technical assistance. The work in the laboratory of S.Y.N. is supported by Academic Sinica and National Science Council (ROC).

REFERENCES

- H. Alder, M. Yoshinouchi, M. B. Prystowsky, P. Appasamy, and R. Baserga (1992), *Nucleic Acids Res* 20, 1769-1775.
- B. J. Andrews and I. Hershowitz (1990), *J Biol Chem* 265, 14057-14060.
- V. Baldin, J. Lukas, M. J. Marcote, M. Pagano, and G. Draetta (1993), *Genes Dev* 7, 812-821.
- L. R. Bandara, V. M. Buck, M. Zamanian, L. H. Johnson, and N. B. La Thangue (1993), *EMBO J* 12, 4317-4324.
- R. Baserga (1991), *J Cell Sci* 98, 433-436.
- R. Bravo, R. Frank, P. A. Blundell, and H. Macdonald-Bravo (1977), *Nature* 326, 515-517.
- C.-D. Chang, L. Ottavio, S. Travali, K. E. Lipson, and R. Baserga (1990), *Mol Cell Biol* 10, 3289-3296.
- R.-H. Charollais, H. Alder, A. Ferber, J. Koniecki, C. Sell, and R. Baserga (1992), *Gene Exp* 2, 285-296.
- P. Chomczynski and N. Sacchi (1987), *Anal Biochem* 162, 156-159.
- D. Cobrinik, S. F. Dowdy, P. W. Hinds, S. Mittnacht, and R. A. Weinberg (1992), *Trends Biochem Sci* 17, 312-315.
- S. Dalton (1992), *EMBO J* 11, 1797-1804.
- S. F. Dowdy, P. W. Hinds, K. Louie, S. I. Reed, A. Arnold, and R. A. Weinberg (1993), *Cell* 73, 499-511.
- M. E. Ewen, H. K. Sluss, C. J. Sherr, H. Matsushime, J.-y. Kato, and D. A. Livingston (1993), *Cell* 73, 487-497.
- E. K. Flemington, S. H. Speck, and W. G. Kaelin, Jr. (1993), *Proc Natl Acad Sci USA* 90, 6914-6918.
- R. Girling, J. F. Partridge, L. R. Banara, N. Burden, N. F. Totty, J. J. Hsuan, and N. B. La Thangue (1993), *Nature* 362, 83-87; 365, 468.
- M. E. Greenberg and E. B. Ziff (1984), *Nature* 311, 433-437.
- C. Hagemeyer, A. Cook, and T. Kouzarides (1993), *Nucleic Acids Res* 21, 4998-5004.
- P. A. Hamel, B. L. Gallie, and R. A. Phillips (1992), *Trends Genet* 8, 180-185.
- S. Heiland, R. Knippers, and N. Kunze (1993), *Eur J Biochem* 217, 813-822.
- K. Helin, J. A. Lees, M. Vidal, N. Dyson, E. Harlow, and A. Fattaey (1992), *Cell* 70, 337-350.
- K. Helin and E. Harlow (1993), *Trends Cell Biol* 3, 43-46.
- K. Helin, E. Harlow, and A. Fattaey (1993a), *Mol Cell Biol* 13, 6501-6508.
- K. Helin, C.-L. Wu, A. R. Fattaey, J. A. Lees, B. D. Dynlacht, C. Ngwu, and E. Harlow (1993b), *Genes Dev* 7, 1850-1861.
- B. Herber, M. Truss, M. Beato, and R. Miller (1994), *Oncogene* 9, 1295-1304.
- D. Hochhauser, C. A. Stanway, A. L. Harris, and I. D. Hickson (1992), *J Biol Chem* 267, 18961-18965.
- H. E. Huber, G. Edwards, P. J. Goodhart, D. R. Patrick, P. S. Huang, M. Ivey-Hoyle, S. F. Barnett, A. Oliff, and D. C. Heimbroom (1993), *Proc Natl Acad Sci USA* 90, 3525-3529.
- T. Inaba, H. Matsushime, M. Valentine, M. F. Roussel, C. J. Sherr, and A. T. Look (1992), *Genomics* 13, 565-574.
- M. Ivey-Hoyle, R. Conroy, H. E. Huber, P. J. Goodhart, A. Oliff, and D. C. Heimbroom (1993), *Mol Cell Biol* 13, 7802-7812.
- D. G. Johnson, J. K. Schwarz, W. D. Cress, and J. R. Nevins (1993), *Nature* 365, 349-352.
- L. H. Johnston (1992), *Trends Cell Biol* 2, 353-357.
- W. G. Kaelin, Jr., W. Krek, W. R. Sellers, J. A. DeCaprio, F. Ajchenbaum, C. S. Fuchs, T. Chittendan, Y. Li, P. J. Farnham, M. A. Blonar, D. M. Livingston, and E. K. Flemington (1992), *Cell* 70, 351-364.
- M. G. Kauffman, S. J. Noga, T. J. Kelly, and A. D. Donnemberg (1990), *Anal Biochem* 191, 41-46.
- S. J. Kim, S. Wagner, F. Liu, M. A. O'Reilly, P. D. Robbins, and M. R. Green (1992), *Nature* 358, 331-334.
- Y. K. Kim and A. S. Lee (1992), *J Biol Chem* 267, 2723-2727.
- W. Krek, D. M. Livingston, and S. Shirodkar (1993), *Science* 262, 1557-1560.
- N. Kunze, G. Yang, M. Dolberg, R. Sundarp, R. Knippers, and A. Richter (1991), *J Biol Chem* 266, 9610-9616.
- C. Labrie, G. F. Morris, and M. B. Mathews (1993), *Mol Cell Biol* 13, 1697-1707.
- E. W.-F. Lam and R. J. Watson (1993), *EMBO J* 12, 2705-2713.
- N. B. La Thangue (1994) *Trends Biochem Sci* 19, 108-114.
- J. A. Lees, M. Saito, M. Vidal, M. Valentine, T. Look,

- E. Harlow, N. Dyson, and K. Helin (1993), *Mol Cell Biol* 13, 7813-7825.
- S.-H. Liu, J.-T. Ma, A. Y. Yueh, S. P. Lees-Miller, C. W. Anderson, and S.-Y. Ng (1993), *J Biol Chem* 268, 21147-21154.
- S.-H. Liu, H.-H. Lee, J.-J. Chen, C.-F. Chuang, and S.-Y. Ng (1994), *Cell Growth Differen* 5, 447-455.
- Y.-C. Liu (1992), GenBank X67329.
- R. F. Margolskee, P. Kavathas, and P. Berg (1988), *Mol Cell Biol* 8, 2837-2847.
- J. D. McKinney and N. Heintz (1991), *Trends Biochem Sci* 16, 430-435.
- T. Motokura and A. Arnold (1993), *Curr Opin Genet Dev* 3, 5-10.
- J. R. Nevins (1992), *Science* 258, 424-429.
- S.-Y. Ng, P. Gunning, S.-H. Liu, J. Leavitt, and L. Kedes (1989), *Nucleic Acids Res* 17, 601-615.
- L. Ottavio, C.-D. Chang, M.-G. Rizzo, S. Travali, C. Casadevall, and R. Baserga (1990), *Mol Cell Biol* 10, 303-309.
- M. Pagano, M. Dürst, S. Joswig, G. Draetta, and P. Jansen-Dürr (1992), *Oncogene* 7, 1681-1686.
- A. B. Pardee and K. Keyomarsi (1992), *Curr Opin Cell Biol* 4, 186-191.
- B. E. Pearson, H.-P. Nasheuer, and T. S.-F. Wang (1991), *Mol Cell Biol* 11, 2081-2095.
- A. Philipp, A. Schneider, I. Vasrik, K. Finke, Y. Xiong, D. Beach, Alitalo, and M. Eilers (1994), *Mol Cell Biol* 14, 4032-4043.
- G. Prelich and B. Stillman (1988), *Cell* 53, 117-126.
- D. E. Quelle, R. A. Ashman, S. A. Shurtleff, J.-y. Kato, D. Bar-Sagi, M. F. Roussel, and C. J. Sherr (1993), *Genes Dev* 7, 1559-1571.
- A. K. Rustgi, N. Dyson, and R. Bernards (1991), *Nature* 352, 541-544.
- J.-L. Salier and K. Kurachi (1989), *Biotechniques* 7, 30-31.
- E. E. Schmidt, R. A. Owen, and G. F. Merrill (1990), *J Biol Chem* 265, 17397-17400.
- J. K. Schwarz, S. H. Devoto, E. J. Smith, S. P. Chelappan, L. Jakoi, and J. R. Nevins (1993), *EMBO J* 12, 1013-1020.
- B. Shan, X. Zhu, P.-L. Chen, T. Durfee, Y. Yang, D. Sharp, and W.-H. Lee (1992), *Mol Cell Biol* 12, 5260-5631.
- B. Shan, C.-Y. Chang, D. Jones, and W.-H. Lee (1994), *Mol Cell Biol* 14, 299-309.
- C. J. Sherr (1991), *Trends Genet* 7, 398-402.
- C. J. Sherr (1993), *Cell* 73, 1059-1065.
- J. E. Slansky, Y. Li, W. G. Kaelin, Jr., and P. G. Farnham (1993), *Mol Cell Biol* 13, 1610-1618.
- D. A. Spandidos (1987), *Gene Anal Technol* 4, 50-56.
- S. Travali, D.-H. Ku, M. G. Rizzo, L. Ottavio, R. Baserga, and B. Calabretta (1989), *J Biol Chem* 264, 7466-7472.
- A. J. Udvardia, K. T. Rogers, P. D. R. Higgins, Y. Murata, K. H. Martin, P. A. Humphrey, and J. M. Horowitz (1993), *Proc Natl Acad Sci USA* 90, 3265-3269.
- L. L. Vindelov and I. J. Christensen (1990), *Cytometry* 11, 753-770.
- S. Waga and B. Stillman (1994), *Nature* 369, 207-212.
- J. C. Wang (1991), *J Biol Chem* 266, 6659-6662.
- Y. Xiong, J. Menninger, D. Beach, and D. C. Ward (1992a), *Genomics* 13, 575-584.
- Y. Xiong, H. Zhang, and D. Beach (1992b), *Cell* 71, 505-514.
- M. Yamaguchi, Y. Hayashi, F. Hirose, S. Matsuoka, T. Moriuchi, T. Shiroishi, K. Moriwaki, and A. Matsukage (1991), *Nucleic Acids Res* 19, 2403-2410.
- M. Zamanian and N. B. La Thangue (1993), *Mol Biol Cell* 4, 389-396.
- H. Zhang, Y. Xiong, and D. Beach (1993), *Mol Biol Cell* 4, 897-906.
- L. Zhu, S. van den Heuvel, K. Helin, A. Fattaey, M. Ewen, D. Livingston, N. Dyson, and E. Harlow (1993), *Genes Dev* 7, 1111-1125.