



Published in final edited form as:

*Med Sci Sports Exerc.* 2018 October ; 50(10): 2156–2163. doi:10.1249/MSS.0000000000001679.

## Pilot Study of Impact of a Pedal Desk on Postprandial Responses in Sedentary Workers

Ho Han<sup>1,2</sup>, Jongil Lim<sup>1,3</sup>, Richard Viskochil<sup>1</sup>, Elroy J. Aguiar<sup>1</sup>, Catrine Tudor-Locke<sup>1</sup>, and Stuart R. Chipkin<sup>1</sup>

<sup>1</sup>Department of Kinesiology, University of Massachusetts Amherst, Amherst, MA

<sup>2</sup>School of Community Health Sciences, Counselling and Counselling Psychology, Oklahoma State University, Stillwater, OK

<sup>3</sup>Department of Counseling, Heath and Kinesiology, Texas A&M University – San Antonio, San Antonio, TX

### Abstract

Physical inactivity has been linked to rates of obesity, diabetes and heart disease through insulin resistance and other mechanisms. While sedentary workplace environments have unintentionally contributed to the risk for chronic diseases, innovations in the workplace environment could potentially rectify this public and occupational health problem.

**Purpose**—To evaluate the effects of light intensity physical activity using a pedal desk (PD) compared to a standard desk (STD) in a pilot study on postprandial metabolic responses and work skills.

**Methods**—Twelve overweight/obese full-time sedentary office workers (6 men and 6 women; BMI=  $28.7 \pm 3.6$  kg/m<sup>2</sup>) were tested in two conditions: (1) PD: pedaling at self-selected light intensity pace for 2-hr and (2) STD: remaining seated for 2-hr in a conventional workstation set up, while performing scripted computer-based work tasks. Blood samples were analyzed for plasma glucose, insulin, and free-fatty acids (FFA) in response to a standardized meal and work skills were evaluated. Paired samples t-tests were used to examine the differences in metabolic responses and work performance tasks between the conditions.

---

Corresponding Author: Stuart R. Chipkin, MD, Research Professor, School of Public Health and Health Sciences, University of Massachusetts Amherst, 240 Thatcher Road, S323Amherst, MA 01003, schipkin@umass.edu, Phone: +1 413 545 3352, Fax: +1 413 545 2906.

The authors declare they have no financial conflicts of interest. Dr. Catrine Tudor-Locke and Mr. Gerald R. Locke, co- invented and thus own intellectual property for the Pennington Pedal Desk™. A management plan was established for financial conflicts of interest that included electronic data capture, a third party audit of the data and manuscript, and designation of Dr. Stuart Chipkin as the Principal Investigator who managed the study. Dr. Tudor-Locke did not determine study design and had no contact with study participants or study data. None of the other authors have a potential conflict of interest. The results of the resent study do not constitute endorsement by the American College of Sports Medicine. The results of the study are presented clearly, honestly, and without fabrication, falsification, or inappropriate data manipulation.

**Conflict of Interest:** The authors declare they have no financial conflicts of interest. Dr. Catrine Tudor-Locke and Mr. Gerald R. Locke, co- invented and thus own intellectual property for the Pennington Pedal Desk™. A management plan was established for financial conflicts of interest that included electronic data capture, a third party audit of the data and manuscript, and designation of Dr. Stuart Chipkin as the Principal Investigator who managed the study. Dr. Tudor-Locke did not determine study design and had no contact with study participants or study data. None of the other authors have a potential conflict of interest. The results of the resent study do not constitute endorsement by the American College of Sports Medicine. The results of the study are presented clearly, honestly, and without fabrication, falsification, or inappropriate data manipulation.

**Results**—PD use required significantly less insulin to maintain glucose concentrations compared with STD condition (peak insulin concentration 42.1 vs. 66.9  $\mu\text{U}/\text{mL}$ ,  $p = 0.03$ , and AUC 302.6 vs. 441.8  $\mu\text{U}\cdot\text{min}/\text{mL}$ ,  $p < 0.001$ ). No significant changes in plasma glucose and FFA concentrations were observed at any time-points (all  $p > 0.05$ ). In addition, pedaling at a self-paced rate caused no adverse effects on work skills ( $p > 0.05$ ).

**Conclusions**—The PD resulted in lower post-meal insulin concentrations without an overall negative impact on work skills. Thus, the PD could have the potential to achieve public and occupational health goals in sedentary work environments.

### Keywords

Insulin; pre-diabetes; light intensity physical activity

---

### Introduction

Sedentary behavior, defined as waking activities in a seated or reclining posture that require an energy expenditure  $< 1.5$  metabolic equivalents [METs] (1), has been linked to an increased risk of obesity and type 2 diabetes (T2D) through increased insulin resistance and abnormal insulin action when controlled for moderate-to-vigorous physical activity levels (2–4). Contemporary growth in business technology has led to decreases in workplace physical activity (5). In fact, excess sitting time in the office-based workplace environment has become the single largest contributor (52%) to total weekday sitting time (6). A recent report on sedentary behavior in the workplace environment concluded that employees spent about two-thirds of their working hours (equivalent to approximately 5 hours/working day) in prolonged, unbroken periods of sitting lasting  $> 30$  minutes (7, 8).

Just as sedentary behavior induces insulin resistance, physical activity can increase insulin sensitivity and improve insulin action. Even light intensity physical activity (e.g., transitions from sitting to standing) can positively affect postprandial elevations of blood glucose and insulin that occur in insulin-resistant individuals (9, 10). Sedentary office workers are thus a key target group for reducing prolonged sitting time, or replacing it with even light intensity activity, with the goal to decrease insulin resistance.

Workplace innovations may be able to help interrupt sedentary behavior and replace it with light intensity activity and reduce insulin resistance for sedentary employees. Some interventions have demonstrated a benefit of intermittent light intensity activity breaks on diminishing postprandial metabolic responses (9, 11). However, many employees may not have the resources or scheduling autonomy to incorporate light intensity activity breaks into their workflow. Another approach has been to incorporate light intensity activity into workplace settings using specially designed “standing desks” or “treadmill desks”. However, these workstations can have limitations including: 1) a decrease in work performance and motor skills (e.g., typing), 2) limited access for those with musculoskeletal conditions, and 3) feelings of fatigue and discomfort after prolonged standing (12). A recent review concluded that substantial research gaps prevent evaluating the overall effectiveness of standing and treadmill desks to reduce sedentary time and improve health (13).

Desks configured with a seated pedal apparatus (see Figure, SDC 1, figure of pedal desk) could be an important alternative tool for reducing workplace inactivity because they are: 1) self-paced, 2) easier to use for workers with existing musculoskeletal problems or reduced mobility, 3) oriented towards non-weight bearing activities, 4) minimally disruptive to the primary work-related tasks, and 5) require smaller footprints than treadmill desks in the workplace (12, 143). However, their use cannot be advocated without documenting an impact on metabolic consequences of inactivity. The primary purpose of this study was to evaluate the effects of light intensity physical activity using a pedal desk (PD) compared with a control condition using a standard desk (STD) on postprandial glucose, insulin and fat concentrations in a pilot study among overweight/obese sedentary office workers in a simulated work environment. A secondary aim was to determine the effectiveness of a PD (compared to a STD) on work skills performed during routine sedentary office work.

## Methods

### Participants

Participants consisted of 12 (6 men and 6 women) full-time sedentary office workers who were recruited from the Amherst, MA area. The study inclusion criteria required that all participants were overweight or obese (BMI > 25 kg/m<sup>2</sup>), aged 21–64 years, and self-reported employment in a sedentary occupation (i.e., mainly seated during working hours). Individuals were excluded if they self-reported weight > 250 lbs. (limitation of the prototype PD), recent injuries or other major health conditions (e.g., cancer, heart disease, liver or kidney disease, etc.) which would prevent using the PD, self-reported diagnosis of diabetes, or a history of reactive hypoglycemia. The study protocol was approved by the Institutional Review Board at the University of Massachusetts Amherst, and written informed consent was obtained from each participant prior to enrollment in the study.

### Study Procedures

Participants attended two separate randomly assigned visits (at least 6 days in between tests) to the laboratory: 1) PD condition: Participants pedaled at a self-selected light intensity for the duration of the experiment (120 minutes) and 2) STD condition: Participants remained seated throughout the experimental period in a conventional workstation set up (i.e., an office chair with standard height desk). There were no rest breaks permitted during the experimental period. Participants were asked to maintain their usual sedentary lifestyle and to eat similar diets for the two days leading up to testing. At the first visit, additional verbal explanation of the study details was provided as well as an orientation to the use of the PD. Height (cm) was measured on the first visit and weight (kg) was taken on both visits.

At both experimental visits, participants rested for 20 minutes and had an intravenous catheter placed in the forearm, which was used for venous blood sampling. Following baseline fasting sample collection, the participant was given a standardized meal (i.e., cornflakes, heavy cream and whole milk) with known carbohydrate (75 g) and fat (50 g) content (a total of 837.5 kcal) to be ingested over a 10-minute period. Blood samples for glucose, insulin and free fatty acid (FFA) concentrations were collected every 15 minutes

during the remainder of the post-meal test while participants were performing scripted computer-based work tasks (Figure 1).

## Measures

**Anthropometrics**—Participant's height was measured to the nearest 0.1 cm using a portable stadiometer (ShorrBoard®; Weight and Measure, LLC, Olney, MD, USA). A dual-frequency total body composition analyzer (Tanita DC-430U; Tanita Corporation, Tokyo, Japan) was used to measure participant's body weight and body mass index (BMI). All measurements were taken twice, and a third measurement was taken if the first two measurements of height and weight were greater than 0.5 cm and 0.5 kg apart, respectively. The average of two closest measurements was used for analysis.

**Measures of pedaling performance**—An accelerometer-based cadence sensor (Garmin Vector™ 2S, Garmin®, USA) paired with a Garmin EDGE® 820 GPS bike computer (Garmin®, USA) was used to continuously track cadence, power and total time in pedaling performed for the duration of experiments. A fixed level of flywheel resistance ( $\approx 0.30$  kiloponds) was used for enabling prolonged duration of pedaling. Participants were not provided with any feedback on their pedaling performances.

## Work Skills

**Stroop Test**—A computerized version of the Stroop Test (e.g., Stroop Color and Word Test) was used to assess participants' executive processing ability, information processing as well as selective attention capacity and skills (14). The test consisted of 3 sections with 60 items per section including: (1) Black trial – containing 3 color words (i.e., *red*, *blue* and *green*) displayed in black color; (2) Congruent trial – containing the color words displayed in a congruent color (e.g., the word “blue” displayed in blue color); and (3) Incongruent trial – containing the color words displayed in an incongruent color (e.g., the word “green” displayed in red color). The three sections were presented to participants in random order with a 30-second rest between sections. At the beginning of the test, participants practiced responding to the experimental stimuli, which were presented at the center of the computer screen and remained on the screen until a response was made, as quickly and accurately as possible by pressing the key corresponding to either *red*, *blue*, or *green* color. The test was repeated for all participants in the next cycle (figure 1). A custom-written MATLAB program (The MathWorks, Inc., Natick, MA, USA) recorded the accuracy of the performance and time to complete each section. The average of two measurements was retained for analysis.

**Mouse proficiency**—A mouse clicking and a drag-and-drop test was performed using a custom-written MATLAB program to determine participant's computer mouse proficiency (15). In the mouse-clicking test, the participant was instructed to click as quickly as possible on 1 of 25 squares that randomly turned green in color until all squares were clicked. The total time was measured and averaged across tests. Twenty-five squares were located at predetermined-positions to ensure that the movement distance was equal across the tests.

The drag-and-drop test involved dragging and dropping an appeared-square into a larger white box located at the bottom of the computer screen. A square appeared until successfully dropped into the box, after which another square on the screen appeared in a random location. The total amount of time taken to complete each test was recorded.

**Typing**—Typing performance was measured using a typing software program (TypingMaster Pro 10 Premium, TypingMaster, Inc., Helsinki, Finland) to evaluate typing skills (e.g., speed and accuracy) (16, 17). Participants were asked to type a non-technical moderately difficult script for 1 minute as quickly and as accurately as possible. Two of four similarly intense scripts were consecutively presented on the screen in a counterbalanced order. The other two scripts were performed at the second visit. Upon completion of typing the scripts, typing speed and accuracy were recorded and averaged.

**Reading comprehension**—Four graduate record examination (GRE) reading comprehension tests (multiple-choice questions only with one answer choice) were used to assess working cognitive performance. The reading comprehension question of the GRE is designed to test a wide range of abilities to understand, summarize, and analyze information, and has been used in previous studies (15, 18). Each test consisted of seven questions of similar difficulty and was timed for completion within 10 minutes. Two of four tests were randomly administered for the first visit and the remainder of the tests was used in the second visit. Reading speed and accuracy of responses were averaged for the two tests given under each condition.

### **Blood Collection, Storage and Analyses**

All the blood samples were collected in vacutainers containing sodium fluoride and potassium oxalate (5:4) mixture (glucose) and EDTA (insulin and FFA) and immediately centrifuged at 3300 revolutions per minute (RPM) for 15 minutes. Plasma samples were subsequently aliquoted into cryotubes and stored in a –80 degree freezer until all participants had completed both test protocols.

**Glucose**—Plasma glucose concentrations were determined using the glucose oxidase method (Analox instruments, Atlanta GA). PD and STD samples were analyzed together in duplicate with an inter-assay coefficient of variability of < 5%.

**Insulin**—Plasma insulin concentrations were determined using a commercially available radioimmunoassay (RIA, Millipore, Billerica, MA). Samples from both trials for each participant were measured in duplicate on the same assay in order to minimize intra-assay variability, with an inter-assay coefficient of variability of < 10%.

**Free Fatty Acids (FFA)**—Circulating FFA concentrations were determined using a colorimetric assay (Sigma Aldrich, St. Louis MO). Samples were assayed in duplicate with an inter-assay coefficient of variability of < 10%.

## Statistical Analysis

Statistical analyses were conducted using IBM SPSS Statistics 22 for Windows (IBM Corp. Armonk, NY) and SAS/STAT software, Version 9.4 of the SAS System for Windows (SAS Institute Inc., Cary, NC, USA). Descriptive statistics were utilized to summarize participant characteristics, pedaling performance, and work skills presented as means, standard deviations, counts and percentages. Linear Mixed Models (PROC MIXED with Toeplitz covariance structure to account for repeated measures) were used to determine main condition effects (PD vs. STD) over time for serum glucose, insulin and FFA levels (adjusted for baseline measures). Where a significant condition by time interaction was observed, the Tests of Effect Slices allowed for comparison of differences in serum glucose, insulin and FFA levels at each time-point between the respective conditions. For the primary analysis (linear mixed models presented above), no additional adjustments were made for age, sex or BMI due to the small sample size of this pilot study. However, we conducted an additional (preliminary and exploratory) ad hoc examination of the effects of sex, age, and BMI interactions, by adding them to the linear mixed models for glucose, insulin and FFA. These exploratory results are reported in the Discussion section. Appropriate caution is advised concerning the interpretation of these findings given the small sample size and limited statistical power for these analyses. Additionally, insulin, glucose and FFA area under the curve (AUC) and incremental AUC (iAUC) were calculated using the trapezoid rule and compared between the two conditions using a paired t-test. Cohens *d* effect sizes were calculated for differences in AUC between conditions, where effect size magnitudes of  $d = 0.2, 0.5, 0.8$ , were considered small, medium and large, respectively (19). Statistical significance was set at  $p < 0.05$  for all analyses.

## Results

The participants recruited for this pilot study were all sedentary office workers (average age of  $38.9 \pm 10.9$  years) who estimated their average time sitting at work was  $8.7 \pm 2.2$  hours per day. These participants were overweight or obese with an average BMI of  $28.7 \pm 3.6$  kg/m<sup>2</sup>. All participants (6 men, 6 women) completed both PD and STD trials and were included in the analyses. Mean values of participant characteristics and quantified pedaling performance are presented in Table 1. On average, participants pedaled at a cadence of  $60.5 \pm 10.6$  RPM throughout the 2-hr PD trial resulting in a power output of  $40.5 \pm 9.2$  watts which falls within the range of 30–50 watts consistent with stationary bicycling of “very light to light effort” (20). Pedaling cadence and power output generated remained relatively stable throughout the 2-hr PD trial (9.3% and 13.8% of coefficients of variations, respectively) (see Figure, SDC 2, Example of a single participant’s pedaling power output and cadence throughout the 2-hr PD trial).

## Mixed Meal Responses

Baseline fasting glucose, insulin and FFA concentrations were not significantly different between PD or STD conditions (Table 2). Interestingly, nine of the subjects had fasting glucose concentrations between 100–126 mg/dL consistent with impaired fasting glucose (21). There was no main effect of condition (PD vs STD) on glucose ( $p = 0.96$ ), and there were no significant differences in peak glucose concentration (134.4 vs. 132.3 mg/dL for PD



and STD, respectively,  $p = 0.70$ ) or plasma glucose concentrations at any of the examined time-points (all  $p > 0.05$ ). In addition, there was no significant difference in the AUC for glucose (1118.0 vs. 1122.1 mg.min/dL,  $p = 0.87$ ) (Figure 2A & B). Furthermore, a Cohen's  $d$  of 0.02 indicates no differences in glucose AUC between conditions.

A main effect of condition was observed for insulin ( $p = 0.004$ ), as well as a desk\*time interaction ( $p = 0.04$ ). While there were no differences between PD and STD during the initial 30 minutes of the mixed meal tolerance test, insulin concentrations were significantly lower with the PD than the STD beginning at 45 minutes and persisting through all subsequent time-points (Test of Effect Slices for desk\*time from 45–120 min, all  $p < 0.05$ ) (Figure 2C). Additionally, the AUC (302.6 vs. 441.8  $\mu\text{U}\cdot\text{min}/\text{mL}$ ,  $p < 0.001$ ) and the peak insulin concentration (42.1 vs. 66.9  $\mu\text{U}/\text{mL}$ ,  $p = 0.03$ ) were significantly lower for the PD session (Figure 2D). In addition, a Cohen's  $d$  of 0.8 indicates a large effect size for the difference in insulin AUC between conditions.

There was no main effect of condition on FFA ( $p = 0.40$ ) and there were no significant differences between conditions in peak FFA concentration (173.4 vs. 175.1  $\mu\text{mol}/\text{L}$ ,  $p = 0.91$ ) or plasma FFA concentrations at any of the examined time-points (all  $p > 0.05$ ). There was also no significant difference in the FFA AUC (12.9 vs. 14.8  $\text{mmol}\cdot\text{min}/\text{L}$ ,  $p = 0.29$ ) between the two conditions (Figure 2E & F). Furthermore, a Cohen's  $d$  of 0.2 also indicates a small (trivial) effect size for the difference in FFA AUC between conditions.

In addition to the standard AUC results presented above and in Figure 2, we also calculated iAUC values (see Table, SDC 3, AUC and iAUC for responses for glucose, insulin and FFA). The findings for iAUC followed the same pattern as the standard AUC results, with significant changes only observed for insulin iAUC (213.6 vs. 358.8  $\text{mmol}\cdot\text{min}/\text{L}$ ,  $p < 0.01$ ).

## Work Skills

The work skill measures are summarized in Table 2. Overall, there were no significant differences in any of the measures of work skills between the PD and STD conditions ( $p > 0.05$ ) except one; the mouse clicking response was slower for the PD (mean  $\pm$  SD = 59.8  $\pm$  5.7 vs. 57.7  $\pm$  4.5 sec, respectively,  $p = 0.01$ ). The PD was not significantly different than the STD for either time to complete or response accuracy for the Stroop Test (all  $p > 0.05$ ) which assessed cognitive processing. The response time was shortest for congruent stimuli (mean  $\pm$  SD = 41.5  $\pm$  7.8 sec for PD, 43.8  $\pm$  6.9 sec for STD) followed by black (43.8  $\pm$  8.0 vs. 44.5  $\pm$  7.1 sec) and incongruent stimuli (56.3  $\pm$  41.3 vs. 55.5  $\pm$  13.1 sec); the accuracy rates occurred in the reverse order for both conditions. The speed of information processing measured by reaction time test for 1 key or for 8 keys was not different between PD and STD (1 key RT: 305.1  $\pm$  42.8 ms for PD, 293.2  $\pm$  33.2 ms for STD; 8 key RT: 578.9  $\pm$  155.1 ms for PD, 580.5  $\pm$  136.9 ms for STD). Similarly, in the reading comprehension test, the PD was not worse than the STD for reading speed or accuracy ( $p = 0.43$  and 0.58, respectively).

The PD was associated with a significantly longer response time in mouse clicking compared with the STD condition ( $p = 0.01$ ). However, the average difference between PD and STD was only 2.2 seconds over approximately one minute of testing. There were no

significant differences (all  $p > 0.05$ ) for other tests assessing computer mouse and typing skills (i.e., Mouse Drag-and-Drop and typing tests).

## Discussion

The primary goal of this study was to determine whether use of a PD in a high-risk group of sedentary workers would positively affect postprandial metabolic parameters without negatively impacting work-related activities. PD use was associated with significant decreases in insulin concentrations beginning at 45 minutes after consuming a mixed carbohydrate/fat meal relative to STD condition, although no significant changes in plasma glucose and FFA concentrations were observed. In terms of work skills, self-selected light intensity pedaling while working generally demonstrated no significant differences in ability to complete work-related activities compared to the STD condition.

The significant decreases in insulin levels from 45–120 minutes and decrease in AUC (large effect size) suggest a benefit of the light intensity activity resulting from use of the PD. Such a continued light intensity physical activity has been shown to be effective on improving insulin sensitivity. For instance, an improvement in insulin sensitivity has been reported after replacing sitting time with a long duration minimal intensity physical activity (e.g., 2-hr standing and 4-hr walking at a leisure pace) compared to energy-matched 1-hr moderate to vigorous intensity physical activity in a day (22). An approximately 5% increase in insulin sensitivity can be achieved by replacing 30 minutes of sedentary time with light intensity physical activity in those with a high risk of T2D (23). Herein, participants were instructed at the beginning of the PD visit to pedal at a speed that was comfortable for them. They were not coached or reminded to pedal during the test period. Nonetheless, participants pedaled steadily (i.e., without break) for the entire 2-hr at a continuous pace consistent with light intensity physical activity. These current findings suggest that using a PD in a workplace environment is appropriate for sedentary office workers at high risk for T2D to lower postprandial insulin levels.

It is likely that lower insulin concentrations in the postprandial period have the potential to benefit sedentary workers. In addition to potential improvements in insulin sensitivity induced by light intensity physical activity, lower insulin concentrations realized during the meal tolerance test while using the PD may also be beneficial for beta cell function. Insulin resistance is often recognized as the first step in the pathophysiology linking obesity to T2D (24), however the transition from normoglycemia to hyperglycemia is characterized by inadequate pancreatic insulin secretion (25, 26). The preservation of beta cell function, or the ability to maintain appropriate insulin supply (e.g., secretion) to match increases to insulin demand (i.e., resistance), thus represents a key component of diabetes prevention (27).

The observation that glucose concentrations were similar between trials despite significantly lower insulin concentrations when using the PD suggests that glycemic control during light intensity pedaling was partially accomplished through skeletal muscle contraction. Contraction-mediated glucose uptake represents a potential mechanism by which insulin secretion can be “spared” while still maintaining normal blood glucose control (28), and has



been linked to cardiometabolic health improvements in sedentary adults at risk for T2D (29, 30) as well as reduced hyperglycemia in adults with T2D (31, 32). Studies evaluating contraction-mediated glucose uptake have often used a moderate intensity exercise protocol (e.g. 30 minute walk following meals), which may not be feasible in an office setting, or may negatively impact work productivity. Results from this study suggest that light intensity pedaling is sufficient to induce contraction-mediated glucose uptake, without any practical decrements in selected workplace skills. If muscle contractions result in lower circulating insulin concentrations through reduced insulin secretion, light intensity postprandial cycling may represent a potent means to preserve insulin secretory capacity and beta cell function. Interestingly, the pedal desk may also be able to assess the direct consequences of muscle contractility independent of other factors affected by changes in posture such as vascular flow.

The lack of significant change in glucose and FFA may have several possible explanations. First, the intensity of the activity performed may have been insufficient to lower glucose concentrations. These participants were not known to have glucose abnormalities and would not have been expected to respond to a meal differently between conditions. The decrease in insulin concentrations associated with use of a PD in a cohort comprised largely of sedentary workers suggests an improvement in insulin sensitivity. We cannot exclude increased clearance as an alternative explanation for the lower insulin concentrations. However, since insulin is not cleared by skeletal muscle, there is no reason to expect light intensity physical activity to cause an increase in insulin clearance. A second reason for lower insulin concentrations without impacting glucose or FFA might be that the duration of a single 2-hr bout of light intensity activity was insufficient. Greater durations of PD use or repeated bouts may be needed to impact postprandial changes in glucose concentrations. Previous studies have documented postprandial changes after repeated bouts versus a single bout of activity (33). This is feasible to study in the future because the PD could be used throughout the day and on repeated days in an office environment. Third, a greater sample size may be needed to demonstrate a change in glucose or FFA. Review of the FFA data (Figure 2E) note that samples during minutes 15–75 of the PD were apparently (but not significantly) lower for the PD visit compared to the STD. The study of additional participants might provide sufficient analytical power to demonstrate a further separation in these parameters. Lastly, the test meal contained fixed amounts of carbohydrate and fat; the impact of the PD on meals with higher concentrations of fat or carbohydrate might produce different outcomes.

In addition to the primary analyses discussed above, we also conducted an additional ad hoc exploration of sex, age and BMI covariate interactions in the linear mixed model analyses for glucose, insulin and FFA. For glucose, although there was a main effect for age ( $p = 0.02$ ), there was no desk\*age interaction ( $p = 0.42$ ). There were no main or interaction effects for sex or BMI (all  $p > 0.05$ ). For insulin, there were no main effects for sex, age or BMI (all  $p > 0.05$ ). However, upon closer inspection of the data [Test of Effect Slices for desk\*time\*sex; and figures (see Figure, SDC 4, sex-dependent effects of PD and STD conditions)], females displayed significantly higher insulin values in the STD vs PD condition from 45–120 min (all  $p < 0.05$ ). In contrast, men only displayed a significantly higher insulin value in the STD vs PD condition at 45 min ( $p = 0.0055$ ) and trended towards significance at 60 min ( $p = 0.0543$ ), with all time-points thereafter not significantly different.

For FFA, there was a main effect for sex ( $p = 0.02$ ), and an interaction effect for desk\*sex ( $p = 0.04$ ). More specifically, females tended to display higher FFA values overall (sex main effect), but particularly so for the STD condition (desk\*sex interaction), as compared to males (Test of Effect Slices for desk\*time\*sex: STD condition, men vs women from -10 to 105 min, all  $p < 0.05$ ; for PD condition, men vs women from -10 to 15 min, all  $p < 0.05$ , time-points thereafter not significantly different). This may explain, at least in part, the aforementioned apparent (but not statistically significant) differences between STD and PD observed in Fig 2E from 15–75 min. In addition, although there was an overall main effect for age ( $p = 0.03$ ), there was no desk\*age interaction ( $p = 0.11$ ), hence age was not retained in the model for further analysis. Finally, there was no main or interaction effect for BMI ( $p > 0.05$ ). Again, we re-iterate that these additional ad hoc analyses were exploratory and should be interpreted with caution given that this was a pilot study with a relatively small sample size. Further studies (with more adequately powered samples) should be conducted to more definitively assess sex, age and BMI covariate interactions.

The PD compared favorably with a standard desk in regards to work-related activities. We chose different skills: executive processing via the Stroop Word Color Test; computer proficiency using mouse click, drop-and-drag and typing; and reading comprehension via GRE questions. All measures except the mouse click test (2.2 seconds out of approximately 1 minute) were equivalent between the PD and the STD conditions. The lack of differences in the multiple work skills between PD and STD are consistent with previous results that a seated active workstation (e.g., pedaling) does not interfere with work performance and/or cognitive function (33, 34). Interestingly, slightly slower responses ( $< 1$  second) on mouse performance tests were also observed while performing the seated active workstations compared to sedentary workstation in previous studies although the differences were negligible (17, 33). The equivalent results for work skills between PD and STD in this study would suggest that the PD could be a viable option for occupational health without interfering with work performance.

We recognize the limitations of this study. First, the sample size was small and may have limited detecting impacts on parameters besides insulin. Second, although we asked participants to maintain their usual lifestyle and diet for two days prior to the testing sessions, we did not document those behaviors. However, participants were required to fast for 12 hours, refrain from caffeine and other stimulants, and avoid moderate-vigorous intensity exercise prior to attending the laboratory. Third, we did not adjust the standardized meal quantity based on individual BMI, i.e., a fixed quantity meal was provided. However, we do not believe this was a major limitation given the small variance in BMI. Fourth, we only examined a single 2-hr bout of PD use (without breaks). Fifth, we tested specific work tasks as markers of work performance. We did this to allow comparisons between conditions but recognize these simulated work-related tasks may not fully represent daily routine sedentary office work activities. Although the tests were performed multiple times throughout the 2-hr test trials, the short task durations (4–10 min) may not be sufficient to simulate true work responsibilities. In addition, our sample size would likely not detect small changes in assessments of work skills. Nonetheless, these results support the concept that light intensity physical activity using a PD can positively impact the adverse impact of sedentary work behaviors with apparently minimal impact on work-related skills. Future

studies will need to consider longer exposures to the PD (>2-hr and including breaks to simulate real-world scenarios) and examination of other metabolic indices and additional measures of insulin sensitivity in real workplace environments.

In summary, this pilot study demonstrated that 2 hr of light activity using a PD resulted in lower insulin concentrations following a mixed meal. Our subjects were overweight/obese office workers who reported sedentary lifestyles and had characteristics suggesting increased risk for diabetes and cardiovascular disease. These results have established a proof-of-concept that participants can perform consistent light intensity physical activity at a self-selected cadence without an obvious negative impact on work skills. If this initial metabolic benefit persists with extended use, habitual use of the PD could have a significant impact on the health of sedentary workers. Further controlled studies and feasibility trials are needed to more fully understand the dose-response effects of the PD at varying intensities of pedaling and/or over longer durations among individuals with newly diagnosed, as well as more long-standing diabetes.

## Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

## Acknowledgments


This publication was supported by the CPH-NEW pilot grant program, funded by Grant Number 1 U19 OH008857 from the U.S. National Institute for Occupational Safety and Health (NIOSH). Its contents are solely the responsibility of the authors and do not necessarily represent the official views of NIOSH. The authors gratefully acknowledge assistance from Chris Moore and Scott Ducharme towards the completion of this project.

## References

1. Matthews CE, Chen KY, Freedson PS, Buchowski MS, Beech BM, Pate RR, et al. Amount of time spent in sedentary behaviors in the United States, 2003–2004. *Am J Epidemiol*. 2008; 167(7):875–81. [PubMed: 18303006]
2. Helmerhorst HJ, Wijndaele K, Brage S, Wareham NJ, Ekelund U. Objectively measured sedentary time may predict insulin resistance independent of moderate-and vigorous-intensity physical activity. *Diabetes*. 2009; 58(8):1776–9. [PubMed: 19470610]
3. Mayer-Davis EJ, D'Agostino R Jr, Karter AJ, Haffner SM, Rewers MJ, Saad M, et al. Intensity and amount of physical activity in relation to insulin sensitivity: the Insulin Resistance Atherosclerosis Study. *JAMA*. 1998; 279(9):669–74. Epub 1998/03/13. [PubMed: 9496984]
4. Larsen RN, Kingwell BA, Robinson C, Hammond L, Cerin E, Shaw JE, et al. Breaking up of prolonged sitting over three days sustains, but does not enhance, lowering of postprandial plasma glucose and insulin in overweight and obese adults. *Clin Sci (Lond)*. 2015; 129(2):117–27. Epub 2015/03/04. DOI: 10.1042/cs20140790 [PubMed: 25731923]
5. Church TS, Thomas DM, Tudor-Locke C, Katzmarzyk PT, Earnest CP, Rodarte RQ, et al. Trends over 5 decades in U.S. occupation-related physical activity and their associations with obesity. *PLoS One*. 2011; 6(5):e19657. Epub 2011/06/08. doi: 10.1371/journal.pone.0019657 [PubMed: 21647427]
6. Miller R, Brown W. Steps and sitting in a working population. *Int J Behav Med*. 2004; 11(4):219–24. Epub 2005/01/20. DOI: 10.1207/s15327558ijbm1104\_5 [PubMed: 15657022]
7. Ryan CG, Dall PM, Granat MH, Grant PM. Sitting patterns at work: objective measurement of adherence to current recommendations. *Ergonomics*. 2011; 54(6):531–8.
8. Thorp AA, Healy GN, Winkler E, Clark BK, Gardiner PA, Owen N, et al. Prolonged sedentary time and physical activity in workplace and non-work contexts: a cross-sectional study of office,


- customer service and call centre employees. *Int J Behav Nutr Phys Act.* 2012; 9(1):128. [PubMed: 23101767]
9. Dunstan DW, Kingwell BA, Larsen R, Healy GN, Cerin E, Hamilton MT, et al. Breaking up prolonged sitting reduces postprandial glucose and insulin responses. *Diabetes Care.* 2012; 35(5): 976–83. Epub 2012/03/01. DOI: 10.2337/dc11-1931 [PubMed: 22374636]
  10. Bailey DP, Locke CD. Breaking up prolonged sitting with light-intensity walking improves postprandial glycemia, but breaking up sitting with standing does not. *J Sci Med Sport.* 2015; 18(3):294–8. Epub 2014/04/08. DOI: 10.1016/j.jsams.2014.03.008 [PubMed: 24704421]
  11. Henson J, Davies MJ, Bodicoat DH, Edwardson CL, Gill JM, Stensel DJ, et al. Breaking Up Prolonged Sitting With Standing or Walking Attenuates the Postprandial Metabolic Response in Postmenopausal Women: A Randomized Acute Study. *Diabetes Care.* 2016; 39(1):130–8. Epub 2015/12/03. DOI: 10.2337/dc15-1240 [PubMed: 26628415]
  12. Tudor-Locke C, Schuna JM Jr, Frensham LJ, Proenca M. Changing the way we work: elevating energy expenditure with workstation alternatives. *Int J Obes (Lond).* 2014; 38(6):755–65. Epub 2013/11/29. DOI: 10.1038/ijo.2013.223 [PubMed: 24285335]
  13. MacEwen BT, MacDonald DJ, Burr JF. A systematic review of standing and treadmill desks in the workplace. *Prev Med.* 2015; 70:50–8. Epub 2014/12/03. DOI: 10.1016/j.ypmed.2014.11.011 [PubMed: 25448843]
  14. Stroop JR. Studies of interference in serial verbal reactions. *J Exp Psychol.* 1935; 18(6):643.
  15. John D, Bassett D, Thompson D, Fairbrother J, Baldwin D. Effect of using a treadmill workstation on performance of simulated office work tasks. *J Phys Act Health.* 2009; 6(5):617–24. Epub 2009/12/04. [PubMed: 19953838]
  16. Larson MJ, LeCheminant JD, Hill K, Carbine K, Masterson T, Christenson E. Cognitive and typing outcomes measured simultaneously with slow treadmill walking or sitting: implications for treadmill desks. *PLoS One.* 2015; 10(4):e0121309. Epub 2015/04/16. doi: 10.1371/journal.pone.0121309 [PubMed: 25874910]
  17. Straker L, Levine J, Campbell A. The effects of walking and cycling computer workstations on keyboard and mouse performance. *Hum Factors.* 2009; 51(6):831–44. Epub 2010/04/27. [PubMed: 20415158]
  18. Baumeister RF, Twenge JM, Nuss CK. Effects of social exclusion on cognitive processes: anticipated aloneness reduces intelligent thought. *J Pers Soc Psychol.* 2002; 83(4):817–27. Epub 2002/10/11. [PubMed: 12374437]
  19. Cohen J. *Statistical Power Analysis for the Behavioral Sciences.* Hillsdale, NJ: Lawrence Earlbaum Associates; 1988.
  20. Ainsworth BE, Haskell WL, Herrmann SD, Meckes N, Bassett DR Jr, Tudor-Locke C, et al. 2011 Compendium of Physical Activities: a second update of codes and MET values. *Med Sci Sports Exerc.* 2011; 43(8):1575–81. DOI: 10.1249/MSS.0b013e31821ece12 [PubMed: 21681120]
  21. American Diabetes Association. Diagnosis and classification of diabetes mellitus. *Diabetes Care.* 2013; 36(Supplement 1):S67–S74. [PubMed: 23264425]
  22. Duvivier BM, Schaper NC, Bremers MA, van Crombrugge G, Menheere PP, Kars M, et al. Minimal intensity physical activity (standing and walking) of longer duration improves insulin action and plasma lipids more than shorter periods of moderate to vigorous exercise (cycling) in sedentary subjects when energy expenditure is comparable. *PLoS One.* 2013; 8(2):e55542. Epub 2013/02/19. doi: 10.1371/journal.pone.0055542 [PubMed: 23418444]
  23. Henson J, Dunstan DW, Davies MJ, Yates T. Sedentary behaviour as a new behavioural target in the prevention and treatment of type 2 diabetes. *Diabetes Metab Res Rev.* 2016; 32(Suppl 1):213–20. Epub 2016/01/28. DOI: 10.1002/dmrr.2759 [PubMed: 26813615]
  24. Kahn SE. The relative contributions of insulin resistance and beta-cell dysfunction to the pathophysiology of Type 2 diabetes. *Diabetologia.* 2003; 46(1):3–19. Epub 2003/03/15. DOI: 10.1007/s00125-002-1009-0 [PubMed: 12637977]
  25. Faerch K, Vaag A, Holst JJ, Hansen T, Jorgensen T, Borch-Johnsen K. Natural history of insulin sensitivity and insulin secretion in the progression from normal glucose tolerance to impaired fasting glycemia and impaired glucose tolerance: the Inter99 study. *Diabetes Care.* 2009; 32(3): 439–44. Epub 2008/12/06. DOI: 10.2337/dc08-1195 [PubMed: 19056613]

26. Cnop M, Vidal J, Hull RL, Utzschneider KM, Carr DB, Schraw T, et al. Progressive loss of beta-cell function leads to worsening glucose tolerance in first-degree relatives of subjects with type 2 diabetes. *Diabetes Care*. 2007; 30(3):677–82. Epub 2007/03/01. DOI: 10.2337/dc06-1834 [PubMed: 17327340]
27. DeFronzo RA, Abdul-Ghani MA. Preservation of beta-cell function: the key to diabetes prevention. *J Clin Endocrinol Metab*. 2011; 96(8):2354–66. Epub 2011/06/24. DOI: 10.1210/jc.2011-0246 [PubMed: 21697254]
28. Lin P-J, Borer KT. Third Exposure to a Reduced Carbohydrate Meal Lowers Evening Postprandial Insulin and GIP Responses and HOMA-IR Estimate of Insulin Resistance. *PLoS One*. 2016; 11(10):e0165378. [PubMed: 27798656]
29. Skov-Jensen C, Skovbro M, Flint A, Helge JW, Dela F. Contraction-mediated glucose uptake is increased in men with impaired glucose tolerance. *Appl Physiol Nutr Metab*. 2007; 32(1):115–24. Epub 2007/03/03. DOI: 10.1139/h06-098 [PubMed: 17332788]
30. Christ-Roberts CY, Pratipanawatr T, Pratipanawatr W, Berria R, Belfort R, Kashyap S, et al. Exercise training increases glycogen synthase activity and GLUT4 expression but not insulin signaling in overweight nondiabetic and type 2 diabetic subjects. *Metabolism*. 2004; 53(9):1233–42. Epub 2004/08/31. [PubMed: 15334390]
31. Oberlin DJ, Mikus CR, Kearney ML, Hinton PS, Manrique C, Leidy HJ, et al. One bout of exercise alters free-living postprandial glycemia in type 2 diabetes. *Med Sci Sports Exerc*. 2014; 46(2): 232–8. Epub 2013/07/23. DOI: 10.1249/MSS.0b013e3182a54d85 [PubMed: 23872939]
32. van Dijk JW, Venema M, van Mechelen W, Stehouwer CD, Hartgens F, van Loon LJ. Effect of moderate-intensity exercise versus activities of daily living on 24-hour blood glucose homeostasis in male patients with type 2 diabetes. *Diabetes Care*. 2013; 36(11):3448–53. Epub 2013/09/18. DOI: 10.2337/dc12-2620 [PubMed: 24041682]
33. Carr LJ, Maeda H, Luther B, Rider P, Tucker SJ, Leonhard C. Acceptability and effects of a seated active workstation during sedentary work: A proof of concept study. *Int J Workplace Health Manag*. 2014; 7(1):2–15.
34. Pilcher JJ, Baker VC. Task Performance and Meta-Cognitive Outcomes When Using Activity Workstations and Traditional Desks. *Front Psychol*. 2016; 7:957. Epub 2016/07/23. doi: 10.3389/fpsyg.2016.00957 [PubMed: 27445921]



**Figure 1.** Study protocol. STD = standard desk; PD = pedal desk. M1 represents the time point of the measurement 1 including tests for Stroop task and motor speed and accuracy. M2 represents the time point of measurement 2 including tests for typing and reading comprehension.





**Figure 2.**

The effect of PD and STD conditions on postprandial blood glucose concentration (A); glucose AUC (B); postprandial blood insulin concentration (C); insulin AUC (D); postprandial blood free fatty acid concentrations (E); free fatty acid AUC (F). Data are presented as means  $\pm$  standard error of the mean. PD = pedal desk; STD = standard desk. \* represents a significant difference between the conditions at  $p < 0.05$ .

**Table 1**

Participant characteristics and measurements for 2-hr pedaling performance

<b>Characteristic</b>	<b>Mean (SD) or %</b>
Age (yrs)	38.9 (10.9)
Sex	6 women, 6 men
Height (cm)	170.8 (11.9)
Weight (kg)	83.4 (12.4)
BMI (kg/m <sup>2</sup> )	28.7 (3.6)
Weight classification	8 overweight/4 obese
Work hours seated/day (self-reported)	8.7 (2.2)
<b>Pedaling Performance</b>	
Average Total Time in Pedaling (min)	120.3 (1.9)
Cadence (rev/min)	60.50 (10.6)
Power (watts)	40.5 (9.2)
Cadence CV in 1-min epoch (%)	9.3
Power CV in 1-min epoch (%)	13.8

*Note.* CV: coefficient of variation; min: minute; rev/min: revolution per minute; SD: standard deviation; yrs: years

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Table 2

Outcome measures.

Task	Pedal Desk		Standard Desk		P	
	Mean	SD	Mean	SD		
<b>Baseline Concentrations</b>						
Glucose (mg/dL)	113.5	15.8	110.6	13.8	0.31	
Insulin ( $\mu$ U/mL)	10.0	5.4	9.2	4.8	0.32	
Free Fatty Acids (mmol/L)	185	47.6	181	53.8	0.79	
<b>Stroop Test</b>						
<sup>a</sup> Time to Complete (sec)	Black	43.77	7.99	44.52	7.13	0.39
	Congruent	41.50	7.84	43.79	6.86	0.12
	Incongruent	56.31	14.33	55.51	13.05	0.59
<sup>b</sup> Accuracy (%)	Black	96.73	2.54	96.95	2.02	0.33
	Congruent	97.15	2.25	98.26	1.40	0.14
	Incongruent	95.76	4.30	95.82	3.10	0.96
<b>Mouse Proficiency Test</b>						
<sup>a</sup> Time to Complete (sec)	Clicking	59.83	5.66	57.65	4.47	0.01
	Drag-and-Drop	70.64	8.47	69.24	6.51	0.21
<b>Choice Reaction Test</b>						
<sup>a</sup> 1 key RT (ms)		305.11	42.81	293.22	33.27	0.23
<sup>a</sup> 8 keys RT (ms)		573.98	155.10	580.51	136.93	0.69
<b>Typing</b>						
<sup>b</sup> Typing Speed (Net WPM)		49.31	16.23	46.65	13.55	0.15
<sup>b</sup> Typing Accuracy (%)		94.07	5.09	92.50	10.18	0.58
<b>Reading Comprehension</b>						
<sup>a</sup> Reading Speed (sec)		373.42	141.18	395.50	146.72	0.43
<sup>b</sup> Reading Accuracy (%)		54.11	17.54	46.99	15.65	0.58

ms: millisecond; n: number; RT: reaction time; SCWT: Stroop Color and Work Test; SD: standard deviation; sec: second; Net WPM: word per minute excluding any errors made.

high score is considered worse in terms of work performance  
high score is considered better in terms of work performance

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript