



## Botryosphaeriaceae from *Eucalyptus* plantations and adjacent plants in China

G.Q. Li<sup>1</sup>, F.F. Liu<sup>1</sup>, J.Q. Li<sup>1</sup>, Q.L. Liu<sup>1</sup>, S.F. Chen<sup>1,\*</sup>

### Key words

Botryosphaeria  
Cophiniforma  
Lasiodiplodia  
Neofusicoccum  
pathogenicity  
plant pathogen

**Abstract** The Botryosphaeriaceae is a species-rich family that includes pathogens of a wide variety of plants, including species of *Eucalyptus*. Recently, during disease surveys in China, diseased samples associated with species of Botryosphaeriaceae were collected from plantation *Eucalyptus* and other plants, including *Cunninghamia lanceolata*, *Dimocarpus longan*, *Melastoma sanguineum* and *Phoenix hanceana*, which were growing adjacent to *Eucalyptus*. In addition, few samples from *Araucaria cunninghamii* and *Cedrus deodara* in two gardens were also included in this study. Disease symptoms observed mainly included stem canker, shoot and twig blight. In this study, 105 isolates of Botryosphaeriaceae were collected from six provinces, of which 81 isolates were from *Eucalyptus* trees. These isolates were identified based on comparisons of the DNA sequences of the internal transcribed spacer regions and intervening 5.8S nrRNA gene (ITS), and partial translation elongation factor 1-alpha (*tef1*), β-tubulin (*tub*), DNA-directed RNA polymerase II subunit (*rpb2*) and calmodulin (*cmdA*) genes, the nuclear ribosomal large subunit (LSU) and the nuclear ribosomal small subunit (SSU), and combined with their morphological characteristics. Results showed that these isolates represent 12 species of Botryosphaeriaceae, including *Botryosphaeria fusiclora*, *Cophiniforma atrovirens*, *Lasiodiplodia brasiliense*, *L. pseudotheobromae*, *L. theobromae* and *Neofusicoccum parvum*, and six previously undescribed species of *Botryosphaeria* and *Neofusicoccum*, namely *B. pseudoramosa* sp. nov., *B. qingyuanensis* sp. nov., *B. wangensis* sp. nov., *N. hongkongense* sp. nov., *N. microconidium* sp. nov. and *N. sinoeucalypti* sp. nov. Aside from *B. wangensis*, *C. atrovirens* and *N. hongkongense*, the other nine Botryosphaeriaceae species were isolated from *Eucalyptus* trees in South China. *Botryosphaeria fusiclora* (26 % of the isolates from *Eucalyptus*) is the dominant species, followed by *L. pseudotheobromae* (23 % of the isolates from *Eucalyptus*). In addition to species found on *Eucalyptus* trees, we also found *B. pseudoramosa* on *M. sanguineum*; *B. wangensis* on *C. deodara*; *C. atrovirens* on *D. longan*; *L. theobromae* on *C. lanceolata*, *D. longan* and *P. hanceana*; and *N. hongkongense* on *A. cunninghamii*. Pathogenicity tests showed that the 12 species of Botryosphaeriaceae are pathogenic to three *Eucalyptus* clones and that *Lasiodiplodia* species are the most aggressive. The results of our study suggest that many more species of the Botryosphaeriaceae remain to be discovered in China. This study also provides confirmation for the wide host range of Botryosphaeriaceae species on different plants.

**Article info** Received: 30 April 2016; Accepted: 12 September 2017; Published: 31 October 2017.

### INTRODUCTION

The Botryosphaeriaceae includes a range of phylogenetically and morphologically diverse fungi with a broad host range and geographic distribution globally (Punithalingam 1980, Slippers & Wingfield 2007, Liu et al. 2012, Phillips et al. 2013). These fungi occur primarily on woody plants including both economically important crops and native trees (Slippers & Wingfield 2007). Many species of Botryosphaeriaceae are well-known pathogens that can cause stem canker, shoot blight and dieback on woody plants; however, some species of Botryosphaeriaceae have been described as latent pathogens or endophytes that cause disease when the plant is under stress conditions (Slippers & Wingfield 2007).

Species of *Eucalyptus* are widely planted in more than 100 countries, and because of the rapid growth of some *Eucalyptus* trees, they represent one of the most widely planted genera for commercial forestry worldwide, with approximately 20 million hectares (Mha) established in plantations (Iglesias-Trabado et al. 2009). In China, *Eucalyptus* plantations have expanded substantially during the past 30 years, with more than 4.5 Mha

of *Eucalyptus* established in South China by the end of 2013 (Chen & Chen 2013). Industrial *Eucalyptus* plantations in China are typically single species or hybrid plantings, often from a few clones that share a common parentage (Wei 2005, Turnbull 2007, Zhou & Wingfield 2011). The model of large-scale plantations with few clones greatly increases the threat from pests and diseases (Wingfield 2003, Wingfield et al. 2008). In recent years, the sustainable development of *Eucalyptus* plantations in China has been increasingly threatened by pathogens and pests (Zhou & Wingfield 2011). The important diseases in Chinese *Eucalyptus* plantations include stem canker/wilt caused by species of Botryosphaeriaceae (Chen et al. 2011c), *Ceratostysis* (Chen et al. 2013, Liu et al. 2015), *Chrysoporthea* (Chen et al. 2010) and *Teratosphaeria* (Chen et al. 2011a); leaf blight/spot caused by species of *Teratosphaeriaceae* (Burgess et al. 2006), *Mycosphaerellaceae* (Burgess et al. 2007), *Calonectria* (Lombard et al. 2010, Chen et al. 2011b) and *Quambalaria* (Zhou et al. 2007); and bacterial wilt associated with *Ralstonia solanacearum* (Cao 1982, Old et al. 2003).

Relatively little research has been conducted on diseases caused by Botryosphaeriaceae on *Eucalyptus* trees in China (Chen et al. 2011c, Li et al. 2015a). Based on DNA sequence comparisons and morphological features, five species of Botryosphaeriaceae have been identified from *Eucalyptus* in China to date, including *Botryosphaeria fabicerciana* from Fujian,

<sup>1</sup> China Eucalypt Research Centre (CERC), Chinese Academy of Forestry (CAF), ZhanJiang 524022, GuangDong Province, China;  
corresponding author e-mail: shuaifei.chen@gmail.com.



**Fig. 1** Disease symptoms on *Eucalyptus* trees caused by *Botryosphaeriaceae*. a. Typical dieback of a *Eucalyptus grandis* clone in Funjian Province; b. dieback of *Eucalyptus globulus*; c–e. stem cankers and lesions on main stems of different *Eucalyptus* clones/genotypes; f. branch and twig blight of a *Eucalyptus grandis* clone; g. fruiting structures with abundant mature dark conidia on a *Eucalyptus* branch; h. new branches germinated after main stem infection.

GuangXi and HaiNan Provinces, *Lasiodiplodia pseudotheobromae* from GuangXi Province, *L. theobromae* from GuangDong and GuangXi Provinces, *Neofusicoccum parvum* from Fujian and GuangXi Provinces and *N. ribis* s.lat. from Fujian Province (Chen et al. 2011c, Li et al. 2015a). These species were collected from cankered stems and blighted branches or twigs, and pathogenicity tests showed that all five species could produce lesions on *Eucalyptus* seedlings or trees (Chen et al. 2011c, Li et al. 2015a).

In China, species of *Botryosphaeriaceae* also have been isolated from a number of other woody and horticultural plants, including *Acacia confusa* (Zhao et al. 2010), *Actinidia chinensis* (Zhou et al. 2015), *Bougainvillea spectabilis*, *Polyscias balfouriana* (Li et al. 2015a), *Juglans regia* (Li et al. 2015b, Yu et al. 2015), *Malus domestica* (Tang et al. 2012, Xu et al. 2015a), *Rosa rugosa* (Chen et al. 2016), *Vitis vinifera* (Yan et al. 2012, 2013) and *Vaccinium corymbosum* (Xu et al. 2015b). *Botryosphaeriaceae* species identified from these plants resided in *Botryosphaeria*, *Lasiodiplodia* and *Neofusicoccum*. These *Botryosphaeriaceae* were all isolated from diseased tissue of the respective plant hosts.

From 2013–2014, surveys were conducted on *Eucalyptus* in plantations and some plants adjacent to *Eucalyptus*, and diseases with symptoms typical of those caused by *Botryosphaeriaceae* were observed. Diseased samples were collected and the putative *Botryosphaeriaceae* fungi (based on microscopic morphology) were isolated. In addition, few samples previously collected from *Araucaria cunninghamii* and *Cedrus deodara* were also included in this study. The aims of this study are to:

- identify these species of *Botryosphaeriaceae* based on phylogenetic analyses and morphological characteristics;
- clarify the geographic distribution of these *Botryosphaeriaceae* species; and
- evaluate pathogenicity of the identified *Botryosphaeriaceae* species on different *Eucalyptus* clones.

## MATERIALS AND METHODS

### Disease symptoms, sample collection and fungal isolation

Disease surveys were mainly conducted on species of *Eucalyptus* in plantations distributed in Fujian, GuangDong, GuangXi and HaiNan Provinces. Disease symptoms typically caused by *Botryosphaeriaceae* include tree dieback, stem canker, branch canker and twig blight (Fig. 1). Other plants, including *Cunninghamia lanceolata*, *Dimocarpus longan*, *Melastoma sanguineum* and *Phoenix hanceana*, which were growing in close proximity to *Eucalyptus* trees, were also randomly surveyed in this study. These surveys were conducted during 2013–2014. Samples of diseased materials, including stems, branches and twigs that showed typical symptoms of *Botryosphaeriaceae* infection, were collected and taken to the laboratory for fungal isolation. Diseased branches of *C. deodara* in HeNan Province and *A. cunninghamii* in Hong Kong Region with similar symptoms typical of *Botryosphaeriaceae* collected previously, were also added in this study (Fig. 1).

Fungi were isolated from diseased stems, branches and twigs, as well as from pycnidia produced on diseased tissues of *Eucalyptus* and other plants. When pycnidia formed on the surface of diseased tissue, the pycnidia were scratched lightly with a sterile scalpel and transferred with a sterile steel needle to 2 % malt extract agar (MEA) media containing 20 g of malt extract powder (Beijing Shuangxuan Microbial Culture Medium Products Factory, Beijing, China) and 20 g of agar per litre of water (Beijing Solarbio Science & Technology Co., Ltd., Beijing, China) under a stereomicroscope (Carl Zeiss Ltd., Munchen, Germany). For diseased tissues that did not produce pycnidia,

small tissue pieces (approximately 0.25 cm<sup>2</sup>) were cut from inner wood and transferred to 2 % MEA. Pieces of pycnidia and wood were incubated at room temperature for 2–5 d until colonies formed. Colonies with morphological characteristics typical of *Botryosphaeriaceae* were transferred to fresh 2 % MEA plates. Pure cultures were obtained by transferring single hyphal tips from colonies to 2 % MEA. Cultures were deposited in the culture collection of the China Eucalypt Research Centre (CERC), Chinese Academy of Forestry (CAF), ZhanJiang, GuangDong Province, China. Isolates linked to type specimens of the fungal species were deposited in the China General Microbiological Culture Collection Center (CGMCC), Beijing, China. The specimens were deposited in the Collection of Central South Forestry Fungi of China (CSFF), GuangDong Province, China.

### DNA extraction, PCR amplification and sequencing

DNA extractions and sequence comparisons were conducted on selected isolates collected from different trees and different regions (Table 1). For the selected isolates, mycelia were scraped from 7-d-old cultures using sterile scalpels and transferred to 2 mL Eppendorf tubes. A CTAB-based protocol, 'Method 5' described by Van Burik et al. (1998), was used to extract the DNA samples. The resulting DNA was checked for purity and concentration using a NanoDrop 2000 Spectrometer (Thermo Fisher Scientific Inc. Waltham, MA, USA). Prior to PCR amplification, each DNA sample was diluted to approximately 100 ng/µL with DNase/RNase-free ddH<sub>2</sub>O (Sangon Biotech Co., Ltd., Shanghai, China). The internal transcribed spacer (ITS) region was amplified using the primers ITS1/ITS4 (White et al. 1990), a part of the translation elongation factor 1-alpha (*tef1*) gene was amplified using the primers EF1-728F/EF1-986R (Carbone & Kohn 1999) or EF1F/EF2R (Jacobs et al. 2004), a part of the β-tubulin (*tub*) gene was amplified using the primers BT-2a/BT-2b (Glass & Donaldson 1995), a part of DNA-directed RNA polymerase II subunit (*rpb2*) gene was amplified using the primers fRPB2-5F/fRPB2-7cR for *Botryosphaeria* and *Cophinaforma* (Liu et al. 1999), *rpb2*-LasF/*rpb2*-LasR for *Lasiodiplodia* (Cruywagen et al. 2017) and RPB2bot6F/RPB2bot7R for *Neofusicoccum* (Pavlic et al. 2009a, Sakalidis et al. 2011), the nuclear ribosomal large subunit (LSU) region was amplified using the primers LR0R/LR5 (Vilgalys & Hester 1990, Cubeta et al. 1991), the nuclear ribosomal small subunit (SSU) region was amplified using the primers NS1/NS4 (White et al. 1990). For the isolates of *Lasiodiplodia*, a portion of the calmodulin (*cmdA*) gene was amplified using the primers CAL-228F/CAL-737R (Carbone & Kohn 1999). All primers were synthesised by Life Technologies (Thermo Fisher Scientific Inc., Shanghai, China). The PCR mixtures to amplify the ITS, *tef1*, *tub*, *rpb2*, *cmdA*, LSU, SSU regions used the TopTaq™ Master Mix Kit (Qiagen Inc., Hilden, Germany). All amplification reactions consisted of 25 µL TopTaq™ Master Mix (contain 1.25 U TopTaq™ DNA Polymerase, 200 µM of each dNTP and 1.5 mM MgCl<sub>2</sub>), 0.2 mM of each primer and 50 ng template DNA (made up to a total volume of 50 µL with RNase-free water). The amplification conditions consisted of an initial denaturation step at 94 °C for 3 min, 35 cycles of 94 °C for 1 min, 55 °C (except 45 °C for SSU) for 1 min, and 72 °C for 1 min, followed by a final elongation step at 72 °C for 10 min.

PCR amplifications were carried out in a thermocycler (Bio-Rad Laboratories, Inc., Berkeley, California, USA). The PCR products were separated by electrophoresis in 1.5 % agarose gels with SYBR Safe DNA Gel Stain (Thermo Fisher Scientific Inc., USA) in 1× Tris-acetate-EDTA (TAE) buffer at a constant voltage (80 V) for 30 min. All PCR products were sequenced in both directions using the primers specified above by Beijing Genomics Institution, Guangzhou, GuangDong Province, China. The

**Table 1** Isolates sequenced and used for phylogenetic analyses, morphological studies and pathogenicity tests in this study.

Species <sup>1</sup>	Isolate No. <sup>2,3</sup>	Genotype <sup>4</sup>	Host	Location	GPS information	Collector	ITS	tef1	tub	rpb2	cndA	LSU	SSU
<i>Botryosphaeria fusispora</i>	CERC1997	AAAA-AA	<i>Eucalyptus</i> hybrid	Beihai Region, GuangXi Province, China	N21°35'41"E109°43'01"	S.F. Chen & G.Q. Li	KX277967	KX278072	KX278177	MF410116	N/A	MF410007	MF410205
	CERC2273	AAAA-AA	<i>Eucalyptus</i> hybrid	Fuzhou Region, Fujian Province, China	N26°13'39"E119°10'51"	S.F. Chen & G.Q. Li	KX277968	KX278073	KX278178	MF410117	N/A	MF410008	MF410206
<b>CERC2274<sup>6,7</sup></b>	AAAA-AA	<i>Eucalyptus</i> hybrid	<i>Eucalyptus</i> hybrid	Fuzhou Region, Fujian Province, China	N26°13'39"E119°10'51"	S.F. Chen & G.Q. Li	KX277969	KX278074	KX278179	MF410118	N/A	MF410009	MF410207
<b>CERC2910</b>	AAAA-AA	<i>Eucalyptus</i> hybrid	<i>Eucalyptus</i> hybrid	Beihai Region, GuangXi Province, China	Unknown	S.F. Chen & G.Q. Li	KX277970	KX278075	KX278180	MF410119	N/A	MF410010	MF410208
<b>CERC2912</b>	AAAA-AA	<i>Eucalyptus</i> hybrid	<i>Eucalyptus</i> hybrid	Beihai Region, GuangXi Province, China	Unknown	S.F. Chen & G.Q. Li	KX277971	KX278076	KX278181	MF410120	N/A	MF410011	MF410209
<b>CERC2913</b>	AAAA-AA	<i>Eucalyptus</i> hybrid	<i>Eucalyptus</i> hybrid	Beihai Region, GuangXi Province, China	Unknown	S.F. Chen & G.Q. Li	KX277972	KX278077	KX278182	MF410121	N/A	MF410012	MF410210
<b>CERC3441<sup>6</sup></b>	AAAA-AA	<i>Eucalyptus</i> hybrid	<i>Eucalyptus</i> hybrid	Zhanjiang Region, GuangDong Province, China	N20°41'20"E110°01'17"	S.F. Chen & G.Q. Li	KX277974	KX278079	KX278184	MF410123	N/A	MF410014	MF410212
<b>CERC3469</b>	AAAA-AA	<i>Eucalyptus</i> hybrid	<i>Eucalyptus</i> hybrid	Zhanjiang Region, GuangDong Province, China	N20°41'20"E110°01'17"	S.F. Chen & G.Q. Li	KX277975	KX278080	KX278185	MF410124	N/A	MF410015	MF410213
<b>CERC3474</b>	AAAA-AA	<i>Eucalyptus</i> hybrid	<i>Eucalyptus</i> hybrid	Zhanjiang Region, GuangDong Province, China	N20°41'20"E110°01'17"	S.F. Chen & G.Q. Li	KX277976	KX278081	KX278186	MF410125	N/A	MF410016	MF410214
<b>CERC3426</b>	AAA-AB	<i>Eucalyptus</i> hybrid	<i>Eucalyptus</i> hybrid	Beihai Region, GuangXi Province, China	N21°35'49"E109°43'49"	S.F. Chen & G.Q. Li	KX277973	KX278078	KX278183	MF410122	N/A	MF410013	MF410211
<b>CERC1998<sup>7</sup></b>	ABAA-AA	<i>Eucalyptus</i> hybrid	<i>Eucalyptus</i> hybrid	Beihai Region, GuangXi Province, China	N21°35'41"E109°43'01"	S.F. Chen & G.Q. Li	KX277977	KX278082	KX278187	MF410126	N/A	MF410017	MF410215
<b>CERC2006</b>	ABAA-AA	<i>Eucalyptus</i> hybrid	<i>Eucalyptus</i> hybrid	Zhanjiang Region, GuangDong Province, China	N21°15'26"E110°07'00"	S.F. Chen & G.Q. Li	KX277978	KX278083	KX278188	MF410127	N/A	MF410018	MF410216
<b>CERC2911<sup>6</sup></b>	ABAA-AA	<i>Eucalyptus</i> hybrid	<i>Eucalyptus</i> hybrid	Beihai Region, GuangXi Province, China	Unknown	S.F. Chen & G.Q. Li	KX277979	KX278084	KX278189	MF410128	N/A	MF410019	MF410217
<b>CERC2918<sup>6</sup></b>	ABAA-AA	<i>Eucalyptus</i> hybrid	<i>Eucalyptus</i> hybrid	Beihai Region, GuangXi Province, China	N21°35'41"E109°43'01"	S.F. Chen & G.Q. Li	KX277980	KX278085	KX278190	MF410129	N/A	MF410020	MF410218
<b>CERC2921</b>	ABAA-AA	<i>Eucalyptus</i> hybrid	<i>Eucalyptus</i> hybrid	Beihai Region, GuangXi Province, China	N21°35'41"E109°43'01"	S.F. Chen & G.Q. Li	KX277981	KX278086	KX278191	MF410130	N/A	MF410021	MF410219
<b>CERC2925</b>	ABAA-AA	<i>Eucalyptus</i> hybrid	<i>Eucalyptus</i> hybrid	Beihai Region, GuangXi Province, China	N21°35'41"E109°43'01"	S.F. Chen & G.Q. Li	KX277982	KX278087	KX278192	MF410131	N/A	MF410022	MF410220
<b>CERC2948</b>	ABAA-AA	<i>Eucalyptus</i> hybrid	<i>Eucalyptus</i> hybrid	QingYuan Region, GuangDong Province, China	N23°51'44"E113°10'56"	S.F. Chen & G.Q. Li	KX277983	KX278088	KX278193	MF410132	N/A	MF410023	MF410221
<b>CERC2949</b>	ABAA-AA	<i>Eucalyptus</i> hybrid	<i>Eucalyptus</i> hybrid	QingYuan Region, GuangDong Province, China	N23°51'44"E113°10'56"	S.F. Chen & G.Q. Li	KX277984	KX278089	KX278194	MF410133	N/A	MF410024	MF410222
<b>CERC2954</b>	ABAA-AA	<i>Eucalyptus</i> hybrid	<i>Eucalyptus</i> hybrid	QingYuan Region, GuangDong Province, China	N23°51'44"E113°10'56"	S.F. Chen & G.Q. Li	KX277985	KX278090	KX278195	MF410134	N/A	MF410025	MF410223
<b>CERC3446<sup>7</sup></b>	ABAA-AA	<i>Eucalyptus</i> hybrid	<i>Eucalyptus</i> hybrid	Zhanjiang Region, GuangDong Province, China	N20°41'20"E110°01'17"	S.F. Chen & G.Q. Li	KX277986	KX278091	KX278196	MF410135	N/A	MF410026	MF410224
<b>CERC2830<sup>7</sup></b>	ACAA-AA	<i>Eucalyptus</i> hybrid	<i>Eucalyptus</i> hybrid	Beihai Region, GuangXi Province, China	N21°35'41"E109°43'01"	S.F. Chen & G.Q. Li	KX277987	KX278092	KX278197	MF410136	N/A	MF410027	MF410225
<b>CERC1999<sup>7</sup></b>	AAAA-AA	<i>Eucalyptus</i> hybrid	<i>Eucalyptus</i> hybrid	Beihai Region, GuangXi Province, China	N21°35'41"E109°43'01"	S.F. Chen & G.Q. Li	KX277988	KX278093	KX278198	MF410139	N/A	MF410030	MF410226
<b>CERC2001</b>	AAAA-AA	<i>Eucalyptus</i> hybrid	<i>Eucalyptus</i> hybrid	Beihai Region, GuangXi Province, China	N21°35'41"E109°43'01"	S.F. Chen & G.Q. Li	KX277989	KX278094	KX278198	MF410140	N/A	MF410031	MF410229
= CGMCC3.18739 <sup>6,8,9</sup>				Beihai Region, GuangXi Province, China	N21°35'41"E109°43'01"	S.F. Chen & G.Q. Li	KX277990	KX278095	KX278199	MF410141	N/A	MF410032	MF410230
<b>CERC2004<sup>9</sup></b>	AAAA-AA	<i>Eucalyptus</i> hybrid	<i>Eucalyptus</i> hybrid	Beihai Region, GuangXi Province, China	Unknown	S.F. Chen & G.Q. Li	KX277991	KX278096	KX278200	MF410142	N/A	MF410033	MF410231
<b>CERC2019</b>	AAAA-AA	<i>Melastoma sanguineum</i>	<i>Melastoma sanguineum</i>	Zhanjiang Region, GuangDong Province, China	N21°13'24"E110°24'04"	S.F. Chen	KX277992	KX278097	KX278201	MF410143	N/A	MF410034	MF410232
<b>CERC2833</b>	AAAA-AA	<i>M. sanguineum</i>	<i>M. sanguineum</i>	Zhanjiang Region, GuangDong Province, China	N21°13'24"E110°24'04"	S.F. Chen	KX277993	KX278098	KX278202	MF410144	N/A	MF410035	MF410233
<b>CERC2885</b>	AAAA-AA	<i>M. sanguineum</i>	<i>M. sanguineum</i>	Zhanjiang Region, GuangDong Province, China	N21°13'24"E110°24'04"	S.F. Chen	KX277994	KX278099	KX278203	MF410145	N/A	MF410036	MF410234
<b>CERC2887<sup>9</sup></b>	AAAA-AA	<i>M. sanguineum</i>	<i>M. sanguineum</i>	Zhanjiang Region, GuangDong Province, China	N21°13'24"E110°24'04"	S.F. Chen	KX277995	KX278100	KX278204	MF410146	N/A	MF410037	MF410235
<b>CERC2888<sup>6</sup></b>	AAAA-AA	<i>M. sanguineum</i>	<i>M. sanguineum</i>	Zhanjiang Region, GuangDong Province, China	N20°41'20"E110°01'17"	S.F. Chen & G.Q. Li	KX277996	KX278101	KX278205	MF410147	N/A	MF410038	MF410236
<b>CERC3455</b>	AAAA-AA	<i>Eucalyptus</i> hybrid	<i>Eucalyptus</i> hybrid	Zhanjiang Region, GuangDong Province, China	N20°41'20"E110°01'17"	S.F. Chen & G.Q. Li	KX277997	KX278102	KX278206	MF410148	N/A	MF410039	MF410237
<b>CERC3462</b>	AAAA-AA	<i>Eucalyptus</i> hybrid	<i>Eucalyptus</i> hybrid	Zhanjiang Region, GuangDong Province, China	N20°41'20"E110°01'17"	S.F. Chen & G.Q. Li	KX277998	KX278103	KX278207	MF410149	N/A	MF410040	MF410238
<b>CERC3472</b>	AAAA-AA	<i>Eucalyptus</i> hybrid	<i>Eucalyptus</i> hybrid	Zhanjiang Region, GuangDong Province, China	N20°41'20"E110°01'17"	S.F. Chen & G.Q. Li	KX277999	KX278104	KX278208	MF410150	N/A	MF410041	MF410239
<b>CERC2946</b>	AAAA-AA	<i>Eucalyptus</i> hybrid	<i>Eucalyptus</i> hybrid	Zhanjiang Region, GuangDong Province, China	N23°44'30"E112°48'49"	S.F. Chen & G.Q. Li	KX278000	KX278105	KX278209	MF410151	N/A	MF410042	MF410240
<b>CERC3455</b>	AAAA-AA	<i>Eucalyptus</i> hybrid	<i>Eucalyptus</i> hybrid	QingYuan Region, GuangDong Province, China	N23°44'30"E112°48'49"	S.F. Chen & G.Q. Li	KX278001	KX278106	KX278210	MF410152	N/A	MF410043	MF410241
<b>CERC2947</b>	AAAA-AA	<i>Eucalyptus</i> hybrid	<i>Eucalyptus</i> hybrid	QingYuan Region, GuangDong Province, China	N34°04'09."8	S.F. Chen	KX278002	KX278107	KX278211	MF410153	N/A	MF410044	MF410242
<b>CERC298</b>	AAAA-AA	<i>C. deodara</i>	<i>C. deodara</i>	XinZhuang, MangChuan, RuZhou Region, HeNan Province, China	E112°49'00."7	S.F. Chen	KX278003	KX278108	KX278212	MF410154	N/A	MF410045	MF410243
<b>CERC299</b>	AAAA-AA	<i>C. deodara</i>	<i>C. deodara</i>	XinZhuang, MangChuan, RuZhou Region, HeNan Province, China	N34°04'09."8	S.F. Chen	KX278004	KX278109	KX278213	MF410155	N/A	MF410046	MF410244
<b>CERC300</b>	AAAA-AA	<i>C. deodara</i>	<i>C. deodara</i>	XinZhuang, MangChuan, RuZhou Region, HeNan Province, China	N34°04'09."8	S.F. Chen	KX278005	KX278110	KX278214	MF410156	N/A	MF410047	MF410245
<b>CERC3487</b>	AAAA-AA	<i>D. longan</i>	<i>D. longan</i>	Zhanjiang Region, GuangDong Province, China	E112°49'00."7	S.F. Chen	KX278006	KX278111	KX278215	MF410157	N/A	MF410048	MF410246
<b>CERC3488</b>	AAAA-AA	<i>D. longan</i>	<i>D. longan</i>	Zhanjiang Region, GuangDong Province, China	E112°49'00."7	S.F. Chen	KX278007	KX278112	KX278216	MF410158	N/A	MF410049	MF410247
<b>CERC3490</b>	AAAA-AA	<i>D. longan</i>	<i>D. longan</i>	Zhanjiang Region, GuangDong Province, China	E112°49'00."7	S.F. Chen	KX278008	KX278113	KX278217	MF410159	N/A	MF410050	MF410248
<b>CERC2844<sup>6,7</sup></b>	AAAAAA	<i>Eucalyptus</i> hybrid	<i>Eucalyptus</i> hybrid	ZhangZhou Region, GuangDong Province, China	N24°45'06"E117°51'02"	S.F. Chen & G.Q. Li	KX278009	KX278114	KX278218	MF410160	N/A	MF410051	MF410249
<b>Lasiostiploida brasiliense</b>	CERC2845 <sup>6,7</sup>	AAAAAA	<i>Eucalyptus</i> hybrid	ZhangZhou Region, GuangDong Province, China	N22°08'12"E110°12'08"	S.F. Chen & G.Q. Li	KX278010	KX278115	KX278219	MF410163	N/A	MF410054	MF410250
<i>L. pseudothecophorae</i>	CERC2862	AAAAAA	<i>Eucalyptus</i> hybrid	Yulin Region, GuangXi Province, China	N24°45'06"E117°51'02"	S.F. Chen & G.Q. Li	KX278011	KX278116	KX278220	MF410164	N/A	MF410055	MF410251
<b>B. wangensis</b>	CERC2280	AAAAAA	<i>Eucalyptus</i> hybrid	ZhangZhou Region, Fujian Province, China	N24°45'06"E117°51'02"	S.F. Chen & G.Q. Li	KX278012	KX278117	KX278221	MF410165	N/A	MF410056	MF410252
<b>B. qingyuanensis</b>													
<b>Copiphorma atrovirens</b>	CERC2300	AAAA-AA	<i>Dimocarpus longan</i>	XinZhuang, MangChuan, RuZhou Region, HeNan Province, China	N34°04'09."8	S.F. Chen	KX278013	KX278118	KX278224	MF410166	N/A	MF410047	MF410253
	CERC3484	AAAA-AA	<i>D. longan</i>	Zhanjiang Region, GuangDong Province, China	E112°49'00."7	S.F. Chen	KX278014	KX278119	KX278225	MF410167	N/A	MF410048	MF410254
	CERC3489	AAAA-AA	<i>D. longan</i>	Zhanjiang Region, GuangDong Province, China	E112°49'00."7	S.F. Chen	KX278015	KX278120	KX278226	MF410168	N/A	MF410049	MF410247
	CERC3491	AAAA-AA	<i>D. longan</i>	Zhanjiang Region, GuangDong Province, China	E112°49'00."7	S.F. Chen	KX278016	KX278121	KX278227	MF410169	N/A	MF410050	MF410248
	CERC3492	AAAA-AA	<i>D. longan</i>	Zhanjiang Region, GuangDong Province, China	E112°49'00."7	S.F. Chen	KX278017	KX278122	KX278228	MF410170	N/A	MF410051	MF410249
	CERC3493	AAAA-AA	<i>D. longan</i>	Zhanjiang Region, GuangDong Province, China	E112°49'00."7	S.F. Chen	KX278018	KX278123	KX278229	MF410171	N/A	MF410052	MF410250
	CERC3494	AAAA-AA	<i>D. longan</i>	Zhanjiang Region, GuangDong Province, China	E112°49'00."7	S.F. Chen	KX278019	KX278124	KX278230	MF410172	N/A	MF410053	MF410251
	CERC3495	AAAA-AA	<i>D. longan</i>	Zhanjiang Region, GuangDong Province, China	E112°49'00."7	S.F. Chen	KX278020	KX278125	KX278231	MF410173	N/A	MF410054	MF410252
	CERC3496	AAAA-AA	<i>D. longan</i>	Zhanjiang Region, GuangDong Province, China	E112°49'00."7	S.F. Chen	KX278021	KX278126	KX278232	MF410174	N/A	MF410055	MF410253
	CERC3497	AAAA-AA	<i>D. longan</i>	Zhanjiang Region, GuangDong Province, China	E112°49'00."7	S.F. Chen	KX278022	KX278127	KX278233	MF410175	N/A	MF410056	MF410254
	CERC3498	AAAA-AA	<i>D. longan</i>	Zhanjiang Region, GuangDong Province, China	E112°49'00."7	S.F. Chen	KX278023	KX278128	KX278234	MF410176	N/A	MF410057	MF410255
	CERC3499	AAAA-AA	<i>D. longan</i>	Zhanjiang Region, GuangDong Province, China	E112°49'00."7	S.F. Chen	KX278024	KX278129	KX278235	MF410177	N/A	MF410058	MF410256
	CERC3500	AAAA-AA	<i>D. longan</i>	Zhanjiang Region, GuangDong Province, China	E112°49'00."7	S.F. Chen	KX278025	KX278130	KX278236	MF410178	N/A	MF410059	MF410257
	CERC3501	AAAA-AA	<i>D. longan</i>	Zhanjiang Region, GuangDong Province, China	E112°49'00."7								

Table 1 (cont.)

Species <sup>1</sup>	Isolate No. <sup>2,3</sup>	Genotype <sup>4</sup>	Host	Location	GPS information	Collector	GenBank accession No. <sup>5</sup>				
							ITS	tel1	tub	rpD2	cndA
<i>L. pseudotrichocephala</i> (cont.)	CERC2281	AAAAAAA	Eucalyptus hybrid	ZhangZhou Region, Fujian Province, China	N24°46'06"E 117°51'02"	S.F. Chen & G.Q. Li	KX278013	KX278118	KX278222	MF410166	MF410057
	CERC2282	AAAAAAA	Eucalyptus hybrid	ZhangZhou Region, Fujian Province, China	N24°46'06"E 117°51'02"	S.F. Chen & G.Q. Li	KX278014	KX278119	KX278223	MF410167	MF410058
	CERC2283	AAAAAAA	Eucalyptus hybrid	ZhangZhou Region, Fujian Province, China	N24°46'06"E 117°51'02"	S.F. Chen & G.Q. Li	KX278015	KX278120	KX278224	MF410168	MF410256
	<b>CERC2286<sup>6,7</sup></b>	AAAAAAA	Eucalyptus hybrid	ZhangZhou Region, Fujian Province, China	N24°46'06"E 117°51'02"	S.F. Chen & G.Q. Li	KX278016	KX278121	KX278225	MF410169	MF410257
	CERC2287 <sup>6</sup>	AAAAAAA	Eucalyptus hybrid	ZhangZhou Region, Fujian Province, China	N24°46'06"E 117°51'02"	S.F. Chen & G.Q. Li	KX278017	KX278122	KX278226	MF410170	MF410258
	CERC2288	AAAAAAA	Eucalyptus hybrid	ZhangZhou Region, Fujian Province, China	N24°46'06"E 117°51'02"	S.F. Chen & G.Q. Li	KX278018	KX278123	KX278227	MF410171	MF410259
	CERC2289	AAAAAAA	Eucalyptus hybrid	ZhangZhou Region, Fujian Province, China	N24°46'06"E 117°51'02"	S.F. Chen & G.Q. Li	KX278019	KX278124	KX278228	MF410172	MF410260
	CERC2290	AAAAAAA	Eucalyptus hybrid	YunFu Region, GuangDong Province, China	N23°15'12"E 111°14'51"	S.F. Chen & G.Q. Li	KX278020	KX278125	KX278229	MF410173	MF410261
	CERC2291	AAAAAAA	Eucalyptus hybrid	YunFu Region, GuangDong Province, China	N23°15'12"E 111°14'51"	S.F. Chen & G.Q. Li	KX278021	KX278126	KX278230	MF410174	MF410262
	CERC3417 <sup>6</sup>	AAAAAAA	Eucalyptus hybrid	Baihai Region, GuangXi Province, China	N21°35'49"E 109°43'49"	S.F. Chen & G.Q. Li	KX278023	KX278128	KX278237	MF410175	MF410263
<i>L. theobromae</i>	CERC3432 <sup>6</sup>	AAAAAAA	Eucalyptus hybrid	Baihai Region, GuangXi Province, China	N21°35'49"E 109°43'49"	S.F. Chen & G.Q. Li	KX278024	KX278129	KX278233	MF410176	MF410264
	CERC3434	AAAAAAA	Eucalyptus hybrid	Baihai Region, GuangXi Province, China	N21°35'49"E 109°43'49"	S.F. Chen & G.Q. Li	KX278025	KX278130	KX278234	MF410178	MF410265
	CERC3438	AAAAAAA	Eucalyptus hybrid	Baihai Region, GuangXi Province, China	N21°35'49"E 109°43'49"	S.F. Chen & G.Q. Li	KX278026	KX278131	KX278235	MF410179	MF410266
	CERC3475	AAAAAAA	Eucalyptus hybrid	Baihai Region, GuangXi Province, China	N21°35'49"E 109°43'49"	S.F. Chen & G.Q. Li	KX278027	KX278132	KX278236	MF410180	MF410267
	<b>CERC3495<sup>7</sup></b>	AAAAAAA	<i>E. urophylla</i> × <i>E. grandis</i>	ZhanJiang Region, GuangDong Province, China	N21°10'23"1"E 110°23"4"	S.F. Chen & G.Q. Li	KX278028	KX278133	KX278237	MF410181	MF410270
	CERC3496	AAAAAAA	<i>E. urophylla</i> × <i>E. grandis</i>	ZhanJiang Region, GuangDong Province, China	N21°10'23"1"E 110°23"4"	S.F. Chen & G.Q. Li	KX278029	KX278134	KX278238	MF410182	MF410271
	<b>CERC2562</b>	AAAAABA	Eucalyptus hybrid	YunFu Region, GuangDong Province, China	N23°15'12"E 111°14'51"	S.F. Chen & G.Q. Li	KX278030	KX278135	KX278239	MF410175	MF410268
	<b>CERC2024<sup>6</sup></b>	AAAAAAA	<i>Phoenix canescens</i>	ZhanJiang Region, GuangDong Province, China	N21°15'26"E 110°07'01"	S.F. Chen & G.Q. Li	KX278031	KX278136	KX278240	MF410184	MF410269
	<b>CERC3420<sup>6,7</sup></b>	AAAAAAA	Eucalyptus hybrid	Baihai Region, GuangXi Province, China	N21°35'49"E 109°43'49"	S.F. Chen & G.Q. Li	KX278032	KX278137	KX278241	MF410185	MF410273
	CERC3424 <sup>6</sup>	AAAAAAA	<i>E. urophylla</i> × <i>E. grandis</i>	Baihai Region, GuangXi Province, China	N21°35'49"E 109°43'49"	S.F. Chen & G.Q. Li	KX278033	KX278138	KX278242	MF410186	MF410274
<i>L. lanceolata</i>	CERC225	ABAAAAA	<i>P. canescens</i>	ZhanJiang Region, GuangDong Province, China	N22°09'12"E 110°12'08"	S.F. Chen & G.Q. Li	KX278034	KX278139	KX278243	MF410187	MF410275
	CERC2264	ABAAAAA	<i>E. urophylla</i> × <i>E. grandis</i>	Yulin Region, GuangXi Province, China	N26°01'40"E 111°27'11"	S.F. Chen & G.Q. Li	KX278035	KX278140	KX278244	MF410188	MF410276
	CERC2275	ABAAAAA	<i>E. urophylla</i> × <i>E. grandis</i>	DingAn County, Hainan Province, China	N19°36'41"E 110°17'16"	S.F. Chen & G.Q. Li	KX278036	KX278141	KX278245	MF410189	MF410277
	CERC2334	ABAAAAA	Eucalyptus hybrid	ShaoGuan Region, GuangDong Province, China	N24°31'32"E 113°37'40"	S.F. Chen & G.Q. Li	KX278037	KX278142	KX278246	MF410190	MF410278
	CERC2957	ABAAAAA	<i>Cunninghamia lanceolata</i>	ShaoGuan Region, GuangDong Province, China	N24°31'32"E 113°37'40"	S.F. Chen & G.Q. Li	KX278038	KX278143	KX278247	MF410191	MF410279
	CERC2958	ABAAAAA	<i>C. lanceolata</i>	YunFu Region, GuangDong Province, China	N23°15'12"E 111°14'51"	S.F. Chen & G.Q. Li	KX278039	KX278144	KX278248	MF410192	MF410280
	CERC2963	ABAAAAA	Eucalyptus hybrid	Baihai Region, GuangXi Province, China	N21°35'49"E 109°43'49"	S.F. Chen & G.Q. Li	KX278040	KX278145	KX278249	MF410193	MF410281
	CERC3418	ABAAAAA	Eucalyptus hybrid	Baihai Region, GuangXi Province, China	N21°35'49"E 109°43'49"	S.F. Chen & G.Q. Li	KX278041	KX278146	KX278250	MF410194	MF410282
	<b>CERC3422</b>	ABAAAAA	Eucalyptus hybrid	ZhanJiang Region, GuangDong Province, China	Unknown	S.F. Chen	KX278042	KX278147	KX278251	MF410195	MF410283
	CERC3485	ABAAAAA	<i>D. longan</i>	ZhanJiang Region, GuangDong Province, China	Unknown	S.F. Chen	KX278043	KX278148	KX278252	MF410196	MF410284
<i>Neofusicoccum hongkongense</i>	CERC3486	ABAAAAA	<i>D. longan</i>	ZhanJiang Region, GuangDong Province, China	Unknown	S.F. Chen	KX278044	KX278149	KX278253	MF410197	MF410285
	CERC3487	ABAAAAA	<i>D. longan</i>	ZhanJiang Region, GuangDong Province, China	Unknown	S.F. Chen	KX278045	KX278150	KX278254	MF410198	MF410286
	CERC3491	ABAAAAA	<i>D. longan</i>	ZhanJiang Region, GuangDong Province, China	Unknown	S.F. Chen	KX278046	KX278151	KX278255	MF410199	MF410287
	CERC3493	ABAAAAA	<i>D. longan</i>	ZhanJiang Region, GuangDong Province, China	N21°10'33"E 110°23'47"	S.F. Chen & G.Q. Li	KX278047	KX278152	KX278256	MF410200	MF410288
	CERC3513 <sup>8</sup>	ABAAAAA	<i>E. urophylla</i> × <i>E. grandis</i>	ZhanJiang Region, GuangDong Province, China	N21°10'33"E 110°23'47"	S.F. Chen & G.Q. Li	KX278048	KX278153	KX278257	MF410201	MF410289
	CERC3514	ABAAAAA	<i>E. urophylla</i> × <i>E. grandis</i>	ZhanJiang Region, GuangDong Province, China	N21°10'33"E 110°23'47"	S.F. Chen & G.Q. Li	KX278049	KX278154	KX278258	MF410202	MF410290
	<b>CERC3516<sup>7</sup></b>	ABAAA	<i>E. urophylla</i> × <i>E. grandis</i>	Hong Kong, China	Unknown	S.F. Chen	KX278050	KX278155	KX278259	N/A	MF410291
	CERC3867	AABAA-AA	<i>Araucaria cunninghamii</i>	Hong Kong, China	Unknown	S.F. Chen	KX278051	KX278156	KX278260	KX278281	MF410292
	= CGMCC3.18747	AABAA-AA	<i>A. cunninghamii</i>	Hong Kong, China	Unknown	S.F. Chen	KX278052	KX278157	KX278261	N/A	MF410293
	CERC2688	AABAA-AA	<i>E. urophylla</i> × <i>E. grandis</i>	ZhanJiang Region, GuangDong Province, China	N21°13'31"E 110°23'47"	S.F. Chen & G.Q. Li	KX278053	KX278158	KX278263	MF410204	MF410294
<i>N. microconidium</i>	CERC3497	AAAAA-AA	<i>E. urophylla</i> × <i>E. grandis</i>	ZhanJiang Region, GuangDong Province, China	N21°13'31"E 110°23'47"	S.F. Chen & G.Q. Li	KX278054	KX278159	KX278264	N/A	MF410095
	= CGMCC3.18748 <sup>6,7,8,9</sup>	6,7,8,9	<i>E. urophylla</i> × <i>E. grandis</i>	QingYuan Region, GuangDong Province, China	N23°51'44"E 113°10'58"	S.F. Chen & G.Q. Li	KX278055	KX278160	KX278265	N/A	MF410096
	CERC3498	AAAAA-AA	<i>E. urophylla</i> × <i>E. grandis</i>	ZhanJiang Region, GuangDong Province, China	N21°13'31"E 110°23'47"	S.F. Chen & G.Q. Li	KX278056	KX278161	KX278266	N/A	MF410097
	= CGMCC3.18751 <sup>6,7,8,9</sup>	6,7,8,9	<i>E. urophylla</i> × <i>E. grandis</i>	ZhanJiang Region, GuangDong Province, China	N21°13'31"E 110°23'47"	S.F. Chen & G.Q. Li	KX278057	KX278162	KX278267	N/A	MF410098
	N. parvum			ZhanJiang Region, GuangDong Province, China	N21°13'31"E 110°23'47"	S.F. Chen & G.Q. Li	KX278058	KX278163	KX278268	N/A	MF410099

Table 1 (cont.)

Species <sup>1</sup>	Isolate No. <sup>2,3</sup>	Genotype <sup>4</sup>	Host	Location	GPS information	Collector	GenBank accession No. <sup>5</sup>					
							ITS	tef1	tub	rpb2	cmdA	LSU
<b>N. sinoeucalyptii</b>	CERC2005 = CGMCC3.18752 <sup>6,7,8,9</sup>	AAAA-AA	E. urophylla × E. grandis	ZhanJiang Region, GuangDong Province, China	N21°15'26"E110°07'00"	S.F. Chen & G.Q. Li	KX278061	KX278166	KX278290	N/A	MF410105	MF410303
CERC3415 <sup>6</sup>	AAAA-AA	Eucalyptus hybrid	BeiHai Region, GuangXi Province, China	N21°35'49"E109°43'49"	S.F. Chen & G.Q. Li	KX278063	KX278168	KX278292	N/A	MF410107	MF410305	
CERC3416 = CGMCC3.18754 <sup>6</sup>	AAAA-AA	Eucalyptus hybrid	BeiHai Region, GuangXi Province, China	N21°35'49"E109°43'49"	S.F. Chen & G.Q. Li	KX278064	KX278169	KX278293	N/A	MF410108	MF410306	
CERC3457	AAAA-AA	E. urophylla × E. grandis	ZhanJiang Region, GuangDong Province, China	N20°41'20"E110°01'17"	S.F. Chen & G.Q. Li	KX278066	KX278171	KX278295	N/A	MF410110	MF410308	
CERC3458	AAAA-AA	E. urophylla × E. grandis	ZhanJiang Region, GuangDong Province, China	N20°41'20"E110°01'17"	S.F. Chen & G.Q. Li	KX278067	KX278172	KX278296	N/A	MF410111	MF410309	
CERC3463 <sup>7</sup>	AAAA-AA	E. urophylla × E. grandis	ZhanJiang Region, GuangDong Province, China	N20°41'20"E110°01'17"	S.F. Chen & G.Q. Li	KX278068	KX278173	KX278297	N/A	MF410112	MF410310	
CERC3464	AAAA-AA	E. urophylla × E. grandis	ZhanJiang Region, GuangDong Province, China	N20°41'20"E110°01'17"	S.F. Chen & G.Q. Li	KX278069	KX278174	KX278298	N/A	MF410113	MF410311	
CERC3467	AAAA-AA	E. urophylla × E. grandis	ZhanJiang Region, GuangDong Province, China	N20°41'20"E110°01'17"	S.F. Chen & G.Q. Li	KX278070	KX278175	KX278299	N/A	MF410114	MF410312	
CERC3517	AAAA-AA	E. urophylla × E. grandis	ZhanJiang Region, GuangDong Province, China	N21°13'31"E110°23'47"	S.F. Chen & G.Q. Li	KX278071	KX278176	KX278300	N/A	MF410115	MF410313	
CERC2265 = CGMCC3.18753 <sup>6,8</sup>	AAA-A-B	E. urophylla × E. grandis	YuLin Region, GuangXi Province, China	N22°08'55"E110°12'00"	S.F. Chen & G.Q. Li	KX278062	KX278167	KX278291	N/A	MF410106	MF410304	
CERC3451	AAAA-AB	E. urophylla × E. grandis	ZhanJiang Region, GuangDong Province, China	N20°41'20"E110°01'17"	S.F. Chen & G.Q. Li	KX278065	KX278170	KX278274	N/A	MF410109	MF410307	

<sup>1</sup> Species names in bold are novel species described in this study.<sup>2</sup> Isolates in bold are in the phylogenetic trees.<sup>3</sup> CERC: Culture Collection of China Eucalypt Research Centre, Chinese Academy of Forestry, ZhanJiang, GuangDong Province, China; CGMCC: China General Microbiological Culture Collection Center, Beijing, China.<sup>4</sup> Genotype within each identified species, determined by ITS, tef1, tub, rpb2, cmdA, LSU and SSU regions; '-' means not available.<sup>5</sup> ITS, internal transcribed spacer region and intervening 5.8S nrRNA gene; tef1, translation elongation factor 1-alpha; tub, β-tubulin; rpb2, DNA-directed RNA polymerase II subunit; cmdA, calmodulin; LSU, nuclear ribosomal large subunit; SSU, nuclear ribosomal small subunit; N/A = not available.<sup>6</sup> Isolates used for morphological studies.<sup>7</sup> Isolates used for pathogenicity tests on three Eucalyptus clones.<sup>8</sup> Isolates represent ex-type.<sup>9</sup> Isolates used for culture growth studies.

nucleotide sequences were edited with MEGA v. 6.0.5 software (Tamura et al. 2013). Sequences obtained in this study were all deposited in GenBank (<http://www.ncbi.nlm.nih.gov>) (Table 1).

### Phylogenetic analyses

The preliminary identities of the isolates sequenced in this study were obtained by conducting a standard nucleotide BLAST search using the ITS, tef1, tub, rpb2, cmdA, LSU, SSU sequences. The sequences of the ex-type strains that were closely related to the *Botryosphaeriaceae* isolates sequenced in this study were downloaded from NCBI (<http://www.ncbi.nlm.nih.gov>) and used for polygenetic analyses (Table 2). Sequences were aligned using MAFFT online v. 7 (<http://mafft.cbrc.jp/alignment/server/>) (Katoh & Standley 2013), with the iterative refinement method (FFT-NS-i setting). The alignments were further edited manually with MEGA v. 6.0.5 software (Tamura et al. 2013). Resulting alignments and phylogenetic trees for all the datasets were deposited in TreeBASE (<http://treebase.org>).

The BLAST results showed that the isolates collected in this study were grouped in the genera *Botryosphaeria*, *Copiphorma*, *Lasiodiplodia* and *Neofusicoccum*. Phylogenetic analyses were conducted for each of the ITS, tef1, tub, rpb2, cmdA, LSU and SSU datasets for genera *Botryosphaeria/Copiphorma*, *Lasiodiplodia* and *Neofusicoccum*, respectively. As the cmdA sequences are only available for *Lasiodiplodia*, and not for *Botryosphaeria*, *Copiphorma* and *Neofusicoccum*, the analyses for cmdA sequences were only conducted for the genus *Lasiodiplodia*.

Phylogenetic analyses were also conducted for combined datasets, as the LSU and SSU sequences are not available for some of the previously described species of *Botryosphaeria*, *Copiphorma*, *Lasiodiplodia* and *Neofusicoccum*, and the rpb2 sequences are not available for some species of *Botryosphaeria*. The ITS, tef1 and tub sequences were combined for phylogenetic analyses of *Botryosphaeria/Copiphorma* isolates, the ITS, tef1, tub, rpb2 and cmdA sequences were combined for *Lasiodiplodia* isolates, and ITS, tef1, tub and rpb2 sequences were combined for *Neofusicoccum* isolates.

Two phylogenetic analysis methods were used: PAUP v. 4.0b10 (Swofford 2003) for the maximum parsimony (MP) analyses and PhyML v. 3.0 (Guindon et al. 2010) for maximum likelihood (ML) tests. For MP analyses, gaps are treated as a fifth character and the characters are unordered and of equal weight with 1 000 random addition replicates. The equally most parsimonious trees were obtained using the heuristic search function and tree bisection and reconstruction (TBR) as the branch swapping algorithms. MAXTREES were limited to 5 000, and branch lengths of zero were collapsed. A bootstrap analysis (50 % majority rule, 1 000 replicates) was performed to determine the confidence levels of the tree-branching points (Felsenstein 1985). Tree length (TL), consistency index (CI), retention index (RI), rescaled consistency index (RC) and homoplasy index (HI) were used to evaluate the trees (Hillis & Huelsenbeck 1992).

For ML analyses of each dataset, the best models of nucleotide substitution were determined using jModelTest v. 2.1.5 (Darriba et al. 2012). Additional ML parameters in PhyML include the retention of the maximum number of 1 000 trees and the determination of nodal support by non-parametric bootstrapping with 1 000 replicates. All phylogenetic trees were viewed using MEGA v. 6.0.5 (Tamura et al. 2013). *Neofusicoccum parvum* (ATCC 58191) was used as the outgroup taxon for analyses of *Botryosphaeria* and *Copiphorma*; *Botryosphaeria dothidea* (CBS 115476) was used as the outgroup taxon for analyses of *Lasiodiplodia* and *Neofusicoccum* (Table 2).

**Table 2** Isolates from other studies used in the phylogenetic analyses for this study.

Species	Isolate numbers <sup>1</sup>	Host	Location	Collector	ITS	tEF1	tUB	rPB2	cMDA	LSU	SSU	Reference
<i>Botryosphaeria agaves</i>	MFLUCC 11-0125 = CBS 133952 <sup>3</sup>	<i>Agave</i> sp.	Thailand	R. Phookamsak	JX646791	JX646556	JX646841	N/A	N/A	JX646508	JX646252	Liu et al. (2012)
<i>B. australmontanum</i>	MFLUCC 10-0051 CMW 25413 = CBS 121769 <sup>3</sup>	<i>Agave</i> sp. <i>Acacia mellifera</i>	Thailand Namibia	P. Chommutti F.J.J. van der Walt & J. Roux	JX646790	JX646555	JX646840	N/A	N/A	JX646807	JX646824	Liu et al. (2012)
<i>B. corticis</i>	CBS 119047 <sup>3</sup> ATCC 22927	<i>Vaccinium corymbosum</i> <i>Vaccinium</i> sp.	USA USA	P.V. Oudemans R.D. Millholland	DQ299245	EU017539	EU673107	N/A	N/A	EU673244	EU673175	Phillips et al. (2006, 2008); Lazzizza et al. (2008)
<i>B. dothidea</i>	CBS 115476 = CMW 8000 <sup>3</sup>	<i>Prunus</i> sp.	Switzerland	B. Slippers	DQ299247	EU673291	EU673108	N/A	N/A	EU673245	EU673176	Phillips et al. (2006, 2008)
<i>B. fabicerciana</i>	CBS 110302 CMW 27094 = CBS 127193 <sup>3</sup>	<i>Vitis vinifera</i> <i>Eucalyptus</i> sp.	Portugal China	A.J.L. Phillips M.J. Wingfield	AY236949	AY236989	AY236927	EU339577	N/A	AY928047	EU673173	Slippers et al. (2004a); Phillips et al. (2008)
<i>B. fuscispora</i>	CMW 27121 = CBS 127194	<i>Eucalyptus</i> sp.	China	M.J. Wingfield	HQ332197	AY259092	AY573218	EU673106	N/A	EU673243	EU673174	Alves et al. (2004); Phillips et al. (2008)
<i>B. kuwatsukai</i>	MFLUCC 10-0098 <sup>3</sup> MFLUCC 11-0507	<i>Entada</i> sp. <i>Entada</i> sp.	Thailand Thailand	S. Boonmee R. Cheewangkoon	JX646789	JX646854	JX646839	N/A	N/A	JX646506	JX646823	Liu et al. (2012)
<i>B. minutispermatia</i>	CBS 135219 = PG 2 <sup>3</sup>	<i>Malus domestica</i>	China	C.S. Wang	KJ433888	KJ646853	JX646838	N/A	N/A	JX646805	JX646822	Liu et al. (2012)
<i>B. ramosa</i>	LSP 5 GZCC 16-0013 <sup>3</sup> GZCC 16-0014	<i>Pyrus</i> sp. Dead wood Dead wood	China Guizhou, China Guizhou, China	C.S. Wang H.A. Ariyawansa H.A. Ariyawansa T.I. Burgess	KJ433410	KJ433410	N/A	N/A	N/A	N/A	N/A	Xu et al. (2015a)
<i>B. rosaceae</i>	CBS 122069 = CMW 26167 <sup>3</sup>	<i>Eucalyptus camaldulensis</i>	Australia		KJ433417	KJ433417	N/A	N/A	N/A	N/A	N/A	Xu et al. (2015a)
<i>B. scharffii</i>	CGMCC 3.18007 <sup>3</sup> CGMCC 3.18008	<i>Malus</i> sp. <i>Amigdalus</i> sp.	Shandong, China Shandong, China	Y. Zhang & J.Q. Zhang Y. Zhang, J.Q. Zhang & Z.P. Dou	KX197074	KX197094	KX197101	N/A	N/A	KX197083	N/A	Zhou et al. (2017)
<i>B. sinensis</i>	IRAN 159C = CBS 124703 <sup>3</sup>	<i>Mangifera indica</i>	Iran	J. Abdollahzadeh	KX197075	KX197095	KX197102	N/A	N/A	KX197084	N/A	Zhou et al. (2017)
<i>Coelophialiforma atrovirens</i>	CBS 124934 = CMW 22674 <sup>3</sup>	<i>Morus</i> sp. <i>Juglans regia</i>	Iran	J. Abdollahzadeh & A. Javadi	JQ772056	JQ772056	N/A	N/A	N/A	N/A	N/A	Abdollahzadeh et al. (2013)
<i>L. brasiliense</i>	CBS 124702 CGMCC 3.17723 CGMCC 3.17724	<i>Pterocarpus angolensis</i>	Henan, China Henan, China	Z.P. Dou Z.P. Dou	KT343254	KU221233	KX197107	N/A	N/A	KX197090	N/A	Zhou et al. (2016, 2017)
<i>L. caatinguensis</i>	CBS 117445 = CMW 13425	<i>Acacia mangium</i>	South Africa	J. Mehl & J. Roux	KT343256	KU221234	KX197108	N/A	N/A	KU587380	N/A	Zhou et al. (2016, 2017)
<i>L. americana</i>	CBS 117446 = CMW 13429	<i>Eucalyptus hybrid</i>	Venezuela	S. Mohali	FJ888456	FJ888456	N/A	N/A	N/A	N/A	N/A	Mehl et al. (2011)
<i>L. laetipidioides aviciniae</i>	CMW 41467 <sup>3</sup> LAS 199 (DNA)	<i>Acicennia marina</i> <i>Acicennia marina</i>	South Africa South Africa	J.A. Osorio & J. Roux J.A. Osorio & J. Roux	KP860835	KP860680	KP860758	KU587878	N/A	N/A	N/A	Osorio et al. (2017)
<i>L. americana</i>	CERC 1961 = CFCC50065 <sup>3</sup>	<i>Pistachia vera</i>	Arizona, USA	T.J. Michalides	KU587957	KU587947	KU587868	KU587880	N/A	N/A	N/A	Osorio et al. (2017)
<i>L. brasiliense</i>	CMM 4015 <sup>3</sup> CMW 35884	<i>Mangifera indica</i> <i>Adansonia madagascariensis</i>	Brazil	M.W. Marques	KP217058	KP217066	KP217074	KP217075	MF410161	MF409965	MF410052	Chen et al. (2015); This study
<i>L. bruguierae</i>	CMW 41470 <sup>3</sup> CMW 41614	<i>Bruguiera gymnorhiza</i> <i>Bruguiera gymnorhiza</i>	South Africa South Africa	J.A. Osorio & J. Roux J.A. Osorio & J. Roux	JX464063	KU886972	KU887466	KU886935	N/A	N/A	N/A	Netto et al. (2014)
<i>L. caatinguensis</i>	CMM 1325 <sup>3</sup>	<i>Citrus sinensis</i>	Itarema, Ceará, Brazil	I.B.L. Coutinho & J.S. Lima	KT008006	KT154760	KT154767	N/A	N/A	KP860678	KP860757	Coutinho et al. (2017)

Table 2 (cont.)

Species	Isolate numbers <sup>1</sup>	Host	Location	Collector	GenBank accession numbers <sup>2</sup>						Reference	
					ITS	tef1	tub	rpb2	cmrA	LSU	SSU	
<i>L. caatinguensis</i> (cont.)	IBL40	<i>Spondias mombin</i>	Itarema, Ceará, Brazil	J.S. Lima & J.E. Cardoso	KT154762	KT154755	KT154769	N/A	N/A	N/A	N/A	Coutinho et al. (2017)
<i>L. chinensis</i>	CGMCC3.18061 <sup>3</sup> CGMCC3.18066	Unknown <i>Hevea brasiliensis</i>	China	W. He & Z.P. Dou Y. Zhang & Y.P. Zhou	KX499889 KX499899	KX499827 KX499837	KX500002 KX500012	KX499865 KX499874	N/A	N/A	N/A	Dou et al. (2017a) Dou et al. (2017a)
<i>L. citricola</i>	CBS 124707 = IRAN 1522C <sup>3</sup>	<i>Citrus</i> sp.	Iran	J. Abdollahzadeh & A. Javadi	GU945354	GU945340	KU887505	KU886351	KU886760	N/A	N/A	Abdollahzadeh et al. (2010), Cruywagen et al. (2017)
<i>L. citricola</i>	CBS 124706 = IRAN 1521C	<i>Citrus</i> sp.	Iran	A. Shekari	GU945353	GU945339	KU887504	KU886350	KU886759	N/A	N/A	Abdollahzadeh et al. (2010), Cruywagen et al. (2017)
<i>L. crassispora</i>	CBS 118741 = WAC12533 <sup>3</sup>	<i>Santalum album</i>	Kununurra, Australia	T.I. Burgess & B. Dell	DQ103550	EU673303	KU887506	KU886353	KU886761	DQ377901	N/A	Burgess et al. (2006), Phillips et al. (2008), Cruywagen et al. (2017)
<i>L. euphorbiola</i>	CBS 110492 CMM 3609 <sup>3</sup> CMW 3350	Unknown <i>Jatropha curcas</i> <i>Adansonia digitata</i>	Unknown Brazil Botswana	Unknown A.R. Machado & O.L. Pereira	EF622086 KF234543	EF622066 KF226689	EU673134 KF254926	N/A	N/A	EU673251	N/A	Alves et al. (2008), Phillips et al. (2008), Machado et al. (2014)
<i>L. exigua</i>	CBS 137785 <sup>3</sup>	<i>Retama raetam</i>	Tunisia	B.T. Linaldeedu	KU887149	KU887026	KU887455	KU886346	KU886754	N/A	N/A	Cruywagen et al. (2017)
<i>L. gilanensis</i>	BL 184 CBS 124704 = IRAN 1523C <sup>3</sup>	<i>Retama raetam</i> Unknown	Tunisia Iran	B.T. Linaldeedu J. Abdollahzadeh & A. Javadi	KU638317 GU945351	KU638336 GU945342	KU887509 KU887511	KU886355 KU886357	KU886764 KU886765	N/A	N/A	Linaldeedu et al. (2015), Abdollahzadeh et al. (2010), Cruywagen et al. (2017)
<i>L. gonoubiensis</i>	CBS 115812 = CMW 14077 <sup>3</sup>	<i>Syzygium cordatum</i>	South Africa	D. Pavlic	AY639595	DQ103566	DQ458860	KU886359	KU886768	DQ377902	EU673193	Pavlic et al. (2004), Burgess et al. (2006), Phillips et al. (2008), Cruywagen et al. (2017)
<i>L. gravistriata</i>	CBS 116355 = CMW 14078	<i>Syzygium cordatum</i>	South Africa	D. Pavlic	AY639594	DQ103567	EU673126	KU886358	KU886767	EU673252	EU673194	Pavlic et al. (2004), Burgess et al. (2006), Phillips et al. (2008), Cruywagen et al. (2017)
<i>L. hormozganensis</i>	CMM 4564 <sup>3</sup> CMM 4565 = IRAN 1500C <sup>3</sup>	<i>Anacardium humile</i> <i>Anacardium humile</i> <i>Olea</i> sp.	Brazil Brazil Iran	M.S.B. Netto M.S.B. Netto J. Abdollahzadeh & A. Javadi	KT250949 KT250947	KT250950 KT266812	N/A N/A	N/A	N/A	N/A	N/A	Netto et al. (2017), Abdollahzadeh et al. (2010), Cruywagen et al. (2017)
<i>L. hyalina</i>	CBS 124708 = IRAN 1498C CGMCC3.17975 <sup>3</sup> CGMCC3.18383	<i>Mangifera indica</i>	Iran	J. Abdollahzadeh & A. Javadi	GU945356	GU945344	KU887514	KU886360	KU886769	N/A	N/A	Abdollahzadeh et al. (2010), Cruywagen et al. (2017)
<i>L. iheringii</i>	IPB 01 <sup>3</sup> = B 6180	<i>Acacia confusa</i> Unknown tree	China	Y. Zhang & Y.P. Zhou Z.P. Dou & Z.C. Liu	KX499879 KY767661	KX499917 KY751302	KX499952 KY751299	KX499955 KY751296	N/A	N/A	N/A	Dou et al. (2017b)
<i>L. indica</i>	IRAN 1520C <sup>3</sup> IRAN 1502C	<i>Angiospermous tree</i>	India	I.B. Prasher & G. Singh	KM376151	N/A	N/A	N/A	N/A	N/A	N/A	Prasher & Singh (2014)
<i>L. iranensis</i>	CBS 167.28 <sup>3</sup> LAEREP1	<i>Salvadora persica</i> <i>Laeliocattleya</i>	Iran	J. Abdollahzadeh & A. Javadi	GU945348 GU945347	GU945336 GU945335	KU887516 KU887517	KU886363 KU886362	KU886771 KU886772	N/A	N/A	Abdollahzadeh et al. (2010), Cruywagen et al. (2017)
<i>L. laeliocattleyae</i>	MFLUCC.11-0435 = CBS 134112 <sup>3</sup>	<i>Mangifera indica</i>	Italy Repatriador, Peru	C. Sibilia P. Guerrero	KU507487 KU507451	KU507454 KU507451	N/A N/A	N/A	N/A	DQ377892	N/A	Crous et al. (2006), Rodriguez-Salvarez et al. (2017)
<i>L. lignicola</i>	IPB 01 <sup>3</sup> CMM 3833 <sup>3</sup> CBS 124925 = CMW 27801 <sup>3</sup>	<i>Jatropha curcas</i> <i>Terminalia catappa</i>	Brazil Madagascar	J. Roux	JX646797	KU887003	JX646845	KU896364	N/A	JX646814	JX646830	Liu et al. (2012), Cruywagen et al. (2017)
<i>L. macrospora</i>	CBS 122519 = CMW 26162 <sup>3</sup>	<i>Terminalia catappa</i>	Madagascar	J. Roux	FJ900596	FJ900642	KU887519	KU896366	KU886774	N/A	N/A	Machado et al. (2014)
<i>L. margaritacea</i>	CBS 137783 <sup>3</sup>	<i>Quercus ilex</i>	WA, Tunnel Creek Gorge	J. Roux	EU144050	EU144065	KU887520	KU896367	KU886775	KX464354	N/A	Begoude et al. (2010), Cruywagen et al. (2017)
<i>L. mediterranea</i>	CBS 124926 = CMW 27820	<i>Adansonia gibbosa</i>	Italy	B.T. Linaldeedu	KJ638312	KJ638331	KU887521	KU896368	KU886776	N/A	N/A	Linaldeedu et al. (2015)
<i>L. missouriana</i>	CBS 128311 = UCD2193MO <sup>3</sup>	<i>Vitis vinifera</i> <i>Vitis</i> sp. ×	Missouri, USA	S. Serra	HQ288225	HQ288267	HQ288304	KU886369	KU886777	N/A	N/A	Urbez-Torres et al. (2012), Cruywagen et al. (2017)
<i>L. parva</i>	CBS 128312 = UCD2199MO CBS 456.78 <sup>3</sup>	<i>Vitis</i> sp. × <i>Vitis fabruscana</i>	Missouri, USA	K. Striegler & G.M. Leavitt	HQ288226	HQ288268	HQ288305	KU896371	KU886779	N/A	N/A	Urbez-Torres et al. (2012), Cruywagen et al. (2017)
			Colombia	O. Rangel	EF622083	EF622063	KU887523	KU896372	KU886780	KF766362	N/A	Alves et al. (2008), Cruywagen et al. (2017)

Table 2 (cont.)

**Table 2** (cont.)

Species	Isolate numbers <sup>1</sup>	Host	Location	Collector	GenBank accession numbers <sup>2</sup>						Reference	
					ITS	tef1	tub	rpb2	cmrA	LSU	SSU	
<i>N. batangarum</i> (cont.)	CBS 124923 = CMW 28320	<i>Terminalia catappa</i>	Cameroon	D. Begoude & J. Roux	FJ900608	FJ900654	FJ900635	FJ900616	N/A	KX464400	N/A	Begoude et al. (2010), Yang et al. (2017)
<i>N. brasiliense</i>	CMM 1338 <sup>3</sup> CMM 1285	<i>Mangifera indica</i> <i>Mangifera indica</i>	Brazil	M.W. Marques	JX513630	KC794031	N/A	N/A	N/A	N/A	N/A	Marques et al. (2013)
<i>N. buxi</i>	CBS 11675 <sup>3</sup> CBS 113714	<i>Buxus sempervirens</i> <i>Buxus sempervirens</i>	France Sweden	H.A. van der Aa O. Constantinescu	KX464165 KX464164	KX464678 KX464677	N/A	KX464010 KX464009	N/A	KX464406 KX464405	N/A	Yang et al. (2017) Yang et al. (2017)
<i>N. cordaticola</i>	CBS 123634 = CMW 13992 <sup>3</sup>	<i>Syzygium cordatum</i>	South Africa	D. Pavlic	EU821898	EU821868	EU821838	EU821928	N/A	KX464409	N/A	Pavlic et al. (2009b), Yang et al. (2017)
<i>N. cryptoaustrale</i>	CBS 123635 = CMW 14056	<i>Syzygium cordatum</i>	South Africa	D. Pavlic	EU821903	EU821873	EU821843	EU821933	N/A	KX464410	N/A	Pavlic et al. (2009b), Yang et al. (2017)
<i>N. eucalypticola</i>	CMS 23785 = CBS 122813 <sup>3</sup>	<i>Eucalyptus trees</i>	South Africa	H.M. Maleme	FJ752742	FJ752713	FJ752756	KX464014	N/A	KX464416	N/A	Crous et al. (2013), Yang et al. (2017)
<i>N. eucalypticola</i>	CBS 115679 = CMW 6539 <sup>3</sup>	<i>Eucalyptus grandis</i>	Orbost, Victoria, Tidbinbilla, NSW, Australia	M.J. Wingfield	AY615141	AY615133	AY615125	N/A	N/A	KF766368	KF766288	Slippers et al. (2004c, 2013)
<i>N. eucalypticola</i>	CBS 115766 = CMW 6217	<i>Eucalyptus rossii</i>	M.J. Wingfield	AY615143	AY615135	AY615127	N/A	N/A	N/A	N/A	N/A	Slippers et al. (2004c, 2013)
<i>N. eucahyptorum</i>	CBS 115791 = CMW 10125 <sup>3</sup>	<i>Eucalyptus grandis</i>	South Africa	H. Smith	AF283686	AY236891	AY236820	N/A	N/A	N/A	N/A	Smith et al. (2001), Slippers et al. (2004b)
<i>N. grevilleae</i>	CBS 115766 = CPC 16999 <sup>3</sup>	<i>Eucalyptus grandis</i> <i>Grevillea aurea</i>	South Africa Australia	H. Smith P.W. Crous & R.G. Shivas	AF283687 JF951137	AY236892	AY236821	N/A	N/A	N/A	N/A	Smith et al. (2001), Slippers et al. (2004b)
<i>N. hellenicum</i>	CERC1947 = CFCC50067 <sup>3</sup>	<i>Pistachia vera</i>	Thessaloniki, Greece	T.J. Michaelides	KP217053	KP217061	KP217069	N/A	N/A	N/A	N/A	Chen et al. (2015)
<i>N. illicii</i>	CERC1948 = CFCC50068	<i>Pistachia vera</i>	Aghios Mamas, Chalkidiki, Greece	T.J. Michaelides	KP217054	KP217062	KP217070	N/A	N/A	N/A	N/A	Chen et al. (2015)
<i>N. kwambonambicense</i>	CGMCC3-18310 <sup>3</sup> CGMCC3-18311	<i>Illicium verum</i> <i>Illicium verum</i>	Guangxi, China Guangxi, China	L. Wang L. Wang	KY350149 KY350150	KY351756	KY350155 KY350156	N/A	N/A	N/A	N/A	Zhang et al. (2017) Zhang et al. (2017)
<i>N. lumnitzerae</i>	CBS 123639 = CMW 14023 <sup>3</sup>	<i>Syzygium cordatum</i>	South Africa	D. Pavlic	EU821900	EU821870	EU821840	EU821930	N/A	KX464422	N/A	Pavlic et al. (2009b), Yang et al. (2017)
<i>N. luteum</i>	CBS 123641 = CMW 14140	<i>Syzygium cordatum</i>	South Africa	D. Pavlic	EU821919	EU821889	EU821859	EU821949	N/A	KX464424	N/A	Pavlic et al. (2009b), Yang et al. (2017)
<i>N. macroclavatum</i>	CMW 41469 <sup>3</sup> CMW 41228	<i>Lumnitzera racemosa</i>	South Africa	J.A. Osorio & J. Roux	KP860724	KP860801	KU587925	N/A	N/A	N/A	N/A	Osorio et al. (2017)
<i>N. luteum</i>	CBS 562.92 = ATCC 58193 <sup>3</sup>	<i>Actinidia deliciosa</i> , lesion on ripe fruit	South Africa New Zealand	J.A. Osorio & J. Roux S.R. Pennycook	KP860882	KP860725	KP860803	KU587926	N/A	N/A	N/A	Osorio et al. (2017)
<i>N. luteum</i>	CBS 118223 = WAC 12444 <sup>3</sup>	<i>Eucalyptus globulus</i>	Western Australia	T. Burgess	DQ093196	DQ093217	DQ093206	KX464022	N/A	KX464436	N/A	Burgess et al. (2005), Yang et al. (2017)
<i>N. mangiferae</i>	CBS 118531 = CMW 7024 <sup>3</sup>	<i>Mangifera indica</i>	Australia	G.I. Johnson	AY615185	DQ093221	AY615172	N/A	N/A	DQ377920	EU673153	Slippers et al. (2005), Phillips et al. (2008)
<i>N. mangroviorum</i>	CBS 118532 = CMW 7779 <sup>3</sup>	<i>Mangifera indica</i>	Australia	G.I. Johnson	AY615186	DQ093220	AY615173	KX464023	N/A	DQ377921	EU673154	Slippers et al. (2005), Phillips et al. (2008), Yang et al. (2017)
<i>N. mediterraneum</i>	CMW 41365 <sup>3</sup> CMW 42481	<i>Avicennia marina</i> <i>Bruguiera gymnorhiza</i>	South Africa South Africa	J.A. Osorio & J. Roux J.A. Osorio & J. Roux	KP860859	KP860702	KP860779	KU587905	N/A	N/A	N/A	Osorio et al. (2017)
<i>N. nonquaesitum</i>	CBS 121718 = CPC 13137 <sup>3</sup>	<i>Eucalyptus sp.</i>	Greece	P.W. Crous, M.J. Wingfield & A.J.L. Phillips	GU251176	GU251308	GU251836	KX464024	N/A	KX464437	N/A	Crous et al. (2007), Yang et al. (2017)
<i>N. occultum</i>	CBS 128008 = MUCC 227 <sup>3</sup>	<i>Umbellularia californica</i> <i>Vaccinium corymbosum</i>	USA Chile	F.P. Trouillas cv. Elliot	GU251163	GU251295	GU251823	KX464025	N/A	KX464438	N/A	Inderbitzin et al. (2010), Yang et al. (2017)
		<i>Eucalyptus grandis</i> hybrid	Australia	T.I. Burgess	EU301030	EU339509	EU339472	EU339558	N/A	KX464438	N/A	Sakalidis et al. (2011), Yang et al. (2017)

**Table 2** (cont.)

Species	Isolate numbers <sup>1</sup>	Host	Location	Collector	GenBank accession numbers <sup>2</sup>						Reference	
					ITS	tef1	tub	rpb2	cmtA	LSU	SSU	
<i>N. occultatum</i> (cont.)	MUCC 286 = WAC 12395	<i>Eucalyptus pellita</i>	Australia	T.I. Burgess	EU736947	EU339511	EU339474	EU339560	N/A	N/A	N/A	Sakalidis et al. (2011)
<i>N. parvum</i>	ATCC 58191 = CMW 9080 = ICMP 8002	<i>Populus nigra</i>	New Zealand	G.J. Samuels	AY236943	AY236888	AY236917	EU821963	N/A	AY928045	EU673151	Slippers et al. (2004a); Alves et al. (2005); Phillips et al. (2008); Pavlic et al. (2009b)
<i>N. pennatisporum</i>	WAC 13453 = MUC 510 <sup>3</sup>	<i>Allocasuarina fraseriana</i>	Western Australia	K.M. Taylor	EF591925	EF591976	EF591959	N/A	N/A	N/A	N/A	Taylor et al. (2009)
<i>N. pistaciae</i>	CBS 595-76 <sup>3</sup>	<i>Pistacia vera</i>	Greece	D.G. Zachos	KX464676	KX464953	KX464008	N/A	KX464404	N/A	Yang et al. (2017)	
<i>N. pistaciarum</i>	CBS 113083 = CPC 5263 <sup>3</sup>	<i>Pistacia vera</i>	USA	T.J. Michailides	KX464186	KX464712	KX464998	KX464027	N/A	KX464465	N/A	Yang et al. (2017)
	CBS 113084 = CPC 5284	Redwood	USA	T.J. Michailides	KX464187	KX464713	KX464999	KX464028	N/A	KX464466	N/A	Yang et al. (2017)
<i>N. proteaeum</i>	CBS 114176 = STE-U 1775 <sup>3</sup>	<i>Leucadendron salignum</i>	South Africa	S. Denman	AF452539	KX464720	KX465006	KX464029	N/A	JX556245	N/A	Denman et al. (2003); Yang et al. (2017)
	CBS 111200 = CPC 1357	<i>Leucadendron</i> sp.	South Africa	P.W. Crous	KX464193	KX464719	KX465005	N/A	N/A	KX464472	N/A	Yang et al. (2017)
<i>N. ribis</i>	CBS 115475 = CMW 7772 <sup>3</sup>	<i>Ribes</i> sp.	USA	B. Slippers & G. Hudler	AY236935	AY236877	AY236906	EU821958	N/A	AY928044	KF762922	Slippers et al. (2004a); Alves et al. (2005); Pavlic et al. (2009b)
	CBS 12126 = CMW 7054	<i>Ribes rubrum</i>	USA	N.E. Stevens	AF241177	AY236879	AY236908	EU821960	N/A	KX464473	N/A	Slippers et al. (2004a); Pavlic et al. (2009b); Yang et al. (2017)
<i>N. sinense</i>	CGMCC3-3; 8315 <sup>3</sup>	Unknown woody plant	Guizhou, China	J.J. Gan	KY350148	KY817755	KY350154	N/A	N/A	N/A	N/A	Zhang et al. (2017)
<i>N. stellaboliviachiana</i>	CBS 110864 = CPC 4598 <sup>3</sup>	<i>Vitis vinifera</i>	South Africa	F. Halleen	AY343407	AY243348	KX465047	KX464042	N/A	KX464513	N/A	Van Niekerk et al. (2004); Yang et al. (2017)
<i>N. terminaliae</i>	CBS 125253 = CMW 26679 <sup>3</sup>	<i>Terminalia sericea</i>	South Africa	D. Begoude & J. Roux	GG471802	GG471780	KX465052	KX464045	N/A	KX464518	N/A	Begoude (2010); Yang et al. (2017)
	CBS 125264 = CMW 26683	<i>Terminalia sericea</i>	South Africa	D. Begoude & J. Roux	GG471804	GG471782	KX465053	KX464046	N/A	KX464519	N/A	Begoude (2010); Yang et al. (2017)
<i>N. umdonicola</i>	CBS 123645 = CMW 14058 <sup>3</sup>	<i>Syzygium cordatum</i>	South Africa	D. Pavlic	EU821904	EU821874	EU821844	EU821934	N/A	KX464522	N/A	Pavlic et al. (2009b); Yang et al. (2017)
	CBS 123646 = CMW 14060	<i>Syzygium cordatum</i>	South Africa	D. Pavlic	EU821905	EU821875	EU821845	EU821935	N/A	KX464523	N/A	Pavlic et al. (2009b); Yang et al. (2017)
<i>N. ursonum</i>	CMW 24480 = CBS 12811 <sup>3</sup>	<i>Eucalyptus</i> trees	South Africa	H.M. Maleme	F.J752746	F.J752709	KX465056	KX464047	N/A	N/A	N/A	Crouse et al. (2013); Yang et al. (2017)
	CMW 23790	<i>Eucalyptus</i> trees	South Africa	H.M. Maleme	F.J752745	F.J752708	KX465057	N/A	N/A	N/A	N/A	Crouse et al. (2013); Yang et al. (2017)
	CBS 112378 = STE-U 5044 <sup>3</sup>	<i>Vitis vinifera</i>	South Africa	F. Halleen	AV343381	AV343342	KX465058	KX464048	N/A	KX464527	N/A	Phillips et al. (2013); Yang et al. (2017)
	CBS 112977 = STE-U 5041	<i>Vitis vinifera</i>	South Africa	F. Halleen	AV343380	AV343341	KX465059	N/A	N/A	KX464528	N/A	Phillips et al. (2013); Yang et al. (2017)
	CBS 110887 = STE-U 5252 <sup>3</sup>	<i>Vitis vinifera</i>	South Africa	J.M. van Niekerk	AY343383	AY343343	KX465061	KX464049	N/A	KX464530	N/A	Van Niekerk et al. (2004); Yang et al. (2017)
<i>N. viticlavatum</i>	CBS 110880 = STE-U 5050	<i>Vitis vinifera</i>	South Africa	J.M. van Niekerk	AY343382	AY343344	KX465008	N/A	N/A	KX464475	N/A	Van Niekerk et al. (2004); Yang et al. (2017)

<sup>1</sup> ALG: Personal culture collection A. Berarf-Tebbaj; ATCC: American Type Culture Collection, Virginia, USA; BL: Personal number of B.T. Linakbeddu; CAA: Personal culture collection Artur Alves, Universidade de Aveiro, Portugal; CBS: CBS-KNAW Fungal Biodiversity Centre, Utrecht, The Netherlands; CERC: Culture collection of China Eucalypt Research Centre, Chinese Academy of Forestry, Zhanjiang, GuangDong, China; CGMCC: China General Microbiological Culture Collection Center, Beijing, China; CMM: Culture collection of Phyllopathogenic Fungi Prof. Maria Menezes, Universidade Federal Rural de Pernambuco, Recife, Brazil; GZCC: Guizhou Academy of Agricultural Sciences Culture Collection, GuiZhou, China; IB: Personal culture collection, I.B.L. Coutinho; IPB: Personal culture collection, I.B.P. Brasheer; ICMP: International Collection of Microorganisms from Plants, Auckland, New Zealand; IRAN: Iranian Fungal Culture Collection, Chiang Rai, Thailand; MUCC: Culture collection of Murdoch University, Perth, Australia; STE-U: Culture collection of the Department of Plant Pathology, University of California, Davis, Plant Pathology Department Culture Collection; WAC: Department of Agriculture, Western Australia Plant Pathogen Collection, South Perth, Western Australia.  
<sup>2</sup> ITS, internal transcribed spacer region and intervening 5.8S rRNA gene; tef1, translation elongation factor 1-alpha; tub, beta-tubulin; rpb2, DNA-directed RNA polymerase II subunit; cmtA, calmodulin; LSU, nuclear ribosomal large subunit; SSU, nuclear ribosomal small subunit; N/A = not available.  
<sup>3</sup> Isolates represent ex-type or are from samples that have been linked morphologically to type materials of the species.

## Morphology

Representative isolates for each genotype of *Botryosphaeriaceae* species identified by DNA sequence comparisons were selected for morphological study. To induce sporulation, selected isolates were transferred to 2 % water agar (WA) media (20 g of agar per litre of water) with double-sterilised pine needles placed on the surface of the media (Smith et al. 1996). These cultures were incubated at 25 °C under near-ultraviolet light for 4–6 wk. Conidia in the pycnidia were mounted in one drop of 80 % lactic acid on glass slides and examined under a stereomicroscope (Carl Zeiss Ltd., Munchen, Germany). Conidia and other structures were examined and recorded using a Zeiss Axio Imager A1 microscope and a Zeiss AxioCam MRc digital camera with Zeiss Axio Vision v. 4.8 software (Carl Zeiss Ltd.). Measurements of conidiomata, conidiophores and conidiogenous cells were made to determine the smallest and the largest values. For the isolates selected as a holotype, the lengths and widths of 100 conidia per isolate were measured, as well as 25 measurements of the remaining isolates of each taxon. Average (mean), standard deviation (SD), minimum (min) and maximum (max) measurements are presented as (min–(mean–SD)–(mean+SD)–max). The average length/average width ratio (L/W) of the conidial measurements was calculated. Colony morphology was characterised by cultures grown on 2 % MEA for 7 d and colony colour was determined using the colour charts of Rayner (1970). For growth studies, a 5-mm-diam plug from the growing margin of 7-d-old colonies of each representative isolate was placed in the centre of 90-mm-diam Petri plates containing 2 % MEA. These cultures were incubated in the dark at 5 °C intervals from 5–40 °C. Five replicate plates of each isolate at each temperature were conducted. Two diameter measurements, orthogonally, were recorded daily until the fastest growing culture reached the edge of the Petri plate. The experiment was repeated once and the average for each of the eight temperatures was calculated.

## Pathogenicity tests

To determine the pathogenicity of the identified species on *Eucalyptus* seedlings, representative isolates of all *Botryosphaeriaceae* species identified in this study were selected to inoculate on *Eucalyptus* seedlings. Three *Eucalyptus* clones, CEPT-11 (*Eucalyptus urophylla* × *E. grandis*), CEPT-12 (*E. urophylla*) and CEPT-13 (*E. urophylla* × *E. tereticornis*), were used for inoculations. The *Eucalyptus* seedlings were 1-yr-old, approximately 1.7 m in height, and had a 2.0 cm diam at the root collar. For each clone, 10 seedlings were inoculated with each isolate. On each inoculated seedling, a 5-mm-diam wound was made on the tree stem using a cork borer to remove the bark and expose the xylem. The wounds are located approximately 30 cm above the root collar. For inoculation, 5-mm-diam plugs of mycelia from the margins of colonies grown on 2 % MEA for 7 d in the dark were taken and placed into the wounds with the mycelia facing the cambium. Inoculated wounds were encased with masking tape to prevent contamination and desiccation. Ten seedlings of each *Eucalyptus* clone were inoculated with sterile MEA plugs to serve as negative controls. One month after inoculation, the bark of inoculated seedlings was removed and the internal lesion/wound length on the cambium was measured. The inoculated fungi were re-isolated by cutting small pieces of wood from the edges of the lesions and cultivating them in 2 % MEA at 25 °C. Re-isolations were made from the seedlings inoculated by mycelium plugs and MEA plugs. The data were analysed by one-way analyses of variance (ANOVA) using SAS v. 9.3 (SAS Institute Inc. 2011).

## RESULTS

### Fungal isolation

In this study, 105 isolates from *Eucalyptus* and other plants that show typical morphology of *Botryosphaeriaceae* were isolated. Eighty-one isolates were collected from *Eucalyptus* trees: 12 from FuJian Province, 39 from GuangDong Province, 29 from GuangXi Province and one from HaiNan Province. Eighteen isolates with typical characteristics of *Botryosphaeriaceae* were collected from other plants which were growing in close proximity to *Eucalyptus*: two from *C. lanceolata*, 10 from *D. longan*, four from *M. sanguineum*, and two from *P. hanceana*. In addition, three isolates were collected from *A. cunninghamii* and *C. deodara*, respectively (Table 1).

### Phylogenetic analyses

For all the 105 isolates in this study, ITS, *tef1*, *tub*, *rpb2*, *cmdA*, LSU and SSU sequence data were generated and deposited in GenBank (Table 1). The PCR fragments are approximately 520 bps for the ITS region, 280 bps for the *tef1* region, 430 bps for the *tub* region, 610 bps for the *rpb2* region, 850 bps for the LSU region and 1040 bps for the SSU region. The genotype for each isolate was determined by the ITS, *tef1*, *tub*, *rpb2*, LSU, SSU sequences for isolates in the genera *Botryosphaeria*, *Copriniforma* and *Neofusicoccum*, and by ITS, *tef1*, *tub*, *rpb2*, *cmdA*, LSU, SSU sequences for isolates in the genus *Lasiodiplodia* (Table 1). The preliminary identities of the isolates were determined from conducting a standard nucleotide BLAST with the sequences of ITS, *tef1*, *tub*, *rpb2*, *cmdA*, LSU and SSU, the results consistently showed that the isolates sequenced in this study resided in *Botryosphaeria*, *Copriniforma*, *Lasiodiplodia* or *Neofusicoccum*. One to two isolates of each genotype were selected and used for phylogenetic analyses, depending on the number of isolates of each genotype (Table 1). Based on the comparisons for six to seven region sequences generated in this study and published sequences from ex-type strains of *Botryosphaeriaceae* downloaded from NCBI, sequences of *Botryosphaeria*, *Copriniforma*, *Lasiodiplodia* or *Neofusicoccum* related to species emerging from this study were used for analyses (Table 2). The aligned sequences of each region of ITS, *tef1*, *tub*, *rpb2*, *cmdA*, LSU, SSU, as well as the combined sequences of three to five (*Botryosphaeria/Copriniforma*: three; *Lasiodiplodia*: five; *Neofusicoccum*: four) regions were deposited in TreeBASE (No. 21430). These datasets for genera *Botryosphaeria/Copriniforma*, *Lasiodiplodia* and *Neofusicoccum*, as well as statistical values for the trees for the MP analyses and parameters for the best-fit substitution models of ML analyses, are provided in Table 3.


### Species residing in *Botryosphaeria*

For the isolates grouping in the genus *Botryosphaeria*, isolates clustered into four phylogenetic groups (Groups A–D) for each of the ITS, *tef1*, *tub*, *rpb2* and ITS/*tef1/tub* datasets (Fig. 2a–d, g). For each of the LSU and SSU datasets, Groups A, C and D clustered together (Fig. 2e–f). The ITS sequences of *Botryosphaeria fabicerciana*, *B. fusispora*, *B. kuwatsukai*, *B. roseae* and the six Chinese isolates (CERC2274, CERC2911, CERC2918, CERC2930, CERC3426 and CERC3441) in Group A are consistent, and all of them grouped into one phylogenetic clade (Fig. 2a). For the *tef1* sequence analyses, the isolates in Group A clustered closely to *B. fabicerciana* and *B. fusispora* (Fig. 2b). For the *tub* sequences, the isolates in Group A resided in the same phylogenetic clade with *B. fusispora* (Fig. 2c). For the *rpb2*, LSU and SSU sequences, the isolates in Group A clustered to the same clade with *B. fabicerciana* and *B. fusispora* (*rpb2* is not available to *B. fusispora*) (Fig. 2d–f). The phylogenetic analyses for ITS, *tef1*, *tub*, *rpb2*, LSU and SSU sequences

**Table 3** Datasets used and statistics resulting from phylogenetic analyses.


Genus	Dataset	No. of taxa		No. of bp <sup>1</sup>	Maximum parsimony											
		PIC <sup>2</sup>	No. of trees		Tree length	CI <sup>3</sup>	RI <sup>4</sup>	RC <sup>5</sup>	HI <sup>6</sup>							
<i>Botryosphaeria/Copriniforma</i>																
<i>Lasiodiplodia</i>	ITS	45	543	42	18	0.8209	0.9506	0.7804	0.1791							
	<i>tef1</i>	45	356	147	875	0.8932	0.9677	0.8643	0.1068							
	<i>tub</i>	34	415	43	8	0.9138	0.9688	0.8852	0.0862							
	<i>rpb2</i>	22	718	92	2	0.9397	0.9809	0.9217	0.0603							
	LSU	34	847	21	3	0.9583	0.9885	0.9454	0.0417							
	SSU	31	1024	9	342	9	1.0000	1.0000	1.0000							
	ITS/ <i>tef1/tub</i>	45	1314	232	5000	337	0.8665	0.9555	0.8305	0.1335						
	ITS	76	526	48	5000	87	0.6897	0.8945	0.6169	0.3103						
	<i>tef1</i>	75	332	142	852	402	0.6219	0.9067	0.5639	0.3781						
	<i>tub</i>	67	409	41	5000	60	0.7667	0.9352	0.7170	0.2333						
<i>Neofusicoccum</i>	<i>rpb2</i>	57	532	104	861	190	0.6421	0.8761	0.5626	0.3579						
	<i>cmdA</i>	45	521	74	496	91	0.9121	0.9776	0.8916	0.0879						
	LSU	28	835	20	3	27	0.7407	0.9247	0.6850	0.2593						
	SSU	19	1020	17	2	23	0.8261	0.9048	0.7474	0.1739						
	ITS/ <i>tef1/tub/rpb2/cmdA</i>	76	2320	409	5000	948	0.5918	0.8713	0.5156	0.4082						
	ITS	77	532	81	1404	187	0.5615	0.8707	0.4889	0.4385						
	<i>tef1</i>	75	307	147	5000	303	0.7426	0.9321	0.6922	0.2574						
	<i>tub</i>	75	424	71	1430	141	0.6170	0.8784	0.5420	0.3830						
	<i>rpb2</i>	55	607	114	652	191	0.7173	0.9069	0.6505	0.2827						
	LSU	55	841	33	3260	70	0.5714	0.8324	0.4757	0.4286						
<i>Genus</i>	SSU	21	1027	3	1	3	1.0000	1.0000	1.0000	0.3700						
	ITS/ <i>tef1/tub/rpb2</i>	77	1870	413	336	884	0.6267	0.8824	0.5530	0.3733						
	Maximum likelihood															
	Dataset															
Genus	Dataset	Subst. model <sup>7</sup>		NST <sup>8</sup>		Rate matrix		Ti/Tv ratio <sup>9</sup>		p-inv		Gamma		Rates		
		ITS		TN+I		1.4207		1.0000		8.3891		0.8010		—		
<i>Botryosphaeria/Copriniforma</i>		<i>tef1</i>		TPM2uf+G		4.6965		1.7386		4.6965		—		0.4470		
<i>Lasiodiplodia</i>		<i>tub</i>		HKY+G		—		—		4.1414		—		0.0220		
<i>Neofusicoccum</i>		<i>rpb2</i>		TN+G		3.1864		1.0000		10.4238		—		0.2610		
<i>Genus</i>		LSU		TN+I		5.1213		1.0000		14.2668		—		0.8440		
<i>Genus</i>		SSU		TM2		0.2353		1.0000		3.3084		—		—		
<i>Genus</i>		ITS/ <i>tef1/tub</i>		TN+G		0.397		0.2353		5.9498		—		0.0970		
<i>Genus</i>		ITS		TPM1uf+I+G		3.2977		1.0000		3.1185		8.3075		—		
<i>Genus</i>		<i>tef1</i>		TN+G		8.3075		3.1185		3.1234		1.0000		5.5149		
<i>Genus</i>		<i>tub</i>		6		1.0000		3.6014		3.8909		1.0000		2.6761		
<i>Genus</i>		<i>rpb2</i>		TM3+G		2.6761		1.0000		4.8566		1.0000		13.8753		
<i>Genus</i>		<i>cmdA</i>		TN+G		—		—		—		—		2.7918		
<i>Genus</i>		LSU		TN+I		1.0000		7.7385		1.0000		16.1138		—		
<i>Genus</i>		SSU		TN+I		3.1234		1.0000		15.5731		—		0.7400		
<i>Genus</i>		ITS/ <i>tef1/tub/rpb2/cmdA</i>		TN3+I+G		3.2898		1.0000		0.6986		5.4586		—		
<i>Genus</i>		ITS		TM1+I+G		11.6895		3.1944		3.1944		22.131		—		
<i>Genus</i>		<i>tef1</i>		HKY+G		2		—		—		2.8135		—		
<i>Genus</i>		<i>tub</i>		TN+G		1.0000		4.0352		1.0000		7.8348		—		
<i>Genus</i>		<i>rpb2</i>		TM3+G		6		1.9463		7.0524		1.0000		19.4804		
<i>Genus</i>		LSU		TN+I		6		1.0000		6.0800		1.0000		25.6908		
<i>Genus</i>		SSU		TN		6		1.0000		0.9096		1.0000		7.9272		
<i>Genus</i>		ITS/ <i>tef1/tub/rpb2</i>		TN+I+G		6		4.8874		1.0000		1.0000		9.1711		


<sup>1</sup> bp = base pairs.  
<sup>2</sup> PIC = number of parsimony informative characters.  
<sup>3</sup> CI = consistency index.  
<sup>4</sup> RI = retention index.  
<sup>5</sup> RC = rescaled consistency index.  
<sup>6</sup> HI = homoplasy index.  
<sup>7</sup> Subst. model = best fit substitution model.  
<sup>8</sup> NST = number of substitution rate categories.  
<sup>9</sup> Ti/Tv ratio = transition/transversion ratio.



**Fig. 2** Phylogenetic trees based on maximum likelihood (ML) analyses for species in *Botryosphaeria* and *Cophiniforma*. a. ITS region; b. tef1 gene region; c. tub gene region; d. rpb2 gene region; e. LSU region; f. SSU region; g. combination of ITS, tef1 and tub regions. Isolates sequenced in this study are in **bold**. Bootstrap support values  $\geq 60\%$  for ML and MP are presented above branches as follows: ML/MP, bootstrap support values  $< 60\%$  are marked with ‘-’, and absent ( $< 50\%$ ) are marked with ‘\*’. Isolates representing ex-type sequences are marked with ‘T’. *Neofusicoccum parvum* (ATCC 58191) was used as the outgroup taxon.


Fig. 2 (cont.)






**Fig. 3** Phylogenetic trees based on maximum likelihood (ML) analyses for species in *Lasiodiplodia*. a. ITS region; b. tef1 gene region; c. tub gene region; d. rpb2 gene region; e. cmdA gene region; f. LSU region; g. SSU region; h. combination of ITS, tef1, tub, rpb2 and cmdA regions. Isolates sequenced in this study are in **bold**. Bootstrap support values  $\geq 60\%$  for ML and MP are presented above branches as follows: ML/MP, bootstrap values  $< 60\%$  are marked with ‘-’, and absent ( $< 50\%$ ) are marked with ‘\*\*’. Isolates representing ex-type sequences are marked with ‘T’. *Botryosphaeria dothidea* (CBS 115476) was used as the outgroup taxon.


Fig. 3 (cont.)





**Fig. 4** Phylogenetic trees based on maximum likelihood (ML) analyses for species in *Neofusicoccum*. a. ITS region; b. *tef1* gene region; c. *tub* gene region; d. *rpb2* gene region; e. LSU region; f. SSU region; g. combination of ITS, *tef1*, *tub* and *rpb2* regions. Isolates sequenced in this study are in **bold**. Bootstrap support values  $\geq 60\%$  for ML and MP are presented above branches as follows: ML/MP, bootstrap support values < 60 % are marked with '–', and absent (< 50 %) are marked with '\*'. Isolates representing ex-type sequences are marked with 'T'. *Botryosphaeria dothidea* (CBS 115476) was used as the outgroup taxon.

Fig. 4 (cont.)



showed that the six Chinese isolates in Group A are most closely related to *B. fabicerciana* and *B. fusispora* (Fig. 2a–f). The analyses of the combination of ITS, *tef1* and *tub* sequences indicated that the six isolates are not forming a well-resolved clade, but are phylogenetically more closely related to *B. fusispora* than to *B. fabicerciana* (Fig. 2g). Based on the phylogenetic analyses for ITS, *tef1*, *tub*, *rpb2*, LSU, SSU and the combination of the ITS, *tef1* and *tub* sequences, the six isolates were identified as *B. fusispora*.

Isolates in Group B (CERC2001, CERC2983 and CERC3452) and Group C (CERC2946 and CERC2947) were found to be consistently distinct from other known phylogenetically related species of *Botryosphaeria* by congruent distinction in the multiple datasets (Group B: ITS, *tef1*, *rpb2* and LSU datasets; Group C: ITS, *tef1* and *rpb2* datasets) (Fig. 2a–f). The analyses of the combination of ITS, *tef1* and *tub* sequences indicated that the isolates in Group B and Group C form two well-resolved clades supported by relatively high bootstrap values (Fig. 2g). Isolates in Groups B and C represent two previously undescribed species of *Botryosphaeria*.

The phylogenetic analyses based on ITS, *tef1*, *tub*, *rpb2*, LSU and SSU sequences consistently showed that three isolates (CERC2298, CERC2299 and CERC2300) in Group D were phylogenetically most closely related to *B. auasmontanum*, *B. dothidea*, *B. minutispermatia* and *B. sinensis* (Fig. 2a–f). The analyses of combined ITS, *tef1* and *tub* sequences showed that isolates in Group D form one well-resolved clade (Fig. 2g). Isolates in Group D were identified as a new species of *Botryosphaeria*.

#### *Species residing in Cophiniforma*

The BLAST results for ITS sequences show that isolates CERC3482, CERC3484, CERC3489 and CERC3490 (Group E) are related to the genus *Cophiniforma*. Only two species of *Cophiniforma* have previously been described, *Cophiniforma atrovirens* (Mehl et al. 2011, Phillips et al. 2013) and *C. mamane* (Gardner 1997, Phillips et al. 2013). The two species of *Cophiniforma* are morphologically very similar, but can be distinguished based on ITS sequence data. BLAST results of the ITS sequences indicate that the four Chinese isolates are more closely related to *C. atrovirens* than to *C. mamane*. A BLAST search of the *tef1* sequences show that the Chinese isolates and the ex-type isolate of *C. atrovirens* (CBS 124934) are identical. *Cophiniforma mamane* does not have a *tef1* sequence and cultures are not available (Phillips et al. 2013). Based on the sequence comparisons of the ITS and *tef1* regions (*tub* gene sequences are not available for species of *Cophiniforma*), isolates in Group E were identified as *C. atrovirens* (Fig. 2a–b, g).

#### *Species residing in Lasiodiplodia*

The isolates in our study that clustered in the genus *Lasiodiplodia* grouped into three phylogenetic groups for the *tef1* dataset (Group F: CERC2284; Group G: CERC2024, CERC3420, CERC3513, CERC3516; Group H: CERC2286, CERC2962, CERC3495) (Fig. 3b), and two clades (Group F = Group G; Group H) for the ITS, *tub*, *rpb2*, *cmdA*, LSU and SSU datasets (Fig. 3a, c–g). For Group F, the sequence analyses of the ITS, *tef1*, *tub*, *rpb2*, *cmdA* datasets showed that the Chinese isolates clustered into the same (ITS, *tef1*, *rpb2* and *cmdA*) clade or close (*tub*) to *L. brasiliense* (LSU and SSU sequences are not available to *L. brasiliense*) (Fig. 3a–g). The analyses indicated that isolates in Group G and Group H clustered into the same (ITS, *tub*, *rpb2*, *cmdA*, LSU and SSU) clade or close (*tef1*) to *L. theobromae* and *L. pseudotheobromae*, respectively (Fig. 3a–g). The analyses of the combination of the ITS, *tef1*, *tub*, *rpb2* and *cmdA* sequences

indicated that the isolates in Groups F, G and H are phylogenetically most closely related to *L. brasiliense*, *L. theobromae* and *L. pseudotheobromae*, respectively (Fig. 3h). Altogether, the results of these phylogenetic analyses identified isolates in Groups F, G and H as *L. brasiliense*, *L. theobromae* and *L. pseudotheobromae*, respectively.

#### *Species residing in Neofusicoccum*

For the Chinese isolates that grouped in the genus *Neofusicoccum*, the isolates in this study clustered into four phylogenetic groups for the ITS, *tef1*, *tub* and *rpb2* datasets (Group I: CERC3497, CERC3498; Group J: CERC2967, CERC2968, CERC2973; Group K: CERC2005, CERC2265, CERC3416, CERC3451; Group L: CERC2951, CERC3503, CERC3504, CERC3508) (Fig. 4a–d). The Chinese *Neofusicoccum* isolates clustered into three groups (Group I, Group J = Group K, Group L) for the LSU dataset, and two groups (Group I, Group J = Group K = Group L) for the SSU dataset (Fig. 4e–f).

Previous studies have shown that phylogenetic analyses of the ITS, *tef1*, *tub* and *rpb2* sequences, especially a combination of the four gene sequences, is an efficient method for species identification in *Neofusicoccum* (Pavlic et al. 2009a, Sakalidis et al. 2011, Osorio et al. 2017). The isolates in each of Group I and J were found to be consistently distinct from other known phylogenetically closely related species of *Neofusicoccum* by congruent distinction in all the ITS, *tef1*, *tub* and *rpb2* datasets (Fig. 4a–d). Isolates in Group K formed a single independent clade that was distinct from any known *Neofusicoccum* species in the *tef1* and *rpb2* datasets (Fig. 4b, d). The analyses of the combination of the ITS, *tef1*, *tub* and *rpb2* sequences indicated that isolates in each of Groups I, J and K formed a well-resolved clade that was distinct from any described *Neofusicoccum* species which are supported by high bootstrap values (Fig. 4g). Therefore, we considered isolates in Groups I, J and K to represent three undescribed species of *Neofusicoccum*.

The Chinese isolates in Group L grouped in the same clade as *N. parvum* based on the ITS, *tub*, *rpb2*, LSU and SSU sequence analyses (Fig. 4a, c–f), and close to *N. parvum* on the *tef1* sequence analysis (Fig. 4b). In the phylogenetic analyses combining four gene regions, isolates in Group L were identified as *N. parvum* (Fig. 4g).

#### *Morphology and taxonomy*

Representative isolates (Table 1, 4) selected for morphological studies produced asexual fruiting structures on pine needles on WA media within 4–6 wk. No sexual structures were observed during the same period of time. For the 12 phylogenetic groups of *Botryosphaeriaceae* which were distinguished by DNA sequences, morphological studies, including culture and conidia characteristics, show that isolates in each of Group A, E, F, G, H and L were morphologically similar to the type specimens linked to it via sequence data, especially in the morphological characterisation of conidia (Table 4), namely *B. fusispora*, *C. atrovirens*, *L. brasiliense*, *L. theobromae*, *L. pseudotheobromae* and *N. parvum*, respectively. For phylogenetic groups B, C, D, I, J and K, morphological differences were observed compared to the phylogenetically most closely related species based on sequence data, and consequently each of the six groups were considered as new species. Based on the phylogenetic analyses and the morphological characteristics, the fungi collected from *Eucalyptus* and other plants in this study represent 12 species of *Botryosphaeriaceae*, including six previously undescribed species. These new species are described as follows.

**Table 4** Conidial measurements of *Botryosphaeriaceae* species examined in this study and comparison with measurements described in previous studies.

Species <sup>1</sup>	Conidial size ( $\mu\text{m}$ ) ( $L \times W$ ) <sup>2</sup>	Mean ( $\mu\text{m}$ ) ( $L \times W$ ) <sup>3</sup>	$L/W^4$	Reference
<i>Botryosphaeria auasmontanum</i>	(8.1–)8.8–11.3(–13) × (2.5–)2.9–3.9(–5)	10.1 × 3.4	3.0	Slippers et al. (2014)
<i>B. corticis</i>	(20.5–)23.5–32.5(–34.5) × (5.0–)5.5–7(–7.5)	28.9 × 6.4	4.5	Phillips et al. (2006)
<i>B. dothidea</i>	(20–)23–27(–30) × 4–5(–6)	26.2 × 5.4	4.9	Slippers et al. (2004a)
<i>B. fabicerciana</i>	(16.5–)19–19.5–24.5(–26) × (4.5–)5–6.5(–7.5)	22.0 × 5.8	3.8	Chen et al. (2011c)
<b><i>B. fusispora</i></b>	<b>(16.5–)19–23.5(–28.5) × 5–6(–8)</b>	<b>21.2 × 5.6</b>	<b>3.8</b>	<b>This study</b>
<i>B. fusispora</i>	16–22 × 4–5.5	20.0 × 5.0	4.0	Liu et al. (2012)
<i>B. kuwatsukai</i>	(18.5–)20–24.5(–26) × 5–7(–8)	22.3 × 6.2	3.6	Xu et al. (2015a)
<i>B. minutispermatia</i>	8–14 × 3–4	13.0 × 3.5	3.7	Ariyawansa et al. (2016)
<b><i>B. pseudoramosa</i><sup>5</sup></b>	<b>(8–)10–13(–16) × (4–)4.5–5(–6)</b>	<b>11.5 × 4.6</b>	<b>2.5</b>	<b>This study</b>
<b><i>B. qingyuanensis</i><sup>5</sup></b>	<b>(15–)19.5–24.5(–28.5) × (5–)6–6.5(–7.5)</b>	<b>22.0 × 6.2</b>	<b>3.5</b>	<b>This study</b>
<i>B. ramosa</i>	(11–)12–15(–16) × (4.7–)5–6(–7)	13.5 × 5.5	2.3	Pavlic et al. (2008)
<i>B. rosaceae</i>	20–31 × 6–8	26.2 × 6.7	3.9	Zhou et al. (2017)
<i>B. scharifii</i>	(11.5–)13–17(–19) × 4–6.5	15.4 × 5.2	2.7	Abdollahzadeh et al. (2013)
<i>B. sinensis</i>	(15–)19–29 × 5–7	24.3 × 5.9	4.1	Zhou et al. (2016)
<b><i>B. wangensis</i><sup>5</sup></b>	<b>(20.5–)22–26(–29) × (4.5–)5.5–6.5(–7.5)</b>	<b>23.8 × 6.0</b>	<b>3.9</b>	<b>This study</b>
<i>Lasiodiplodia brasiliense</i>	22.7–29.2 × 11.7–17	26.0 × 14.6	1.8	Netto et al. (2014)
<b><i>L. brasiliense</i></b>	<b>(22–)25–27(–28) × (12–)13.5–15(–15.5)</b>	<b>26.0 × 14.4</b>	<b>1.8</b>	<b>This study</b>
<i>L. pseudotheobromae</i>	(22.5–)23.5–32(–33) × (13.5–)14–18(–20)	28.0 × 16.0	1.7	Alves et al. (2008)
<b><i>L. pseudotheobromae</i></b>	<b>(22.5–)24.5–28.5(–31.5) × (12–)13–15(–16)</b>	<b>26.5 × 13.8</b>	<b>1.9</b>	<b>This study</b>
<i>L. theobromae</i>	(19–)21–31(–32.5) × (12–)13–15.5(–18.5)	26.2 × 14.2	1.9	Alves et al. (2008)
<b><i>L. theobromae</i></b>	<b>(21–)24–26.5(–29.5) × (11–)12.5–14(–16)</b>	<b>25.3 × 13.1</b>	<b>1.9</b>	<b>This study</b>
<i>Neofusicoccum algeriense</i>	(14.5–)17–18(–21) × (4.5–)5.5–5.7(–6.5)	17.6 × 5.6	3.1	Berraf-Tebbal et al. (2014)
<i>N. batangarum</i>	(12–)14–17.5(–20) × (4–)4.5–6(–6.5)	15.5 × 5.5	2.9	Begoude et al. (2010)
<i>N. cordatula</i>	18–28 × 4.5–7	23.3 × 5.3	4.3	Pavlic et al. (2009b)
<b><i>N. hongkongense</i><sup>5</sup></b>	<b>(11.5–)13–15.5(–17.5) × (4–)4.5–5(–5.5)</b>	<b>14.1 × 4.7</b>	<b>3.0</b>	<b>This study</b>
<i>N. kwambonambiense</i>	16–28 × 5–8	22.3 × 6.3	3.6	Pavlic et al. (2009b)
<b><i>N. microconidium</i><sup>5</sup></b>	<b>(10–)11.5–13(–14.5) × (4–)4.5–5.5(–6)</b>	<b>12.3 × 5.0</b>	<b>2.5</b>	<b>This study</b>
<i>N. mangiferae</i>	(11–)12–15(–17.5) × 5–6.6	13.6 × 5.4	2.0–2.5	Slippers et al. (2005)
<i>N. occulatum</i>	14–22 × 3.5–7.5	18.3 × 5.2	3.5	Sakalidis et al. (2011)
<i>N. parvum</i>	(12–)13.5–21(–24) × 4–6(–10)	17.1 × 5.5	3.2	Phillips et al. (2013)
<b><i>N. parvum</i></b>	<b>(15.5–)16.5–19(–21) × (4.5–)5–6(–6.5)</b>	<b>17.9 × 5.5</b>	<b>3.3</b>	<b>This study</b>
<i>N. ribis</i>	(16–)19–23(–24) × 5–6(–7)	20.8 × 5.5	3.8	Slippers et al. (2004a)
<i>N. sinense</i>	(15.2–)17.6–20.4(–23) × (6.9–)7.4–8(–9)	18.7 × 7.7	2.4	Zhang et al. (2017)
<b><i>N. sinoeucalypti</i><sup>5</sup></b>	<b>(13–)15–20.5(–25.5) × (4–)5–5.5(–6.5)</b>	<b>17.7 × 5.2</b>	<b>3.4</b>	<b>This study</b>
<i>N. umdonicola</i>	15–23.5 × 4.5–6.5	19.4 × 5.5	3.5	Pavlic et al. (2009b)

<sup>1</sup> Isolates and measurements in **bold** were examined in this study.<sup>2</sup> Minimum–(average – standard deviation)–(average + standard deviation)–maximum or minimum–maximum, L × W = length × width.<sup>3</sup> L × W = average length × average width.<sup>4</sup> L/W = average length/average width.<sup>5</sup> Novel species described in this study.

## TAXONOMY

### *Botryosphaeria pseudoramosa* G.Q. Li & S.F. Chen, sp. nov.

— MycoBank MB822323; Fig. 5

**Etymology.** Named for its phylogenetic resemblance to *B. ramosa*.

**Sexual morph unknown.** *Conidiomata* pycnidial, produced on pine needles on WA within 2–4 wk, globose to ovoid, dark brown to black, up to 698  $\mu\text{m}$  wide, sometimes with a neck up to 1660  $\mu\text{m}$  long, arising from the substrate, covered by hyphal hairs, embedded in needle tissue, semi-immersed to superficial, unilocular, with a central ostiole. *Conidiophores* reduced to conidiogenous cells. *Conidiogenous cells* holoblastic, discrete, hyaline, cylindrical to lageniform, phialidic with periclinal thickening, (10–)11–16(–22.5) × (1–)2–3.5(–4)  $\mu\text{m}$ . *Paraphyses* not seen. *Conidia* hyaline, thin-walled, smooth with granular contents, unicellular, aseptate ellipsoid to fusoid, base subtruncate to bluntly rounded, (8–)10–13(–16) × (4–)4.5–5(–6)  $\mu\text{m}$  (av. = 11.5 × 4.6  $\mu\text{m}$ , n = 100; L/W = 2.5) (Table 4).

**Culture characteristics** — Colonies on MEA have fluffy mycelia with an uneven margin and a few cottony aerial mycelia reaching to the lid of the Petri plate, with an appressed mycelial mat that is sparse to moderately dense. Colony mycelia initially white, becoming smoke grey (21<sup>”f</sup>) to pale mouse grey (15<sup>”d</sup>) at the surface and olivaceous (21<sup>”k</sup>) to iron grey (23<sup>”m</sup>) at the reverse within 10–14 d. Optimal growth temperature is 30 °C, covering the 90 mm plates after 5 d. No growth at 5 °C. After 5 d, colonies at 10 °C, 15 °C, 20 °C, 25 °C, 30 °C, 35 °C and 40 °C reached 17 mm, 20 mm, 64 mm, 80 mm, 87 mm, 33 mm and 8 mm, respectively.

**Specimens examined.** CHINA, GuangXi, from twigs of one *Eucalyptus* tree, fruiting structures induced on needles of *Pinus* sp. on water agar, 24 May 2014, S.F. Chen & G.Q. Li (holotype CSFF2025, culture ex-type CERC2001 = CGMCC3.18739); GuangDong, from twigs of one *Eucalyptus* tree, 24 May 2014, S.F. Chen & G.Q. Li (CSFF2026, culture CERC3455 = CGMCC3.18741); GuangDong, from twigs of one *Melastoma sanguineum* plant, 17 Mar. 2014, S.F. Chen (CSFF2027, culture CERC2983 = CGMCC3.18740).

**Notes** — *Botryosphaeria pseudoramosa* is phylogenetically closely related to *B. ramosa* and *B. scharifii*. *Botryosphaeria pseudoramosa* can be distinguished from *B. ramosa* and *B. scharifii* based on the morphology of their conidia. Conidia of *B. pseudoramosa* (av. 11.5 × 4.6; L/W = 2.5) are smaller than *B. ramosa* (av. 13.5 × 5.5; L/W = 2.3) (Pavlic et al. 2008) and *B. scharifii* (av. 15.4 × 5.2; L/W = 2.7) (Abdollahzadeh et al. 2013) (Table 4).

### *Botryosphaeria qingyuanensis* G.Q. Li & S.F. Chen, sp. nov.

— MycoBank MB822324; Fig. 6

**Etymology.** Named for the QingYuan Region where the fungus was isolated for the first time.

**Sexual morph unknown.** *Conidiomata* pycnidial, produced on pine needles on WA within 2–4 wk, solitary, globose to ovoid, dark brown to black, up to 317  $\mu\text{m}$  wide, 229  $\mu\text{m}$  high, embedded in needle tissue, semi-immersed to superficial, unilocular, with a central ostiole. *Conidiophores* absent. *Conidiogenous cells* holoblastic, discrete, hyaline, cylindrical to lageniform, phialidic with periclinal thickening, (7–)7.5–12(–14.5) × (2–)2.5–3.5  $\mu\text{m}$ . *Paraphyses* not seen. *Conidia* hyaline, thin-walled, smooth with

granular contents, unicellular, aseptate narrowly fusiform, base subtruncate to bluntly rounded, (15–)19.5–24.5(–28.5) × (5–)6–6.5(–7.5) µm (av. = 22 × 6.2 µm, n = 100; L/W = 3.5) (Table 4).

**Culture characteristics** — Colonies on MEA have fluffy mycelia with an uneven margin and few cottony aerial mycelia reaching to the lid of the Petri plate, with an appressed mycelial mat that is sparse to moderately dense. Colony mycelia initially white, becoming smoke grey (21<sup>\*\*\*f</sup>) to pale mouse grey (15<sup>\*\*\*d</sup>) at the surface and smoke grey (21<sup>\*\*\*f</sup>) to iron grey (23<sup>\*\*\*k</sup>) at the reverse within 10–14 d. Optimal growth temperature is (25–)30 °C, reaching the edge of the 90 mm plates after 5 d. No growth is observed at 5 °C and 40 °C. After 5 d, colonies at 10 °C, 15 °C, 20 °C, 25 °C, 30 °C and 35 °C reach 14 mm, 22 mm, 52 mm, 73 mm, 74 mm and 12 mm, respectively.

**Specimens examined.** CHINA, GuangDong, from twigs of one *Eucalyptus* tree, fruiting structures induced on needles of *Pinus* sp. on water agar, 4 Dec. 2013, S.F. Chen & G.Q. Li (holotype CSFF2028, culture ex-type CERC2946 = CGMCC3.18742); GuangDong, from twigs of one *Eucalyptus* hybrid tree, fruiting structures induced on needles of *Pinus* sp. on water agar, 4 Dec. 2013, S.F. Chen & G.Q. Li (CSFF2029, culture CERC2947 = CGMCC3.18743).


**Notes** — *Botryosphaeria qingyuanensis* is phylogenetically closely related to *B. corticis*, *B. fabicerciana*, *B. fusispora*, *B. kuwatsukai* and *B. rosaceae*, but can be distinguished from these species based on morphological or growth characteristics. Conidia of *B. qingyuanensis* (av. 22 × 6.2; L/W = 3.5) are wider than these of *B. fabicerciana* (av. 22 × 5.8; L/W = 3.8) and the optimal growth temperature of *B. qingyuanensis* ((25–)30 °C) is different from that of *B. fabicerciana* (25(–30) °C) (Chen et al. 2011c). Conidia of *B. qingyuanensis* are longer and wider than *B. fusispora* (av. 20 × 5; L/W = 4) (Liu et al. 2012). Conidia of *B. qingyuanensis* are smaller than *B. corticis* (av. 28.9 × 6.4;

L/W = 4.5) (Phillips et al. 2006) and *B. rosaceae* (av. 26.2 × 6.7; L/W = 3.9) (Zhou et al. 2017). Conidia of *B. qingyuanensis* are slightly shorter than *B. kuwatsukai* (av. 22.3 × 6.2; L/W = 3.6) (Xu et al. 2015a) and no conidia or microconidia are observed for *B. qingyuanensis*, but conidia with 1–3 septa before germination and microconidia (3–8 × 1–2 µm) have been found for *B. kuwatsukai* (Xu et al. 2015a) (Table 4).

***Botryosphaeria wangensis* G.Q. Li & S.F. Chen, sp. nov.** — MycoBank MB822325; Fig. 7

**Etymology.** Named after the Wang village where the fungus was isolated for the first time.

**Sexual morph** unknown. **Conidiomata** pycnidial, produced on pine needles on WA within 2–4 wk, solitary, globose to ovoid, dark brown to black, up to 698 µm wide, 484 µm high, embedded in needle tissue, semi-immersed to superficial, unilocular, with a central ostiole, exuding conidia in a yellow mucoid mass. **Conidiophores** absent. **Conidiogenous cells** holoblastic, discrete, hyaline, cylindrical to lageniform, phialidic with periclinal thickening, (6–)8.5–13.5(–15) × 2–3(–3.5) µm. **Paraphyses** not seen. **Conidia** hyaline, thin-walled, smooth with granular contents, unicellular, aseptate, becoming 1-septate before germination, narrowly fusiform, base subtruncate to bluntly rounded, (20.5–)22–26(–29) × (4.5–)5.5–6.5(–7.5) µm (av. = 23.8 × 6 µm, n = 100; L/W = 3.9) (Table 4). **Spermatophores** hyaline, smooth, branched, cylindrical to subcylindrical (Fig. 7f). **Spermatogenous cells** discrete or integrated, hyaline, smooth, cylindrical, producing spermatia on their tips, holoblastic or proliferating via phialides with periclinal thickenings, 6.5–16 × 1.5–2.5 µm. **Spermatia** unicellular, aseptate, hyaline, thin-walled, allantoid to rod-shaped, 3.5–4.5 × 1–1.5 µm, L/W = 2.9.



**Fig. 5** *Botryosphaeria pseudoramosa*. a–b. Conidiomata formed on pine needle culture; c–d. conidiogenous cells and developing conidia; e. conidia; f. living culture after 10 d on 2 % MEA (front). — Scale bars: a–b = 500 µm; c–e = 10 µm; f = 1 cm.

**Culture characteristics** — Colonies on MEA have fluffy mycelia with an uneven margin and a few cottony aerial mycelia reaching to the lid of the Petri plate, with an appressed mycelial mat that is sparse to moderately dense. Colony mycelia initially white, becoming smoke grey (21<sup>\*\*\*</sup>f) to mouse grey (13<sup>\*\*\*</sup>i) at the surface and olivaceous grey (21<sup>\*\*\*</sup>i) to iron grey (23<sup>\*\*\*</sup>k) at the reverse within 10–14 d. Optimal growth temperature is 30 °C, covering the 90 mm plates after 5 d. No growth at 5 °C and 40 °C. After 5 d, colonies at 10 °C, 15 °C, 20 °C, 25 °C, 30 °C and 35 °C reach 15 mm, 21 mm, 50 mm, 69 mm, 89 mm and 24 mm, respectively.

**Specimens examined.** CHINA, HeNan, from twigs of one *Cedrus deodara* tree, fruiting structures induced on needles of *Pinus* sp. on water agar, 26 Nov. 2013, S.F. Chen (holotype CSFF2030, culture ex-type CERC2298 = CGMCC3.18744); HeNan, from twigs of one *Cedrus deodara* tree, 26 Nov. 2013, S.F. Chen (CSFF2031, culture CERC2300 = CGMCC3.18746).

**Notes** — *Botryosphaeria wangensis* is phylogenetically closely related to *B. auasmontanum*, *B. dothidea*, *B. minutispermatica* and *B. sinensis*. *Botryosphaeria wangensis* can be distinguished from its phylogenetically closely related species by the size of their conidia. Conidia of *B. wangensis* (av. 23.8 × 6; L/W = 3.9) are longer and wider than those of *B. auasmontanum* (av. 10.1 × 3.4; L/W = 3) (Slippers et al. 2014) and *B. minutispermatica* (av. 13 × 3.5; L/W = 3.7) (Ariyawansa et al. 2016) and shorter and wider than those of *B. dothidea* (av. 26.2 × 5.4; L/W = 4.9) (Slippers et al. 2004a) and *B. sinensis* (av. 24.3 × 5.9; L/W = 4.1) (Zhou et al. 2016) (Table 4).



**Fig. 6** *Botryosphaeria qingyuanensis*. a. Conidiomata formed on pine needle culture; b–c. conidiogenous cells and developing conidia; d. conidia; e. living culture after 10 d on 2 % MEA (front). — Scale bars: a = 500 µm; b–d = 10 µm; e = 1 cm.

***Neofusicoccum hongkongense* G.Q. Li & S.F. Chen, sp. nov.**

— MycoBank MB822328; Fig. 8

**Etymology.** Named after the Hong Kong Region where it was isolated for the first time.


**Sexual morph unknown.** *Conidiomata* pycnidial, produced on pine needles on WA within 2–4 wk, solitary, globose to ovoid, dark brown to black, up to 694 µm wide, up to 776 µm high, embedded in needle tissue, semi-immersed to superficial, unilocular, with a central ostiole. *Conidiophores* reduced to conidiogenous cells. *Conidiogenous cells* holoblastic, discrete, hyaline, cylindrical, phialidic with periclinal thickening, (9.5–) 12–18.5(–22) × (1.5–)2–2.5(–3) µm. *Paraphyses* not seen. *Conidia* hyaline, thin-walled, smooth with granular contents, unicellular, aseptate narrowly fusiform, base subtruncate to bluntly rounded, (11.5–)13–15.5(–17.5) × (4–)4.5–5(–5.5) µm (av. = 14.1 × 4.7 µm, n = 100; L/W = 3) (Table 4).

**Culture characteristics —** Colonies on MEA have fluffy mycelia with an uneven margin and a few cottony aerial mycelia

reaching to the lid of the Petri plate, with an appressed mycelial mat that is sparse to moderately dense. Colony mycelia initially white, becoming smoke grey (21<sup>\*\*\*</sup>f) to grey olivaceous (21<sup>\*\*\*</sup>b) at the surface and grey olivaceous (21<sup>\*\*\*</sup>b) to olivaceous grey (21<sup>\*\*\*</sup>i) at the reverse within 10–14 d. Optimal growth temperature is 25 °C, covering the 90 mm plates after 5 d. No growth at 5 °C or 40 °C. After 5 d, colonies grown at 10 °C, 15 °C, 20 °C, 25 °C, 30 °C and 35 °C reach 25 mm, 41 mm, 66 mm, 90 mm, 84 mm and 9 mm, respectively.

**Specimens examined.** CHINA, Hong Kong, from twigs of *Araucaria cunninghamii*, fruiting structures induced on needles of *Pinus* sp. on water agar, 11 Mar. 2014, S.F. Chen (holotype CSFF2034, culture ex-type CERC2973 = CGMCC3.18749); Hong Kong, from twigs of *Araucaria cunninghamii*, 11 Mar. 2014, S.F. Chen (CSFF2035, culture CERC2968 = CGMCC3.18748).

**Notes —** Based on phylogenetic analyses, *N. hongkongense* phylogenetically clustered in the *N. parvum*/*N. ribis* species complex. *Neofusicoccum hongkongense* can be distinguished from other species in the *N. parvum*/*N. ribis* complex by the size and shape of their conidia. The conidia of *N. hongkongense*



**Fig. 7** *Botryosphaeria wangensis*. a–b. Conidiomata on pine needle culture; c–d. conidiogenous cells and developing conidia; e. conidia with 1 septum; f. spermatogenous cells; g. spermatia; h. living culture after 10 d on 2 % MEA (front). — Scale bars: a–b = 500 µm; c–g = 10 µm; h = 1 cm.


(av.  $14.1 \times 4.7$ ; L/W = 3) are shorter and narrower than those of *N. algeriense* (av.  $17.6 \times 5.6$ ; L/W = 3.1) (Berraf-Tebbal et al. 2014), *N. batangarum* (av.  $15.5 \times 5.5$ ; L/W = 2.9) (Begoude et al. 2010), *N. cordaticola* (av.  $23.3 \times 5.3$ ; L/W = 4.3) (Pavlic et al. 2009b), *N. kwambonambiense* (av.  $22.3 \times 6.3$ ; L/W = 3.6) (Pavlic et al. 2009b), *N. oculatum* (av.  $18.3 \times 5.2$ ; L/W = 3.5) (Sakalidis et al. 2011), *N. parvum* (av.  $17.1 \times 5.5$ ; L/W = 3.2) (Phillips et al. 2013), *N. ribis* (av.  $20.8 \times 5.5$ ; L/W = 3.8) (Slippers et al. 2004a), *N. sinense* (av.  $18.7 \times 7.7$ ; L/W = 2.4) (Zhang et al. 2017), *N. sinoeucalypti* (av.  $17.7 \times 5.2$ ; L/W = 3.4) (this study) and *N. umdonicola* (av.  $19.4 \times 5.5$ ; L/W = 3.5) (Pavlic et al. 2009b). The conidial size of *N. brasiliense* remains unknown (Marques et al. 2013) (Table 4).

***Neofusicoccum microconidium* G.Q. Li & S.F. Chen, sp. nov.**


— MycoBank MB822326; Fig. 9

**Etymology.** Named for the small conidia of this fungus.

**Sexual morph** unknown. **Conidiomata** pycnidial, produced on pine needles on WA within 2–4 wk, solitary, globose to ovoid, dark brown to black, up to 895 µm wide, 1729 µm high, embedded in needle tissue, semi-immersed to superficial, unilocular, with a central ostiole, exuding conidia in a white mucoid mass. **Conidiophores** reduced to **conidiogenous cells**. **Conidiogenous cells** holoblastic, discrete, hyaline, cylindrical, phialidic with periclinal thickening, (10.5–)12.5–18(–20.5) × (2–)2.5–3(–3.5) µm. **Paraphyses** not seen. **Conidia** hyaline, thin-walled, smooth with granular contents, unicellular, aseptate narrowly fusiform, base subtruncate to bluntly rounded, (10–)11.5–13(–14.5) × (4–)4.5–5.5(–6) µm (av. = 12.3 × 5 µm, n = 100; L/W = 2.5) (Table 4).



**Fig. 8** *Neofusicoccum hongkongense*. a. Conidiomata formed on pine needle culture; b. conidiogenous cells and developing conidia; c. conidiogenous cells; d. conidia; e. living culture after 10 d on 2 % MEA (front). — Scale bars: a = 500 µm; b–d = 10 µm; e = 1 cm.



**Fig. 9** *Neofusicoccum microconidium*. a. Conidiomata formed on pine needle culture; b–c. conidiogenous cells and developing conidia; d. conidia; e. living culture after 10 d on 2 % MEA (front). — Scale bars: a = 500 µm; b–d = 10 µm; e = 1 cm.

**Culture characteristics** — Colonies on MEA have fluffy mycelia with an uneven margin and a few cottony aerial mycelia reaching to the lid of the Petri plate, with an appressed mycelial mat that is sparse to moderately dense. Colony mycelia initially white, becoming pale mouse grey (15<sup>”””d</sup>) to olivaceous grey (21<sup>”””i</sup>) at the surface and olivaceous grey (21<sup>”””i</sup>) to iron grey (23<sup>”””k</sup>) at the reverse within 10–14 d. Optimal growth temperature is 30 °C, reaching the edge of the 90 mm plates after 5 d. No growth at 5 °C. After 5 d, colonies at 10 °C, 15 °C, 20 °C, 25 °C, 30 °C, 35 °C and 40 °C reach 24 mm, 34 mm, 66 mm, 74 mm, 86 mm, 36 mm and 8 mm, respectively.

**Specimens examined.** CHINA, GuangDong, from twigs of *E. urophylla* × *E. grandis*, fruiting structures induced on needles of *Pinus* sp. on water agar, 22 July 2014, S.F. Chen & G.Q. Li (holotype CSFF2032, culture ex-type CERC3497 = CGMCC3.18750); GuangDong, from twigs of *E. urophylla* × *E. grandis*, 22 July 2014, S.F. Chen & G.Q. Li (CSFF2033, culture CERC3498 = CGMCC3.18751).

**Notes** — *Neofusicoccum microconidium* is phylogenetically closely related to *N. mangiferae*. The two species can be distinguished from each other based on conidial morphology. Conidia of *N. microconidium* (av. 12.3 × 5; L/W = 2.5) are smaller than those of *N. mangiferae* (av. 13.6 × 5.4; L/W = 2–2.5) (Slippers et al. 2005) (Table 4).

***Neofusicoccum sinoeucalypti* G.Q. Li & S.F. Chen, sp. nov.** — MycoBank MB822327; Fig. 10

**Etymology.** Named after the host genus *Eucalyptus* from which it was isolated for the first time.

**Sexual morph** solitary, globose to ovoid, dark brown to black, up to 1 007 µm wide, 685 µm high, embedded in needle tissue, semi-immersed to superficial, unilocular, with a central ostiole. **Conidiophores** absent. **Conidiogenous cells** holoblastic, discrete, hyaline, cylindrical to lageniform, phialidic with periclinal thickening, (10–)10.5–11 × 2–3 µm. **Paraphyses** not seen. **Conidia** hyaline, thin-walled, smooth with granular contents, unicellular, aseptate, narrowly fusiform, base subtruncate to bluntly rounded, (13–)15–20.5(–25.5) × (4–)5–5.5(–6.5) µm (av. = 17.7 × 5.2 µm, n = 100; L/W = 3.4). **Spermatophores** hyaline, smooth, cylindrical to subcylindrical. **Spermatogenous cells** discrete or integrated, hyaline, smooth, cylindrical, producing spermatia on their tips, holoblastic or proliferating via phialides with periclinal thickenings, 8.5–15.5 × 1.5–2 µm. **Spermatia** unicellular, aseptate, hyaline, thin-walled, allantoid to rod-shaped, 2.5–4.5 × 1.5 µm, L/W = 2.1.

**Culture characteristics** — Colonies on MEA have fluffy mycelia with an uneven margin and a few cottony aerial mycelia that reach to the lid of the Petri plate, with an appressed mycelial mat that is sparse to moderately dense. Colony mycelia are initially



**Fig. 10** *Neofusicoccum sinoeucalypti*. a. Conidiomata formed on pine needle culture; b. conidiogenous cells and developing conidia; c. immature conidia; d–e. mature conidia with 1–2 septa; f. spermatogenous cells; g. spermatia; h. living culture after 10 d on 2 % MEA (front). — Scale bars: a = 500 µm; b–g = 10 µm; h = 1 cm.

white, becoming pale mouse grey (15<sup>”””d</sup>) to mouse grey (13<sup>”””i</sup>) at the surface and olivaceous buff (21<sup>””d</sup>) to iron grey (23<sup>”””k</sup>) at the reverse within 10–14 d. Optimal growth temperature is 30 °C, reaching the edge of the 90 mm plates after 5 d. No growth at 5 °C or 40 °C. After 5 d, colonies at 10 °C, 15 °C, 20 °C, 25 °C, 30 °C and 35 °C reach 25 mm, 31 mm, 53 mm, 78 mm, 90 mm and 11 mm, respectively.

*Specimens examined.* CHINA, GuangDong, from twigs of *E. urophylla* × *E. grandis*, fruiting structures induced on needles of *Pinus* sp. on water agar, 30 July 2013, S.F. Chen & G.Q. Li (holotype CSFF2036, culture ex-type CERC2005 = CGMCC3.18752); GuangXi, from twigs of *E. urophylla* ×

*E. grandis*, 25 Oct. 2013, S.F. Chen & G.Q. Li (CSFF2037, culture CERC2265 = CGMCC3.18753); GuangXi, from twigs of *Eucalyptus* hybrid, 22 May 2014, S.F. Chen & G.Q. Li (CSFF2038, culture CERC3416 = CGMCC3.18754).

*Notes* — *Neofusicoccum sinoeucalypti* clustered in the *N. parvum*/*N. ribis* species complex. Other species in this complex include *N. algeriense*, *N. batangarum*, *N. brasiliense*, *N. cordaticola*, *N. hongkongense* (this study), *N. kwambonambiense*, *N. occultatum*, *N. parvum*, *N. ribis* and *N. umdonicola*. For these species, except *N. brasiliense* (morphological data not available) (Marques et al. 2013), spermatia have been reported only in *N. sinoeucalypti* and are allantoid to rod-shaped. Conidia of

*N. sinoeucalypti* (av.  $17.7 \times 5.2$ ; L/W = 3.4) are longer and wider than those of *N. hongkongense* (av.  $14.1 \times 4.7$ ; L/W = 3), longer and narrower than those of *N. batangarum* (av.  $15.5 \times 5.5$ ; L/W = 2.9) (Begoude et al. 2010) and *N. parvum* (av.  $17.1 \times 5.5$ ; L/W = 3.2) (Phillips et al. 2013), shorter and narrower than those of *N. cordaticola* (av.  $23.3 \times 5.3$ ; L/W = 4.3) (Pavlic et al. 2009b), *N. kwambonambiense* (av.  $22.3 \times 6.3$ ; L/W = 3.6) (Pavlic et al. 2009b), *N. ribis* (av.  $20.8 \times 5.5$ ; L/W = 3.8) (Slippers et al. 2004a), *N. sinense* (av.  $18.7 \times 7.7$ ; L/W = 2.4) (Zhang et al. 2017) and *N. umdonicola* (av.  $19.4 \times 5.5$ ; L/W = 3.5) (Pavlic et al. 2009b), shorter than those of *N. occulatum* (av.  $18.3 \times 5.2$ ; L/W = 3.5) (Sakalidis et al. 2011), and narrower than those of *N. algeriense* (av.  $17.6 \times 5.6$ ; L/W = 3.1) (Berraf-Tebbal et al. 2014). The optimal growth temperature of *N. sinoeucalypti* (30 °C) is different compared to *N. algeriense* (25 °C) (Berraf-Tebbal et al. 2014), *N. batangarum* (25 °C) (Begoude et al. 2010), *N. brasiliense* (27.7 °C) (Marques et al. 2013), *N. hongkongense* (25 °C) (this study), *N. occulatum* (25 °C) (Sakalidis et al. 2011) and *N. ribis* (25 °C) (Slippers et al. 2004a) (Table 4).

#### Distribution of Botryosphaeriaceae

According to the phylogenetic and morphological analyses of the 105 isolates collected in this study, twelve species of Botryosphaeriaceae were identified from seven hosts in the FuJian, GuangDong, GuangXi, HaiNan and HeNan Provinces

and the Hong Kong Region of China (Fig. 11). These species include *B. fusispora* (21 isolates: all from *Eucalyptus* hybrids), *B. pseudoramosa* (12 isolates: 8 from *Eucalyptus* hybrids, 4 from *M. sanguineum*), *B. qingyuanensis* (2 isolates: both from one *Eucalyptus* hybrid), *B. wangensis* (3 isolates: all from *C. deodara*), *C. atrovirens* (5 isolates: all from *D. longan*), *L. brasiliense* (1 isolate: from a *Eucalyptus* hybrid), *L. pseudotheobromae* (19 isolates: 17 from unknown *Eucalyptus* hybrids, two from *E. urophylla* × *E. grandis*), *L. theobromae* (20 isolates: six from unknown *Eucalyptus* hybrids, five from *E. urophylla* × *E. grandis*, 2 from *C. lanceolata*, 5 from *D. longan*, 2 from *P. hanceana*), *N. hongkongense* (3 isolates: all from *A. cunninghamii*), *N. microconidium* (2 isolates: both from *E. urophylla* × *E. grandis*), *N. parvum* (6 isolates: all from *E. urophylla* × *E. grandis*) and *N. sinoeucalypti* (11 isolates: nine from *E. urophylla* × *E. grandis*, two from *Eucalyptus* hybrids) (Table 1, Fig. 11). The 81 isolates collected from *Eucalyptus* trees include nine species (except for *B. wangensis*, *C. atrovirens* and *N. hongkongense*) of Botryosphaeriaceae. Of these nine species from *Eucalyptus*, *B. fusispora* (26 % of the isolates), *L. pseudotheobromae* (23 % of the isolates) and *L. theobromae* (14 % of the isolates) are dominant and are distributed throughout the surveyed Provinces of South China. Of the 12 species of Botryosphaeriaceae, *L. theobromae* (isolated from *C. lanceolata*, *D. longan*, a *Eucalyptus* hybrid and *P. hanceana*) and *B. pseudoramosa* (isolated from a *Eucalyptus* hybrid and *M. sanguineum*) were collected from more than one plant host (Fig. 11).




Fig. 11 Map showing the 12 species of Botryosphaeriaceae detected from different regions and plant hosts. The different Botryosphaeriaceae species are indicated as numbers 1 to 12; the plant hosts are shown as letters A to G. For example, A8 indicates *L. theobromae* (number 8 of fungal species) isolated from *Eucalyptus* spp. (letter A of plant species) in HaiNan Province. The pies in colours indicate Botryosphaeriaceae isolated from different plant hosts in this study, the pies without colour indicate Botryosphaeriaceae species reported from *Eucalyptus* in previous studies (Chen et al. 2011c, Li et al. 2015a).

**Table 5** Average lesion length (mm) on seedlings of three *Eucalyptus* clones inoculated with *Botryosphaeriaceae*.

Species	Isolates	<i>Eucalyptus</i> clones		
		CEPT-11	CEPT-12	CEPT-13
<i>Botryosphaeria fusicpora</i>	CERC1998	17.6 ± 1.8 m-p <sup>1</sup>	27.1 ± 9.3 k-p	10.6 ± 0.7 op
	CERC2274	10.3 ± 0.4 op	10.7 ± 0.4 op	9.2 ± 0.2 op
	CERC2930	12.3 ± 1.2 op	13.9 ± 2.4 m-p	14.5 ± 1.6 m-p
	CERC3446	12.0 ± 0.6 op	17.5 ± 4.5 m-p	15.5 ± 2.8 m-p
<i>B. pseudoramosa</i>	CERC2001	13.5 ± 0.8 n-p	15.5 ± 4.0 m-p	11.1 ± 0.5 op
	CERC3452	16.8 ± 2.0 m-p	26.9 ± 7.3 k-p	13.9 ± 1.9 m-p
<i>B. qingyuanensis</i>	CERC2946	10.4 ± 0.5 op	16.2 ± 5.6 m-p	11.1 ± 0.8 op
	CERC2947	11.0 ± 0.5 op	18.2 ± 5.2 l-p	10.4 ± 0.5 op
<i>B. wangensis</i>	CERC2298	10.6 ± 0.5 op	12.3 ± 0.8 op	10.0 ± 0.2 op
	CERC2299	9.3 ± 1.1 op	11.1 ± 0.3 op	9.8 ± 0.1 op
<i>Copriniforma atrovirens</i>	CERC3484	11.0 ± 1.5 op	10.0 ± 1.5 op	8.8 ± 1.1 op
	CERC3489	12.2 ± 0.7 op	9.2 ± 0.2 op	9.4 ± 0.3 op
<i>Lasiodiplodia brasiliense</i>	CERC2284	41.7 ± 5.6 j-m	139.7 ± 27.3 cd	95.8 ± 15.7 f
	CERC2286	90.7 ± 16.9 fg	84.5 ± 12.6 f-h	100.1 ± 21.8 ef
<i>L. pseudotheobromae</i>	CERC3417	78.4 ± 9.7 fg	121.0 ± 12.6 de	128.4 ± 12.2 d
	CERC3495	120.9 ± 11.5 de	138.1 ± 12.8 cd	85.9 ± 9.5 fg
	CERC3420	26.7 ± 2.3 k-p	46.6 ± 7.5 i-l	26.8 ± 5.6 k-p
	CERC3513	123.9 ± 16.0 d	150.7 ± 21.9 b	219.5 ± 19.8 a
<i>Neofusicoccum hongkongense</i>	CERC3516	126.3 ± 13.1 d	142.0 ± 21.6 bc	173.5 ± 18.7 b
	CERC2968	17.7 ± 1.2 m-p	12.9 ± 2.2 op	14.6 ± 1.2 m-p
<i>N. microconidium</i>	CERC2973	21.6 ± 1.2 k-p	31.8 ± 5.6 k-p	18.7 ± 1.3 m-p
	CERC3497	32.7 ± 2.2 k-p	47.7 ± 7.5 i-k	40.8 ± 6.8 j-n
<i>N. parvum</i>	CERC3498	16.6 ± 1.2 m-p	17.2 ± 1.7 m-p	20.8 ± 4.3 l-p
	CERC2951	10.3 ± 0.5 op	11.9 ± 0.9 op	10.3 ± 0.3 op
<i>N. sinoeucalypti</i>	CERC3504	17.3 ± 0.7 m-p	30.1 ± 5.2 k-p	22.1 ± 3.2 k-p
	CERC3509	16.0 ± 1.5 m-p	17.5 ± 4.0 k-p	15.3 ± 2.4 m-p
Control	CERC2005	27.5 ± 4.7 k-p	68.0 ± 9.0 g-i	62.3 ± 8.7 h-j
	CERC3463	30.9 ± 4.8 k-p	24.0 ± 4.3 k-p	39.3 ± 7.7 j-o
Control		10.5 ± 0.6 op	10.0 ± 0.2 op	9.4 ± 0.3 op

<sup>1</sup> Mean ± SE followed by different lowercase letters indicates treatments that are significantly different ( $P < 0.05$ ); Mean = average lesion length; SE = standard error of mean.

### Pathogenicity tests

Twenty-eight isolates representing the 12 species of *Botryosphaeriaceae* identified in this study were used for inoculations on three different *Eucalyptus* clones (different parents) (Table 1, 5). Pathogenicity tests indicate that all of the *Botryosphaeriaceae* isolates tested produce lesions on stems of the three *Eucalyptus* clones, while MEA unclonised plugs produced only wounds. Overall, isolates in species of *Lasiodiplodia* produce relatively longer lesions than that of *Botryosphaeria*, *Copriniforma* and *Neofusicoccum*. For all three tested *Eucalyptus* clones, the lesions produced by *Lasiodiplodia* isolates are all significantly longer than the wounds caused by negative controls, except isolate CERC3420 (*L. theobromae*) on CEPT-11 and CEPT-13 ( $P < 0.05$ ) (Table 5). For isolates in the genera of *Botryosphaeria*, *Copriniforma* and *Neofusicoccum*, isolates CERC3497 (*N. microconidium*) and CERC2005 (*N. sinoeucalypti*) also produce significantly longer lesions on CEPT-11 and CEPT-13 ( $P < 0.05$ ) (Table 5). Analysis of variance shows significant differences in the susceptibility of the three *Eucalyptus* clones to some of the isolates we tested. For example, the lesions produced by isolate CERC2284 (*L. brasiliense*) on three *Eucalyptus* clones are significantly different from each other ( $P < 0.05$ ) (Table 5). Analysis of results also show that not all the isolates of the same species of *Botryosphaeriaceae* react in the same manner to the *Eucalyptus* clones. For example, lesions produced by isolate CERC3420 (*L. theobromae*) on clone CEPT-12 are significantly longer than those on CEPT-13, whereas lesions produced by isolate CERC3513 (*L. theobromae*) on CEPT-12 are significantly shorter than those on CEPT-13 ( $P < 0.05$ ) (Table 5). In addition, based on the lesions caused by all *Botryosphaeriaceae* isolates in this study, CEPT-11 (average lesion length:  $33.0 \pm 2.4$  mm) is more tolerant than CEPT-12 (average lesion length:  $44.2 \pm 3.2$  mm) and CEPT-13 (average lesion length:  $42.0 \pm 3.4$  mm). All 12 species of *Botryosphaeriaceae* were re-isolated successfully

from the lesions, and no *Botryosphaeriaceae* were isolated from the negative controls, thus fulfilling Koch's postulates.

### DISCUSSION

In this study, disease samples from symptomatic trees with stem cankers, shoot and twig blight were collected mainly from *Eucalyptus* and six other plant hosts in China. *Botryosphaeriaceae* was isolated from these diseased samples. Based on phylogenetic analyses and morphological characteristics, 12 species of *Botryosphaeriaceae* were isolated from these samples and the genera *Botryosphaeria*, *Copriniforma*, *Lasiodiplodia* and *Neofusicoccum* were identified from among a relatively large collection of isolates. These species include *Botryosphaeria fusicpora*, *Copriniforma atrovirens*, *Lasiodiplodia brasiliense*, *L. pseudotheobromae*, *L. theobromae*, *Neofusicoccum parvum* and each of three previously undescribed species of *Botryosphaeria* and *Neofusicoccum*, namely *B. pseudoramosa* sp. nov., *B. qingyuanensis* sp. nov., *B. wangensis* sp. nov., *N. hongkongense* sp. nov., *N. microconidium* sp. nov. and *N. sinoeucalypti* sp. nov.

In this study, ITS, *tef1*, *tub*, *rpb2*, *cmdA*, LSU and SSU sequences were generated to distinguish and describe new species of *Botryosphaeria*, *Copriniforma*, *Lasiodiplodia* and *Neofusicoccum*. For the six to seven regions used for analyses of *Botryosphaeria*, *Lasiodiplodia* and *Neofusicoccum*, phylogenetic analyses based on sequence comparisons show that polymorphic nucleotides exist between some isolates collected in this study and other closely related species. Sequences of the ITS, *tef1* and *tub* regions are widely used to distinguish and describe new species of *Botryosphaeria*, *Lasiodiplodia* and *Neofusicoccum* of *Botryosphaeriaceae* (Phillips et al. 2013, Chen et al. 2015, Linaldeddu et al. 2015, Coutinho et al. 2017), except ITS, *tef1* and *tub*, *rpb2* genes are also used for the species identification of *Neofusicoccum* (Pavlic et al. 2009a,

Sakalidis et al. 2011, Osorio et al. 2017, Yang et al. 2017) and *rpb2* and *cmdA* are also used for *Lasiodiplodia* (Cruywagen et al. 2017, Dou et al. 2017a, b, Osorio et al. 2017). The phylogenetic analyses based on a combination of the three to five regions (*Botryosphaeria*: ITS, *tef1* and *tub*; *Lasiodiplodia*: ITS, *tef1*, *tub*, *rpb2* and *cmdA*; *Neofusicoccum*: ITS, *tef1*, *tub* and *rpb2*) indicated that these isolates form independent phylogenetic clades supported by high bootstrap values, which are identified and described as six new species. In the other Chinese isolates, the differences we did find occurred only in one of the two (*Cophiniforma*), six (*Botryosphaeria* and *Neofusicoccum*) or seven (*Lasiodiplodia*) regions, these isolates reside in the same clade to previously identified species or form independent phylogenetic clades but not supported by high bootstrap values, and they were identified as *B. fusiclava*, *C. atrovirens*, *L. brasiliense*, *L. pseudotheobromae*, *L. theobroma* and *N. parvum*.

The identification of 12 *Botryosphaeriaceae* species is also supported by morphological and/or biological characteristics. For each of the six species that have been described previously, their culture morphology and conidial characteristics are very similar to that of the type specimens. For the six newly described species in this study, morphological differences exist among them and other phylogenetically closely related species, especially in terms of the size and shape of conidia, as well as conidium septum characteristics. We also observed biological differences, for example optimal growth temperatures, among some of the species. For the six new species, *B. pseudoramosa*, *B. wangensis*, *N. hongkongense* and *N. microconidium* are easily distinguished from other phylogenetically close species based on conidial morphology. Although some overlap in conidial shape and size is observed among some species, such as *B. fabicerciana*, *B. kuwatsukai* and *B. qingyuanensis*, these species can be distinguished from each other by the presence of a conidial septum (older conidia) and microconidia, as well as the optimal growth temperature. The newly described species *N. sinoeucalypti* can be distinguished from other species with similar conidia in the *N. parvum*/*N. ribis* complex by conidial morphology and the optimal growth temperature.

Except for *B. wangensis*, *C. atrovirens* and *N. hongkongense*, the other nine species were isolated from *Eucalyptus* trees in South China. Of the *Botryosphaeriaceae* species isolated from *Eucalyptus*, *B. fusiclava*, *L. pseudotheobromae* and *L. theobromae* are dominant and distributed in the GuangDong, GuangXi and HaiNan Provinces; *L. pseudotheobromae* and *L. theobromae* have also been found in previous studies (Chen et al. 2011c, Li et al. 2015a), suggesting that they may be widely distributed on *Eucalyptus* trees in other areas in South China. Four new species, *B. pseudoramosa*, *B. qingyuanensis*, *N. microconidium* and *N. sinoeucalypti*, were isolated from *Eucalyptus* in China. This study also presents the first report of *L. brasiliense* on *Eucalyptus* in the world. Species of *Botryosphaeriaceae* are distributed in all the areas surveyed where *Eucalyptus* is planted. The results of our study suggest that the species diversity of *Botryosphaeriaceae* on *Eucalyptus* in China may be higher than what was previously expected (Chen et al. 2011c).

In addition to *Botryosphaeriaceae* species identified on *Eucalyptus*, we also identified *B. pseudoramosa* from *Melastoma sanguineum*, *B. wangensis* from *C. deodara*, *C. atrovirens* from *D. longan*, *L. theobromae* from *C. lanceolata*, *D. longan* and *P. hanceana*, and *N. hongkongense* from *A. cunninghamii*. Aside from *L. theobromae* from *P. hanceana* (Lu et al. 2000), which has been reported previously, these *Botryosphaeriaceae* species are reported from their respective plant hosts for the first time. Disease materials were collected randomly from limited areas, including the areas which were adjacent to *Eucalyptus* plantations, and further work is needed to better understand

the biodiversity and distribution of *Botryosphaeriaceae* on their hosts.

Based on sequence comparisons of the seven gene regions, the same genotype of *L. theobromae* was shared by species of *Eucalyptus* in all the surveyed provinces in South China, and *C. lanceolata*, *D. longan* and *P. hanceana* planted in GuangDong Province (Table 1). We isolated the newly described species *B. pseudoramosa* from both *Eucalyptus* trees and *M. sanguineum*, and isolates from different hosts in geographically close areas do share the same genotype (Table 1). These results provide confirmation for the wide host range of *L. theobromae* and *B. pseudoramosa* on different plants. Previous studies used genetic diversity and geographic distribution comparisons to show the wide host range of *N. mediterraneum* on different crop trees in California (Chen et al. 2014a, b). The results of our current study further show that some *Botryosphaeriaceae* have wide geographic and host ranges.

Inoculation experiments revealed that all species of *Botryosphaeriaceae* identified in this study are pathogenic to the tested *Eucalyptus* clones, which is consistent with previous work showing that *Botryosphaeriaceae* species include important pathogens of *Eucalyptus* (Pavlic et al. 2007, Mohali et al. 2009, Rodas et al. 2009, Chen et al. 2011c). Pathogenicity tests in this study showed that species of *Lasiodiplodia* are more aggressive than *Botryosphaeria* and *Neofusicoccum* on three *Eucalyptus* clones, including one clone of *E. urophylla* × *E. grandis*, which is consistent with results in previous studies (Chen et al. 2011c). Results in Mohali et al. (2009) showed that some species of *Neofusicoccum* were more aggressive than *Lasiodiplodia* on clones of *E. urophylla* × *E. grandis*, which indicated that resistance of different genotypes of *E. urophylla* × *E. grandis* can be significantly different. Therefore, the identification of commercially available *Eucalyptus* genotypes resistant to *Botryosphaeriaceae* will promote the selection of resistant materials for wide-scale planting.

Of the fungal species we found, *L. theobromae* and *L. pseudotheobromae* are the most aggressive and are also widely distributed on *Eucalyptus* trees in different regions; it is essential that these pathogens be monitored carefully to help make decisions regarding disease management. Except for species of *Lasiodiplodia*, other fungi of the genera *Botryosphaeria*, *Cophiniforma* and *Neofusicoccum* also produce lesions on inoculated seedlings; although these species are not highly virulent to *Eucalyptus* and are not widespread, these fungi still need to be monitored carefully because some of them may be highly aggressive to their original hosts or may spread and act as important pathogens in a suitable environment.

Our results in this study indicate that some species of *Botryosphaeriaceae* are widely distributed in different geographic regions on different hosts. These fungal species have significant potential to cause diseases of *Eucalyptus*. Management of the diseases on *Eucalyptus* reported in this study will need to rely on sound breeding programs to select *Eucalyptus* genotypes to match climatic and edaphic factors and silvicultural practices (spacing and thinning) as part of an integrated management strategy (Old et al. 2003). Further study is needed to better understand the genetic diversity of the species at the population level and to understand the biological and epidemiological characteristics of these species to help with long-term disease management.

**Acknowledgements** This study was supported by the National Key R&D Program of China (project no. 2017YFD0600103), the National Natural Science Foundation of China (NSFC) (project nos 31622019 and 31400546). We thank Ms CaiYun Xu and Mr ShengLong Zhang for their assistance in collecting disease samples in GuangDong and GuangXi Provinces. We thank WILEY for its linguistic assistance during the preparation of this manuscript.

## REFERENCES

- Abdollahzadeh J, Javadi J, Mohammadi Goltepeh E, et al. 2010. Phylogeny and morphology of four new species of *Lasiodiplodia* from Iran. *Persoonia* 25: 1–10.
- Abdollahzadeh J, Zare R, Phillips AJL. 2013. Phylogeny and taxonomy of *Botryosphaeria* and *Neofusicoccum* species in Iran, with description of *Botryosphaeria scharifii* sp. nov. *Mycologia* 105: 210–220.
- Alves A, Correis A, Luque J, et al. 2004. *Botryosphaeria corticola* sp. nov. on *Quercus* species, with notes and description of *Botryosphaeria stevensii* and its anamorph, *Diplodia mutila*. *Mycologia* 96: 598–613.
- Alves A, Crous PW, Correis A, et al. 2008. Morphological and molecular data reveal cryptic speciation in *Lasiodiplodia theobromae*. *Fungal Diversity* 28: 1–13.
- Alves A, Phillips AJL, Henriques I, et al. 2005. Evaluation of amplified ribosomal DNA restriction analysis as a method for the identification of *Botryosphaeria* species. *FEMS Microbiology Letters* 245: 221–229.
- Ariyawansa HA, Hyde KD, Liu JK, et al. 2016. Additions to Karst Fungi 1: *Botryosphaeria minutispermatia* sp. nov., from Guizhou Province, China. *Phytotaxa* 275: 35–44.
- Begoué BAD. 2010. Characterization of *Botryosphaeriaceae* and *Cryptonectriaceae* associated with *Terminalia* spp. in Africa. PhD thesis, Department of Microbiology and Plant Pathology, Forestry and Agricultural Biotechnology Institute, University of Pretoria, South Africa.
- Begoué BAD, Slippers B, Wingfield MJ, et al. 2010. *Botryosphaeriaceae* associated with *Terminalia catappa* in Cameroon, South Africa and Madagascar. *Mycological Progress* 9: 101–123.
- Berraf-Tebbal A, Guereiro MA, Phillips AJL. 2014. Phylogeny of *Neofusicoccum* species associated with grapevine trunk diseases in Algeria, with description of *Neofusicoccum algeriense* sp. nov. *Phytopathologia Mediterranea* 53: 416–427.
- Burgess TI, Andjic V, Hardy GESTJ, et al. 2006. First report of *Phaeophleospora destructans* in China. *Journal of Tropical Forest Science* 18: 144–146.
- Burgess TI, Barber PA, Hardy GESTJ. 2005. *Botryosphaeria* spp. associated with eucalypts in Western Australia, including the description of *Fusicoccum macroclavatum* sp. nov. *Australasian Plant Pathology* 34: 557–567.
- Burgess TI, Barber PA, Sufaati S, et al. 2007. *Mycosphaerella* spp. on Eucalyptus in Asia: new species, new hosts and new records. *Fungal Diversity* 24: 135–157.
- Cao JD. 1982. Investigation of bacterial wilt in *Eucalyptus saligna* and *E. grandis* introduced from Brazil. *Guangxi Forestry Science and Technology* 4: 30–31. [In Chinese.]
- Carbone I, Kohn LM. 1999. A method for designing primer sets for speciation studies in filamentous ascomycetes. *Mycologia* 91: 553–556.
- Chen SF, Barnes I, Chungu D, et al. 2011a. High population diversity and increasing importance of the *Eucalyptus* stem canker pathogen, *Teratosphaeria zuluensis*, in South China. *Australasian Plant Pathology* 40: 407–415.
- Chen SF, Gryzenhout M, Roux J, et al. 2010. Identification and pathogenicity of *Chrysoporthea cubensis* on *Eucalyptus* and *Syzygium* spp. in South China. *Plant Disease* 94: 1143–1150.
- Chen SF, Li GQ, Liu FF, et al. 2015. Novel species of *Botryosphaeriaceae* associated with shoot blight of pistachio. *Mycologia* 107: 780–792.
- Chen SF, Li GQ, Liu QL, et al. 2016. Characteristics of *Lasiodiplodia theobromae* from *Rosa rugosa* in South China. *Crop Protection* 79: 51–55.
- Chen SF, Lombard L, Roux J, et al. 2011b. Novel species of *Calonectria* associated with *Eucalyptus* leaf blight in Southeast China. *Persoonia* 26: 1–12.
- Chen SF, Morgan DP, Hasey JK, et al. 2014a. Phylogeny, morphology, distribution, and pathogenicity of *Botryosphaeriaceae* and *Diaporthaceae* from English walnut in California. *Plant Disease* 98: 636–652.
- Chen SF, Morgan DP, Michailides TJ. 2014b. *Botryosphaeriaceae* and *Diaporthaceae* associated with panicle and shoot blight of pistachio in California, USA. *Fungal Diversity* 67: 157–179.
- Chen SF, Pavlic D, Roux J, et al. 2011c. Characterization of *Botryosphaeriaceae* from plantation-grown *Eucalyptus* species in South China. *Plant Pathology* 60: 739–751.
- Chen SF, Van Wyk M, Roux J, et al. 2013. Taxonomy and pathogenicity of *Ceratocystis* species on *Eucalyptus* trees in South China, including *C. chinaeucensis* sp. nov. *Fungal Diversity* 58: 267–279.
- Chen SX, Chen XF. 2013. Technical problems and thinking on *Eucalypt* plantation management in China. *Eucalypt Science & Technology* 30: 52–59. [In Chinese.]
- Coutinho IBL, Freire FCO, Lima CS, et al. 2017. Diversity of genus *Lasiodiplodia* associated with perennial tropical fruit plants in northeastern Brazil. *Plant Pathology* 66: 90–104.
- Crous PW, Groenewald JZ, Shivas RG, et al. 2011. Fungal Planet description sheets: 69–91. *Persoonia* 26: 108–156.
- Crous PW, Slippers B, Wingfield MJ, et al. 2006. Phylogenetic lineages in the *Botryosphaeriaceae*. *Studies in Mycology* 55: 235–253.
- Crous PW, Wingfield MJ, Guarro J, et al. 2013. Fungal Planet description sheets: 154–213. *Persoonia* 31: 188–296.
- Crous PW, Wingfield MJ, Phillips AJL. 2007. *Neofusicoccum mediterraneum*. Fungal Planet 19. CBS-KNAW Fungal Biodiversity Centre, Utrecht, The Netherlands.
- Crugwagen EM, Slippers B, Roux J, et al. 2017. Phylogenetic species recognition and hybridisation in *Lasiodiplodia*: a case study on species from baobabs. *Fungal Biology* 121: 420–436.
- Cubeta MA, Echandi E, Abernethy T, et al. 1991. Characterization of anastomosis groups of binucleate *Rhizoctonia* species using restriction analysis of an amplified ribosomal RNA gene. *Phytopathology* 81: 1395–1400.
- Damm U, Crous PW, Fourie PH. 2007. *Botryosphaeriaceae* as potential pathogens of *Prunus* species in South Africa, with descriptions of *Diplodia africana* and *Lasiodiplodia plurivora* sp. nov. *Mycologia* 99: 664–680.
- Darriba D, Taboada GL, Doallo R, et al. 2012. jModelTest 2: more models, new heuristics and parallel computing. *Nature Methods* 9: 772.
- Denman S, Crous PW, Groenewald JZ, et al. 2003. Circumscription of *Botryosphaeria* species associated with *Proteaceae* based on morphology and DNA sequence data. *Mycologia* 95: 294–307.
- Dou ZP, He W, Zhang Y. 2017a. *Lasiodiplodia chinensis*, a new holomorphic species from China. *Mycosphere* 8: 521–532.
- Dou ZP, He W, Zhang Y. 2017b. Does morphology matter in taxonomy of *Lasiodiplodia*? An answer from *Lasiodiplodia hyalina* sp. nov. *Mycosphere* 8: 1014–1027.
- Farr DF, Elliott M, Rossman AY, et al. 2005. *Fusicoccum arbuti* sp. nov. causing cankers on Pacific madrone in western North America with notes on *Fusicoccum dimidiatum*, the correct name for *Scytalidium dimidiatum* and *Natrassia mangiferae*. *Mycologia* 97: 730–741.
- Felsenstein J. 1985. Confidence intervals on phylogenetics: an approach using bootstrap. *Evolution* 39: 783–791.
- Gardner DE. 1997. *Botryosphaeria mamane* sp. nov. associated with witches'-brooms on the endemic forest tree *Sophora chrysophylla* in Hawaii. *Mycologia* 89: 298–303.
- Glass NL, Donaldson GC. 1995. Development of primer sets designed for use with the PCR to amplify conserved genes from filamentous ascomycetes. *Applied and Environmental Microbiology* 61: 1323–1330.
- Guindon S, Dufayard JF, Lefort V, et al. 2010. New algorithms and methods to estimate maximum-likelihood phylogenies: assessing the performance of PhyML 3.0. *Systematic Biology* 59: 307–321.
- Hillis DM, Huelsenbeck JP. 1992. Signal, noise and reliability in molecular phylogenetic analyses. *Journal of Heredity* 83: 189–195.
- Iglesias-Trabado G, Carbaeira-Tenreiro R, Folgueira-Lozano J. 2009. *Eucalyptus universalis*. Global cultivated eucalypt forest map Version 1.2 In: GIT Forestry Consulting's EUCALYPTOLOGICS: Information resources on Eucalyptus cultivation worldwide. Retrieved from <http://www.git-forestry.com> [accessed: 19 Oct. 2009].
- Inderbitzin P, Bostock RM, Trouillas FP, et al. 2010. A six locus phylogeny reveals high species diversity in *Botryosphaeriaceae* from California almond. *Mycologia* 102: 1350–1368.
- Jacobs K, Bergdahl DR, Wingfield MJ, et al. 2004. *Leptographium wingfieldii* introduced into North America and found associated with exotic *Tomicus piniperda* and native bark beetles. *Mycological Research* 108: 411–418.
- Katoh K, Standley DM. 2013. MAFFT multiple sequence alignment software version 7: improvements in performance and usability. *Molecular Biology and Evolution* 30: 772–780.
- Lazzizera C, Frisullo S, Alves A, et al. 2008. Morphology, phylogeny and pathogenicity of *Botryosphaeria* and *Neofusicoccum* species associated with drupe rot of olives in southern Italy. *Plant Pathology* 57: 948–956.
- Li GQ, Arnold RJ, Liu FF, et al. 2015a. Identification and pathogenicity of *Lasiodiplodia* species from *Eucalyptus urophylla* × *grandis*, *Polyscias balfouriana* and *Bougainvillea spectabilis* in Southern China. *Journal of Phytopathology* 163: 956–967.
- Li GQ, Liu FF, Li JQ, et al. 2015b. Characterization of *Botryosphaeria dothidea* and *Lasiodiplodia pseudotheobromae* from English walnut in China. *Journal of Phytopathology* 164: 348–353.
- Linaldeddu BT, Dedda A, Scanu B, et al. 2015. Diversity of *Botryosphaeriaceae* species associated with grapevine and other woody hosts in Italy, Algeria and Tunisia, with descriptions of *Lasiodiplodia exigua* and *Lasiodiplodia mediterranea* sp. nov. *Fungal Diversity* 71: 201–214.
- Liu FF, Mbenoun M, Barnes I, et al. 2015. New *Ceratocystis* species from *Eucalyptus* and *Cunninghamia* in South China. *Antonie van Leeuwenhoek* 107: 1451–1473.
- Liu JK, Phookamsak R, Doilom M, et al. 2012. Towards a natural classification of *Botryosphaeraiales*. *Fungal Diversity* 57: 149–210.

- Liu YJ, Whelen S, Hall BD. 1999. Phylogenetic relationships among Ascomycetes: evidence from an RNA polymerase II subunit. *Molecular Biology and Evolution* 16: 1799–1808.
- Lombard L, Zhou XD, Crous PW, et al. 2010. Calonectria species associated with cutting rot of Eucalyptus. *Persoonia* 24: 1–11.
- Lopes A, Phillips AJL, Alves A. 2017. Mating type genes in the genus *Neofusicoccum*: Mating strategies and usefulness in species delimitation. *Fungal Biology* 121: 394–404.
- Lu B, Hyde KD, Ho WH, et al. 2000. Checklist of Hong Kong fungi. Fungal Diversity Press, Hong Kong.
- Machado AR, Pinho DB, Pereira OL. 2014. Phylogeny, identification and pathogenicity of the Botryosphaeriaceae associated with collar and root rot of the biofuel plant *Jatropha curcas* in Brazil, with a description of new species of *Lasiodiplodia*. *Fungal Diversity* 67: 231–247.
- Marques MW, Lima NB, De Moraes Jr MA, et al. 2013. *Botryosphaeria*, *Neofusicoccum*, *Neoscyclodium* and *Pseudofusicoccum* species associated with mango in Brazil. *Fungal Diversity* 61: 195–208.
- Mehl JWM, Slippers B, Roux J, et al. 2011. Botryosphaeriaceae associated with *Pterocarpus angolensis* (kiaat) in South Africa. *Mycologia* 103: 534–553.
- Mohali S, Slippers B, Wingfield MJ. 2006. Two new *Fusicoccum* species from *Acacia* and *Eucalyptus* in Venezuela, based on morphology and DNA sequence data. *Mycological Research* 110: 405–413.
- Mohali SR, Slippers B, Wingfield MJ. 2007. Identification of Botryosphaeriaceae from *Eucalyptus*, *Acacia* and *Pinus* in Venezuela. *Fungal Diversity* 25: 103–125.
- Mohali SR, Slippers B, Wingfield MJ. 2009. Pathogenicity of seven species of the Botryosphaeriaceae on *Eucalyptus* clones in Venezuela. *Australasian Plant Pathology* 38: 135–140.
- Netto MSB, Assunção IP, Lima GSA, et al. 2014. Species of *Lasiodiplodia* associated with papaya stem-end rot in Brazil. *Fungal Diversity* 67: 127–141.
- Netto MSB, Lima GSA, Correia KC, et al. 2017. Analysis of phylogeny, distribution, and pathogenicity of Botryosphaeriaceae species associated with gummosis of *Anacardium* in Brazil, with a new species of *Lasiodiplodia*. *Fungal Biology* 121: 437–451.
- Old KM, Wingfield MJ, Yuan ZQ. 2003. A manual of diseases of Eucalypts in South-East Asia. Centre for International Forestry Research, Indonesia.
- Osorio JA, Crous CJ, De Beer ZW, et al. 2017. Endophytic Botryosphaeriaceae, including five new species, associated with mangrove trees in South Africa. *Fungal Biology* 121: 361–393.
- Pavlic D, Slippers B, Coutinho TA, et al. 2007. Botryosphaeriaceae occurring on native *Syzygium cordatum* in South Africa and their potential threat to *Eucalyptus*. *Plant Pathology* 56: 624–636.
- Pavlic D, Slippers B, Coutinho TA, et al. 2009a. Multiple gene genealogies and phenotypic data reveal cryptic species of the Botryosphaeriaceae: A case study on the *Neofusicoccum parvum*/N. *ribis* complex. *Molecular Phylogenetics and Evolution* 51: 259–268.
- Pavlic D, Slippers B, Coutinho TA, et al. 2009b. Molecular and phenotypic characterization of three phylogenetic species discovered within the *Neofusicoccum parvum*/N. *ribis* complex. *Mycologia* 101: 636–647.
- Pavlic D, Slippers B, Teresa A, et al. 2004. *Lasiodiplodia gonubiensis* sp. nov., a new Botryosphaeria anamorph from native *Syzygium cordatum* in South Africa. *Studies in Mycology* 50: 313–322.
- Pavlic D, Wingfield MJ, Barber P, et al. 2008. Seven new species of the Botryosphaeriaceae from baobab and other native trees in Western Australia. *Mycologia* 100: 851–866.
- Phillips AJL, Alves A, Abdollahzadeh J, et al. 2013. The Botryosphaeriaceae: genera and species known from culture. *Studies in Mycology* 76: 51–167.
- Phillips AJL, Alves A, Pennycook SR, et al. 2008. Resolving the phylogenetic and taxonomic status of dark-spored teleomorph genera in the Botryosphaeriaceae. *Persoonia* 21: 29–55.
- Phillips AJL, Oudemans PV, Correia A, et al. 2006. Characterisation and epitypification of *Botryosphaeria corticis*, the cause of blueberry cane canker. *Fungal Diversity* 21: 141–155.
- Prasher IB, Singh G. 2014. *Lasiodiplodia indica* – A new species of coelomycetous mitosporic fungus from India. *Kavaka* 43: 64–69.
- Punithalingam E. 1980. Plant diseases attributed to *Botryodiplodia theobromae* Pat. Cramer, Vaduz, Liechtenstein.
- Rayner RW. 1970. A mycological colour chart. Commonwealth Mycological Institute and British Mycological Society, Kew, Surrey, UK.
- Rodas CA, Slippers B, Gryzenhout M, et al. 2009. Botryosphaeriaceae associated with Eucalyptus canker diseases in Colombia. *Forest Pathology* 39: 110–123.
- Rodríguez-Gálvez E, Guerrero P, Barradas C, et al. 2017. Phylogeny and pathogenicity of *Lasiodiplodia* species associated with dieback of mango in Peru. *Fungal Biology* 121: 452–465.
- Sakalidis ML, Hardy GEStJ, Burgess TI. 2011. Use of the Genealogical Sorting Index (GSI) to delineate species boundaries in the *Neofusicoccum parvum*–*Neofusicoccum ribis* species complex. *Molecular Phylogenetics and Evolution* 60: 333–344.
- SAS Institute Inc. 2011. SAS® 9.3 System Options: Reference, 2nd edn. Cary, North California, USA.
- Slippers B, Boissin E, Phillips AJL, et al. 2013. Phylogenetic lineages in the Botryosphaerales: a systematic and evolutionary framework. *Studies in Mycology* 76: 31–49.
- Slippers B, Crous PW, Denman S, et al. 2004a. Combined multiple gene genealogies and phenotypic characters differentiate several species previously identified as *Botryosphaeria dothidea*. *Mycologia* 96: 82–101.
- Slippers B, Fourie G, Crous PW, et al. 2004b. Multiple gene sequences delimit *Botryosphaeria australis* sp. nov. from *B. lutea*. *Mycologia* 96: 1030–1041.
- Slippers B, Fourie G, Crous PW, et al. 2004c. Speciation and distribution of *Botryosphaeria* spp. on native and introduced *Eucalyptus* trees in Australia and South Africa. *Studies in Mycology* 50: 343–358.
- Slippers B, Johnson GI, Crous PW, et al. 2005. Phylogenetic and morphological re-evaluation of the *Botryosphaeria* species causing diseases of *Mangifera indica*. *Mycologia* 97: 99–110.
- Slippers B, Roux J, Wingfield MJ, et al. 2014. Confronting the constraints of morphological taxonomy in the Botryosphaerales. *Persoonia* 33: 155–168.
- Slippers B, Wingfield MJ. 2007. Botryosphaeriaceae as endophytes and latent pathogens of woody plants: diversity, ecology and impact. *Fungal Biology Reviews* 21: 90–106.
- Smith H, Crous PW, Wingfield MJ, et al. 2001. *Botryosphaeria eucalyptorum* sp. nov., a new species in the *B. dothidea*-complex on *Eucalyptus* in South Africa. *Mycologia* 93: 277–285.
- Smith H, Wingfield MJ, Crous PW, et al. 1996. *Sphaeropsis sapinea* and *Botryosphaeria dothidea* endophytic in *Pinus* spp. and *Eucalyptus* spp. in South Africa. *South African Journal of Botany* 62: 86–88.
- Swofford DL. 2003. PAUP\*. Phylogenetic Analysis Using Parsimony (\*and other methods). Version 4.0b10. Sinauer Associates, Sunderland, MA, USA.
- Tamura K, Stecher G, Peterson D, et al. 2013. MEGA6: Molecular evolutionary genetics analysis version 6.0. *Molecular Biology and Evolution* 30: 2725–2729.
- Tang W, Ding Z, Zhou ZQ, et al. 2012. Phylogenetic and pathogenic analyses show that the causal agent of apple ring rot in China is *Botryosphaeria dothidea*. *Plant Disease* 96: 486–496.
- Taylor K, Barber PA, Hardy GEStJ, et al. 2009. Botryosphaeriaceae from tuart (*Eucalyptus gomphocephala*) woodland, including descriptions of four new species. *Mycological Research* 113: 337–353.
- Trakunyingcharoen T, Lombard L, Groenewald JZ, et al. 2015. Caulicolous Botryosphaerales from Thailand. *Persoonia* 34: 87–99.
- Turnbull JW. 2007. Development of sustainable forestry plantations in China: a review. ACIAR Impact Assessment Series Report No. 45. Australian Centre for International Agricultural Research, Canberra.
- Úrbez-Torres JR, Peduto F, Striegler RK, et al. 2012. Characterization of fungal pathogens associated with grapevine trunk diseases in Arkansas and Missouri. *Fungal Diversity* 52: 169–189.
- Van Burik JAH, Schreckhise RW, White TC, et al. 1998. Comparison of six extraction techniques for isolation of DNA from filamentous fungi. *Medical Mycology* 36: 299–303.
- Van Niekerk JM, Crous PW, Groenewald JZ, et al. 2004. DNA phylogeny, morphology and pathogenicity of *Botryosphaeria* species on grapevines. *Mycologia* 96: 781–798.
- Vilgalys R, Hester M. 1990. Rapid genetic identification and mapping of enzymatically amplified ribosomal DNA from several *Cryptococcus* species. *Journal of Bacteriology* 172: 4238–4246.
- Wei RP. 2005. Genetic diversity and sustainable productivity of Eucalypt plantations in China. In: Wang H (ed), *Changing patterns: tree introduction and phytogeography*: 19–27. China Forestry Publishing House, Beijing, China. [In Chinese.]
- White TJ, Bruns T, Lee S, et al. 1990. Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. In: Innis MA, Gelfand DH, Sninsky JJ, et al. (eds), *PCR Protocols: a guide to methods and applications*: 315–322. Academic Press, San Diego, CA, USA.
- Wingfield MJ. 2003. Increasing threat of diseases to exotic plantation forests in the Southern Hemisphere: lessons from *Cryphonectria* canker. *Australasian Plant Pathology* 32: 133–139.
- Wingfield MJ, Slippers B, Hurley BP, et al. 2008. Eucalypt pests and diseases: growing threats to plantation productivity. *Southern Forests* 70: 139–144.
- Xu C, Wang CS, Ju LL, et al. 2015a. Multiple locus genealogies and phenotypic characters reappraise the causal agents of apple ring rot in China. *Fungal Diversity* 71: 215–231.

- Xu C, Zhang H, Zhou Z, et al. 2015b. Identification and distribution of Botryosphaeriaceae species associated with blueberry stem blight in China. *Europe Journal of Plant Pathology* 143: 737–752.
- Yan JY, Xie Y, Yao SW, et al. 2012. Characterization of *Botryosphaeria dothidea*, the causal agent of grapevine canker in China. *Australasian Plant Pathology* 41: 351–357.
- Yan JY, Xie Y, Zhang W, et al. 2013. Species of Botryosphaeriaceae involved in grapevine dieback in China. *Fungal Diversity* 61: 221–236.
- Yang T, Groenewald JZ, Cheewangkoon R, et al. 2017. Families, genera and species of Botryosphaerales. *Fungal Biology* 121: 322–346.
- Yu ZD, Tang GH, Peng SB, et al. 2015. *Neofusicoccum parvum* causing canker of seedlings of *Juglans regia* in China. *Journal of Forestry Research* 26: 1019–1024.
- Zhang M, Lin S, He W, et al. 2017. Three species of *Neofusicoccum* (Botryosphaeriaceae, Botryosphaerales) associated with woody plants from southern China. *Mycosphere* 8: 797–808.
- Zhao JP, Lu Q, Liang J, et al. 2010. *Lasiodiplodia pseudotheobromae*, a new record of pathogenic fungus from some subtropical and tropical trees in southern China. *Cryptogamie Mycologie* 31: 431–439.
- Zhou XD, De Beer ZW, Xie YJ, et al. 2007. DNA-based identification of *Quambalaria pitereka* causing severe leaf blight of *Corymbia citriodora* in China. *Fungal Diversity* 25: 245–254.
- Zhou XD, Wingfield MJ. 2011. Eucalypt diseases and their management in China. *Australasian Plant Pathology* 40: 339–345.
- Zhou Y, Gong GS, Cui YL, et al. 2015. Identification of Botryosphaeriaceae species causing kiwifruit rot in Sichuan Province, China. *Plant Disease* 99: 699–708.
- Zhou YP, Dou ZP, He W, et al. 2016. *Botryosphaeria sinensis* sp. nov., a new species from China. *Phytotaxa* 245: 43–50.
- Zhou YP, Zhang M, Dou ZP, et al. 2017. *Botryosphaeria rosaceae* sp. nov. and *B. ramosa*, new botryosphaeriaceous taxa from China. *Mycosphere* 8: 162–171.