

Article

Selenium Speciation in the Fountain Creek Watershed (Colorado, USA) Correlates with Water Hardness, Ca and Mg Levels

James S. Carsella ^{1,2}, Irma Sánchez-Lombardo ^{3,4}, Sandra J. Bonetti ¹ and Debbie C. Crans ^{2,3,*}

¹ Department of Chemistry, Colorado State University–Pueblo, 2200 Bonforte Blvd, Pueblo, CO 81001, USA; jim.carsella@csupueblo.edu (J.S.C.), sandra.bonetti@csupueblo.edu (S.J.B.)

² Cell and Molecular Biology Program, Colorado State University, Fort Collins, CO 80523, USA

³ Department of Chemistry, Colorado State University, Fort Collins, CO 80523, USA

⁴ División de Ciencias Básicas, Universidad Juárez Autónoma de Tabasco, Cunduacán, Tabasco 86690, Mexico; irmas_l@yahoo.com.mx

* Correspondence: debbie.crans@colostate.edu; Tel.: +1-970-491-7635

Academic Editors: Claudio Santi and Luana Bagnoli

Received: 4 March 2017; Accepted: 19 April 2017; Published: 30 April 2017

Abstract: The environmental levels of selenium (Se) are regulated and strictly enforced by the Environmental Protection Agency (EPA) because of the toxicity that Se can exert at high levels. However, speciation plays an important role in the overall toxicity of Se, and only when speciation analysis has been conducted will a detailed understanding of the system be possible. In the following, we carried out the speciation analysis of the creek waters in three of the main tributaries—Upper Fountain Creek, Monument Creek and Lower Fountain Creek—located in the Fountain Creek Watershed (Colorado, USA). There are statistically significant differences between the Se, Ca and Mg levels in each of the tributaries and seasonal swings in Se, Ca and Mg levels have been observed. There are also statistically significant differences between the Se levels when grouped by Pierre Shale type. These factors are considered when determining the forms of Se present and analyzing their chemistry using the reported thermodynamic relationships considering Ca^{2+} , Mg^{2+} , SeO_4^{2-} , SeO_3^{2-} and carbonates. This analysis demonstrated that the correlation between Se and water hardness can be explained in terms of formation of soluble CaSeO_4 . The speciation analysis demonstrated that for the Fountain Creek waters, the Ca^{2+} ion may be mainly responsible for the observed correlation with the Se level. Considering that the Mg^{2+} level is also correlating linearly with the Se levels it is important to recognize that without Mg^{2+} the Ca^{2+} would be significantly reduced. The major role of Mg^{2+} is thus to raise the Ca^{2+} levels despite the equilibria with carbonate and other anions that would otherwise decrease Ca^{2+} levels.

Keywords: speciation; selenium determination; acidity; toxicity; thermodynamics

1. Introduction

The environmental levels of Se are regulated and strictly enforced in the USA by the Environmental Protection Agency (EPA) [1–4] because even though low levels are beneficial high levels are toxic [5–10]. The high Se toxicity on aquatic life and particularly fish reproduction [11–14] has led to the development of a chronic exposure criterion [11]. High Se levels are often found when waterways run over the Se-rich shale deposits [3] or when agricultural runoff carries Se into waterways from Se rich soils [15–18]. Specifically, the Lower Fountain Creek (LF) in southeastern Colorado is a known and major contributor to the Se levels downstream from its confluence with the Arkansas River (AR); this confluence is in the eastern part of the city of Pueblo, Colorado (for a map see [3,19]). These high Se levels in the Fountain

Creek and in the Arkansas River increase the potential of damaging environmental effects downstream in the Arkansas River basin [3]. While Se toxicity is linked to particular species [5,12–14,16,20] and the presence of selenite (SeO_3^{2-}) [21,22], many environmental studies only measure and report the total amount of elemental Se. When analyzing and investigating Se speciation, a better understanding of the interaction of Se species in a natural environment is gained [5,7,21,23–28]. Recent studies of fish in the Fountain Creek Watershed demonstrated that high numbers of fish species can be observed at sites with high levels of Se [21]. This work led to the hypothesis that the toxicity is dependent on the presence of toxic Se species and not the total Se level [21] and support the need for additional studies that consider the effects of Se species on different types of fish. The following manuscript analyzes the chemistry of Se speciation in these waters and that of two major cations, Ca^{2+} and Mg^{2+} , using the thermodynamic parameters.

Total Se concentrations in aqueous environments generally vary between 0.06 to 400 $\mu\text{g/L}$ [29–32]. The most common forms of Se in surface waters are Se(IV) and Se(VI) with some reports also including Se(II) in either organic or inorganic form [8,15,33–35]. Selenium is most soluble in aqueous solution under oxidizing conditions and the distribution of the oxidized forms of Se, selenite (SeO_3^{2-}) and selenate (SeO_4^{2-}) depend on specific conditions [8,15,34,35]. The less soluble and reduced form of Se, Se^{2-} , exists complexed in the earth's crust, and may also be found at low levels in reducing water environments [8]. Pourbaix diagrams are important to describe the speciation of Se in idealized systems [8,15,21,34,35]. However, considering that the Pourbaix diagrams are calculated assuming specific conditions, for example, where the concentration and ionic strength do not affect the speciation; such representations are at best, a first approximation [15,21,34,35]. From the reported Pourbaix diagrams, it is clear that selenous acid (H_2SeO_3) is favored at the conditions normally found in oxidizing natural stream waters ($\text{pE} \approx 13.5$) at a $\text{pH} < 3.0$ [36]. Hydrogen selenite (HSeO_3^-) is favored in the same oxidizing conditions up to about $\text{pH} 5.0$. Above $\text{pH} 5.0$ selenate (SeO_4^{2-}) is the most favored Se species. Using this information, it follows that in less aerated waters ($\text{pE} \approx 13.5$) at pH values between 3 and 8, hydrogen selenite, HSeO_3^- is the favored species [21,34]. However, recently we demonstrated that for Fountain Creek the Se was about 90% Se(VI) and 10% Se(IV) even through the Pourbaix diagrams predicted more Se(IV) [21]. In the following, we explore the following hypothesis: *classical speciation plots are effective in describing the Se speciation when considering two influential cations, Ca^{2+} and Mg^{2+}* . The conversions between the different oxidation states of Se are reported to be slow [35] and it is possible that the speciation in the watershed does not represent equilibrium conditions. Previously a correlation between total elemental Se with water hardness [19] was reported. However, the detailed chemistry of this correlation involving a combination of Ca^{2+} and Mg^{2+} was not investigated [19]. Specifically, the possibility of the formation of the soluble and insoluble forms of CaSeO_4 correlate with the observed aqueous Se levels and equilibrium conditions.

In the following, the Se speciation chemistry of waters in the Fountain Creek Watershed sites was explored. Since the two most common cations determining water hardness are calcium (Ca^{2+}) and magnesium (Mg^{2+}), in addition to Se, Ca, and Mg levels were measured at the Fountain Creek sites. Furthermore, the speciation chemistry and its relationship to Se content with Ca^{2+} and Mg^{2+} contents were explored. Samples were collected from three different Fountain Creek reaches: Upper Fountain Creek (UF), Monument Creek (MC) and the previously mentioned LF portion of Fountain Creek. The Se concentrations, the pH , and the concentrations of Ca^{2+} and Mg^{2+} were determined. Since correlations have been reported between Se levels and water hardness, the current manuscript explores the chemical speciation of Se, Ca, and Mg. The goal was to examine if specific chemical relationships between Se, Ca^{2+} , and Mg^{2+} combined can explain the reported correlation with water hardness. Speciation diagrams show that the Se content indeed correlates with the soluble CaSeO_4 levels even though the thermodynamic equilibria of Se speciation alone are not sufficient to describe all the properties of this system.

Table 1. Se levels in the water along the Fountain Creek Watershed [21,37] ^a.

Sample Sites	Spring									Fall								
	T (°C)	pH	Water Se (µg/L)	Std. Error Water Se	Water Ca (µg/L)	Std. Error Water Ca	Water Mg (µg/L)	Std. Error Water Mg	Alkalinity (mg CaCO ₃ /L)	T (°C)	pH	Water Se (µg/L)	Std. Error Water Se	Water Ca (µg/L)	Std. Error Water Ca	Water Mg (µg/L)	Std. Error Water Mg	Alkalinity (mg CaCO ₃ /L)
UF-1	4.6	7.2	0.193	0.003	28,440	892.5	5354	140.2	72	5.8	8.2	0.063	0.007	19,307	582.3	5481	261.7	76
UF-2	4.9	7.2	0.140	0.017	28,867	1993.3	5474	341.9	50	5.7	8.1	0.000	0.007	13,003	1355.7	3481	450.2	63
UF-3	5.7	7.6	0.673	0.090	34,863	2409.8	6890	456.9	81	6.0	8.1	0.043	0.009	13,897	291.7	4041	61.7	146
UF-4	6.6	7.6	1.323	0.313	39,017	2989.2	10,134	1504.9	83	7.0	8.1	0.340	0.010	15,303	205.1	5490	109.3	154
MC-1	4.2	7.4	0.207	0.012	12,227	357.3	1742	25.1	66	5.5	7.9	0.177	0.009	15,757	99.4	4772	102.3	86
MC-2	3.9	7.5	0.277	0.007	19,380	875.6	3400	253.9	74	5.3	8.0	0.330	0.023	21,547	394.7	7599	223.0	89
MC-3	4.1	7.2	0.343	0.017	21,623	1374.0	3818	283.5	73	4.5	8.1	0.597	0.038	22,450	283.6	8043	143.3	88
MC-4	4.1	7.5	0.443	0.023	26,327	1978.4	4365	324.2	78	5.0	8.0	0.860	0.021	27,047	91.3	8450	26.5	86
MC-5	4.8	7.4	1.863	0.216	38,503	3260.0	6712	538.8	110	6.4	8.1	5.283	0.269	41,080	172.1	13,217	416.0	105
LF-1	7.7	7.5	2.050	0.202	42,303	2751.5	8135	538.8	104	9.9	8.1	2.970	0.072	29,947	312.5	10,960	573.0	108
LF-2	12.2	7.2	2.780	0.008	53,465	698.1	14,540	714.4	141	15.2	8.1	3.773	0.060	38,537	520.7	18,177	137.8	179
LF-3	10.6	7.6	3.290	0.125	64,447	2316.7	17,703	514.4	138	15.1	8.1	3.800	0.057	41,837	806.5	19,770	182.5	180
LF-4	11.3	7.7	9.687	0.617	69,083	827.5	22,943	627.7	140	15.2	8.1	18.59	1.131	46,683	254.6	29,830	531.5	183
LF-5	11.8	7.8	7.910	0.377	69,373	1624.5	23,183	793.0	-	15.8	8.2	14.05	0.677	47,137	542.6	29,900	517.3	-
Ref.	^b	[37]	[37]	^b	^b		^b		[38–43]	^b	[37]	[37]		^b		^b		[38–43]

^a Measurements are averages of three as indicated in the experimental section. Abbreviations: UF, Upper Fountain; MC, Monument Creek; LF, Lower Fountain. ^b Information provided in this work not previously published.

2. Results

2.1. Characterization of the Se-Levels and Speciation in the Fountain Creek

Fountain and Monument Creeks were sampled for dissolved Se in the past [3,4,18,19,21,37,44–46]. High levels of Se were reported (3.4–12 $\mu\text{g/L}$) [3] with up to 64 $\mu\text{g/L}$ [3] between Colorado Springs and Pueblo, Colorado. Selenium originates from the Pierre Shale located in southeastern Colorado [2] which is underlying both Fountain and Monument Creeks. Fourteen sites were monitored in the spring and fall of 2007 (Table 1, Figure 1) and part of this data was reported previously [21,37]. Se concentrations and cation concentrations were measured using inductively coupled plasma–mass spectrometry (ICP–MS) and the results with Ca^{2+} and Mg^{2+} are now reported in this manuscript. In Table 1 the Se, pH, and alkalinity reported previously are listed for all water samples. The Ca and Mg concentrations and temperature are reported here. As previously shown, the temperature and pH influence speciation of different elements and these parameters were investigated here.




Figure 1. Total Se concentrations in the creek water (solid symbol) spring and (open symbol) fall 2007 in $\mu\text{g/L}$. UF stands for Upper Fountain Creek (squares), MC Monument Creek (circles) and LW, Lower Fountain Creek (triangles) [21]. Data points were measured in triplicates and details of the sampling site was described.

Several Analyses of Variances (ANOVA) tests were conducted on the Se levels from all reaches to determine which Se measurements are statistically different between site, reach, shale type and Ca, and Mg levels. In Figure 1, the observed Se concentrations are plotted at each site for both spring and fall.

Due to the exponential increase in the Se, Ca, and Mg concentrations in the reaches of the sampled creeks, natural log transformations were used to fit the data to a normal distribution to carry out the ANOVA analysis. An ANOVA test was also conducted on the Se, Ca, and Mg levels from the Pierre shales (Pierre Shale (PS), Upper Pierre Shale (UPS), Lower Pierre Shale (LPS), Continuous Pierre Shale (CPS) and No Pierre Shale (NPS)). A statistically significant difference in Se levels exists between LF and both the UF and MC reaches ($p < 0.0005$). No statistically significant difference exists between the UF and MC reaches of the Fountain Creek Study area. The ANOVA with Tukey's pairwise comparisons results also show a statistically significant difference between the Se levels and the Shale type ($p < 0.0005$). The results indicate that a statistically significant difference exists between the Se levels in the water exposed to LPS and the rest of the PS formations as well as NPS area of the study area. In addition, there is a statistically significant difference between the Se levels in the water in the

UPS and the area of NPS. No statistically significant differences were found between the NPS and CPS. A statistically significant difference exists between all reaches for Ca and Mg levels ($p < 0.0005$) with the lowest levels in MC. The ANOVA between shale type and Mg indicates that a statistically significant difference exists between LPS and UPS and NPS. There is an overlap between CPS and all shale types with respect to Mg levels. The Ca levels are statistically significantly different in all shale types except for the CPS in the spring data set. In the fall Ca dataset, there is an overlap between the levels in the CPS and NPS as well as LPS and UPS. A statistically significant difference exists between CPS-NPS and UPS-LPS ($p < 0.0005$).

The highest Se concentrations were observed at the MC-5 site that is in Colorado Springs and at the LF-4 and LF-5 sites. The latter sites are in the city of Pueblo (during both, spring and fall), see Figure 2, and for a map see [3,19]. These sites with very high Se levels are located where the human use is becoming more prevalent and thus include both contributions from natural and human waste sources [21,37]. The highest Se concentration in Monument Creek at MC-5 could be attributed to the presence of UPS under Monument Creek, while the elevated Se concentrations in the LF-4 and LF-5 sites are due to the LPS formation under the creek [19].




Figure 2. Total Se level ([Se]/µg/L) vs Total Hardness (A), Total Ca level (B) and Total Mg level (C) in the creek water (solid symbol) spring and (open symbol) fall 2007 in µg/L. UF is the Upper Fountain Creek (squares), MC Monument Creek (circles) and LF Lower Fountain Creek (triangles) [21]. Data points were measured in triplicates and details of the sampling sites were described.

2.2. Se, Ca and Mg Levels and Hardness in the Fountain Creek Water

Because correlations were reported between Se levels and water hardness by Herrmann and coworkers [19], the current manuscript explores the chemical speciation of Se, Ca and Mg to examine

if specific chemical relationships between Se and Ca or Mg can explain the reported correlation with water hardness. In Figure 2, the Se concentrations are plotted as a function of total hardness, and a correlation was suggested for the final sites at the Fountain Creek Watershed. Because water hardness is generally attributed to the amount of calcium (Ca^{2+}) and magnesium ions (Mg^{2+}) dissolved in the water, we investigated the relationships by plotting Se level as a function of the hardness (in Figure 2A), the concentrations of Ca^{2+} (Figure 2B) and the concentration of Mg^{2+} (Figure 2C). The plots show that more Ca was measured in the water in spring as compared to the fall and that this difference correlates with the difference observed in the hardness.

Based on Figure 2 plots of Se versus Ca^{2+} and Se versus Mg^{2+} as well as plots of each cation versus collection site shown in Figure 3, the Se levels increase as the creek runs downstream and across the PS formation. Variations in Se levels exist for each cation and water sample location and some different patterns emerge between the three parameters. Specifically, in the fall the Se concentrations are higher than in the spring for most samples. Linear relationships can be suggested between Se levels and hardness, and between Se and Mg^{2+} levels for both fall and spring. Linearity is also observed for Se levels and Ca^{2+} levels measured in the spring, but not in the fall. This change in the pattern is of interest because it could be related to uptake of Se by plants in the creek waters, and this will be the topic of a future investigation factoring in measurements of Se(IV) and Se(VI) [19]. Most UF and MC sites that have not been exposed to the PS formation, exhibit characteristically low Se concentrations in the water. The Se levels begin to rise at the UF-4 and MC-5 sites which are at the intersections of the UF reach with the CPS formation and the MC reach with the UPS formation. The water from both the UF and MC reaches confluence with the LF reach and flow into the LF-1 site. The most interesting pattern shown here is the correlation of the Se levels with Ca^{2+} concentrations in the spring measurements and with the Mg^{2+} concentrations in both spring and fall seasons.




Figure 3. (A) Total calcium concentrations (presumed Ca^{2+}) in the creek water (solid symbols) spring and (open symbols) fall year in mg/L and (B) total magnesium concentration (presumed Mg^{2+}) in the creek water (solid symbols) spring and (open symbols) fall year in $\mu\text{g/L}$. UF is the Upper Fountain Creek (symbol squares), MC Monument Creek (symbol circles) and LF Lower Fountain Creek (symbol triangles). Data points were measured in triplicates and details of the sampling sites were described [21].

In the case of the Mg^{2+} , there is less seasonal variation (Figure 2C). In Figure 3B the concentrations of Mg at the different sample sites were plotted, and the same trend for the sample locations was found for the Mg levels in samples collected in the fall or spring (see Figures 2 and 3). The UF-4, MC-5, LF-4 and LF-5 sites contain the highest levels of Ca in spring. Therefore, the fact that the Se levels correlate differently with the hardness in the fall versus the spring streams can be attributed to the seasonal variations in Ca^{2+} levels shown in Figure 2B. We suggest that although Mg^{2+} is important for defining the environment in these systems because the concentrations of Mg^{2+} remain constant, it influences hardness by facilitating dissolution of insoluble Ca^{2+} species as has been reported previously for

MgCO₃-CO₂ systems [47,48]. We suggest that similar effects exist for the MgSeO₃-SeO₂ system and that this can explain at least in part the concentrations of SeO₃²⁻ and SeO₄²⁻ systems above those expected by the solubility products.

2.3. Speciation Profile for Sites with Low and High Levels of Se

In Figure 4 we show the distribution of Se species that would be anticipated if the most of the oxidized Se species present are in a thermodynamic equilibrium using the reported formation constants (see the Materials and Methods section for formation constants). The figures show that under oxidizing conditions the major species is predicted to be SeO₄²⁻ from pH 5 to 8, but that at higher concentrations the contribution of Se(IV) to Se(VI) is smaller. However, it must be noted, that these calculations are done at oxidizing conditions, and as reported previously using Pourbaix diagrams, when the pE measured at these sites along the Fountain Creek is used, the predicted major Se component was Se(IV) [21]. However, speciation studies showed that regardless, about 90% of the Se was found to be in oxidation state VI [21] and using these results we calculated the speciation that would be observed for the UF-2 and LF-4 sites (Figure 4).




Figure 4. Speciation diagrams were calculated for two different sites along the Fountain Creek in the stream water: (A) The low concentration UF-2 sites with $[\text{Se}]_{\text{tot}} 3.36 \times 10^{-10}$ M, and (B) LF-4 with $[\text{Se}]_{\text{tot}} 1.22 \times 10^{-7}$ M. Recently the speciation was determined to around 90% Se(VI) for several of the sites studied in this work [25].

Ca²⁺ is known to form both soluble and insoluble Ca²⁺-SeO₄²⁻ species and the distribution of these species was investigated using the LF-4 site concentrations of Se (Figure 5A). The levels of the soluble form of CaSeO₄ was determined previously in other waters in studies using isotopically pure materials. These studies confirmed the presence of a significant amount of oxidized Se(VI) in such waters [47,48].

Studies with the Fountain Creek waters have previously shown that the formation of CaSeO₄ is competing with the formation of Ca²⁺ complexes with CaSeO₃, CO₃²⁻, and with HCO₃⁻; and the latter will be discussed below. Figure 5A shows the speciation of Se(IV) and Se(VI) in the presence of Ca²⁺ showing more CaSeO₄ than CaSeO₃. As in the case of the Ca²⁺, the Mg²⁺ speciation was considered in Figure 5. The speciation diagram was constructed for the Mg and Se species using the LF-4 site concentrations, showing that the most favorable species in the water under these conditions were MgSeO₄ and the insoluble MgSeO₃.

As elevated concentrations of Ca²⁺ and Mg²⁺ were measured, the speciation diagrams for both cations were plotted in Figure 5A and by comparison, we include the speciation diagram for the Se, HCO₃⁻ and CO₃²⁻ species with both Ca²⁺ and Mg²⁺ together in Figure 6. As expected, the carbonate species with both cations are the most abundant since their equilibrium constants are the highest.




Figure 5. Speciation diagrams were calculated for LF-4 site along the Fountain Creek in the stream water with concentrations (A) of [Se(IV)] 1.2×10^{-8} M, [Se(VI)] 1.1×10^{-7} M and $[Ca]_{\text{tot}}$ 1.7×10^{-3} M and (B) [Se(IV)] 1.2×10^{-8} M, [Se(VI)] 1.1×10^{-7} M and $[Mg]_{\text{tot}}$ 9.4×10^{-4} M.

The carbonates thus shift the selenite equilibrium for the Mg^{2+} since no insoluble $MgSeO_3$ is present; however, this is not the case for the $CaSeO_3$ that continues to have insoluble material. The presence of carbonates with their large equilibrium constants and the higher concentration of Ca^{2+} versus Mg^{2+} would support the formation of insoluble calcium selenite. This behavior could be a consequence of a cycle whereby carbonates are consumed by organisms present in the water leading to more release of Ca^{2+} . This is a logical explanation given that the pH of the Fountain Creek water (Table 1) is slightly alkaline and the alkalinity measurements indicate that under these conditions the waters are oversaturated with Ca^{2+} according to the Langelier saturation index (LSI) [49]. The LSI was calculated for the LF-4 site in the summer using the alkalinity value of 140 mg/L of $CaCO_3$ and the K_{sp} for $CaCO_3$ (6.92×10^{-9}) where $LSI = \log([Ca^{2+}][CO_3^{2-}]/K_{\text{sp}}) = 2.45$; this indicates that the water is saturated with $CaCO_3$ which would allow precipitation of some $CaCO_3$ particles in the water.




Figure 6. Speciation diagram was calculated for the LF-4 site along the Fountain Creek in the stream water with concentrations of [Se(IV)] 1.2×10^{-8} M, [Se(VI)] 1.1×10^{-7} M, $[Ca]_{\text{tot}}$ 1.7×10^{-3} M and $[Mg]_{\text{tot}}$ 9.4×10^{-4} M.

Although the formation constant for $CaCO_3$ is large, at pH values near neutral the concentration of CO_3^{2-} in the creek waters is very low. The protonated form of HCO_3^- is present, but the complex that it forms with Ca^{2+} is much less stable. The methods used in this study specifically preclude measurements of carbonate and the values are thus not routinely measured. However, carbonate levels in river waters have been reported previously and are in the range of 20–200 mg/L $CaCO_3$ [49,50]. To illustrate the levels of calcium carbonates in these systems, we assume a carbonate level of 140 mg/L

in the speciation diagram calculated in Figure 6. Since the formation constant for the soluble CaSeO_4 is high ($K = 10^{2.68}$) a significant portion of the soluble Se presumably exists as a soluble form of CaSeO_4 , however, the solubility product for the insoluble CaSeO_4 may be reached at some of the sites with high Se concentrations. The possibility that the CaSeO_4 species is a major species involved in the correlation between Se levels and hardness was considered, and the speciation diagrams including the Ca^{2+} cations were examined at the LF-4 site concentrations and are shown in Figure 6. As seen from Figure 6 the major species in solution are indeed CaSeO_4 and $\text{Ca}(\text{HCO}_3)_2$.

At the pH values ($5 < \text{pH} < 8$) of the water that has been investigated, most of the carbonate would be protonated and not as prone to precipitation. In addition, Castanier and coworkers show that production of carbonate particles by heterotrophic bacteria follows different paths and these induce a pH increase and an accumulation of CO_3^{2-} and HCO_3^- ions in the system [51]. This is important because the high concentration of Ca^{2+} found in the water correlates with the total Se and because in our previous work [21] we found that the major Se species is present in the nontoxic form of Se(VI) as SeO_4^{2-} . However, additional analysis is needed and will be the topic of future communications. The subject of other cations will be considered particularly when the processing of the Se due to uptake by the plants in the creek bed is examined. Thus, speciation of iron and other metal ions become important [25,52–57].

2.4. Exploration of Other Physical Parameters and Their Impact on the Speciation

To further explore the effects of other physical parameters the Se, Ca and Mg levels were measured in addition to pH, temperature, and discharge. In this study fourteen sites were monitored in spring and fall of 2007 (Table 1). We found that the highest Se concentrations in each reach were observed at the UF-4 and MC-5 sites in Colorado Springs and LF-4 and LF-5 sites in Pueblo in both seasons as shown in Figure 2.

The water temperature values are more consistent within a season (Table 1) and the water temperatures are colder during the spring. The discharge rates are higher during the spring runoff and the lower discharge during the fall resulted in higher Se concentrations (Figure 7A). This is consistent with the observation of higher Se levels when there are higher water temperatures and smaller discharge rates; this allows for more dissolution of Se into the creek water. The total Se content in the water during spring and fall correlates with the water temperature and the creek discharge rate as shown from the plots in Figure 7.




Figure 7. Total Se concentration in nM vs. Creek discharge in m^3/s (A) and Se concentration in nM vs. water temperature ($^{\circ}\text{C}$) (B). Selenium concentrations in the creek water (solid symbol) spring and (open symbol) fall 2007 in nM. UF stands for Upper Fountain Creek (squares), MC Monument Creek (circles) and LW, Lower Fountain Creek (triangles) Regression lines are shown as solid for spring and dashed for fall. Details of the sampling sites were described previously [21].

The best relationship to dissolved Se in the creek exists with flow rate (Figure 7A, $R^2_{\text{spring}} = 0.8094$, $R^2_{\text{Fall}} = 0.7941$). Temperature is also correlated with dissolved Se but the correlation is not as strong as the flow rate (Figure 7B). Together the data for temperature and flow rate suggest that there are symbiotic effects on dissolved Se and that these parameters assist in increasing the amount of dissolved Se, and because these parameters affect the Ca^{2+} (Figure 8A,C) and Mg^{2+} (Figure 8B,D) in a similar manner, they contribute to increasing the Ca^{2+} level above that expected for the solubility product (Figure 8).




Figure 8. Total Ca concentration in nM plotted vs. creek discharge in m^3/s (A). For comparison, total Mg concentration in nM plotted vs. creek discharge in m^3/s (B). Total Ca concentration in nM plotted vs. water temperature ($^{\circ}\text{C}$) (C). Total Mg concentration in nM plotted vs. water temperature ($^{\circ}\text{C}$) (D). concentrations in the creek water (solid symbol) spring and (open symbol) fall 2007 in $\mu\text{g}/\text{L}$. UF stands for Upper Fountain Creek (squares), MC Monument Creek (circles) and LW, Lower Fountain Creek (triangles) Regression lines are shown as solid for spring and dashed for fall. Details of the sampling sites were described previously [21].

3. Materials and Methods

3.1. Materials and Reagents

The standards for EPA method 200.8 were purchased as a NIST traceable multi-element custom made standard from Inorganic Ventures (IV, Christiansburg, VA, USA, CSTU-STD-1) and from SPEX CertiPrep (Metuchen, NJ, USA, part# CL-CAL-2A). The IV standard was made to contain $1000 \mu\text{g}/\text{mL}$ of Ca, Mg, Na, Fe, K. Furthermore, it contains $10 \mu\text{g}/\text{mL}$ Al, Sb, As, Ba, Cd, Se, Ag, Pb, Be, Co, Cr, Zn, Cu, Mn, Mo, Ni, V, U, Th, Tl. The SPEX standard was diluted to 50 ppb and used as a check

standard to verify calibration. The internal standards used were (Inorganic Ventures 2008ISS-125, ^6Li , Sc, Y, In, Tb, Bi, Ho) and Germanium (Ge) (Inorganic Ventures MSGE-10PPM-125ML). The nitric acid (A467-500, Fisher Scientific, Waltham, MA, USA) and hydrochloric acid (Fisher A466-500) used in these procedures were purchased as Optima grade from Fisher Scientific.

3.2. Fountain Creek Sampling

Water quality measurements of surface waters included measuring temperature, pH, specific conductance, and dissolved oxygen per the protocols described by the United States Geological Survey (USGS) protocols [19,58]. Details of the sampling site was described previously including the Upper Fountain creek, the Monument Creek and the Lower Mountain Creek [21]. Each parameter was measured three times at each site over a 10-day period. The measurements were made at 0, 5 and 10-day intervals. Water samples were taken at equivalent distance points on a line across the creek at each sampling site. The samples were collected in 250 mL low density polyethylene (LDPE) plastic bottles and stored on ice for transport. The collection was performed per limnologic sampling protocol by holding the bottle with the mouth of the container facing downstream at an angle near 45° . The container was rinsed with creek water 3 times prior to taking the sample. Triplicate samples were collected at each site during each sampling interval and filtered through a 0.45 micron reconstituted cellulose membrane syringe filter (Phenex AF0-8103-12) purchased from Phenomenex (Torrance, CA, USA) [58]. After filtering, water samples including field blanks for each trip were preserved with 1% Optima grade HNO_3 . The sample was divided in two, one was analyzed and the other stored at 4°C . Six milliliters were used for cation determination by ICP-MS.

3.3. Analysis for Se, Ca and Mg

Water samples were analyzed for Se, Ca and Mg on a 7500ce ICP-MS (Agilent, Santa Clara, CA, USA) following the ICP-MS drinking water method, EPA Method 200.8 [59]. Since these samples were not intended for evaluation for municipal drinking water, the ICP-MS was run with the Octopole Reaction System (ORS). Hydrogen was used as a reaction gas for ^{40}Ca and ^{78}Se measurements. The elements ^{24}Mg and ^{44}Ca were measured using helium as a collision gas, to reduce interferences. Multi-element environmental external calibration standard (CSTU-STD-1) and internal standards were diluted in 1% nitric and 0.5% hydrochloric acid prior to analysis. The internal standard was diluted to a final concentration of 20 ppb from a 10-ppm stock. Germanium was added to the final internal standard mix before dilution to make a final concentration of 20 ppb. The Ge was used as an internal standard for Se because of Ge having a closer ionization potential to Se. Scandium was used as the internal standard for Ca and Mg.

3.4. Alkalinity


Alkalinity was measured by titration on all samples collected using the protocols listed in the book *Standard Methods for the Examination of Water and Wastewater* for alkalinity [60]. Details of each sampling site was described previously including the Upper Fountain creek, the Monument Creek and the Lower Mountain Creek [21].

3.5. Statistical Calculations

Analysis of Variance was used to determine if statistically significant differences exist between the water components (Ca, Mg, and Se) exposed to Pierre Shale type, creek reach, and sites. Creek discharge rates were obtained from USGS monitoring sites via web interfaces [19,58]. The sites nearest the monitoring stations (UF-1, UF-4, MC-1, MC-2, MC-4, LF-1, LF-2, and LF-4) used the data directly from the monitoring stations. The sites between the stations (UF-2, UF-3, MC-3, MC-5, LF-3, and LF-5) used a weighted average of the flow rates measured at the USGS stations above and below the sampling site used in this study.

3.6. Speciation Calculations

Species distribution diagrams were calculated by using HYSS 2003 software [61]. The concentration used for the consideration described here for the UF-2 site was $[\text{Se}]_{\text{tot}} 3.36 \times 10^{-10}$ M, and for LF-4 site was $[\text{Se}]_{\text{tot}} 1.22 \times 10^{-7}$ M. When the speciation diagrams with Ca and Mg cations were constructed the concentrations used were $[\text{Se(IV)}] 1.2 \times 10^{-8}$ M, $[\text{Se(VI)}] 1.1 \times 10^{-7}$ M, $[\text{Ca}]_{\text{tot}} 1.7 \times 10^{-3}$ M and $[\text{Mg}]_{\text{tot}} 9.4 \times 10^{-4}$ M. The label Cr designate crystalline phase, see below constants for $\text{CaSeO}_3 \cdot \text{H}_2\text{O}_{(\text{cr})}$ and $\text{MgSeO}_3 \cdot 6\text{H}_2\text{O}_{(\text{cr})}$ at 25 °C and $I = 0$ [62]. The speciation diagrams were constructed using the following equilibrium reactions and constants [62]:



4. Conclusions

A speciation analysis was carried out using Se and pH data on creek samples from 14 sites in the Fountain Creek Watershed which were supplemented with new measurements of Ca, Mg and temperature at these same sites. Details of the sampling sites were described previously including the Upper Fountain Creek, the Monument Creek and the Lower Mountain Creek locations [21]. Several ANOVA analyses were conducted on the Se levels from all reaches to consider which Se measurements are statistically different between site, reach, the type of shale and Ca, and Mg levels. An ANOVA was also performed on the Pierre shales (Pierre Shale (PS), Upper Pierre Shale (UPS), Lower Pierre Shale (LPS), Continuous Pierre Shale (CPS) and No Pierre Shale (NPS)) and a statistically significant difference in Se levels exists between LF and both the UF and MC reaches ($p < 0.0005$). The subsequent analysis supports the interpretation that the Ca^{2+} ion is mainly responsible for the observed high Se levels in these waters presumably by the formation of the soluble and insoluble CaSeO_4 . Although the Mg^{2+} levels also correlate with the Se level, this cation is not present at the same high level and thus presumably serves mainly to increase the Ca^{2+} levels beyond the solubility limit through ion-pair formation as reported previously [52]. These findings are important because several of these sites are found to have high levels of Se. However, previously it was demonstrated that there are no apparent toxicity effects associated with the fish population and these high Se levels [19,25,52–57]. The data presented here thus further characterize the Se-rich waters in which most of the Se is in the Se(VI)

oxidation state. The studies here validate the observation that Se toxicity is reduced because Se(VI) is the predominant species and the studies presented here suggest that a major part of the Se(VI) is in the form of CaSeO_4 .

The detailed analysis of the Se species and its ability to explain observations underline the importance of consideration of the speciation chemistry in these environmental systems. The studies demonstrate that measurement of Se species allows for additional insights into the processes in a hydrological system. The detailed considerations including Mg^{2+} and Ca^{2+} allowed for an insight into the form of Se and the environmental system at hand. These considerations do require measurements of the oxidation state of the Se species and demonstrate the need for continued development of advanced methods for detection and measurements of components in complex matrices. We illustrate in this manuscript that the use of speciation analysis significantly enhances the understanding of previously reported results for other systems [25,52–57].

Acknowledgments: J.S.C. and S.J.B. would like to thank the following for funding this research: The Board of Pueblo, Colorado, County Commissioners, the Lower Arkansas Valley Water Conservancy District, Colorado, and the Board of Water Works of Pueblo and J.S.C., S.J.B. and D.C.C. would like to thank Colorado State University-Pueblo for funding from 2014–2015 and 2016–2017 Faculty SEED Grants.

Author Contributions: J.S.C. collected the samples, measured the Se, Ca and Mg levels, analyzed data for Table 1 and ran the references; I.S.L. carried out the speciation analysis and prepared most of the figures; S.J.B. wrote a grant to obtain funding and supervised J.S.C. at CSU-Pueblo; D.C.C. wrote most of the manuscript and supervised J.S.C. and I.S.L.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. United States Environmental Protection Agency National Recommended Water Quality Criteria-Aquatic Life Criteria Table. Available online: <https://www.epa.gov/wqc/national-recommended-water-quality-criteria-aquatic-life-criteria-table> (accessed on 18 April 2016).
2. Colorado Department of Public Health and Environment. *The Basic Standards and Methodologies for Surface Water*; Colorado Department of Public Health and Environment Water Quality Control Commission: Denver, CO, USA, 2013; Volume 31, p. 218.
3. Divine, C.E.; Gates, T.K. *Sources and Occurrence of Selenium in the Arkansas River and Fountain Creek near Pueblo, Colorado*; Arcadis G&M, Inc.: Highlands Ranch, CO, USA, 2006; pp. 1–41.
4. Edelmann, P.; Ferguson, S.A.; Stogner, R.W., Sr.; August, M.; Payne, W.F.; Bruce, J.F. *Evaluation of Water Quality, Suspended Sediment, and Stream Morphology with an Emphasis on Effects of Stormflow on Fountain and Monument Creek Basins, Colorado Spring and Vicinity, Colorado, 1981 Through 2001*; United States Geological Survey: Denver, CO, USA, 2005.
5. Bird, S.M.; Ge, H.; Uden, P.C.; Tyson, J.F.; Block, E.; Denoyer, E. High-Performance liquid chromatography of selenoamino acids and organo selenium compounds. Speciation by inductively coupled plasma mass spectrometry. *J. Chromatogr. A* **1997**, *789*, 349–359. [[CrossRef](#)]
6. Bird, S.M.; Tyson, J.F. Speciation of selenoamino acids and organoselenium compounds in selenium-Enriched yeast using high-Performance liquid chromatography–Inductively coupled plasma mass spectrometry. *J. Anal. At. Spectrom.* **1997**, *12*, 785–788. [[CrossRef](#)]
7. Block, E.; Booker, S.J.; Flores-Penalba, S.; George, G.N.; Gundala, S.; Landgraf, B.J.; Liu, J.; Lodge, S.N.; Pushie, M.J.; Rozovsky, S.; et al. Trifluoroselenomethionine: A New Unnatural Amino Acid. *Chembiochem* **2016**, *17*, 1738–1751. [[CrossRef](#)] [[PubMed](#)]
8. Combs, G.F., Jr.; Gray, W.P. Chemopreventive agents: Selenium. *Pharmacol. Ther.* **1998**, *79*, 179–192. [[CrossRef](#)]
9. Mousa, S.A.; O'Connor, L.; Rossman, T.G.; Block, E. Pro-Angiogenesis action of arsenic and its reversal by selenium-Derived compounds. *Carcinogenesis* **2007**, *28*, 962–967. [[CrossRef](#)] [[PubMed](#)]
10. Weekley, C.M.; Harris, H.H. Which form is that? The importance of selenium speciation and metabolism in the prevention and treatment of disease. *Chem. Soc. Rev.* **2013**, *42*, 8870–8894. [[CrossRef](#)] [[PubMed](#)]
11. United States Environmental Protection Agency. *Draft Aquatic Life Ambient Water Quality Criterion for Selenium–Freshwater 2015*; United States Environmental Protection Agency: Washington, DC, USA, 2015; pp. 1–740.

12. Hamilton, S.J.; Holley, K.M.; Buhl, K.J.; Bullard, F.A. Selenium impacts on razorback sucker, Colorado: Colorado River: III. Larvae. *Ecotoxicol. Environ. Saf.* **2005**, *61*, 168–189. [[CrossRef](#)] [[PubMed](#)]
13. Hamilton, S.J.; Holley, K.M.; Buhl, K.J.; Bullard, F.A. Selenium impacts on razorback sucker, Colorado River, Colorado II. Eggs. *Ecotoxicol. Environ. Saf.* **2005**, *61*, 32–43. [[CrossRef](#)] [[PubMed](#)]
14. Hamilton, S.J.; Holley, K.M.; Buhl, K.J.; Bullard, F.A.; Ken Weston, L.; McDonald, S.F. Selenium impacts on razorback sucker, Colorado River, Colorado I. Adults. *Ecotoxicol. Environ. Saf.* **2005**, *61*, 7–31. [[CrossRef](#)] [[PubMed](#)]
15. Mast, M.A.; Mills, T.J.; Paschke, S.S.; Keith, G.; Linard, J.I. Mobilization of selenium from the Mancos Shale and associated soils in the lower Uncompahgre River Basin, Colorado. *Appl. Geochem.* **2014**, *48*, 16–27. [[CrossRef](#)]
16. Kennedy, C.J.; McDonald, L.E.; Loveridge, R.; Stroscher, M.M. The effect of bioaccumulated selenium on mortalities and deformities in the eggs, larvae, and fry of a wild population of cutthroat trout (*Oncorhynchus clarki lewisi*). *Arch. Environ. Contam. Toxicol.* **2000**, *39*, 46–52. [[CrossRef](#)] [[PubMed](#)]
17. Lemly, A.D. A teratogenic deformity index for evaluating impacts of selenium on fish populations. *Ecotoxicol. Environ. Saf.* **1997**, *37*, 259–266. [[CrossRef](#)] [[PubMed](#)]
18. Presser, T.S.; Sylvester, M.A.; Low, W.H. Bioaccumulation of selenium from natural geologic sources in western states and its potential consequences. *Environ. Manag.* **1994**, *18*, 423–436. [[CrossRef](#)]
19. Herrmann, S.J.; Turner, J.A.; Carsella, J.S.; Lehmpuhl, D.W.; Nimmo, D.R. Bioaccumulation of selenium by the Bryophyte *Hygrohypnum ochraceum* in the Fountain Creek Watershed, Colorado. *Environ. Manag.* **2012**, *50*, 1111–1124. [[CrossRef](#)] [[PubMed](#)]
20. Mathews, T.J.; Fortner, A.M.; Jett, R.T.; Morris, J.; Gable, J.; Peterson, M.J.; Carriker, N. Selenium bioaccumulation in fish exposed to coal ash at the Tennessee Valley Authority Kingston spill site. *Environ. Toxicol. Chem.* **2014**, *33*, 2273–2279. [[CrossRef](#)] [[PubMed](#)]
21. Carsella, J.S.; Melnykov, I.; Bonetti, S.J.; Sánchez-Lombardo, I.; Crans, D.C. Selenium speciation in the Fountain Creek watershed and its effects on fish diversity. *J. Biol. Inorg. Chem.* **2017**. [[CrossRef](#)] [[PubMed](#)]
22. Rosenfeld, I.; Beath, O.A. *Selenium: Geobotany, Biochemistry, Toxicity, and Nutrition*; Academic Press: New York, NY, USA, 1964; p. 411.
23. Block, E. Fifty years of smelling sulfur. *J. Sulfur Chem.* **2013**, *34*, 158–207. [[CrossRef](#)]
24. Crans, D.C. Antidiabetic, Chemical, and Physical Properties of Organic Vanadates as Presumed Transition-State Inhibitors for Phosphatases. *J. Org. Chem.* **2015**, *80*, 11899–11915. [[CrossRef](#)] [[PubMed](#)]
25. Crans, D.C.; Woll, K.A.; Prusinskas, K.; Johnson, M.D.; Norkus, E. Metal speciation in health and medicine represented by iron and vanadium. *Inorg. Chem.* **2013**, *52*, 12262–12275. [[CrossRef](#)] [[PubMed](#)]
26. Doucette, K.A.; Hassell, K.N.; Crans, D.C. Selective speciation improves efficacy and lowers toxicity of platinum anticancer and vanadium antidiabetic drugs. *J. Inorg. Biochem.* **2016**, *165*, 56–70. [[CrossRef](#)] [[PubMed](#)]
27. Kiss, T.; Jakusch, T.; Hollender, D.; Dörnyei, Á.; Enyedy, É.A.; Pessoa, J.C.; Sakurai, H.; Sanz-Medel, A. Biospeciation of antidiabetic VO(IV) complexes. *Coord. Chem. Rev.* **2008**, *252*, 1153–1162. [[CrossRef](#)]
28. Kiss, T.; Odani, A. Demonstration of the importance of metal ion speciation in bioactive systems. *Bull. Chem. Soc. Jpn.* **2007**, *80*, 1691–1702. [[CrossRef](#)]
29. Fawell, J.K.; Combs, G.F. Guidelines for drinking-water quality. In *Selenium in Drinking Water: Background Document for Development of WHO Guidelines for Drinking Water Quality*; World Health Organization: Geneva, Switzerland, 2011; Volume 2011, p. 22.
30. Lindberg, P.; Bingefors, S. Selenium levels of forages and soils in different regions of Sweden. *Acta Agric. Scand.* **1970**, *20*, 133–136. [[CrossRef](#)]
31. Scott, R.C.; Voegeli, P.T. *Radiochemical Analyses of Ground and Surface Water in Colorado, 1954–1961*; Colorado Water Conservation Board: Denver, CO, USA, 1961.
32. Smith, M.; Westfall, B. Further field studies on the selenium problem in relation to public health. *Public Health Rep. (1896–1970)* **1937**, *52*, 1375–1384. [[CrossRef](#)]
33. Kulp, T.R.; Pratt, L.M. Speciation and weathering of selenium in Upper Cretaceous chalk and shale from South Dakota and Wyoming, USA. *Geochim. Cosmochim. Acta* **2004**, *68*, 3687–3701. [[CrossRef](#)]
34. Torres, J.; Pintos, V.; Domínguez, S.; Kremer, C.; Kremer, E. Selenite and selenate speciation in natural waters: Interaction with divalent metal ions. *J. Solut. Chem.* **2010**, *39*, 1–10. [[CrossRef](#)]

35. Torres, J.; Pintos, V.; Gonzatto, L.; Domínguez, S.; Kremer, C.; Kremer, E. Selenium chemical speciation in natural waters: Protonation and complexation behavior of selenite and selenate in the presence of environmentally relevant cations. *Chem. Geol.* **2011**, *288*, 32–38. [CrossRef]
36. Baes, C.F.; Mesmer, R.E. *Hydrolysis of Cations*; John Wiley & Sons: New York, NY, USA, 1976; p. 489.
37. Nimmo, D.R.; Herrmann, S.J.; Carsella, J.S.; McGarvy, C.M.; Foutz, H.P.; Herrmann-Hoesing, L.M.; Gregorich, J.M.; Turner, J.A.; Vanden Heuvel, B.D. Mercury and selenium in fish of Fountain Creek, Colorado (USA): Possible sources and implications. *Springerplus* **2016**, *5*, 437. [CrossRef] [PubMed]
38. United States Geological Survey National Water Information System: Web Interface. USGS Site 07106500. Available online: http://waterdata.usgs.gov/co/nwis/uv/?site_no=07106500 (accessed on 26 January 2017).
39. United States Geological Survey National Water Information System: Web Interface. USGS Site 07106200. Available online: http://waterdata.usgs.gov/co/nwis/uv/?site_no=07106200 (accessed on 26 January 2017).
40. United States Geological Survey National Water Information System: Web Interface. USGS Site 07103755. Available online: http://waterdata.usgs.gov/co/nwis/uv/?site_no=07103755 (accessed on 26 January 2017).
41. United States Geological Survey National Water Information System: Web Interface. USGS Site 07103970. Available online: http://waterdata.usgs.gov/co/nwis/uv/?site_no=07103970 (accessed on 26 January 2017).
42. United States Geological Survey National Water Information System: Web Interface. USGS Site 07104000. Available online: http://waterdata.usgs.gov/co/nwis/uv/?site_no=07104000 (accessed on 26 January 2017).
43. United States Geological Survey National Water Information System: Web Interface. USGS Site 07106300. Available online: http://waterdata.usgs.gov/co/nwis/uv/?site_no=07106300 (accessed on 26 January 2017).
44. United States Environmental Protection Agency. *Methods for Measuring the Acute Toxicity of Effluents to Freshwater and Marine Organisms*, 3rd ed. United States Environmental Protection Agency: Washington, DC, USA, 1985.
45. Herrmann, S.; Nimmo, D.; Carsella, J.; Herrmann-Hoesing, L.; Turner, J.; Gregorich, J.; Heuvel, B.V.; Nehring, R.; Foutz, H. Differential Accumulation of Mercury and Selenium in Brown Trout Tissues of a High-Gradient Urbanized Stream in Colorado, USA. *Arch. Environ. Contam. Toxicol.* **2016**, *70*, 204–218. [CrossRef] [PubMed]
46. Herrmann, S.J.; Sublette, J.E.; Helland, L.K.; Nimmo, D.W.R.; Carsella, J.S.; Herrmann-Hoesing, L.M.; Heuvel, B.D.V. Species richness, diversity, and ecology of Chironomidae (Diptera) in Fountain Creek: A Colorado Front Range sandy-Bottom watershed. *West. N. Am. Nat.* **2016**, *76*, 186–252. [CrossRef]
47. Nakayaam, F. Calcium Activity, Complex and Ion-Pair in Saturated CaCO₃. *Soil Sci.* **1968**, *106*, 429–434. [CrossRef]
48. Nakayama, F. Magnesium complex and ion-Pair in MgCO₃–CO₂ solution system. *J. Chem. Eng. Data* **1971**, *16*, 178–181. [CrossRef]
49. Panthi, S. Carbonate Chemistry and Calcium Carbonate Saturation State of Rural Water Supply Projects in Nepal. In Proceedings of the Seventh International Water Technology Conference, Cairo, Egypt, 1–3 April 2003; pp. 545–560.
50. Wurts, W.A.; Durborow, R.M. Interactions of pH, carbon dioxide, alkalinity and hardness in fish ponds. *SRAC Publ.* **1992**, 1–3.
51. Castanier, S.; Le Métayer-Levrel, G.; Perthuisot, J.-P. Ca-Carbonates precipitation and limestone genesis—The microbiogeologist point of view. *Sediment. Geol.* **1999**, *126*, 9–23. [CrossRef]
52. Ip, C.; Birringer, M.; Block, E.; Kotrebai, M.; Tyson, J.F.; Uden, P.C.; Lisk, D.J. Chemical speciation influences comparative activity of selenium-enriched garlic and yeast in mammary cancer prevention. *J. Agric. Food Chem.* **2000**, *48*, 2062–2070. [CrossRef] [PubMed]
53. Kotrebai, M.; Birringer, M.; Tyson, J.F.; Block, E.; Uden, P.C. Selenium speciation in enriched and natural samples by HPLC–ICP–MS and HPLC–ESI–MS with perfluorinated carboxylic acid ion-Pairing agents. *Analyst* **2000**, *125*, 71–78. [CrossRef] [PubMed]
54. Fleming, D.E.; Groves, J.W.; Gherase, M.R.; George, G.N.; Pickering, I.J.; Ponomarenko, O.; Langan, G.; Spallholz, J.E.; Alauddin, M.; Ahsan, H. Soft tissue measurement of arsenic and selenium in an animal model using portable X-ray fluorescence. *Radiat. Phys. Chem.* **2015**, *116*, 241–247. [CrossRef]
55. MacDonald, T.C.; Korbas, M.; James, A.K.; Sylvain, N.J.; Hackett, M.J.; Nehzati, S.; Krone, P.H.; George, G.N.; Pickering, I.J. Interaction of mercury and selenium in the larval stage zebrafish vertebrate model. *Metallomics* **2015**, *7*, 1247–1255. [CrossRef] [PubMed]

56. Choudhury, S.; Thomas, J.K.; Sylvain, N.J.; Ponomarenko, O.; Gordon, R.A.; Heald, S.M.; Janz, D.M.; Krone, P.H.; Coulthard, I.; George, G.N. Selenium preferentially accumulates in the eye lens following embryonic exposure: A confocal X-ray fluorescence imaging study. *Environ. Sci. Technol.* **2015**, *49*, 2255–2261. [[CrossRef](#)] [[PubMed](#)]
57. Hamada, Y.Z.; Bayakly, N.; Peipho, A.; Carlson, B. Accurate Potentiometric Studies of Chromium–Citrate and Ferric–Citrate Complexes in Aqueous Solutions at Physiological and Alkaline pH Values. *Synth. React. Inorg. Met. Org. Nano Met. Chem.* **2006**, *36*, 469–476. [[CrossRef](#)]
58. Fishman, M.J. *Methods of Analysis by the US Geological Survey National Water Quality Laboratory; Determination of Inorganic and Organic Constituents in Water and Fluvial Sediments*; U.S. Geological Survey: Denver, CO, USA, 1993.
59. United States Environmental Protection Agency. In *Method 200.8 Determination of Trace Elements in Water and Wastes by Inductively Coupled Plasma–Mass Spectrometry*; rev. 5.4; Environmental Protection Agency: Cincinnati, OH, USA, 1994; pp. 1–57.
60. American Public Health Association (APHA). *Standard Methods for the Examination of Water and Wastewater*; APHA: Washington, DC, USA, 2005.
61. Alderlghi, L.; Gans, P.; Ienco, A.; Peters, D.; Sabatini, A.; Vacca, A. Hyperquad simulation and speciation (HySS): A utility program for the investigation of equilibria involving soluble and partially soluble species. *Coord. Chem. Rev.* **1999**, *184*, 311–318. [[CrossRef](#)]
62. Olin, Å.; Noläng, B.; Öhman, L.-O.; Osadchii, E.; Rosén, E. *Chemical Thermodynamics of Selenium*; Elsevier Science: Amsterdam, The Netherlands, 2005.

Sample Availability: Not Available.



© 2017 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).