

# Author Correction: An intercross population study reveals genes associated with body size and plumage color in ducks

Zhengkui Zhou<sup>1</sup>, Ming Li<sup>2</sup>, Hong Cheng<sup>2</sup>, Wenlei Fan<sup>1</sup>, Zhengrong Yuan<sup>3</sup>, Qiang Gao<sup>4</sup>, Yaxi Xu<sup>1,2</sup>, Zhanbao Guo<sup>1</sup>, Yunsheng Zhang<sup>1</sup>, Jian Hu<sup>1</sup>, Hehe Liu<sup>1</sup>, Dapeng Liu<sup>1</sup>, Weihuang Chen<sup>2</sup>, Zhuqing Zheng<sup>2</sup>, Yong Jiang<sup>1</sup>, Zhiguo Wen<sup>1</sup>, Yongming Liu<sup>5</sup>, Hua Chen<sup>5</sup>, Ming Xie<sup>1</sup>, Qi Zhang<sup>1</sup>, Wei Huang<sup>1</sup>, Wen Wang<sup>6</sup>, Shuisheng Hou<sup>1</sup> & Yu Jiang<sup>2</sup>

Correction to: *Nature Communications*; <https://doi.org/10.1038/s41467-018-04868-4>; published online 17 July 2018

In the original version of this Article, there was an error in the legend for Fig. 2, whereby the descriptions of panels a, b and c were presented in a different order to the corresponding figure panels. The text ‘a GWAS of duck plumage color, including 76 colored ducks and 30 white Pekin ducks. The gray horizontal dashed lines indicate the Bonferroni significance threshold of the GWAS ( $1 \times 10^{-9}$ ). b Fixation index ( $F_{ST}$ ) of all SNPs along chromosome 13 between mallards and Pekin ducks. Red dots indicate fixed SNPs. c The nucleotide diversity ( $\pi$ ) of mallards (blue line) and Pekin ducks (red line) from 16.0 to 17.0 Mb on chromosome 13.’ should have read ‘a Fixation index ( $F_{ST}$ ) of all SNPs along chromosome 13 between mallards and Pekin ducks. Red dots indicate fixed SNPs. b The nucleotide diversity ( $\pi$ ) of mallards (blue line) and Pekin ducks (red line) from 16.0 to 17.0 Mb on chromosome 13. c GWAS of duck plumage color, including 76 colored ducks and 30 white Pekin ducks. The gray horizontal dashed lines indicate the Bonferroni significance threshold of the GWAS ( $1 \times 10^{-9}$ ).’ This has been corrected in both the PDF and HTML versions of the Article.

Published online: 25 September 2018



**Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2018

<sup>1</sup>State Key Laboratory of Animal Nutrition; Key Laboratory of Animal (Poultry) Genetics Breeding and Reproduction, Ministry of Agriculture; Institute of Animal Science, Chinese Academy of Agricultural Sciences, Beijing 100193, China. <sup>2</sup>College of Animal Science and Technology, Northwest A&F University, Yangling 712100, China. <sup>3</sup>College of Biological Sciences and Technology, Beijing Forestry University, Beijing 100083, China. <sup>4</sup>Institute of Genetics and Developmental Biology, Chinese Academy of Sciences, Beijing 100101, China. <sup>5</sup>Beijing Institute of Genomics, Chinese Academy of Sciences, Beijing 100101, China. <sup>6</sup>Research Center for Ecology and Environmental Sciences, Northwestern Polytechnical University, Xi'an 710129, China. Correspondence and requests for materials should be addressed to S.H. (email: [houss@263.net](mailto:houss@263.net)) or to Y.J. (email: [yu.jiang@nwafu.edu.cn](mailto:yu.jiang@nwafu.edu.cn))